

## Genetic Evidence that *Legionella pneumophila* RpoS Modulates Expression of the Transmission Phenotype in Both the Exponential Phase and the Stationary Phase

Michael A. Bachman and Michele S. Swanson\*

Department of Microbiology and Immunology, University of Michigan Medical School,  
Ann Arbor, Michigan 48109-0620

Received 17 September 2003/Returned for modification 19 November 2003/Accepted 22 January 2004

**The opportunistic pathogen *Legionella pneumophila* alternates between two states: replication within phagocytes and transmission between host amoebae or macrophages. In broth cultures that model this life cycle, during the replication period, CsrA inhibits expression of transmission traits. When nutrients become limiting, the alarmone (p)ppGpp accumulates and the sigma factors RpoS and FliA and the positive activators LetA/S and LetE promote differentiation to the transmissible form. Here we show that when cells enter the postexponential growth phase, RpoS increases expression of the transmission genes *fliA*, *flaA*, and *mip*, factors *L. pneumophila* needs to establish a new replication niche. In contrast, in exponential (E)-phase cells whose (p)ppGpp levels are low, *rpoS* inhibits expression of transmission traits, on the basis of three separate observations. First, *rpoS* RNA levels peak in the E phase, suggestive of a role for RpoS during replication. Second, in multiple copies, *rpoS* decreases the amounts of *csrA*, *letE*, *fliA*, and *flaA* transcripts and inhibits the transmission traits of motility, infectivity, and cytotoxicity. Third, *rpoS* blocks expression of cytotoxicity and motility by E-phase bacteria that have been induced to express the LetA activator ectopically. The data are discussed in the context of a model in which the alarmone (p)ppGpp enables RpoS to outcompete other sigma factors for binding to RNA polymerase to promote transcription of transmission genes, while LetA/S acts in parallel to relieve CsrA posttranscriptional repression of the transmission regulon. By coupling transcriptional and posttranscriptional control pathways, intracellular *L. pneumophila* could respond to stress by rapidly differentiating to a transmissible form.**

*Legionella pneumophila* has the remarkable ability to replicate within two distantly related host cells, freshwater amoebae and alveolar macrophages. However, infection of the human lung is likely fortuitous. Disease outbreaks that occur when *L. pneumophila* contaminates water sources can affect large numbers of people, but no secondary cases due to person-to-person transmission have been reported. Accordingly, the ability of *L. pneumophila* to cause pneumonia in humans is likely the consequence of selective pressures to disrupt bactericidal activities of amoebae that are common to macrophages (reviewed in reference 59).

When ingested by a macrophage or an amoeba, *L. pneumophila* blocks its immediate delivery to lysosomes (6, 31). Instead, its phagosome transiently associates with mitochondria and endoplasmic reticulum, and then the microbe replicates to high numbers in an acidic vacuole that acquires lysosomal traits (18, 30, 58, 60). When it has exhausted the nutrient supply, *L. pneumophila* must exit the host cell and search for new prey. To do so, after the replication period, the progeny not only increase their resistance to extracellular stresses (4, 20, 22, 42) but also express a pore-forming activity to escape the host (2, 8, 35), assemble a unipolar flagellum to disseminate (8, 51), and use a type IV secretion system to inhibit

phagosome maturation in the new host cell (5, 52, 56, 62), where the cycle repeats.

From studies of broth cultures that model the *L. pneumophila* life cycle, several features of the regulatory circuit that governs bacterial differentiation have been deduced. Exponential (E)-phase bacteria respond to nutrient depletion or other stress by producing the alarmone (p)ppGpp (21). Subsequently, as the bacteria transition to the postexponential (PE) phase, the sigma factor RpoS induces motility, sodium sensitivity, and evasion of the endocytic pathway, and it also promotes subsequent replication of the intracellular bacteria (3, 20). In parallel, the LetA/S two-component regulator system also induces motility, sodium sensitivity, and evasion of lysosomes, and it activates cytotoxicity, pigmentation, resistance to heat and osmotic stress, and cell shortening (17, 22, 42). LetA/S likely does so by relieving translational repression by CsrA, thereby activating the transmission phenotype by pathways that are both dependent on and independent of the flagellar sigma factor FliA (17, 44). By an unknown mechanism, the LetE protein enhances expression of LetA/S-dependent transmission traits (22). Thus, by integrating the activity of multiple sigma factors, a two-component regulatory system, and a repressor of translation, *L. pneumophila* can efficiently alternate between an intracellular replicative state and a form fit to spread among populations of amoebae or macrophages.

Among gram-negative bacteria, RpoS, LetA/S, and CsrA are highly conserved global regulators of stationary-phase physiology that act by distinct mechanisms. The RpoS sigma factor directs RNA polymerase to transcribe genes important for

\* Corresponding author. Mailing address: University of Michigan Medical School, 6734 Medical Sciences Building II, Ann Arbor, MI 48109-0620. Phone: (734) 647-7295. Fax: (734) 764-3562. E-mail: mswanson@umich.edu.

survival in stationary phase and under other stressful conditions (24). In contrast, the LetA/S homologues known as GacA/S or BarA/SirA act posttranscriptionally (23). Specifically, these two-component regulatory systems relieve repression by the mRNA binding protein CsrA by inducing the expression of the regulatory RNA *csrB*, which binds CsrA protein to derepress translation of mRNAs (reviewed in reference 50). In a number of human pathogens, including *L. pneumophila*, *Salmonella enterica* serovar Typhimurium, and *Pseudomonas aeruginosa*, a subset of genes in both the RpoS and the LetA/S (GacA/S, BarA/SirA) regulons encode virulence effectors (1, 3, 15, 17, 20, 22, 34, 39, 42, 44, 49). By using both transcriptional and translational control mechanisms, bacteria could respond quickly to environmental fluctuations by altering their composition.

In a number of bacterial species, the RpoS sigma factor and the LetA/S posttranscriptional regulatory system directly interact, although to different effect. In *Escherichia coli* and *Pseudomonas fluorescens*, LetA/S homologues upregulate *rpoS* transcript levels beginning in E phase and are required for RpoS-dependent traits in PE phase (46, 63). Conversely, *Erwinia carotovora* RpoS functionally opposes the LetA/S homologues GacA/S by positively activating RsmA (CsrA), the mRNA binding protein that destabilizes targeted transcripts (45). In *L. pneumophila*, phenotypic analysis of bacteria that lack or overexpress *rpoS* has predicted not only unexpected roles for RpoS but also both cooperative and antagonistic interactions with LetA/S. RpoS is dispensable for *L. pneumophila* to become resistant to oxidative, osmotic, and acidic stress in the PE phase (20). Instead, the *rpoS* locus promotes resistance to osmotic stress during the E phase, whereas LetA/S activates stress resistance in the PE phase (3, 20, 22, 42, 44). Conversely, when present in multiple copies, wild-type *rpoS* inhibits *L. pneumophila* replication in *Acanthamoeba castellanii* (20) and also motility and cytotoxicity of PE-phase *L. pneumophila*, two traits induced by LetA/S (3). On the basis of these paradoxical results, we tested the hypothesis that, depending on the growth phase, RpoS either positively or negatively modulates traits that are induced by LetA/S.

#### MATERIALS AND METHODS

**Bacterial strains, plasmids, and media.** *L. pneumophila* Lp02, a virulent thymine auxotroph (5); MB379 (*rpoS::kan*; 3); and *mariner* Tn mutant strains (Kan<sup>r</sup>; 22) MB414 (*letA-22*), MB417 (*letS-36*), and MB420 (*letE-121*) derived from Lp02 were cultured in *N*-(2-acetamido)-2-aminoethanesulfonic acid (ACES; Sigma)-buffered yeast extract broth supplemented with 100 µg of thymidine per ml (AYET) at 37°C. Bacteria were plated on ACES-buffered charcoal-yeast extract agar supplemented with 100 µg of thymidine per ml (CYET). Where indicated, media was supplemented with gentamicin to a final concentration of 10 µg/ml or with kanamycin to 25 µg/ml.

The RSF1010-based plasmids used in this study were digested with AgeI to delete *mobA*, since conjugation function has been demonstrated to interfere with *L. pneumophila* virulence (57). Plasmids pMB384 (pKB5Δ*mobA* Thy<sup>r</sup>), pLrpoS (pMB391, pKB5Δ*mobA* containing a 6.8-kb *rpoS* genomic locus), and pLrpoSD (3) were transformed into Lp02 by electroporation and selected on CYE, which lacks thymidine. Likewise, pLetA (pMMB-GentΔ*mobA* with a 1.4-kb *letA*-containing fragment colinear with the P<sub>TAC</sub> promoter) was transformed into Lp02 or MB379 and selected with gentamicin. As previously reported, LetA homologues can inhibit growth in laboratory culture (14), a trait we observed as heterogeneity in the colony size of transformants, regardless of their *rpoS* allele. Inocula from small colonies grew poorly in gentamicin-supplemented AYET and displayed heterogeneous colony morphology when colony purified or plated from broth cultures. These small isolates were not studied further. Approximately 1 to 10%

TABLE 1. Primers used in this study

Primer	Sequence
<i>rpoS</i> -int1	5'-CCA AGC GAG GAT TCC GTT TTT C-3'
<i>rpoS</i> -int2 <sup>a</sup>	5'-AAA ACG TCT TGC GAT AAC CTG-3'
<i>mip</i> -int1	5'-AAA TCT CTA TCG CTA ACG CAC AAG-3'
<i>mip</i> -int2 <sup>a</sup>	5'-AAC TCA CTC ACC CCA CGA C-3'
<i>fliA</i> -int1	5'-GGT GCC GCG TTC TGT TTA TC-3'
<i>fliA</i> -int2 <sup>a</sup>	5'-CGG TGC GTT GCT TGA CTT-3'
<i>fliA</i> -int1	5'-CCA ATT TTA CCG GGG CAC TA-3'
<i>fliA</i> -int2 <sup>a</sup>	5'-AAC GGC GCT GAT TGA TAA AG-3'
<i>csrA</i> -int1 <sup>b</sup>	5'-AGC TGC TAT GTT AGG GAG-3'
<i>csrA</i> -int2 <sup>a,c</sup>	5'-TAC TGC TTG TTC CGA ATC-3'
<i>letE</i> Tn1	5'-TAC ATG CAC TAA AAA GCG GTT CTG-3'
<i>letE</i> Tn2 <sup>a</sup>	5'-TGA TCA TGC TAC GAG CTC AAG TAA-3'

<sup>a</sup> Used for ssDNA probe synthesis.

<sup>b</sup> Derived from *csrA* uni139-BamHI (17).

<sup>c</sup> Derived from *csrA* rev340-HindIII (17).

of total transformants, large isolates exhibited a stable colony size when colony purified or plated from broth cultures. Inocula from three large-colony isolates of each strain containing plasmid-borne *letA* replicated in broth, reached PE phase, and became motile and cytotoxic with kinetics similar to that of wild-type Lp02 (data not shown), and these clones were used for subsequent experiments.

**RNA preparation and Northern analysis.** For mutant analysis, overnight cultures were quantified by optical density at 600 nm (OD<sub>600</sub>) and then diluted to 0.3 or 0.03 to generate PE-phase and E-phase cultures after an additional 16 h of incubation. Time course experiments were performed by diluting overnight cultures to an OD<sub>600</sub> of 0.015 in 25 ml and, to allow for equal aeration, dividing them into five 5-ml aliquots. After an additional overnight incubation, cultures were in mid-E phase (time = 0 h). Subsequently, RNA was prepared by collecting one aliquot every 3 h for a 12-h period, at which point the cultures had entered the PE phase. Bacteria from 4.5 ml of E-phase cultures and 1.5 ml of PE-phase cultures were collected by centrifugation, resuspended in 1 ml of Trizol reagent, and purified as directed by the manufacturer (Invitrogen). RNA samples were quantified by OD<sub>260</sub>, and 10 µg of total RNA per sample was electrophoresed on a 1.2% formaldehyde gel. The gel was prepared, electrophoresed, and transferred to BrightStar BioDetect nylon membranes in accordance with the NorthernMax kit protocol (Ambion, Inc.).

For each gene of interest, a segment within the open reading frame was amplified by PCR with the primers listed in Table 1. Biotin-labeled, antisense, single-stranded DNA (ssDNA) probes were synthesized with the Strip EZ PCR kit (Ambion, Inc.) from 10 ng of each PCR product by using primer 2 of each pair and 200 nM biotin 14-ATP (Invitrogen). Membranes were hybridized with NorthernMax Ultrahyb and a final probe concentration of ~1 ng/ml. Probes were initially quantified, and mRNA was subsequently detected with the BrightStar BioDetect kit (Ambion) and Kodak ML chemiluminescence film. The net intensity of hybridization was measured with Kodak Digital Science 1D software and expressed as arbitrary units (data not shown). Probes were stripped by incubation for 10 min at room temperature in probe degradation buffer, 10 min at 68°C, and 10 min in blot reconstitution buffer at 68°C (Strip EZ PCR kit; Ambion).

**Contact-dependent cytotoxicity.** Quantification of contact-dependent cytotoxicity of PE-phase cultures was performed as previously described (21). For LetA induction experiments, overnight cultures were diluted to an OD<sub>600</sub> of ~0.007, divided in two, and incubated overnight with or without 200 µM isopropyl-β-D-thiogalactopyranoside (IPTG). On the following morning, culture density was quantified by OD<sub>600</sub>, motility was assessed by phase microscopy, cultures were diluted to equivalent concentrations in RPMI medium plus 10% fetal bovine serum (R10), and A/J mouse bone marrow macrophages were infected with each sample in triplicate. Inocula were diluted and plated on CYET containing gentamicin. To quantify CFU, after a 1-h incubation with macrophages, bacteria were removed by repeated washing with R10 and macrophage viability was quantified by incubation with 10% Alamar blue. Reduction of the colorimetric dye by viable macrophages was measured by the ratio of OD<sub>570</sub> to OD<sub>600</sub> in a Spectramax 250 plate reader (Molecular Devices).

**Bone marrow-derived macrophage infectivity.** Enumeration of *L. pneumophila* bacteria that bind, enter, and survive within A/J mouse bone marrow-derived macrophages during a 2-h incubation was performed as previously described (8), with the following modifications. Macrophages were infected with *L. pneumophila* at a multiplicity of infection of 0.2 to avoid cytotoxicity, and gentamicin

treatment was omitted since control studies indicate that this antibiotic is only weakly bactericidal against PE-phase *Legionella* (S. Sturgill-Koszycki, unpublished results). Instead, the majority of extracellular bacteria were removed by washing the infected monolayers with  $3 \times 0.5$  ml of R10 immediately before lysis by trituration with  $2 \times 0.5$  ml of phosphate-buffered saline. Infectivity was expressed as (cell-associated CFU at 2 h/CFU added at 0 h)  $\times 100$ .

## RESULTS

*L. pneumophila* expresses *rpoS* RNA maximally in E phase by a mechanism independent of the LetA/S pathway. Consistent with the primary role of RpoS as a stationary-phase sigma factor, *E. coli* expresses *rpoS* RNA and protein maximally in early PE phase (24, 46). As a first step to investigate how RpoS regulates growth phase-dependent differentiation of *L. pneumophila*, we compared *rpoS* expression by wild-type bacteria cultured to either the E or the PE phase. Although the *L. pneumophila* RpoS protein is most abundant in the PE phase (20), its *rpoS* transcripts were maximal in the E phase and undetectable in the PE phase (Fig. 1), a pattern generally consistent with a previous genetic study of *rpoS* promoter activity (42). To a first approximation, its pattern of RNA abundance suggests an atypical E-phase function and/or mode of regulation for *L. pneumophila* RpoS.

In *E. coli*, *rpoS* transcription is induced to a low level in the E phase by BarA, a tripartite sensor kinase that is homologous to LetS (22, 46). Therefore, we examined whether *L. pneumophila* activates *rpoS* expression in the E phase via its cognate LetA/S two-component regulatory system. The effect of a panel of regulatory mutations on *rpoS* RNA accumulation was determined by Northern analysis (Fig. 1). In the E phase, the amount of *rpoS* RNA in *letA* and *letS* mutants approached that of wild-type *L. pneumophila*. Likewise, LetE, which is predicted to cooperate with LetA/S to induce transmission traits (22), had only a modest effect on *rpoS* RNA accumulation by E-phase *L. pneumophila* (Fig. 1). Thus, *rpoS* RNA accumulates during the replication period by a mechanism that is largely independent of LetA/S and LetE, putative posttranscriptional activators of the PE transmission phenotype (22, 23).

**RpoS and LetA/S induce transmission genes by independent mechanisms.** As another approach to investigate how RpoS and LetA/S cooperate to control *L. pneumophila* differentiation, their effects on the expression of three genes known to promote phagocyte infection were compared. The macrophage infectivity potentiator (Mip) is a peptidyl-prolyl *cis-trans* isomerase that enhances *L. pneumophila* invasion and replication within both amoebae and macrophages (10, 11). As expected, the *mip* RNA was present in both E- and PE-phase wild-type cells (36, 64). The LetA/S and LetE proteins are minor inducers of *mip* RNA expression in the PE phase since, compared to the wild type, bacteria that lack *letA*, *letS*, and *letE* contained less but still appreciable levels of *mip* transcripts (Fig. 1). In comparison, RpoS is critical for the accumulation of *mip* RNA in the PE phase, since its transcripts were barely detectable in PE *rpoS* mutant cells (Fig. 1). The observation that *mip* PE-phase expression is controlled primarily by RpoS may account for the similar lag in growth initiation that is characteristic of both *mip* and *rpoS* mutants that have infected either macrophages or amoebae (3, 10, 11).

In addition to Mip, *L. pneumophila* uses two genes of the flagellar regulon to infect phagocytes efficiently. The flagellar




FIG. 1. Accumulation of *rpoS*, *fliA*, *flaA*, and *mip* mRNAs by E- and PE-phase wild-type (WT) and mutant *L. pneumophila*. Northern analysis was performed on 10  $\mu$ g of total RNA collected from E-phase ( $OD_{600}$ , 0.8 to 1.2) and PE-phase ( $OD_{600}$ , 3.1 to 3.6) cultures of wild-type Lp02 bacteria and *letA*-22, *letS*-36, *letE*-121, and *rpoS* mutant bacteria with a biotin-labeled ssDNA probe. Hybridization was detected with streptavidin conjugated to alkaline phosphatase and visualized by CDP-Star (Tropix). The hybridization pattern shown is representative of results obtained from at least two independent sets of RNA samples. The relative amount of total RNA in each sample is demonstrated by ethidium bromide staining of *rrnA* from the formaldehyde gel prior to membrane transfer. Greater hybridization to *mip* in the wild-type PE-phase sample is due to a higher concentration of total RNA, as demonstrated by ethidium bromide staining of *rrnA*.

sigma factor FliA is required for expression of the three PE-phase transmission traits of motility, infectivity, and cytotoxicity and also for replication in amoebae (22, 27, 44). FliA activates several genes needed for the terminal stages of flagellum development, including *flaA*, the structural gene for flagellin

which itself is required for efficient infection of phagocytes (12). Therefore, we next used these two genes of the flagellar regulon as tools to investigate how the RpoS and the LetA/S and LetE pathways cooperate to regulate differentiation of PE-phase *L. pneumophila*.

Replicating wild-type *L. pneumophila* contained a small amount of full-length *fliA* RNA; upon entry into the PE phase, the quantity increased dramatically (Fig. 1). Results of previous studies predict that *fliA* induction in the PE phase requires LetA/S to relieve CsrA repression (17, 44). Indeed, the growth phase regulation of *fliA* expression required LetA/S, as well as RpoS. Full-length *fliA* RNA was rare in E- and PE-phase *letA* and *letS* mutants, diminished in *letE* mutants, and undetectable in *rpoS* mutants (Fig. 1). Therefore, the robust induction of *fliA* RNA levels in the PE phase appears to be accomplished by two mechanisms: the sigma factor RpoS likely regulates transcription initiation, whereas the LetA/S pathway is predicted to relieve posttranscriptional repression mediated by CsrA (17).

Consistent with the paradigm of FliA-directed transcription of the *flaA* flagellin gene and genetic assays of its promoter activity (17, 21, 26, 28), *flaA* RNA levels were induced in the PE phase by LetA/S-, LetE-, and RpoS-dependent pathways (Fig. 1). Compared to wild-type *L. pneumophila*, PE-phase *letA* and *letS* mutants have dramatically reduced amounts of *flaA* RNA, whereas *letE* and *rpoS* mutants contained a low level of *flaA* RNA (Fig. 1), a pattern consistent with the degree of their motility defects (3, 22).

It was notable, however, that the abundance of *flaA* RNA in the *letE* and the *rpoS* mutant strains did not correspond directly to their levels of *fliA* transcripts. Relative to PE-phase *letA* or *letS* mutants, *rpoS* mutants reproducibly harbored less RNA encoding the FliA sigma factor yet accumulated similar amounts of RNA encoding flagellin. In contrast, PE-phase *letE* mutants contained a substantial amount of *fliA* RNA but little *flaA* RNA (Fig. 1). These results are also consistent with a model in which the steady-state level of *flaA* RNA is controlled by two mechanisms: the sigma factor RpoS regulates transcription initiation, whereas LetE and LetA/S act posttranscriptionally as stabilizers of effector mRNAs (23, 24). Accordingly, *letE* mutants contain ample RpoS and FliA sigma factors to transcribe *flaA*, but the mRNA product is destabilized by the constitutive CsrA posttranscriptional repressor. Conversely, an *rpoS* mutant transcribes *fliA* poorly, but the mRNA yield is sufficient for *flaA* expression since LetA/S and LetE relieve mRNA destabilization by CsrA (17, 44, 50).

**In multiple copies, rpoS represses transcription of transmission genes.** Previous studies documented that in multiple copies the wild-type *rpoS* locus inhibits *L. pneumophila* replication in amoebae (20) and also represses two *fliA*-dependent PE transmission traits, motility and cytotoxicity (3). Therefore, as an independent approach to analyze how RpoS functionally interacts with other regulators of *L. pneumophila* differentiation, the effect of multiple copies of the wild-type *rpoS* locus on the transcription of known transmission genes was examined. For this purpose, RNA samples from wild-type bacteria that contained either the *rpoS* plasmid pLrpoS or the control vector pMB384 were collected at 3-h intervals between the E phase and the PE phase for Northern analysis (Fig. 2A). As expected, compared to bacteria containing the control vector, *L. pneumophila* transformed with pLrpoS had increased levels of both

full-length and shorter *rpoS* transcripts throughout the E phase (Fig. 2B).

When *rpoS* RNA was excessive, the expression of a number of transmission genes was inhibited. *L. pneumophila* transformed with pLrpoS contained less RNA for two regulators of transmission that are transcribed predominantly in the E phase, namely, the positive activator *letE* and the repressor *csrA* (17, 44). A similar effect was observed for two genes expressed primarily in the PE phase: multiple copies of *rpoS* significantly decreased the magnitude, but not the timing, of *fliA* and *flaA* expression (Fig. 2B). The observation that excess *rpoS* can inhibit the expression of four different genes, *letE*, *csrA*, *fliA*, and *flaA*, is consistent with the work of Nystrom and colleagues demonstrating that *E. coli* sigma factors compete for binding to core RNA polymerase. By influencing the competition among sigma factors, growth or experimental conditions can alter the cellular pattern of gene expression (16, 33, 40, 43).

**RpoS overexpression represses LetA/S-dependent virulence traits.** To test more rigorously whether RpoS can antagonize LetA/S-dependent activation of the transmission phenotype, the effect of pLrpoS on the expression of three transmission traits by wild-type *L. pneumophila* was analyzed. Whereas ~75% of PE-phase wild-type cells that contained only the chromosomal *rpoS* locus were motile, motility was rare or absent when *L. pneumophila* contained multiple copies of *rpoS*, consistent with the inhibitory effects of pLrpoS on *fliA* and *flaA* expression (3).

In the PE phase, *L. pneumophila* expresses a contact-dependent pore-forming activity that is toxic to macrophages and is thought to promote bacterial escape from phagocytes when nutrients become scarce (2, 8, 35). Induction of cytotoxicity in the PE phase requires LetA/S, LetE, and FliA, but not RpoS (3, 22). Nevertheless, when wild-type *L. pneumophila* was transformed with pLrpoS, cytotoxicity was completely repressed (Fig. 3), a phenotype consistent with the negative effect of pLrpoS on *fliA* expression (Fig. 2B). To verify that the pLrpoS-mediated repression was conferred by the *rpoS* locus and not other plasmid sequences, the effect of its derivative pLrpoSΔ, which contains an *rpoS*:*kan* null allele, was examined. Strains transformed with this control plasmid were at least as cytotoxic as the wild type that contained only the parent vector pMB384, demonstrating *rpoS* specificity (Fig. 3).

When ingested by macrophages, *L. pneumophila* can evade lysosomal degradation (31), provided the LetA/S regulator is active (22). In contrast, RpoS is dispensable for lysosome evasion, but it is necessary to block fusion of the bacterial phagosome with earlier, nonbactericidal endosomal compartments (3). To test whether multicopy *rpoS* also inhibited lysosome evasion, macrophage infection by *rpoS* mutant and wild-type cells with or without pLrpoS was assessed. After a 2-h incubation, similar percentages of wild-type and *rpoS* bacteria were viable and cell associated (Fig. 3), a result that reflects the persistence of *rpoS* mutants in vacuoles that contain LAMP-1 but not lysosomal contents (3). In contrast, carriage of pLrpoS by wild-type *L. pneumophila* lowered infectivity from ~33% to ~2% of the inoculum, a phenotype shared with *letA* and *fliA* mutants (22). The virulence defects conferred by plasmid-borne *rpoS* were not a simple consequence of slowing growth, as judged by replication of bacteria in broth (Fig. 2A) or in




FIG. 2. RpoS overexpression represses *letE*, *csrA*, *fliA*, and *flaA* expression by a mechanism that does not affect *L. pneumophila* growth

macrophages (3). In summary, multiple copies of *rpoS* caused poor motility, cytotoxicity, and infectivity in the PE phase, a spectrum of defects reminiscent of the pattern conferred by mutations in *letA*, *letS*, *letE*, or *fliA* (22). Therefore, in excess, *rpoS* antagonizes activators of the transmission phenotype.

**RpoS prevents LetA induction of transmission traits in the E phase.** On the basis of its E-phase RNA expression and the inhibitory effects of pLrpoS, we postulated that during the replication period, RpoS protein represses transmission directly or indirectly. This hypothesis predicts that genetic inactivation of *rpoS* will allow LetA to induce transmission traits prematurely during the replication period. To test this supposition, wild-type and *rpoS* mutant bacteria were transformed with pLetA, a plasmid that encodes *letA* under the control of an inducible pTlac promoter, and then their ability to respond to LetA by expressing cytotoxicity during the E phase was assessed (Fig. 4).

As demonstrated previously, wild-type bacteria were not cytotoxic in the E phase. Likewise, IPTG induction of *letA* was not sufficient to induce cytotoxicity in E-phase cultures of wild-type *L. pneumophila*. However, as predicted, in the absence of *rpoS*, E-phase cells responded to activating signals by LetA. Indeed, E-phase *rpoS* mutant cells induced to express *letA* resembled PE-phase *L. pneumophila* cells in their ability to kill macrophages (Fig. 4). Likewise, RpoS also repressed the expression of motility during the E phase, since induction of *letA* expression triggered flagellar motility only in the absence of *rpoS* function (data not shown). RpoS repression was relieved as cells entered the PE phase; in the presence or absence of pLetA, wild-type cells became fully cytotoxic, as expected (8). Therefore, the low level of RpoS protein normally present in E-phase *L. pneumophila* (20) blocks expression of transmission traits, even when the *letA* activator is expressed ectopically.

## DISCUSSION

In *L. pneumophila*, RpoS function and regulation are atypical of its homologues in several other species of gram-negative bacteria, on the basis of four observations. First, although *L. pneumophila* substantially increases its resistance to oxidative, osmotic, and acidic stress in the PE phase, none of these fortifications require RpoS (3, 20). Second, RpoS induces osmotic resistance of replicating *L. pneumophila* (20). Third, transcription of *E. coli rpoS* increases ~5- to 10-fold upon transition to the stationary phase (25); in *L. pneumophila*, abundant E-phase transcripts disappear in the PE phase (Fig. 1 and 2). Last, *P. fluorescens* and *E. coli* LetA/S homologues

in broth. (A) OD<sub>600</sub> of cultures of wild-type Lp02 containing the vector pMB384 (open circles) or pLrpoS (pMB391, closed circles). At each time point shown, aliquots were removed, centrifuged, permeabilized with Trizol reagent, extracted for total RNA as directed by the manufacturer (Invitrogen), and analyzed as described for panel B. (B) Northern analysis was performed on 10 µg of total RNA collected as described for panel A. The same membrane was stripped and reprobbed for each transcript. The relative amount of total RNA in each sample is demonstrated by ethidium bromide staining of *rrnA* from the formaldehyde gel prior to membrane transfer. The results shown are representative of two sets of RNA samples collected from independent cultures at similar optical densities. WT, wild type.




FIG. 3. Multiplicity expression of *rpoS* eliminates cytotoxicity and infectivity. (A) Contact-dependent cytotoxicity for macrophages. Macrophages were incubated for 1 h with samples of PE-phase wild-type Lp02 containing the vector pMB384 (circles), pLrpoS harboring a 6.8-kb *rpoS* locus (pMB391, squares), or pLrpoS::kan carrying an *rpoS*::kan mutation in the same locus (pMB392, triangles). Macrophage viability was assessed by quantifying reduction of the colorimetric dye Alamar blue. Shown are means and standard errors of triplicate samples from one representative experiment of three performed. (B) Efficiency of macrophage infection. Macrophages were infected at a multiplicity of infection of 0.2 with the following PE-phase strains: wild-type (WT) strain Lp02 containing the vector pMB384, *rpoS* mutant strain MB379 (*rpoS*::kan) containing pMB384 and the wild type plus pLrpoS. The percentage of each *L. pneumophila* inoculum that was cell associated after a 2-h infection with  $2.5 \times 10^5$  macrophages is shown. The results shown are means of macrophage samples infected in triplicate; error bars represent standard deviations.

induce *rpoS* transcription (46), but *L. pneumophila* *rpoS* RNA species appear to be destabilized in the PE phase by a LetA/S-dependent pathway (Fig. 1). Together, these observations suggest that, in *L. pneumophila*, RpoS is used in pathways or under conditions beyond coordination of stationary-phase physiology.

Studies of *E. coli* and *S. enterica* serovar Typhimurium provide precedents for a role for RpoS during the *L. pneumophila*




FIG. 4. RpoS represses LetA-induced cytotoxicity in the E phase. After overnight growth in the presence (closed symbols) or absence (open symbols) of 200  $\mu$ M IPTG to induce expression from pLetA, samples of E-phase wild-type Lp02 (circles) or *rpoS* mutant MB379 (squares) were incubated for 1 h with macrophages. Macrophage viability was assessed by quantifying reduction of the colorimetric dye Alamar blue. Shown are means and standard errors of triplicate samples from one representative experiment of six performed.

replication period. Although *E. coli* RpoS protein levels are low under nutrient-rich conditions, this alternate sigma factor directs the transcription of *xthA* and *katG*, genes that prevent UV and oxidative damage (32, 41, 53). When macrophages ingest *S. enterica* serovar Typhimurium, another gram-negative bacterium that replicates within intracellular vacuoles, its RpoS expression increases dramatically, approaching levels typical of stationary phase (9). RpoS is required for full virulence of *S. enterica* serovar Typhimurium in mice, as it activates the *katE*-encoded catalase and other *spv*-dependent and -independent factors (15). The biphasic life cycle of *L. pneumophila* may account for the diverse roles postulated for RpoS. To persist as a free-living aquatic microbe, *L. pneumophila* must tolerate nutrient limitation and other environmental stresses. To replicate efficiently within phagocytes, *L. pneumophila* must adapt to another harsh environment, i.e., acidic lysosomal compartments (58). To thrive within this stressful but nutrient-rich site, *L. pneumophila* may have evolved a regulatory circuit in which RpoS confers on replicating bacteria tolerance to lysosomal stresses while also preventing induction of the transmission phenotype.

The inverse pattern of *L. pneumophila* RpoS RNA and protein levels (20) likely indicates that RpoS is subject to post-transcriptional control, which is well documented in *E. coli* (25). A stockpile of *rpoS* mRNA may enable replicating *L. pneumophila* to respond quickly to deteriorating conditions. As *L. pneumophila* differentiates to the transmissible form, a LetA-dependent increase in *rpoS* promoter activity, predicted by a transcriptional *rpoS*-*lacZ* reporter (42), may further increase template levels. However, in the PE phase, full-length *rpoS* transcripts disappear (Fig. 1 and 2) while RpoS protein accumulates (20). LetA/S and LetE are attractive candidates to couple *rpoS* RNA translation to decay, given their predicted role as antagonists of the posttranscriptional repressor CsrA (23, 44). Interestingly, PE-phase *letA*, *letS*, and *letE* mutant cells contained a large quantity of smaller *rpoS* RNAs, which were rare in wild-type PE cells (Fig. 1). Similar *rpoS* species

have been observed in other gram-negative bacteria under conditions of *rpoS* message excess and are thought to be stable degradation intermediates (38, 46). Further studies are required to determine whether the *L. pneumophila* LetA/S regulatory pathway mediates *rpoS* RNA decay after its translation in the PE phase.

This study provides insight into the organization of the *L. pneumophila* RpoS regulon, showing that this sigma factor is required for PE-phase expression of *mip*, the flagellar sigma factor *fliA*, and its target *flaA*, three well-described virulence factors that promote *L. pneumophila* infection of amoebae and macrophages (19, 22, 27). Both *mip* and *rpoS* mutants exhibit a characteristic lag in growth initiation after infection of phagocytes (3, 10, 11). However, unlike a *mip* mutant, an *rpoS* mutant cannot replicate at all in amoebae (20). Its more severe phenotype is consistent with the observation that RpoS regulates additional genes, including *fliA* and *flaA*, factors that together with Mip are required for efficient infection and growth inside protozoa (12, 27).

Several lines of genetic and molecular evidence indicate that RpoS and LetA/S activate the expression of transmission genes by cooperative but distinct mechanisms. In comparison to an *rpoS* mutation, mutations in *letA* or *letS* have a less extreme effect on the levels of *mip* and *fliA* transcripts, yet they impair *flaA* expression more severely (Fig. 1). Thus, a LetA/S- and LetE-dependent mechanism may compensate for low *fliA* message in an *rpoS* mutant to allow expression of FliA-dependent traits. Consistent with their more extreme effects on *fliA* and *flaA* expression, LetA/S and FliA are required for cytotoxicity and infectivity in the PE phase, whereas RpoS is not (3, 22). Furthermore, when *L. pneumophila* is engulfed by a phagocyte, RpoS and LetA/S promote two different activities: immediate arrest of phagosome maturation and subsequent bacterial replication. *letA* or *letS* mutants infect macrophages and amoebae poorly (22, 42). However, even though 50% of internalized *letA* mutant bacteria are degraded by macrophage lysosomes, the survivors replicate as efficiently as the wild type (22). *rpoS* mutants display the opposite pattern: the majority of *rpoS* mutant bacteria remain intact in an intermediate endosomal compartment, but they fail to multiply (3). Also, the requirement for LetA can be bypassed by genetic inactivation of the CsrA repressor, whereas the need for RpoS cannot (44). Therefore, RpoS and LetA/S have specialized roles in the transmissive phase, as postulated previously (3, 42). By analogy to their demonstrated activity in *E. coli* and other gram-negative bacteria, RpoS is expected to control transcription initiation by RNA polymerase, whereas LetA/S is likely to act posttranscriptionally by relieving CsrA repression to stabilize *fliA* and/or *flaA* mRNA and promote synthesis of these and other transmission effectors (23, 24, 50).

By examining bacteria that lack or overexpress *rpoS*, we found that, depending on the growth phase, RpoS either negatively or positively influences expression of LetA/S-dependent transmission traits. In replicating *L. pneumophila*, RpoS not only promotes osmotic resistance (20) but also prevents LetA-induced transmission traits, since ectopically expressed LetA induces cytotoxicity and motility only when *rpoS* is absent (Fig. 4). Likewise, when overexpressed by PE cells, *rpoS* disrupts LetA/S-dependent motility, cytotoxicity, and infectivity (Fig. 3, data not shown) by inhibiting expression of *fliA*, *flaA*, and *letE*

(Fig. 2). Consistent with this experimental condition, in wild-type *L. pneumophila* cultures, *fliA* and *flaA* transcripts are scarce when *rpoS* RNA is abundant (Fig. 1 and 2). Together, the data indicate that, by some direct or indirect mechanism, RpoS prevents activation of the transmission phenotype by replicating *L. pneumophila*.

During the replication period, the low level of RpoS protein (20) could inhibit the transmission phenotype indirectly by inducing a repressor. In *E. coli* and *S. enterica* serovar Typhimurium, RpoS-dependent repression of pilus phase variation is attributed to transcriptional activation of a repressor gene (13, 48). In *E. coli*, RpoS is also known to cooperate with Hfq to repress expression of *mutH* (61) and with Fis to repress a collection of genes at various points in the growth cycle (65). Because *E. carotovora* RpoS induces the posttranscriptional repressor RsmA (45), we tested whether *L. pneumophila* *csrA* RNA levels are regulated by RpoS. To the contrary, our data indicate that RpoS represses transmission genes along with *csrA*, and an *rpoS* mutation had little effect on *csrA* expression (data not shown).

By analogy to *E. coli*, *L. pneumophila* RpoS could also inhibit transmission gene expression by competing with other sigma factors for binding to RNA polymerase (16, 33, 40). For example, when overexpressed, RpoS inhibits RpoD-directed transcription by PE-phase *E. coli* by competing for binding to core RNA polymerase (16). In *L. pneumophila*, the observation that overexpression of *rpoS* inhibits multiple genes, including loci normally transcribed in either the E or the PE phase (Fig. 2), is in keeping with such a mechanism. Furthermore, the observation that, depending on the experimental conditions, RpoS can either repress or activate the expression of the same gene, *fliA* (Fig. 1 and 2B), is consistent with the detailed genetic and biochemical studies of Nystrom and colleagues demonstrating that sigma factors compete for binding to core RNA polymerase (16, 33, 40).

By taking into account the paradigm of sigma factor competition, known transcriptional and posttranscriptional control mechanisms of RpoS activity (25), the nucleotide sequence and genetic analysis of the *L. pneumophila* flagellar regulon by Heuner and colleagues (29), and genetic evidence from our laboratory and others, the following working model of *L. pneumophila* regulation of differentiation can be proposed. During the replication period, the *rpoS* gene is transcribed at high levels (Fig. 1); although most of the RpoS protein is presumably degraded (20, 25), there must be sufficient protein bound to RNA polymerase to mediate osmotic resistance (20) and promote intracellular replication (3, 20). At the same time, although sigma factor RpoN may direct some transcription of *fliA* (29), the posttranscriptional repressor CsrA targets any mRNA encoded by the flagellar regulon for degradation (17). Consequently, when amino acids are abundant and (p)ppGpp levels are low, *fliA* and *flaA* RNAs are rare and *L. pneumophila* cells are not motile, cytotoxic, or infectious. When the basal level of E-phase RpoS protein is eliminated genetically, sufficient RpoN may bind the freed RNA polymerase to induce the flagellar regulon, provided posttranscriptional repression by CsrA is also alleviated by ectopic expression of LetA/S (29, 44).

When conditions deteriorate, we postulate that accumulation of (p)ppGpp (21) increases the amount of RpoS protein (20, 25) and also promotes both RpoS- and RpoN-mediated

TABLE 2. Demand theory<sup>a</sup> of *L. pneumophila* differentiation

Growth phase	Demand for transmission genes	Predicted mode of regulation	Regulators	Observed mode of regulation	Reference(s)
E (replicative)	Low	Negative	RpoS, CsrA	Negative	This work, 17, 43
PE (transmissive)	High	Positive	RpoS, LetA/S, LetE, FliA	Positive	This work, 3, 12, 20, 22, 27, 29, 41, 43

<sup>a</sup> See Discussion and references 47, 54, and 55.

transcription by RNA polymerase (33, 37, 40). By some mechanism, RpoS promotes transcription initiation in the flagellar regulon (Fig. 1), perhaps indirectly by elevating the expression of an RpoN coactivator protein (7). According to this model, excess RpoS could inhibit PE expression of the flagellar regulon (Fig. 2) by outcompeting RpoN for binding to RNA polymerase. In wild-type *L. pneumophila*, (p)ppGpp (21, 22) or some other PE-phase signal (66) concomitantly activates the LetA/S two-component regulatory system to relieve CsrA repression of *fliA* mRNA translation (17, 44). In this scenario, a rapid and robust induction of the flagellar and transmission regulons occurs when (p)ppGpp coordinates a parallel increase in RpoS- and RpoN-dependent transcription initiation and a LetA/S-dependent decrease in posttranscriptional repression by CsrA. Consequently, by coupling both transcriptional and translational controls, intracellular *L. pneumophila* can respond to deteriorating conditions by efficiently differentiating to a cytotoxic, motile, and infectious form that is fit to be transmitted to a new host.

The demand theory of gene regulation states that one can predict a positive mode of control for genes that are in high demand in an organism's natural environment and negative control for those genes in low demand (47, 54, 55). For organisms that alternate between different environments, the mode of regulation is predicted to change in accordance with the current demand for the regulated genes. Consistent with the predictions of the demand theory, when conditions are favorable for growth, *L. pneumophila* transmission genes are in low demand and thus are repressed by CsrA and RpoS. Conversely, when the amino acid supply becomes limiting, transmission genes are in high demand and are thus under a positive mode of control, mediated by LetA/S, LetE, FliA, and RpoS (Table 2 and Fig. 1 to 4) (17, 44). Knowledge of the regulatory circuit that controls *L. pneumophila* differentiation can be used to identify the effectors of the replicative and transmission phenotypes. Once the effector genes are identified, detailed analysis of their transcription initiation and mRNA stability can test current genetic models of how RpoS and the LetA/S- and-LetE system cooperate to govern bacterial physiology to allow *L. pneumophila* to tolerate or exploit its surroundings.

#### ACKNOWLEDGMENTS

We gratefully acknowledge Victor DiRita, N. Cary Engleberg, Alex Ninfa, and Ari Molofsky for insight and critical analysis and also advice on the manuscript.

This project was funded by NIH grant AI 44212-01 and the University of Michigan President's Initiative Fund for Graduate Training in Microbial Pathogenesis.

#### REFERENCES

- Ahmer, B. M., J. van Reeuwijk, P. R. Watson, T. S. Wallis, and F. Heffron. 1999. Salmonella SirA is a global regulator of genes mediating enteropathogenesis. *Mol. Microbiol.* **31**:971-982.
- Alli, O. A. T., L.-Y. Gao, L. L. Pedersen, S. Zink, M. Radulic, M. Doric, and Y. Abu Kwaik. 2000. Temporal pore formation-mediated egress from macrophages and alveolar epithelial cells by *Legionella pneumophila*. *Infect. Immun.* **68**:6431-6440.
- Bachman, M. A., and M. S. Swanson. 2001. RpoS co-operates with other factors to induce *Legionella pneumophila* virulence in the stationary phase. *Mol. Microbiol.* **40**:1201-1214.
- Bandyopadhyay, P., and H. M. Steinman. 1998. *Legionella pneumophila* catalase-peroxidases: cloning of the *katB* gene and studies of KatB function. *J. Bacteriol.* **180**:5369-5374.
- Berger, K. H., and R. R. Isberg. 1993. Two distinct defects in intracellular growth complemented by a single genetic locus in *Legionella pneumophila*. *Mol. Microbiol.* **7**:7-19.
- Bozue, J. A., and W. Johnson. 1996. Interaction of *Legionella pneumophila* with *Acanthamoeba castellanii*: uptake by coiling phagocytosis and inhibition of phagosome-lysosome fusion. *Infect. Immun.* **64**:668-673.
- Buck, M., M.-T. Gallegos, D. J. Studholme, Y. Guo, and J. D. Gralla. 2000. The bacterial enhancer-dependent  $\sigma^{54}$  ( $\sigma^N$ ) transcription factor. *J. Bacteriol.* **182**:4129-4136.
- Byrne, B., and M. S. Swanson. 1998. Expression of *Legionella pneumophila* virulence traits in response to growth conditions. *Infect. Immun.* **66**:3029-3034.
- Chen, C. Y., L. Eckmann, S. J. Libby, F. C. Fang, S. Okamoto, M. F. Kagnoff, J. Fierer, and D. G. Guiney. 1996. Expression of *Salmonella typhimurium* *rpoS* and *rpoS*-dependent genes in the intracellular environment of eukaryotic cells. *Infect. Immun.* **64**:4739-4743.
- Ciacciottio, N. P., and B. S. Fields. 1992. *Legionella pneumophila* mip gene potentiates intracellular infection of protozoa and human macrophages. *Proc. Natl. Acad. Sci. USA* **89**:5188-5191.
- Ciacciottio, N. P., R. Long, B. I. Eisenstein, and N. C. Engleberg. 1989. A *Legionella pneumophila* gene encoding a species-specific surface protein potentiates the initiation of intracellular infection. *Infect. Immun.* **57**:1255-1262.
- Dietrich, C., K. Heuner, B. C. Brand, J. Hacker, and M. Steinert. 2001. Flagellum of *Legionella pneumophila* positively affects the early phase of infection of eukaryotic host cells. *Infect. Immun.* **69**:2116-2122.
- Dove, S. L., S. G. Smith, and C. J. Dorman. 1997. Control of *Escherichia coli* type 1 fimbrial gene expression in stationary phase: a negative role for RpoS. *Mol. Gen. Genet.* **254**:13-20.
- Duffy, B. K., and G. Defago. 2000. Controlling instability in *gacA-gacS* regulatory genes during inoculant production of *Pseudomonas fluorescens* biocontrol strains. *Appl. Environ. Microbiol.* **66**:3142-3150.
- Fang, F. C., S. J. Libby, N. A. Buchmeier, P. C. Loewen, J. Switala, J. Harwood, and D. G. Guiney. 1992. The alternative sigma factor KatF (RpoS) regulates *Salmonella* virulence. *Proc. Natl. Acad. Sci. USA* **89**:11978-11982.
- Farewell, A., K. Kvint, and T. Nystrom. 1998. Negative regulation by RpoS: a case of sigma factor competition. *Mol. Microbiol.* **29**:1039-1051.
- Fettes, P. S., V. Forsbach-Birk, D. Lynch, and R. Marre. 2001. Overexpression of the *Legionella pneumophila* homologue of the *E. coli* regulator *csrA* affects cell size, flagellation, and pigmentation. *Int. J. Med. Microbiol.* **291**:353-360.
- Fields, B. S., J. M. Barbaree, E. B. Shotts, Jr., J. C. Feeley, W. E. Morrill, G. N. Sanden, and M. J. Dykstra. 1986. Comparison of guinea pig and protozoan models for determining virulence of *Legionella* species. *Infect. Immun.* **53**:553-559.
- Fischer, G., H. Bang, B. Ludwig, K. Mann, and J. Hacker. 1992. Mip protein of *Legionella pneumophila* exhibits peptidyl-prolyl-*cis/trans* isomerase (PPIase) activity. *Mol. Microbiol.* **6**:1375-1383.
- Hales, L. M., and H. A. Shuman. 1999. The *Legionella pneumophila* *rpoS* gene is required for growth within *Acanthamoeba castellanii*. *J. Bacteriol.* **181**:4879-4889.
- Hammer, B. K., and M. S. Swanson. 1999. Coordination of *Legionella* pneu-



- mophila* virulence with entry into stationary phase by ppGpp. *Mol. Microbiol.* **33**:721–731.
22. Hammer, B. K., E. Tateda, and M. Swanson. 2002. A two-component regulator induces the transmission phenotype of stationary phase *Legionella pneumophila*. *Mol. Microbiol.* **44**:107–118.
  23. Heeb, S., and D. Haas. 2001. Regulatory roles of the GacS/GacA two-component system in plant-associated and other gram-negative bacteria. *Mol. Plant-Microbe Interact.* **14**:1351–1363.
  24. Hengge-Aronis, R. 1996. Regulation of gene expression during entry into stationary phase, p. 1497–1512. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed., vol. 1. ASM Press, Washington, D.C.
  25. Hengge-Aronis, R. 2002. Signal transduction and regulatory mechanisms involved in control of the  $\sigma^S$  (RpoS) subunit of RNA polymerase. *Microbiol. Mol. Biol. Rev.* **66**:373–395.
  26. Heuner, K., L. Bender-Beck, B. C. Brand, P. C. Luck, K.-H. Mann, R. Marre, M. Ott, and J. Hacker. 1995. Cloning and genetic characterization of the flagellum subunit gene (*flaA*) of *Legionella pneumophila* serogroup 1. *Infect. Immun.* **63**:2499–2507.
  27. Heuner, K., C. Dietrich, C. Skriwan, M. Steinert, and J. Hacker. 2002. Influence of the alternative  $\sigma^{28}$  factor on virulence and flagellum expression of *Legionella pneumophila*. *Infect. Immun.* **70**:1604–1608.
  28. Heuner, K., J. Hacker, and B. C. Brand. 1997. The alternative sigma factor  $\sigma^{28}$  of *Legionella pneumophila* restores flagellation and motility to an *Escherichia coli* *flaA* mutant. *J. Bacteriol.* **179**:17–23.
  29. Heuner, K., and M. Steinert. 2003. The flagellum of *Legionella pneumophila* and its link to the expression of the virulent phenotype. *Int. J. Med. Microbiol.* **293**:133–143.
  30. Horwitz, M. A. 1983. Formation of a novel phagosome by the Legionnaires' disease bacterium (*Legionella pneumophila*) in human monocytes. *J. Exp. Med.* **158**:1319–1331.
  31. Horwitz, M. A. 1983. The Legionnaires' disease bacterium (*Legionella pneumophila*) inhibits phagosome lysosome fusion in human monocytes. *J. Exp. Med.* **158**:2108–2126.
  32. Ivanova, A., C. Miller, G. Glinisky, and A. Eisenstark. 1994. Role of *rpoS* (*katF*) in *oxyR*-independent regulation of hydroperoxidase I in *Escherichia coli*. *Mol. Microbiol.* **12**:571–578.
  33. Jishage, M., K. Kvint, V. Shingler, and T. Nystrom. 2002. Regulation of sigma factor competition by the alarmone ppGpp. *Genes Dev.* **16**:1260–1270.
  34. Johnston, C., D. A. Pegues, C. J. Hueck, A. Lee, S. I. Miller. 1996. Transcriptional activation of *Salmonella typhimurium* invasion genes by a member of the phosphorylated response-regulator superfamily. *Mol. Microbiol.* **22**:715–727.
  35. Kirby, J. E., J. P. Vogel, H. L. Andrews, and R. R. Isberg. 1998. Evidence of pore-forming ability by *Legionella pneumophila*. *Mol. Microbiol.* **27**:323–336.
  36. Köhler, R., A. Bubert, W. Goebel, M. Steinert, J. Havker, and B. Bubert. 2000. Expression and use of the green fluorescent protein as a reporter system in *Legionella pneumophila*. *Mol. Gen. Genet.* **262**:1060–1069.
  37. Kvint, K., A. Farewell, and T. Nystrom. 2000. RpoS-dependent promoters require guanosine tetraphosphate for induction even in the presence of high levels of  $\sigma^S$ . *J. Biol. Chem.* **275**:14795–14798.
  38. Lange, R., D. Fischer, and R. Hengge-Aronis. 1995. Identification of transcriptional start sites and the role of ppGpp in the expression of *rpoS*, the structural gene for the  $\sigma^S$  subunit of RNA polymerase in *Escherichia coli*. *J. Bacteriol.* **177**:4676–4680.
  39. Latifi, A., M. Foglino, K. Tanaka, P. Williams, and A. Lazdunski. 1996. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators LasR and RhIR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol. Microbiol.* **21**:1137–1146.
  40. Laurie, A. D., L. M. Bernardo, C. C. Sze, E. Skarfstad, A. Szalewska-Palasz, T. Nystrom, and V. Shingler. 2003. The role of the alarmone (p)ppGpp in sigma N competition for core RNA polymerase. *J. Biol. Chem.* **278**:1494–1503.
  41. Loewen, P. C., B. Hu, J. Strutinsky, and R. Sparling. 1998. Regulation in the *rpoS* regulon of *Escherichia coli*. *Can. J. Microbiol.* **44**:707–717.
  42. Lynch, D., N. Rieser, K. Glogglar, F. Forsbach-Brik, and R. Marre. 2003. The response regulator LetA regulates the stationary-phase stress response in *Legionella pneumophila* and is required for efficient infection of *Acanthamoeba castellanii*. *FEMS Microbiol. Lett.* **219**:241–248.
  43. Magnusson, L. U., T. Nystrom, and A. Farewell. 2003. Underproduction of sigma 70 mimics a stringent response. A proteome approach. *J. Biol. Chem.* **278**:968–973.
  44. Molofsky, A., and M. Swanson. 2003. *Legionella pneumophila* CsrA is a pivotal repressor of transmission traits and activator of replication. *Mol. Microbiol.* **50**:445–461.
  45. Mukherjee, A., Y. Cui, Y. Liu, A. Ishihama, A. Eisenstark, and A. K. Chatterjee. 1998. RpoS (sigma-S) controls expression of *rsmA*, a global regulator of secondary metabolites, harpin, and extracellular proteins in *Erwinia carotovora*. *J. Bacteriol.* **180**:3629–3634.
  46. Mukhopadhyay, S., J. P. Audia, R. N. Roy, and H. E. Schellhorn. 2000. Transcriptional induction of the conserved alternative sigma factor RpoS in *Escherichia coli* is dependent on BarA, a probable two-component regulator. *Mol. Microbiol.* **37**:371–381.
  47. Neidhardt, F. C., and M. A. Savageau. 1996. Regulation beyond the operon, p. 1310–1324. In F. C. Neidhardt, R. Curtiss III, J. L. Ingraham, E. C. C. Lin, K. B. Low, B. Magasanik, W. S. Reznikoff, M. Riley, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella*: cellular and molecular biology, 2nd ed., vol. 1. ASM Press, Washington, D.C.
  48. Nicholson, B., and D. Low. 2000. DNA methylation-dependent regulation of *pef* expression in *Salmonella typhimurium*. *Mol. Microbiol.* **35**:728–742.
  49. Reimann, C., M. Beyeler, A. Latifi, H. Winteler, M. Foglino, A. Lazdunski, and D. Haas. 1997. The global activator GacA of *Pseudomonas aeruginosa* PAO positively controls the production of the autoinducer *N*-butyryl-homoserine lactone and the formation of the virulence factors pyocyanin, cyanide, and lipase. *Mol. Microbiol.* **24**:309–319.
  50. Romeo, T. 1998. Global regulation by the small RNA-binding protein CsrA and the non-coding RNA molecule CsrB. *Mol. Microbiol.* **29**:1321–1330.
  51. Rowbotham, T. J. 1986. Current views on the relationships between amoebae, legionellae and man. *Isr. J. Med. Sci.* **22**:678–689.
  52. Roy, C. R., K. H. Berger, and R. R. Isberg. 1998. *Legionella pneumophila* DotA protein is required for early phagosome trafficking decisions that occur within minutes of bacterial uptake. *Mol. Microbiol.* **28**:663–674.
  53. Sak, B. D., A. Eisenstark, and D. Touati. 1989. Exonuclease III and the catalase hydroperoxidase II in *Escherichia coli* are both regulated by the *katF* gene product. *Proc. Natl. Acad. Sci. USA* **86**:3271–3275.
  54. Savageau, M. A. 1977. Design of molecular control mechanisms and the demand for gene expression. *Proc. Natl. Acad. Sci. USA* **74**:5647–5651.
  55. Savageau, M. A. 1983. Regulation of differentiated cell-specific functions. *Proc. Natl. Acad. Sci. USA* **80**:1411–1415.
  56. Segal, G., M. Purcell, and H. A. Shuman. 1998. Host cell killing and bacterial conjugation require overlapping sets of genes within a 22-kb region of the *Legionella pneumophila* genome. *Proc. Natl. Acad. Sci. USA* **95**:1669–1674.
  57. Segal, G., and H. A. Shuman. 1998. Intracellular multiplication and human macrophage killing by *Legionella pneumophila* are inhibited by conjugal components of IncQ plasmid RSF1010. *Mol. Microbiol.* **30**:197–208.
  58. Sturgill-Koszycki, S., and M. S. Swanson. 2000. *Legionella pneumophila* replication vacuoles mature into acidic, endocytic organelles. *J. Exp. Med.* **192**:1261–1272.
  59. Swanson, M. S., and B. K. Hammer. 2000. *Legionella pneumophila* pathogenesis: a fateful journey from amoebae to macrophages. *Annu. Rev. Microbiol.* **54**:567–613.
  60. Swanson, M. S., and R. R. Isberg. 1995. Association of *Legionella pneumophila* with the macrophage endoplasmic reticulum. *Infect. Immun.* **63**:3609–3620.
  61. Tsui, H. T., G. Feng, and M. E. Winkler. 1997. Negative regulation of *mutS* and *mutH* repair gene expression by Hfq and RpoS global regulators of *E. coli* K-12. *J. Bacteriol.* **179**:7476–7487.
  62. Vogel, J. P., H. L. Andrews, S. K. Wong, and R. R. Isberg. 1998. Conjugative transfer by the virulence system of *Legionella pneumophila*. *Science* **279**:873–876.
  63. Whistler, C. A., N. A. Corbell, A. Sarniguet, W. Ream, and J. E. Loper. 1998. The two-component regulators GacS and GacA influence accumulation of the stationary-phase sigma factor  $\sigma^S$  and the stress response in *Pseudomonas fluorescens* Pf-5. *J. Bacteriol.* **180**:6635–6641.
  64. Wieland, H., M. Faigle, F. Lang, H. Northoff, and B. Neumeister. 2002. Regulation of the *Legionella mip*-promotor during infection of human monocytes. *FEMS Microbiol. Lett.* **212**:127–132.
  65. Xu, J., and R. C. Johnson. 1995. Identification of genes negatively regulated by Fis: Fis and RpoS comodulate growth-phase-dependent gene expression in *Escherichia coli*. *J. Bacteriol.* **177**:938–947.
  66. Zusman, T., O. Gal-Mor, and G. Segal. 2002. Characterization of a *Legionella pneumophila* *relA* insertion mutant and roles of RelA and RpoS in virulence gene expression. *J. Bacteriol.* **184**:67–75.