

## An antibiotic-altered microbiota provides fuel for the enteric foe

Cell Research (2014) 24:5-6. doi:10.1038/cr.2013.142; published online 29 October 2013

**Antibiotic therapies disrupt the intestinal microbiota and render the host susceptible to enteric infections. A recent report by Ng *et al.* explores the ability of two intestinal pathogens (*Salmonella enterica* serovar Typhimurium and *Clostridium difficile*) to use this disruption to their advantage and consume host carbohydrates that would otherwise be unavailable in the presence of a normal gut microbiota.**

The human intestinal microbiota is an extremely dense mixed microbial population within the human gut. It is composed of trillions of microbial cells forming a complex and competitive ecosystem that significantly impacts 'host' human cells as well as invading bacterial pathogens [1]. Disruption of gut microbiota diversity with oral antibiotics increases the risk of enteric infections and also causes a number of immune disorders (e.g., atopic dermatitis, asthma, and type 1 diabetes) [2-4]. A recent article by Ng *et al.* [5] identifies a mechanism by which invading pathogens exploit this disruption in the intestinal microbiota through the uptake of liberated host carbohydrates. Such a discovery suggests that antibiotics not only negatively affect our indigenous microbiota, but they can also actually promote the proliferation of enteric pathogens.

Enteric pathogens interact extensively with the intestinal microbiota [3], therefore it should be no surprise that these pathogens use an antibiotic-disturbed gut microbiota to their advantage. In 2008, Serkirov *et al.* [6]


used increasing doses of vancomycin and streptomycin to disrupt the murine microbiota prior to infecting these mice with *S. Typhimurium* (the causative agent of enteric salmonellosis/gastroenteritis in humans). They found that these antibiotics altered the gut microbiota in a dose-dependent manner. Additionally, in mice treated with increasing doses of antibiotics prior to infection, *S. Typhimurium* was more capable of colonizing the intestinal tract. Stecher *et al.* [7] found that *S. Typhimurium* actually exploits the host's inflammatory immune response to overcome colonization resistance from the intestinal microbiota. Hence, the mechanisms by which these enteric pathogens evade the protection of our gut microbiota are rather complex and more investigation is required.

In a recent paper published in *Nature*, Ng and colleagues, based in Justin Sonnenburg's Stanford laboratory, identify a mechanism by which antibiotic-associated pathogens exploit the increase in mucosal carbohydrate availability that occurs after disruption of the gut microbiota with oral antibiotics [5]. Many commensal and pathogenic bacteria use sialic acids acquired from their hosts as an energy source [8]. However, some bacteria, such as *Bacteroides thetaiotaomicron* (*Bt*), encode a sialidase required to cleave and release sialic acid from the mucosal glycoconjugates, but lack the catabolic pathway to consume it. *S. Typhimurium* and *C. difficile* conversely possess the *nan* operon necessary to consume sialic acid within the lumen of the intestine but

lack the sialidase to liberate it [9, 10].

Ng *et al.* infected both germ-free (microbiota-free) and *Bt*-monoassociated mice with *S. Typhimurium* and *C. difficile* to assess the uptake of free sialic acid by these pathogens in a microbiota-dependent environment. In the mice infected with *S. Typhimurium*, transcriptional profiling revealed that the *nanE* and *fucI* operons (genes encoding catabolic pathways for sialic acid and another monosaccharide, fucose, respectively) were significantly upregulated in the *Bt*-monoassociated mice relative to the germ-free mice. They constructed a mutant strain of *S. Typhimurium* in which they deleted the *nanA* and *fucI* operons. In competition experiments, the mutant strain showed a significant disadvantage in the *Bt*-monoassociated mice, yet this deletion had no effect on germ-free mice. *C. difficile* encodes the *nan* operon necessary for sialic acid consumption but does not encode any genes related to fucose consumption. Ng *et al.* measured the expression of *nanA* and *nanE* upon infection of germ-free and *Bt*-monoassociated mice with *C. difficile* by quantitative RT-PCR. In *Bt*-monoassociated mice both genes exhibited elevated expression levels and an increased density of *C. difficile* relative to germ-free mice. Such findings show that usage of sialic acid by *S. Typhimurium* and *C. difficile* and fucose consumption by *S. Typhimurium* is microbiota-dependent.

The authors also assessed whether sialic acid use is related to pathogen proliferation in antibiotic-treated microbiota. After treatment of conventional



**Figure 1** Schematic overview of antibiotic-altered microbiota experiments. The caecum of conventional mice (normal microbiota) exhibited low sialic acid levels compared to antibiotic-treated mice (altered microbiota). *C. difficile* mutant expansion was significantly compromised in conventional mice and the *C. difficile nan* operon was highly induced in antibiotic-treated mice. *S. Typhimurium* mutants displayed a competitive defect in antibiotic-treated mice relative to conventional mice. All results are consistent with the concept that liberated sialic acid allows for the growth and proliferation of WT *S. Typhimurium* and *C. difficile* within an antibiotic-disturbed gut microbiota.

mice with streptomycin, they quantified free sialic acids in the caeca of these mice and found lower levels of sialic acid in untreated mice relative to the streptomycin-treated mice. *S. Typhimurium* mutants with deletions in the *nan* and *fuc* operons (*St-AnanA* and *St-AnanAΔfucI*) displayed a competitive defect relative to wild-type (WT) *S. Typhimurium*, which is consistent with sialic acid and fucose being necessary for *S. Typhimurium* proliferation. They reported a similar finding in *C. difficile* mutants lacking *nanT* (*Cd-nanT*), as *Cd-nanT*<sup>-</sup> expansion was significantly

compromised in antibiotic-treated mice relative to conventional mice. Furthermore, the *nan* operon in WT *C. difficile* was highly induced in antibiotic-treated mice relative to the *C. difficile in vitro* model. These models support the idea that antibiotics not only disrupt the gut microbiota but they also allow for the release of carbohydrates necessary for the proliferation of enteric pathogens (Figure 1).

Diseases such as vaginal candidiasis, *Clostridium difficile* colitis, and bacterial urinary tract infections typically arise shortly after antibiotic therapy

[6]. As shown by Ng *et al.*, it is unlikely that simple disruption of the gut microbiota homeostasis is the cause for these enteric infections. Further insight into the mechanisms involved in antibiotic disruption of the intestinal microbiota is necessary to determine what molecules or mechanisms each of these pathogens is exploiting to effectively grow and proliferate inside its host. Moreover, this study could potentially lead to novel therapeutic strategies, possibly through probiotic regimens using bacteria that target these carbohydrates for digestion or via the administration of drugs that might inhibit the enzymes used by the gut microbiota to free these carbohydrates [5].

Leah T Stiemsma<sup>1</sup>,  
Stuart E Turvey<sup>2</sup>, Brett B Finlay<sup>1</sup>

<sup>1</sup>Department of Microbiology & Immunology, University of British Columbia, Vancouver, BC, V6T 1Z4; <sup>2</sup>Department of Pediatrics, Child & Family Research Institute, Vancouver, BC, V5Z 4H4

Correspondence: Brett B Finlay  
E-mail: bfinlay@msl.ubc.ca

## References

- 1 Kau AL, Ahern PP, Griffin NW, *et al.* *Nature* 2011; **474**:327-336.
- 2 Penders J, Gerhold K, Stobberingh EE, *et al.* *J Allergy Clin Immunol* 2013; **132**:601-607.
- 3 Kamada N, Seo SU, Chen GY, *et al.* *Nat Rev Immunol* 2013; **13**:321-335.
- 4 Giongo A, Gano KA, Crabb DB, *et al.* *ISME J* 2011; **5**:82-91.
- 5 Ng KM, Ferreyra JA, Higginbottom SK, *et al.* *Nature* 2013; **502**:96-99.
- 6 Sekirov I, Tam NM, Jogova M, *et al.* *Infect Immun* 2008; **76**:4726-4736.
- 7 Stecher B, Robbiani R, Walker AW, *et al.* *PLoS Biol* 2007; **5**:2177-2189.
- 8 Vimr ER, Kalivoda KA, Deszo EL, *et al.* *Microbiol Mol Biol Rev* 2004; **68**:132-153.
- 9 Hoyer LL, Hamilton AC, Steenbergen SM, *et al.* *Mol Microbiol* 1992; **6**:873-884.
- 10 Sebaihia M, Wren BW, Mullany P, *et al.* *Nat Genet* 2006; **38**:779-786.