

RESEARCH PAPER

Olive seed protein bodies store degrading enzymes involved in mobilization of oil bodies

Agnieszka Zienkiewicz^{1,2}, Krzysztof Zienkiewicz^{1,3}, Juan David Rejón¹, Juan de Dios Alché¹, Antonio Jesús Castro¹ and María Isabel Rodríguez-García^{1,*}

¹ Department of Biochemistry, Cell and Molecular Biology of Plants, Estación Experimental del Zaidín, CSIC, 18008 Granada, Spain

² Chair of Plant Physiology and Biotechnology, Nicolaus Copernicus University, 87-100 Toruń, Poland

³ Department of Cell Biology, Nicolaus Copernicus University, 87-100 Toruń, Poland

* To whom correspondence should be addressed. E-mail: mariaisabel.rodriguez@eez.csic.es

Received 9 May 2013; Revised 8 August 2013; Accepted 30 September 2013

Abstract

The major seed storage reserves in oilseeds are accumulated in protein bodies and oil bodies, and serve as an energy, carbon, and nitrogen source during germination. Here, the spatio-temporal relationships between protein bodies and several key enzymes (phospholipase A, lipase, and lipoxygenase) involved in storage lipid mobilization in cotyledon cells was analysed during *in vitro* seed germination. Enzyme activities were assayed in-gel and their cellular localization were determined using microscopy techniques. At seed maturity, phospholipase A and triacylglycerol lipase activities were found exclusively in protein bodies. However, after seed imbibition, these activities were shifted to the cytoplasm and the surface of the oil bodies. The activity of neutral lipases was detected by using α -naphthyl palmitate and it was associated mainly with protein bodies during the whole course of germination. This pattern of distribution was highly similar to the localization of neutral lipids, which progressively appeared in protein bodies. Lipoxygenase activity was found in both the protein bodies and on the surface of the oil bodies during the initial phase of seed germination. The association of lipoxygenase with oil bodies was temporally correlated with the appearance of phospholipase A and lipase activities on the surface of oil bodies. It is concluded that protein bodies not only serve as simple storage structures, but are also dynamic and multifunctional organelles directly involved in storage lipid mobilization during olive seed germination.

Key words: Lipase, lipoxygenase, oil bodies, olive, phospholipase A, protein bodies.

Introduction

Seed germination is accompanied by intense metabolic activity of cells, including respiration and protein synthesis, as well as multiple cell divisions, cell elongation, and complex structural changes at the subcellular level. Establishment of the seedling is the result of proper seed germination, and seedling growth is initially supported by metabolites produced by the hydrolysis and conversion of the major stored seed reserves, such as proteins and lipids (Bewley and Black, 1994; Zienkiewicz *et al.*, 2011).

Seed storage proteins (SSPs) are degraded during germination and used for seedling growth. These proteins provide

carbon, nitrogen, and sulphur resources for subsequent seedling development (Müntz, 1996; Herman and Larkins, 1999). SSPs are accumulated during seed maturation in the protein storage vacuoles (PSVs) of the embryo and endosperm cells. At the late stage of seed maturation, the PSVs are transformed into protein bodies (PBs), which are surrounded by a membrane derived from the vacuolar membrane (Hara-Nishimura *et al.*, 1987; Strzałka *et al.*, 1995). Seed storage triacylglycerols (TAGs) are accumulated in the embryo and endosperm during seed development and also provide the sources of carbon and energy for germination

and post-germinative growth of seedlings (Napier *et al.*, 1996). The storage lipids are assembled in discrete spherical organelles of 0.5–2.5 µm called oil bodies (OBs) or lipid bodies (Murphy, 2012). The OB consists of a core of TAGs covered by a single layer of phospholipids with proteins embedded. These proteins (oleosin, caleosin, and steroleosin) are responsible for the stability of OBs and prevent the degradation of storage lipids until the seed germinates (Tai *et al.*, 2002; Lin and Tzen, 2004; Poxleitner *et al.*, 2006). Storage lipid mobilization is initiated by the activity of endogenous lipases, which hydrolyse storage TAGs to glycerol and free fatty acids (Quettier and Eastmond, 2009). *SUGAR-DEPENDENT (SDPI)* and *SDPI-LIKE (SDPIL)* encode a TAG lipase with a patatin-like acyl-hydrolase domain that can associate with the OB surface and is capable of hydrolysing TAGs in preference to diacylglycerol or monoacylglycerol (Eastmond, 2006; Kelly *et al.*, 2011). In an alternative pathway for TAG mobilization, a specific lipoxygenase (LOX) enzyme is involved. Matsui *et al.* (1999) found that trypsin *in vitro* digestion of OB-associated proteins led to the oxygenation of TAGs by the action of LOX in cucumber cotyledons. This lipoxygenase is capable of catalysing stereospecific oxygenation of the linoleate moieties of TAGs to (9Z,11E 1,2Z)-13-hydroperoxy octadeca-9,11-dienoic acid (13-HPOD) (Feussner and Wasternack, 1998; Gerhardt *et al.*, 2005). More recently, it was reported that a patatin-like phospholipase (PLA) promotes the LOX-dependent oxygenation of OB phospholipids in cucumber cotyledons (Rudolph *et al.*, 2011).

Seed OBs have often been shown to associate with other organelles such as glyoxysomes, PSVs, or PBs (Fernandez and Staehelin, 1987; Hayashi *et al.*, 2001; Poxleitner *et al.*, 2006). There is agreement that this interaction could help breach the OB membrane in order to trigger the process of TAG degradation (Hayashi *et al.*, 2001; Poxleitner *et al.*, 2006). A direct relationship between OBs and glyoxysomes is thought to facilitate fatty acid release and their transfer between the two organelles. Moreover, this process requires the participation of a lipase active at neutral pH (Hayashi *et al.*, 2001). Other data suggested that PBs store lipases in their membranes; thus, one of the possible functions of the OB–PB association might be to join the enzyme and its substrate (Fernandez and Staehelin, 1987). Previous results showed that in mature olive seed the cotyledon cells contain large PBs surrounded by numerous OBs (Alché *et al.*, 2006; Zienkiewicz *et al.*, 2011). During seed germination, a progressive decrease of OB and PB number was observed, which is accompanied by a close spatial relationship between these two organelles (Zienkiewicz *et al.*, 2011).

To date, despite extensive studies of TAG mobilization during seed germination, little is known about the nature of PB–OB interaction. Here, new evidence is reported on the direct involvement of PBs in breakdown of OBs during seed germination. By analysing the spatio-temporal behaviour of lipase, LOX, and PLA, it was demonstrated that PBs are spatially and functionally connected to storage lipid mobilization.

Materials and methods

Plant material

Mature seeds of *Olea europaea* L. were obtained from olive trees (cv. 'Picual') grown in the Estación Experimental del Zaidín (Granada, Spain).

In vitro germination of olive embryos

In vitro germination of olive embryos was carried out as described by Cañas and Benbadis (1988). Cotyledons were collected from mature and imbibed (24h) seeds and at different times of *in vitro* germination (6h and 3 d) and seedling growth (4, 8, 15, and 26 d).

RNA isolation and cDNA synthesis

Frozen samples were ground in liquid nitrogen using a mortar and pestle. Total RNA was extracted using an RNeasy Plant Total RNA kit (Quiagen, Germany). First-strand cDNA was synthesized with 0.5 µg total of RNA, oligo(dT)₁₉ primer (0.5 µg), and reverse transcriptase (Fermentas, Germany) according to the manufacturer's instructions.

Quantitative real-time PCR (qRT-PCR)

Gene expression analysis was performed by qRT-PCR using an iCycler (Bio-Rad, USA). Primers for gene-specific amplification (Supplementary Table S1 available at *JXB* online) were designed using the Primer3 program (http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi). Target genes (and the olive ubiquitin2 gene as a housekeeping marker; Padilla *et al.*, 2012) were subjected to qRT-PCR in 96-well optical reaction plates in 20 µl mixtures per well, using iQTM SYBR[®] Green Supermix (Bio-Rad) and following the manufacturer's instructions. Cycle threshold (Ct) values were obtained with the iQTM software (Bio-Rad) and data were analysed with the 2^{ΔΔCT} method (Livak and Schmittgen, 2001). The data are presented as means ± standard deviation (SD) of three biological repeats, obtained from three independent experiments.

Isolation of protein and oil bodies

PBs and OBs were isolated from cotyledons at different times of *in vitro* germination according to Torrent *et al.* (1989) and Zienkiewicz *et al.* (2010), respectively.

Protein extraction

Material was powdered in liquid nitrogen and suspended in 1.5 ml of extraction buffer (0.05 M phosphate buffer, pH 7.0). Total proteins and proteins from the PB fraction were eluted under continuous and vigorous stirring at 4 °C for 2h. The samples were then centrifuged at 13 500 g for 30min at 4 °C and the resulting supernatants were used for activity assays and Western blot analysis. OB-associated proteins were extracted as described by Zienkiewicz *et al.* (2010). Protein content in each extract was measured by using a commercial Bradford procedure (Bio-Rad).

SDS–PAGE and immunoblotting

SDS–PAGE was performed according to Laemmli (1970) on 12% (w/v) acrylamide gels with 4.5% stacking gels. Total proteins (50 µg per sample) were mixed with an equal volume of 2× SDS sample buffer (Laemmli, 1970) and boiled for 3min prior to gel loading. After electrophoresis, the resulting gels were stained by Coomassie Brilliant blue (CBB), or were transferred onto PVDF membranes in a Semi-dry Transfer Cell (Bio-Rad). The membranes were blocked in Tris-buffered saline (TBS) buffer containing 0.5% (w/v) non-fat dry milk for 1h. Immunodetection of LOX was carried out

by incubation with a polyclonal anti-*Glycine max* (soybean) LOX antibody (Ab) (Agrisera, Sweden) diluted 1:1000 in TBS buffer for 12 h at 4 °C. A DyLight 488 conjugated anti-rabbit IgG (Agrisera), diluted 1:2000 in TBS buffer for 2 h, served as the secondary Ab. The signal was detected in a Pharos FX molecular imager (Bio-Rad). Densitometric measurements were carried out from images of membranes using Quantity One 4.6.2 software (Bio-Rad).

In-gel lipase and LOX activity

SDS-PAGE was performed as above, but the sample boiling step was omitted. After electrophoresis, SDS was removed from polyacrylamide gels by washing them three times for 30 min each in a solution containing 0.05 M phosphate buffer (pH 7.0) and 2.5% (v/v) Triton X-100. Lipase activity was measured as previously described (Rejón *et al.*, 2012). LOX activity assays were prepared according to Heinisch *et al.* (1996). The gel was incubated for 30 min in 50 ml of solution containing 0.2 M borate buffer (pH 9.0) and 50 µl of α -linolenic acid (dissolved in 50 µl of ethanol). Subsequently, the gel was rinsed briefly in distilled water and incubated with 100 ml of a solution containing 0.5 g of *N,N*-dimethyl-*p*-phenylenediamine, 4.5 ml of methanol, and 0.5 ml of acetic acid.

Histochemical studies

Material was processed for light microscopy as previously described (Zienkiewicz *et al.*, 2011). Protein material and neutral lipids were stained according to Serrano *et al.* (2008). The samples were observed in a LM Zeiss Axioplan microscope (Carl Zeiss, Germany) and images were obtained with a ProGres C3 digital camera using the ProGres CapturePro 2.6 software (Jenoptik AG, Germany).

Microscopic immunolocalization of LOX

Semi-thin sections were incubated overnight at 4 °C with an anti-LOX Ab [diluted 1:50 in 1% bovine serum albumin (BSA) in phosphate-buffered saline (PBS) pH 7.2], following by an anti-rabbit IgG DyLight 488-conjugated secondary Ab (diluted 1:200 in 1% BSA in PBS pH 7.2) for 1 h at 37 °C. Samples were observed in a Zeiss Axioplan microscope using the filter combination: BP365, FT395, and LP397. For immunogold experiments, ultra-thin sections were incubated with an anti-LOX Ab (diluted 1:50 in 1% BSA in PBS pH 7.2), overnight at 4 °C, followed by an anti-rabbit IgG 15 nm gold-conjugated secondary Ab (British Biocell International, UK) diluted 1:100 in 1% BSA in PBS pH 7.2 for 1 h. After washing, sections were stained with 5% (w/v) uranyl acetate for 30 min. Samples were observed in a JEM-1011 transmission electron microscope (JEOL, Japan) operating at 80 kV. Negative controls were prepared by omitting the incubation with the primary Ab or by incubation with a pre-immune rabbit serum.

Light microscopic localization of lipase and LOX activity

Lipase activity was revealed by incubating the semi-thin sections for 30 min at 37 °C in a developing solution containing 40 mg of α -naphthyl palmitate prepared in 16 ml of *N,N*-dimethylformamide and 80 mg of Fast blue BB salt in 144 ml of 0.1 M phosphate buffer, pH 7.0. LOX activity was detected by incubating the semi-thin sections for 30 min at room temperature in a solution containing α -linolenic acid. Then, the sections were stained with a solution containing 0.5 g of *N,N*-dimethyl-*p*-phenylenediamine, 4.5 ml of methanol, and 0.5 ml of acetic acid. Observations were carried out with an Axioplan microscope.

In situ localization of lipase and PLA activity

For whole-mount localization, olive cotyledons were cut into small pieces using a scalpel. Detection of neutral lipids was carried out by using Nile Red. Samples were incubated for 10 min in a solution

containing 0.05 mg ml⁻¹ Nile Red (Sigma-Aldrich) dissolved in acetone. Samples were observed with a Nikon C1 confocal laser scanning microscope (CLSM) (Nikon, Japan) using an argon (488) laser. Z-series images were collected and processed with the software EZ-C1 Gold version 2.10 build 240 (Nikon). Detection of cellular membranes was performed by using FM4-64 dye according to the manufacturer's instructions (Molecular Probes, USA). SYTOX green nucleic acid stain was used for determination of cotyledon cell viability and was performed according to the manufacturer's instructions (Molecular Probes). Heat treatment (95 °C for 10 min) was used as a positive control of cells with permeabilized membranes. In order to detect lipase activity, samples were incubated for 30 min in an aqueous solution of 25 µg ml⁻¹ resorufin ester (lipase substrate, Sigma-Aldrich). Samples were observed with a Nikon C1 CLSM using a He-Ne (549 nm) laser. Phospholipase A₁ and A₂ activities were detected using BODIPY[®] FL C₁₁-PC (Molecular Probes). Samples were incubated for 10 min in 1 mM BODIPY[®] FL C₁₁-PC ethanolic solution and observed with a Nikon C1 CLSM using an argon (488 nm) laser. *In situ* localization of lipase and PLA activity was also performed in isolated OBs as described previously by Zienkiewicz *et al.* (2013)

Immunolocalization experiments in isolated OBs

Purified OBs were incubated in a 1.5 ml tube with the anti-LOX Ab, rabbit anti-ubiquitin (Alché *et al.*, 2000), or the mixture of anti-LOX and chicken anti-Clo (Davids Biotechnologie, Germany) Abs (diluted 1:30 in PBS buffer, pH 7.2, containing 1% BSA), for 2 h at room temperature, followed by incubation with an anti-rabbit IgG DyLight 550-conjugated secondary Ab (Agrisera), and/or an anti-chicken IgY DyLight 448-conjugated secondary Ab (Agrisera) diluted 1:200 in PBS buffer, pH 7.2 for 1 h at 37 °C under gentle agitation. Samples were observed with a Zeiss Axioplan epifluorescence microscope under blue and/or green light irradiation. Negative controls were treated as above, but the primary Ab was omitted or pre-immune serum was used instead of the primary Ab.

Results

In-gel detection of lipase activity

Lipase activity was detected after protein separation on an SDS-polyacrylamide gel and after incubation with α -naphthyl palmitate as the substrate (Fig. 1A–D). At seed maturity, up to seven different lipolytic bands were observed in the total (Fig. 1A) and cytoplasmic fraction (Fig. 1B), with molecular weights ranging from 90 kDa to 30 kDa. After seed imbibition, only the lipolytic bands of 53, 48, 40, 34, and 32 kDa were visible in both fractions (Fig. 1A, B, arrows). The band of 53 kDa showed a higher intensity compared with the other bands and was detectable until the third day of *in vitro* germination, whereas the band of 40 kDa was visible during the whole course of *in vitro* germination and seedling growth. In PB extracts, only one lipolytic band of 53 kDa was detected during the course of *in vitro* germination (Fig. 1C). No lipolytic bands were detected in the OB fraction (Fig. 1D).

Expression and activity of LOXs

OeLOX1 and *OeLOX2* mRNA levels were found to increase significantly after seed imbibition. The highest level of *OeLOX1* was observed at 6 h of *in vitro* germination and was followed by a decrease after 3 d of *in vitro* germination (Fig. 2A). During the next stages of seedling growth,




Fig. 1. In-gel assay of lipase activity in the total protein extract (A) and in proteins associated with the cytoplasmic (B), PB (C), and OB (D) fractions. Major activity bands are indicated with arrows. M, protein markers; MS, mature dry seed; IM, seed after imbibition (24 h); 6 h, and 3 d, time of *in vitro* germination; 4 d, 8 d, 15 d, and 26 d, days of seedling growth.

OeLOX1 mRNA levels remained unchanged. High levels of *OeLOX2* transcripts were observed until the fourth day of seedling growth (Fig. 2B). A significant decrease in *OeLOX2* mRNA levels occurred at the final analysed steps of seedling growth. Western blot experiments showed the presence of two bands of ~100 kDa and 98 kDa, which were cross-recognized by the anti-LOX Ab (Fig. 2C). The specificity of this Ab was demonstrated previously (Zienkiewicz *et al.*, 2013). The LOX levels significantly increased after seed imbibition and during the subsequent steps of *in vitro* germination (Fig. 2C). No LOX proteins could be detected at the last analysed stage of seedling growth. In-gel assays of LOX activity indicated the presence of three LOX isozymes of ~100, 98, and 96 kDa (Fig. 2D). The highest LOX activities were observed just after seed imbibition and during the first days of *in vitro* germination. Later steps of seedling growth were accompanied by a gradual decrease in all LOX activities (Fig. 2D).

Distribution of total proteins, neutral lipids, and lipase activity in olive cotyledons

CBB and Sudan Black B (SBB) stains were used to study the distribution of proteins and storage lipids in olive cotyledons, respectively. To analyse the potential correlation in the distribution of proteins, lipids, and lipase activity, α -naphthyl palmitate was used as a substrate. In mature seeds, proteins were detected in the area of PBs (Fig. 3A), while storage lipids were

accumulated in the form of numerous OBs located around PBs (Fig. 3A1). At this stage, lipase activity was detected mainly in PBs, with a high concentration on their boundaries (Fig. 3A2, arrows). After seed imbibition, protein-free areas appeared in the PB matrix (Fig. 3B, arrows). In some PBs, rounded areas stained by SBB were also found (Fig. 3B1, arrows). Lipase activity was detected mainly in the area of PBs (Fig. 3B2, arrows). After 6 h of *in vitro* germination, a rich pool of proteins was observed in the area of PBs and in the cytoplasm (Fig. 3C). SSB-stained areas in the matrix of the PBs were more numerous and more intensively stained when compared with imbibed seed (Fig. 3C1, arrows). Lipase activity was detected in the area of PBs as well as in the cytoplasm surrounding the OBs (Fig. 3C2). After 3 d of *in vitro* germination, proteins were uniformly distributed in some PBs, whereas in others they were visible as irregular clusters (Fig. 3D). Lipids were observed in the area of PBs in the form of irregular clusters and also as rounded areas after SSB staining (Fig. 3D1, arrows). The lipase activity was found in the matrix of the PBs, especially concentrated around lipid-containing PB areas (Fig. 3D2, insert). On the fourth day of seedling growth, when in most of the cotyledon cells only one, large PB was present, proteins and lipids were found in both the cytoplasm and PB matrix (Fig. 3E, E1). Lipase activity was detected mainly in the cytoplasm surrounding the central PB (Fig. 3E1). After 15 d of seedling growth, cells displayed differentiating chloroplasts, and proteins were detected in the




Fig. 2. Relative expression levels of *OeLOX1* (A) and *OeLOX2* (B) determined by qRT-PCR using the expression level of the genes in mature seed cotyledons as the calibrator. (C) Immunodetection of lipoxigenase (arrows). (D) In-gel assay of lipoxigenase activity (arrows). MS, mature dry seed; IM, seed after imbibition (24h); 6h and 3d, time of *in vitro* germination; 4 d, 8 d, 15 d, and 26 d, days of seedling growth.

chloroplast, cytoplasm, and the lumen of the central vacuole (Fig. 3F). Lipids were found homogeneously distributed in the lumen of the central vacuole and in the cytoplasm (Fig. 3F1). Lipase activity was located mainly in the cytoplasm surrounding the central vacuole. Less intense staining was also observed in the chloroplasts (Fig. 3F2). Control reactions were performed by omitting lipase substrate and did not show any labelling (Supplementary Fig. S1A at JXB online).

Localization of LOX proteins and activity in olive cotyledons

The cellular localization of LOX proteins and LOX activity was examined in olive cotyledons by using fluorescence and light microscopy, respectively. In mature seeds, the fluorescent labelling was detected in the area of PBs (Fig. 4A). Purple colour, indicating LOX activity, was visible mainly in PBs (Fig. 4A, inset). After seed imbibition, the intensity of the fluorescence and of LOX activity staining increased in PBs as well as in the cytoplasm surrounding OBs (Fig. 4B and inset).

After 6h of *in vitro* germination, high fluorescent labelling was observed in the area of PBs and cytoplasm (Fig. 4C). At this stage, significant LOX activity was visible on the boundary of OBs (Fig. 4C, inset). After 3 d of *in vitro* germination, both LOX proteins and activity were found in numerous, rounded areas located in the matrix of the PBs (Fig. 4D, circles). Moreover, intense LOX activity was attached to the boundary of OBs (Fig. 4D, inset). After 4 d of seedling growth, LOX antigen was located inside large PBs and in the surrounding cytoplasm (Fig. 4E). Weak LOX activity was visible in the area of PBs and cytoplasm (Fig. 4E, inset). After 15 d, when a large central vacuole was formed, the fluorescent labelling and LOX activity were detected mainly in the cytoplasm surrounding the central vacuole as well as in the chloroplasts (Fig. 4F and inset). Control reactions did not show any labelling (Supplementary Fig. S1B–D at JXB online).

Gold immunolabelling of LOX proteins provided additional details about its subcellular localization, and confirmed the tissue distribution pattern observed by fluorescence microscopy. In the mature seed, the gold particles were located in the




Fig. 3. (A–F) Coomassie Brilliant blue (CBB) staining of total proteins in sections from olive cotyledons. (A1–F1) Sudan black B (SBB) staining of neutral lipids in sections from olive cotyledons. The arrows denote lipidic masses localized in the protein bodies. (A2–F2) Cellular localization of lipase activity in sections from olive cotyledons. Ch, chloroplast; PB, protein body; V, vacuole. Bars=25 μ m.

area of PBs (Fig. 5A). After seed imbibition, the labelling was found in the area of PBs (Fig. 5B, arrows) as well as on the boundaries of OBs (Fig. 5B, arrowheads). A similar pattern of labelling was observed 6h after imbibition (Fig. 5C). After 3 d of *in vitro* germination, the gold particles were detected

in the electron-dense areas located in the PB matrix (Fig. 5D, circle). In the large PBs formed after 4 d of seedling growth, gold particles were found in the area of the PBs (Fig. 5E, arrows) and OBs (Fig. 5E, arrowhead). As cotyledon development progressed (15th day), the signal was localized in the




Fig. 4. Localization of lipooxygenases (anti-LOX Ab) (A–F) with their activity (A–F, insets) in sections from olive cotyledons. Fluorescent labelling of LOX is located mainly in the area of PBs. After 6 h of *in vitro* germination, the fluorescence appears in the OB boundary (arrows) and in the cytoplasm. An intense LOX activity is found at the boundaries of PBs connected with OBs. Significant LOX activity is observed on the surface of the OBs (insets, arrows). Bars=10 μ m.

chloroplasts as well as in the surrounding cytoplasm (Fig. 5F, arrows). Control reactions did not show any gold labelling (Supplementary Fig. S1E, F at JXB online).

Cellular localization of lipase and PLA activities in living olive cotyledons

The viability of the cotyledon cells, their intact structure, and the integrity of their membranes were confirmed by using FM4-64 dye and SYTOX green (Supplementary Figs S2, S3 at JXB online). After Nile Red staining, the OBs in olive cotyledon cells were visible as numerous red-orange fluorescence

spots surrounding non-labelled PBs (Figs 6A–C, 7A–C). *In situ* lipase activity was detected by using a lipase substrate (resorufin ester). The product of its hydrolysis emitted red fluorescence after excitation with light of 544 nm (Figs 6D–F, 7D–F). BODIPY[®] FL C₁₁-PC was used for detection of PLA activities. The cleavage product generated from this substrate by PLAs can be detected as green fluorescence under the wavelength of 488 nm (Figs 6G–I, 7G–I). In the cotyledons from mature, dry seed, the lipase activity was observed exclusively in the area of PBs (Fig. 6D–F). A similar localization pattern was found for products of PLA activity (Fig. 6G–I). After seed imbibition, the lipase activity was located inside




Fig. 5. (A–F) Ultrastructural localization of LOX in olive cotyledons. Gold particles are located on the OB boundaries (arrowheads) and in the PBs (arrows). Some labelling is also present in the surrounding cytoplasm and chloroplasts (arrows). Ch, chloroplast; OB, oil body; PB, protein body; W, cell wall. Bars=1 μm .

PBs as well as in the area occupied by OBs (Fig. 7D–F). The green fluorescence, indicating PLA activity, was located exclusively in the area occupied by OBs (Fig. 7G–I, arrows). No fluorescent labelling was found in PBs.

Detection of oil body-associated enzymes in olive cotyledons

Purified OBs were visible as spherical structures (0.2–2 μm) that stained positively with Nile Red (Supplementary Fig. S4 at *JXB* online). The purity of isolated OBs was confirmed by the absence of ubiquitin (cytoplasmic marker) on their surface (Supplementary Fig. S5A, B). Additionally, the identity of isolated OBs and the integrity of their membrane were verified by co-localization of LOX and caleosin (Supplementary Fig.

S5C–E). OBs isolated from cotyledons at seed maturity did not show the presence of lipase activity (Fig. 8A). After imbibition, a strong lipase activity appeared on the surface of the OBs as ring-shaped red fluorescence (Fig. 8B). The activity of PLAs was detected only on the surface of OBs isolated from imbibed seeds (Fig. 8C, D). The presence of LOX protein on the surface of OBs was also tested. The LOX was detected only on the surface of OBs isolated from cotyledons of imbibed seeds, but it was absent in OBs isolated from mature, dry seed (Fig. 8E, F).

Discussion

In germinating oilseeds, such as *Arabidopsis* or cucumber, neutral lipids stored in OBs are mobilized to provide the carbon skeletons and energy necessary for their post-germinative




Fig. 6. Localization of OB lipase (resorufin ester) and phospholipase A (BODIPY[®] FL C₁₁-PC) activities in living olive cotyledons dissected out from mature seed by confocal laser scanning microscopy. (A–C) Olive cotyledons stained with Nile Red. Numerous OBs are indicated with arrows. (D–F) Olive cotyledons incubated with resorufin ester. Lipase activity is located mainly in the area of PBs. (G–I) Olive cotyledons incubated with BODIPY[®] FL C₁₁-PC. Green fluorescent labelling is detected exclusively in the PB area. PB, protein body. Bars=25 μm, inset bars=10 μm.

growth (Graham, 2008). This process occurs by the action of hydrolytic enzymes such as phospholipase, LOX, and lipase at different stages of seed germination (Rudolph *et al.*, 2011). Here, the changes in the *in vivo* localization of PLA activity in cotyledons during seed imbibition are reported for the first time. Interestingly, the PLA activity in the cotyledons from mature olive seed was associated with PBs, while after seed imbibition PLA activity was evident in the cytoplasm surrounding the OBs and on their surface. To date PLA activity had been detected only on the OB surface and in the cytoplasm of sunflower cotyledon protoplasts. PLA was expressed or activated on the surface of OBs during the early phase of seedling development, prior to the action of lipase (Bhatla *et al.*, 2009). In the light of the microscopy findings reported here, PLA activity in the olive cotyledons could be shifted from PBs to OBs concomitantly with the progress of seed germination. Moreover, a close temporal correlation was found between PLA and lipase activities during olive seed germination. These data suggest that PLA activity associated with the surface of OBs is responsible for the phospholipid breakdown leading to TAG mobilization by lipase.

Lipase is major enzyme associated with the seed (Eastmond, 2006; Kelly *et al.*, 2011). Lipase is known to be associated with the membranes of OBs, glyoxysomes, and with PBs (Fernandez and Staehelin, 1987; Beisson *et al.*, 2001; Hayashi *et al.*, 2001). Using α -naphthyl palmitate as an enzyme substrate, a group of seven putative lipases were identified in olive cotyledons. Interestingly, only 53 kDa lipase activity was present in PBs during the early steps of seed germination. The lipase activity was detectable in PBs mainly in the areas where lipids were present. Indeed, some studies suggested that lipase could be initially localized at its storage sites in the PBs or PSVs (Fernandez and Staehelin, 1987; Bhatla *et al.*, 2009). Moreover, it was demonstrated that OBs interact with the PSV tonoplast in the germinating *Arabidopsis* seed, and this interaction might be promoted by caleosin of the OBs (Poxleitner *et al.*, 2006). These data suggest that the interaction of OBs with the membrane of another organelle could help breach the phospholipid monolayer of the OB, providing access to lipase. The present results indicate that lipase present in PBs could act by facilitating the metabolism of storage lipids in PBs during seed germination. After using α -naphthyl




Fig. 7. Localization of OB lipase (resorufin ester) and phospholipase A (BODIPY[®] FL C₁₁-PC) activities in living olive cotyledons dissected out from imbibed seed by confocal laser scanning microscopy. (A–C) Optical section of olive cotyledons stained with Nile Red. Numerous OBs are indicated with arrows. (D–F) Olive cotyledons incubated with resorufin ester. Lipase activity is detected in the area of PBs as well as on the OB surface (arrows). (G–I) Olive cotyledons incubated with BODIPY[®] FL C₁₁-PC. Green fluorescent labelling was detected only on the OB surface (arrows). PB, protein body. Bars=25 μm, inset bars=10 μm.

palmitate as the lipase substrate, lipase activity was not shown on the surface of the OBs. However, the presence and activity of an alternative lipase responsible for OB hydrolysis cannot be ruled out. A previous study described evidence of a 95 kDa patatin domain TAG lipase (SDPI) associated with purified OBs and involved in their mobilization in germinating *Arabidopsis* seeds (Eastmond, 2006). In order to detect TAG lipase activity in olive OBs, resorufin ester was used as a marker for lipase intracellular localization. Lipase activity was initially localized in PBs prior to seed germination, this activity becoming progressively transferred to the OB surface during early steps of olive seed germination. This pattern was also observed in sunflower cotyledon protoplasts, where lipase activity shifted from the PSVs to OBs during seed germination (Bhatla *et al.*, 2009). The different substrate-dependent localization patterns of lipase activity in olive cotyledons suggest that some lipases may act only in PBs, whereas others may hydrolyse TAGs in both PBs and OBs.

Fatty acids released from TAGs may be further degraded either via the β -oxidation in glyoxysomes or by an alternative

pathway, which is dependent on a LOX enzyme (Feussner *et al.*, 2001; Graham, 2008). A recent study found evidence that the lipid breakdown by LOX in cucumber cotyledons is promoted by a patatin-like phospholipase (Rudolph *et al.*, 2011). The anti-LOX Ab used in this work recognized two protein bands, of 98 kDa and 100 kDa, on western blots. However, the LOX activity profile revealed the existence of a third LOX isoenzyme with an estimated mol. wt of 96 kDa. The sizes of olive cotyledon LOXs are well within the molecular weight range (95–100 kDa) reported for other plant LOXs (Andreou and Feussner, 2009). The present study also detected a smaller protein band of 63 kDa, which showed enhanced LOX activity. In a previous study (Zienkiewicz *et al.*, 2013) a 63 kDa protein band with LOX activity was also detected in olive pollen. Interestingly, limited proteolysis of soybean LOX1 generated a 60 kDa fragment with enhanced activity and membrane binding ability (Maccarrone *et al.*, 2001). Although LOX expression and activity subsequently increased during the initial steps of *in vitro* germination, a further decrease occurred coincidentally with the reduction in the number of OBs




Fig. 8. Detection of lipase (resorufin ester) and phospholipase A (BODIPY[®] FL C₁₁-PC) activity and lipoxygenase (anti-LOX Ab) on OBs isolated from cotyledons dissected out from mature and imbibed seed. (A and B) Lipase activity visible as red fluorescence is located on the OB surface in imbibed seed (arrows). OBs isolated from mature seed do not show any labelling. (C and D) Green fluorescence is detected only on the surface of OBs (arrows) isolated from cotyledons after seed imbibition. (E and F) LOX is detected on the surface of OBs (arrows) in imbibed seed but it is absent in OBs isolated from cotyledons of mature seed. Bars=10 μ m.

(Zienkiewicz *et al.*, 2011) in the cytoplasm of olive cotyledons. A similar pattern was also observed in barley embryos, in which levels of LOX expression and activity declined during the last phase of germination (Holtman *et al.*, 1996). Olive LOXs were mainly localized in PBs as well as on the surface of OBs. Here, the presence of LOXs in the PBs during *in vitro* germination was detected for the first time. The localization of LOXs in both the cytoplasm and PBs suggests that this enzyme may be shifted from PBs to the cytoplasm and the OB surface. Interestingly, LOXs were specifically localized in rounded intra-PB areas. A previous study reported that these areas do not contain SSPs but do contain peroxidase (POX) (Zienkiewicz *et al.*, 2011). It has been showed that fatty acid peroxides, which are products of LOX activity, subsequently serve as substrates for POX (Koljak *et al.*, 1997; Brash, 1999). A functional relationship between LOX and POX has been previously studied in developing oat plants at four stages:

germination, growth, senescence, and dark-incubated senescence (Yi *et al.*, 2005). LOX and POX were found to exhibit similar activity patterns in all examined stages (Yi *et al.*, 2005). Specific co-localization of LOXs and POX in PBs of olive cotyledon seems to confirm this functional relationship.

Conclusions

PBs from olive cotyledons store PLA and lipase at seed maturity, which are transferred to the cytoplasm and the surface of OBs during early steps of germination. Meanwhile, the activity of neutral lipases is limited mainly to PBs and their distribution is closely correlated with the appearance of neutral lipids in the PB area as germination progresses. LOX enzymes act both in PBs and on the surface of OBs during the initial phase of seed germination. At this time, OB-associated

LOX activity is temporally correlated with the appearance of PLA and lipase activities on the OB surface. To the authors' knowledge, this is the first report indicating the presence of diverse lipolytic enzymes inside the PBs implicated in the mobilization of storage lipids during seed germination. It is proposed that PBs should be considered not only as simple storage structures but also as dynamic and multifunctional organelles.

Supplementary data

Supplementary data are available at *JXB* online.

Figure S1. Negative controls.

Figure S2. Fluorescent labelling of cellular membranes with FM4-64 dye in the olive cotyledon cells.

Figure S3. Analysis of cell viability in fresh cotyledon tissue by using SYTOX green.

Figure S4. Purified oil bodies (arrows) stained with Nile Red and observed under transmitted light (A) and under fluorescence (B).

Figure S5. Localization of ubiquitin on isolated OBs (A) using the anti-UBQ Ab, showing the absence of ubiquitin (cytoplasmic marker) on their surface (B). Co-localization of caleosin (C) and LOX (D) on isolated OBs using the anti-Clo3 and anti-LOX Abs (E).

Table S1. Gene accession numbers and sequences of the primers used in this study.

Acknowledgements

This work was supported by the ERDF-co-financed projects AGL2008-00517 and BFU2011-22779 (Spanish Ministry of Science and Innovation) and P2010-CVI5767, P2010-AGR6274, and P2011-CVI7487 (Andalusian Regional Government). AZ thanks the CSIC for providing JAEdoc grant funding. Microscopy work was carried out at the CTEM facilities of the EEZ-CSIC Institute (Granada, Spain). We also thank Ms C. Martínez-Sierra for her excellent technical support.

References

- Alché JD, Butowt R, Castro AJ, Rodríguez-García MI.** 2000. Ubiquitin and ubiquitin-conjugated proteins in the olive (*Olea europaea* L.) pollen. *Sexual Plant Reproduction* **12**, 285–291.
- Alché JD, Jiménez-López JC, Wang W, Castro AJ, Rodríguez-García MI.** 2006. Biochemical characterization and cellular localization of 11S type storage proteins in olive (*Olea europaea* L.) seeds. *Journal of Agricultural Food Chemistry* **54**, 5562–5570.
- Andreou A, Feussner I.** 2009. Lipoxygenases. Structure and reaction mechanism. *Phytochemistry* **70**, 1504–1510.
- Beisson F, Ferte N, Bruley S, Voultoiry R, Verger R, Arondel V.** 2001. Oil bodies as substrates for lipolytic enzymes. *Biochimica et Biophysica Acta* **1531**, 47–58.
- Bewley JD, Black M.** 1994. *Seeds: physiology of development and germination*. London: Plenum Press.
- Bhatla SC, Vandana S, Kaushik V.** 2009. Recent development in the oil body-associated signaling molecules during lipolysis in oil seeds. *Plant Signaling and Behavior* **4**, 176–182.
- Brash AR.** 1999. Lipoxygenases: occurrence, functions, catalysis, and acquisition of substrate. *Journal of Biological Chemistry* **274**, 23679–23682.
- Cañas LA, Benbadis A.** 1988. *In vitro* plant regeneration from cotyledon fragment of the olive tree (*Olea europaea* L.). *Plant Science* **54**, 65–74.
- Eastmond PJ.** 2006. *SUGAR-DEPENDENT* encodes a patatin domain triacylglycerol lipase that initiates storage oil breakdown in germinating *Arabidopsis* seeds. *The Plant Cell* **18**, 665–675.
- Fernandez DE, Staehelin LA.** 1987. Does gibberellic acid induce the transfer of lipase from protein bodies to lipid bodies in barley aleurone cells. *Plant Physiology* **85**, 487–496.
- Feussner I, Kuhn H, Wasternack C.** 2001. Lipoxygenase-dependent degradation of storage lipids. *Trends in Plant Science* **6**, 268–273.
- Feussner I, Wasternack C.** 1998. Lipoxygenase catalyzed oxygenation of lipids. *Fett/Lipid* **100**, 146–152.
- Gerhardt B, Fischer K, Balkenhohl TJ, Pohnert G, Kühn H, Wasternack C, Feussner I.** 2005. Lipoxygenase-mediated metabolism of storage lipids in germinating sunflower cotyledons and β -oxidation of (9Z,11E,13S)-13-hydroxy-octadeca-9,11-dienoic acid by the cotyledonary glyoxysomes. *Planta* **220**, 919–930.
- Graham IA.** 2008. Seed storage oil mobilization. *Annual Review of Plant Biology* **59**, 115–142.
- Hara-Nishimura I, Hayashi M, Nishimura M, Akazawa T.** 1987. Biogenesis of protein bodies by budding from vacuoles in developing pumpkin cotyledons. *Protoplasma* **136**, 49–55.
- Hayashi Y, Hayashi M, Hayashi H, Hara-Nishimura I, Nishimura M.** 2001. Direct interaction between glyoxysomes and lipid bodies in cotyledons of the *Arabidopsis thaliana ped1* mutant. *Protoplasma* **218**, 83–94.
- Heinisch O, Kowalski E, Ludwig H, Tauscher B.** 1996. Staining for soybean lipoxygenase activity in electrophoretic gels. *European Journal of Lipid Science and Technology* **98**, 183–184.
- Herman EM, Larkins BA.** 1999. Protein storage bodies and vacuoles. *The Plant Cell* **11**, 601–613.
- Holtman WL, van Duijn C, Sedee NJA, Douma AC.** 1996. Differential expression of lipoxygenase isoenzymes in embryos of germinating barley. *Plant Physiology* **111**, 569–576.
- Kelly AA, Quettier AL, Shaw E, Eastmond PJ.** 2011. Seed storage oil mobilization is important but not essential for germination or seedling establishment in *Arabidopsis*. *Plant Physiology* **157**, 866–875.
- Kolijak R, Boutaud O, Shieh BH, Samel N, Brash AR.** 1997. Identification of a naturally occurring peroxidase-lipoxygenase fusion protein. *Science* **26**, 1994–1996.

- Laemmli UK.** 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- Lin LJ, Tzen JTC.** 2004. Two distinct steroleosins are present in seed oil bodies. *Plant Physiology and Biochemistry* **42**, 601–608.
- Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* **25**, 402–408.
- Maccarrone M, Salucci ML, van Zadelhoff G, Malatesta F, Veldink G, Vliegthart JFG, Finazzi-Agrò A.** 2001. Tryptic digestion of soybean lipoxygenase-1 generates a 60kDa fragment with improved activity and membrane binding ability. *Biochemistry* **40**, 6819–6827.
- Matsui K, Hijiya K, Tabuchi Y, Kajiwara T.** 1999. Cucumber cotyledon lipoxygenase during postgerminative growth. Its expression and action on lipid bodies. *Plant Physiology* **119**, 1279–1287.
- Murphy DJ.** 2012. The dynamic roles of intracellular lipid droplets: from archaea to mammals. *Protoplasma* **249**, 541–585.
- Müntz K.** 1996. Proteases and proteolytic cleavage of storage proteins in developing and germinating dicotyledonous seeds. *Journal of Experimental Botany* **47**, 605–622.
- Napier JA, Stobart AK, Shewry PR.** 1996. The structure and biogenesis of plant oil bodies: the role of the ER membrane and the oleosin class of proteins. *Plant Molecular Biology* **31**, 945–956.
- Padilla M, Hernández ML, Sanz C, Martínez-Rivas JM.** 2012. Molecular cloning, functional characterization and transcriptional regulation of a 9-lipoxygenase gene from olive. *Phytochemistry* **74**, 58–68.
- Poxleitner M, Rogers SW, Samuels AL, Browse J, Rogers JC.** 2006. A role of caleosin in degradation of oil-body storage lipids during seed germination. *The Plant Journal* **47**, 917–933.
- Quettier A-L, Eastmond PJ.** 2009. Storage oil hydrolysis during early seedling growth. *Plant Physiology and Biochemistry* **47**, 485–490.
- Rejón JD, Zienkiewicz A, Rodríguez-García MI, Castro AJ.** 2012. Profiling and functional classification of esterases in olive (*Olea europaea*) pollen during germination. *Annals of Botany* **110**, 1035–1045.
- Rudolph M, Schlereth A, Körner M, Feussner K, Berndt E, Melzer M, Hornung E, Feussner I.** 2011. The lipoxygenase-dependent oxygenation of lipid body membranes is promoted by a patatin-type phospholipase in cucumber cotyledons. *Journal of Experimental Botany* **62**, 749–760.
- Serrano I, Suárez C, Olmedilla A, Rapoport HF, Rodríguez-García MI.** 2008. Structural organization and cytochemical features of the pistil in olive (*Olea europaea* L.) cv. Picual at anthesis. *Sexual Plant Reproduction* **21**, 99–111.
- Strzałka K, Hara-Nishimura I, Nishimura M.** 1995. Changes in physical properties of vacuolar membrane during transformation of protein bodies into vacuoles in germinating pumpkin seeds. *Biochimica et Biophysica Acta* **1239**, 103–110.
- Tai SS, Chen MC, Peng CC, Tzen JT.** 2002. Gene family of oleosin isoforms and their structural stabilization in sesame seed oil bodies. *Bioscience, Biotechnology, and Biochemistry* **66**, 2146–2153.
- Torrent M, Geli MI, Ludevid MD.** 1989. Storage-protein hydrolysis and protein-body breakdown in germinated *Zea mays* L. seeds. *Planta* **180**, 90–95.
- Yi H, Yi M, Choe HT.** 2005. Changes in lipoxygenase properties and activity related to postgerminative growth and senescence in oat (*Avena sativa* L. cv. Victory 1). *Journal of Plant Biology* **48**, 429–439.
- Zienkiewicz K, Castro AJ, Alché JD, Zienkiewicz A, Suárez C, Rodríguez-García MI.** 2010. Identification and localization of a caleosin in olive (*Olea europaea* L.) pollen during *in vitro* germination. *Journal of Experimental Botany* **61**, 1537–1546.
- Zienkiewicz A, Jiménez-López JC, Zienkiewicz K, Alché JD, Rodríguez-García MI.** 2011. Development of the cotyledon cells during olive (*Olea europaea* L.) *in vitro* seed germination and seedling growth. *Protoplasma* **248**, 751–765.
- Zienkiewicz A, Zienkiewicz K, Rejón JD, Rodríguez-García MI, Castro AJ.** 2013. New insights into the early steps of oil body mobilization during pollen germination. *Journal of Experimental Botany* **64**, 293–302.