

Predictors of Improvement in Cardiometabolic Risk Factors With Weight Loss in Women

Caitlin A. Dow, PhD; Cynthia A. Thomson, PhD, RD; Shirley W. Flatt, MS; Nancy E. Sherwood, PhD; Bilge Pakiz, EdD; Cheryl L. Rock, PhD, RD

Background—Weight loss is associated with improvements in cardiometabolic risk factors, including serum glucose, insulin, C-reactive protein, and blood lipids. Few studies have evaluated the long-term (>18 months) effect of weight loss on these risk factors or sought to identify factors associated with sustained improvements in these measures.

Methods and Results—In 417 overweight/obese women (mean [SD] age, 44 [10] years) participating in a weight loss trial, we sought to identify predictors of weight loss–associated cardiometabolic risk factors after 12 and 24 months of intervention. Total cholesterol (TC), low-density lipoprotein (LDL)–cholesterol (LDL-C), high-density lipoprotein (HDL)–cholesterol, non-HDL-cholesterol, triglycerides (TG), insulin, glucose, C-reactive protein (CRP), and cardiopulmonary fitness were measured at baseline and at 12 and 24 months. After 24 months, significant reductions in body weight, waist circumference, CRP, TC, HDL-cholesterol, and non-HDL-cholesterol were observed ($P<0.01$). After 24 months, mean TC and non-HDL-cholesterol were reduced regardless of the amount of weight lost, whereas reductions in LDL-cholesterol, CRP, insulin, and TG were observed only in those who lost $\geq 10\%$ body weight. Step-test performance improved only in those who lost $\geq 10\%$ body weight after 24 months. Change in weight demonstrated a positive predictive value for change in cholesterol, insulin, glucose, and triglycerides. Baseline level of the biomarker showed the greatest predictive value for follow-up measures for insulin, cholesterol, glucose, and triglycerides.

Conclusions—Our data extend the results from short-term weight loss trials and suggest that the magnitude of weight loss and baseline values for risk factors are associated with improvements in cardiometabolic risk factors even after 24 months.

Clinical Trials Registration Information—URL: <http://www.clinicaltrials.gov/>. Unique identifier: NCT00640900. (*J Am Heart Assoc.* 2013;2:e000152 doi: 10.1161/JAHA.113.000152)

Key Words: lipids • obesity • risk factors • women

Obesity remains a significant risk factor for disease-associated morbidity and mortality in the United States, including that associated with diabetes, coronary heart disease, and stroke.¹ Obese individuals often present with elevated fasting values of glucose,² lipids,¹ and inflammatory markers,² as well as insulin resistance.³ Even with modest

weight loss, individuals may demonstrate significant improvements in metabolic risk factors for obesity-related disease, particularly cardiovascular disease and diabetes.^{1,4–6} It is estimated that a 10% reduction in body weight in an obese individual can promote significant reductions in metabolic parameters including circulating glucose, insulin, and inflammatory markers, even if the body mass index (BMI) remains within an at-risk category.⁷ Yet few studies have evaluated this relationship beyond 12 months.

The ability to promote weight loss through dietary interventions is well established, and behavioral weight loss programs and counseling remain a primary clinical approach to reduce body weight in overweight and obese adults.⁶ These programs vary in intensity, delivery approach, behavioral constructs applied, dietary plan, and duration. But independent of the approach, energy deficit can be achieved and is associated with modest weight loss.^{3,5,8} Further, the demonstrated weight loss associated with targeted dietary interventions generally translates to improvements in metabolic indices related to obesity-associated disease.^{4,6,9–11}

From the Department of Nutritional Sciences (C.A.D., C.A.T.) and Mel and Enid Zuckerman College of Public Health (C.A.T.), University of Arizona, Tucson, AZ; Department of Family and Preventive Medicine, University of California, San Diego, La Jolla, CA (S.W.F., B.P., C.L.R.); HealthPartners Institute for Education and Research, Minneapolis, MN (N.E.S.).

Correspondence to: Cynthia A. Thomson, PhD, RD, Mel and Enid Zuckerman College of Public Health, University of Arizona, 1295 N. Martin Street, Drachman Building, Room A260, Tucson, AZ 85724. E-mail: cthompson@email.arizona.edu

Received February 11, 2013; accepted October 25, 2013.

© 2013 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

In a recently conducted randomized, controlled weight loss study,¹² women were randomized to receive a structured weight loss program delivered either in person or via telephone or to receive minimal diet counseling, considered usual care, for a period of 24 months. Women assigned to the structured weight loss program arms demonstrated greater reductions in body weight, BMI, and waist circumference over time than those in the usual care condition. More than 70% of women had sustained weight loss at 24 months; the variance in weight loss afforded an opportunity to evaluate the relationship between weight loss and modification of cardiometabolic risk factors. Changes in lipids and C-reactive protein (CRP) according to study arm assignment have been previously reported.¹² Overall, blood lipids did not change in those randomized to the structured weight loss plan, although CRP values were significantly reduced after 12 and 24 months of intervention.¹² The purpose of this analysis was to assess changes in these and additional cardiometabolic risk factors associated with weight loss. Further, this analysis sought to determine which factors (baseline value of biomarker, baseline weight, percent change in body weight, age, race/ethnicity) predicted improvement in cardiometabolic risk factors at 12 and 24 months, independent of treatment arm.

Methods

Study Subjects and Recruitment

The weight loss intervention trial recruited 446 overweight/obese women from a screened sample of 564 women. Women were eligible to participate if they were >18 years of age, had a BMI between 25 and 40 kg/m², and were a minimum of 15 kg above ideal body weight as defined by the 1983 Metropolitan Life tables.¹³ Participants were recruited via listservs and flyers distributed through each of the 4 study sites (University of California, San Diego; University of Arizona, Tucson; University of Minnesota, Minneapolis; and Kaiser Permanente Center Northwest). Any woman who was pregnant, planning a pregnancy, lactating, reported a history of eating disorders, other psychiatric disorders, comorbid conditions, or food allergies, or was unable to perform a 3-minute step test for cardiopulmonary fitness assessment was excluded from enrollment. All study participants completed the consent process and provided written informed consent prior to randomization. The institutional review boards approved this study for Human Subjects' Protection at each recruitment site prior to study initiation.

Intervention

The details of the study participation have been previously described.¹² Briefly, all women enrolled in the trial received

weight loss counseling. Eligible women were randomized in a 3:3:2 allocation to center-based, telephone-based, or usual care that included weight loss counseling by a dietetics professional. Women in the center-based intervention arm visited the weight loss center weekly for brief counseling sessions and weigh-ins with designated trained staff; they also had access to Web-based resource materials. Women in the telephone-based arm similarly received weekly counseling by telephone to reinforce weight loss behaviors and goals. They also had access to Web-based resource materials. Prepackaged meals and snacks provided 42% to 68% of total energy needs of the prescribed eating plan and were provided free of charge for participants randomized to these 2 study arms. Generally, after the initial 12-month period, women were transitioned to fewer prepackaged meals and more self-selected food items. In addition to the prepackaged foods, women assigned to the weight loss program received one-on-one counseling to promote adherence to a low-fat (20% to 30% total energy), reduced-energy (1200 to 2000 kcal), high fruit-and-vegetable eating plan with recommendations for increased physical activity to achieve current guidelines¹⁴ of 30 minutes of planned exercise ≥ 5 days/week. Usual-care counseling sessions took place once at baseline and again at 6 months. Usual-care study participants received written educational materials to support their dietary goals (500 to 1000 kcal/day deficit); regular physical activity was also promoted. For these analyses treatment arms were collapsed.

Outcomes

The primary outcome of this study was change in cardiometabolic risk factors. Subjects provided a fasting blood sample at baseline and after 12 and 24 months for evaluation of cardiometabolic risk factors. Specifically, fasting plasma was sampled to measure lipid profiles including total cholesterol, triglycerides, and high-density lipoprotein (HDL) using enzymatic methods (Kodak Ektachem Analyzer System; Johnson & Johnson Clinical Diagnostics). Low-density lipoprotein (LDL) was calculated using the Friedewald equation.¹⁵ Non-HDL-cholesterol (non-HDL-C) was calculated by subtracting HDL from the total cholesterol value. Blood glucose was determined using enzymatic methods (Kodak Ektachem DT60 Analyzer; Johnson & Johnson). A double-antibody radioimmunoassay, with <0.2% cross-reactivity with human proinsulin, was used for the analysis of fasting serum insulin (Linco Research Inc). The homeostasis model assessment of insulin resistance (HOMA-IR) was calculated by dividing the product of fasting glucose (in milligrams per deciliter) and fasting insulin (in microunits per milliliter) by 405.¹⁶ CRP was quantified using a high-sensitivity polystyrene-enhanced turbidimetric in vitro immunoassay kit (DiaSorin Inc).

Secondary outcomes included weight, height, BMI (weight in kilograms/height in meters squared) as well as waist and hip circumference, which were measured at baseline and every 6 months throughout the 24-month study period. In addition, a 3-minute step test was administered at baseline and repeated at 12 and 24 months. This test measures heart rate during the first 30 seconds of recovery from stepping.

Statistical Analysis

Risk categories for cardiometabolic outcomes, used as cut points in tabulation, were identified as follows: glucose (cut point, 100 mg/dL),¹⁷ insulin (cut point, 15 μ U/mL),¹⁸ HOMA-IR (cut point, 3.0),¹⁶ CRP (low risk, <1 mg/L; medium risk, 1 to 2.99 mg/L; high risk, \geq 3.0 mg/L),¹⁹ total cholesterol (cut point, 200 mg/dL),¹⁷ LDL-cholesterol (cut point, 130 mg/dL),²⁰ non-HDL-C (cut point, 160 mg/dL),²¹ HDL-cholesterol (cut point, 60 mg/dL),¹⁷ triglycerides (cut point, 150 mg/dL),¹⁷ waist (cut point, 88 cm),²² and waist-to-height ratio (cut point, 0.5).²³ Cardiopulmonary fitness recovery heart rate categories for ages <30, 30 to 39, 40 to 59, and \geq 50 years were based on literature.²⁴ Logarithmic transformations were applied in all analyses to improve normality of skewed distributions with the exception of the β coefficients reported in Table 2.

Longitudinal analyses were limited to the 417 women who provided at least 1 follow-up blood sample at either 12 or 24 months. Differences in biochemical risk factors between baseline and each of 12 and 24 months (overall and stratified by a priori biomarker risk category) were analyzed using longitudinal mixed models, controlled for intervention group. Likelihood ratio tests explored interactions between risk strata and time. Each model tested for differences over time and included a subject-specific intercept as a random effect. Regression models predicted laboratory values after 12 and 24 months of study participation as a function of race/ethnicity, age, baseline levels of laboratory analyte, waist circumference, weight change, randomization assignment, cardiorespiratory fitness, and waist-to-height ratio. For each term in each model, we present parameter estimates (β coefficients). To further explore the relationship between change in body weight and cardiometabolic risk factors, stratified analyses, applying a priori clinical cut points for weight loss, were conducted using longitudinal mixed models in which women were classified as either losing \geq 10% of baseline body weight or losing <10% of baseline body weight at each of the 12- and 24-month points. Models were controlled for intervention group assignment and tested for interaction between the weight loss categories and biomarker change. In consideration of multiple comparisons issues, statistical significance was set at $P<0.01$. Analyses were conducted using SAS version 9.3 (SAS Institute, Inc, Cary, NC).

Results

Four hundred and forty-six overweight/obese women were enrolled; 4 were removed from analysis because of postrandomization exclusions and 25 did not provide a blood sample at either 12 or 24 months, leaving 417 in the study sample. Table 1 provides baseline demographic characteristics of the study population. Women enrolled in the study were a mean (SD) age of 44 (10) years, had a mean weight of 92.1 (10.8) kg with a BMI of 33.9 (3.4) kg/m², and had a waist circumference of 108.6 (9.5) cm. At baseline, 159 women (36%) were postmenopausal. At follow-up, mean weight loss as a percentage of baseline weight was 8.6% (0.4%) in 417 subjects with available measured body weight at 12 months and 6.6% (0.5%) in 407 subjects with weight data at 24 months; 85% and 74% of women had any weight loss at 12 and 24 months, respectively. At study entry, 2 women were using hypoglycemic agents, and 1% of the women (n=5) were using them at study end. Twenty-six subjects took lipid-lowering drugs at study entry and 30 women took them at 24 months; 17 women reported estrogen use throughout the study (data not shown).

As shown in Table 2, several predictors of change in cardiometabolic biomarkers were identified although neither age nor race/ethnicity showed an association. Most relevant was the strong and significant association between percent body weight reduction and change in biomarkers, such that for every 10% point reduction in body weight, there was an estimated 4-point reduction in insulin and glucose, a 5.4 mg/dL reduction in cholesterol, and an 18 mg/dL decrease in serum triglycerides. Baseline level of each biomarker was

Table 1. Baseline Characteristics of Overweight/Obese Women Enrolled in a 24-Month Weight Loss Study (n=442)

Characteristic	Baseline Value, Mean (SD)
Age, y	44 (10)
Weight, kg	92.1 (10.8)
Height, cm	164.9 (6.3)
BMI, kg/m ²	33.9 (3.4)
Waist circumference, cm	108.6 (9.5)
Hip circumference, cm	120.2 (8.2)
Waist-to-hip ratio	0.9 (0.06)
Postmenopausal, n (%)	159 (36)
Education, n	
High school or less	52
Some college	189
College graduate	95
Graduate school	106

BMI indicates body mass index; SD, standard deviation.

Table 2. Predictors of Cardiometabolic Biomarker Values After 12 and 24 Months of Weight Loss Study Participation

	Models Predicting 12-Month Biomarker Values (n=406)				
	Insulin ($R^2=0.37$)	Cholesterol ($R^2=0.46$)	Glucose ($R^2=0.26$)	TG ($R^2=0.66$)	CRP ($R^2=0.16$)
Baseline level of biomarker	0.551**	0.597**	0.786**	0.752**	0.470**
Baseline weight	-0.057	-0.093	0.069	0.286	0.041
Percent change in weight	0.274**	0.403*	0.351**	1.514**	0.091*
Baseline waist	0.294	0.409	0.211	-0.585	0.013
Age	-0.025	0.520	0.016	0.301	-0.021
White race/ethnicity	-2.022	5.980	-0.964	1.428	-0.420
Dietary intervention	0.0033	-1.380	-0.597	-5.874*	-0.167
Step-test heart rate	-0.036	0.071	0.058	-0.053	-0.008
Waist-to-height ratio	-24.329	-72.124	-24.980	95.266	4.740
	Models Predicting 24-Month Biomarker Values (n=383)				
	Insulin ($R^2=0.52$)	Cholesterol ($R^2=0.39$)	Glucose ($R^2=0.31$)	TG ($R^2=0.56$)	CRP ($R^2=0.07$)
Baseline level of biomarker	0.770**	0.505**	0.525**	1.150**	0.430*
Baseline weight	0.073	-0.300	0.104	0.673	-0.060
Percent change in weight	0.399**	0.544*	0.406**	1.788**	0.064
Baseline waist	-0.173	0.847	-0.306	-2.147	-0.029
Age	0.067	0.637**	0.064	0.817	-0.098
White race/ethnicity	-0.744	5.592	-1.137	-3.287	-2.545
Dietary intervention	0.264	-2.682	0.423	-6.210	0.917
Step-test heart rate	0.001	0.277	0.069	-0.324	0.872
Waist-to-height ratio	31.76	-127.74	31.81	211.26	0.829

CRP indicates C-reactive protein; TG, triglycerides.

* $P<0.01$, ** $P<0.001$. Values shown are β coefficients in a regression model for each biomarker during each period.

strongly and inversely associated with follow-up measures in the individual marker. Percent change in weight was also associated with modification of biomarkers at 12 and 24 months, particularly for reductions in triglyceride values. Baseline weight or waist circumference was not associated with change in any of the markers.

To further evaluate the relationship between weight change and changes in cardiometabolic biomarkers, models were stratified at a 10% cut point in weight loss percentage and evaluated at the 12 and 24 month follow-up points (Table 3). At 12 months, 167 women had lost at least 10% of initial body weight (48.4% of those randomized to the intervention arms and 13.9% of those in the usual care arm; data not shown). In these women, CRP was reduced from 4.0 to 2.6 mg/L. Waist-to-height ratio (WtHR), total cholesterol, and low-density lipoprotein-cholesterol (LDL-C) were reduced regardless of weight loss percentage (Table 3). Most lipids (except triglycerides) did not show significant interactions between degree of weight loss and degree of lipid change over time (data not shown). All other cardiometabolic risk factors shown in Table 3 had significant interactions between degree of weight loss and time ($P<0.01$ for each).

At 24 months, 130 women had lost $\geq 10\%$ of their initial body weight (28.8% of those in the intervention groups and 17.5% of those randomized to usual care; data not shown). Reductions in triglycerides, glucose, insulin, and HOMA-IR were only evident in women who lost $\geq 10\%$ body weight at follow-up. Cardiopulmonary fitness (based on the 3-minute step test) also improved in study participants. Interestingly, this effect was independent of the degree of weight loss only at 12 months (Table 3). After 24 months, cardiopulmonary fitness significantly improved only in women whose weight loss was $\geq 10\%$, whereas it was significantly reduced in those who did not lose 10% of their initial weight compared with baseline values (Table 3).

To evaluate the clinical relevance of the change in cardiometabolic risk factors, an analysis was completed to assess overall change in metabolic markers over time as well as change within clinical cut-point substrata. Of note, all the biomarkers showed significant interactions between risk level and time ($P<0.001$ for each), and there was a significant interaction between weight loss and time. Generally, the higher the baseline biomarker values, the greater was the reduction with weight loss, as was shown for CRP, chole-

Table 3. Evaluation of Cardiometabolic Risk Factors, Stratified at 10% Weight Loss After 12 and 24 Months Separately

	Lost 10% of Weight After 12 Months (n=167)		Did Not Lose 10% of Weight After 12 Months (n=240)		Lost 10% of Weight After 24 Months (n=130)		Did Not Lose 10% of Weight After 24 Months (n=253)	
	Baseline	12 Months	Baseline	12 Months	Baseline	24 Months	Baseline	24 Months
Weight (kg), mean (SD)	93.2 (11.3)	77.3 (10.7)**	91.0 (10.3)	88.3 (11.2)**	93.3 (10.3)	77.3 (10.1)**	91.0 (10.9)	89.7 (11.9)*
Waist (cm), mean (SD)	109.1 (9.5)	93.8 (8.5)**	107.8 (9.4)	102.7 (9.5)**	109.0 (9.2)	94.8 (8.9)**	107.8 (9.5)	104.2 (9.8)**
Waist/height ratio	0.66 (0.06)	0.58 (0.05)**	0.66 (0.06)	0.63 (0.06)**	0.66 (0.06)	0.57 (0.06)**	0.66 (0.06)	0.63 (0.06)**
BMI, mean (SD)	33.9 (3.4)	28.2 (3.4)**	33.7 (3.4)	32.7 (3.8)**	34.0 (3.3)	28.2 (3.4)**	33.6 (3.4)	33.2 (4.0)*
CRP (mg/L), mean (SD)	4.0 (3.5)	2.6 (5.6)**	4.3 (3.9)	4.0 (5.7)**	3.9 (3.1)	2.2 (2.8)**	4.2 (3.9)	5.0 (11.3)
TG (mg/dL), mean (SD)	107 (48)	94 (39)**	113 (64)	114 (63)	104 (45)	92 (33)**	114 (56)	124 (61)**
Glucose (mg/dL), mean (SD)	94 (11)	93 (24)	94 (11)	96 (13)*	93 (10)	89 (11)**	94 (11)	96 (14)
Insulin (μ U/mL), mean (SD)	16.9 (9.1)	12.9 (10.0)**	18.2 (8.7)	17.5 (9.2)	16.3 (8.1)	13.3 (6.2)**	18.1 (8.6)	21.1 (12.2)**
HOMA-IR, % insulin resistant [†]	62%	35%**	68%	65%	61%	39%**	67%	78%**
Step-test heart rate/30 s, mean (SD)	54 (10)	46 (9)**	54 (9)	50 (9)**	54 (9)	46 (8)**	54 (10)	50 (9)**
Step test poor to fair, %	68.5	30.7**	70.2	58.4**	68.5	36.8**	46.1	87.1**
Step test good to outstanding, %	31.5	69.3**	29.8	41.6**	31.5	63.8**	53.9	12.9**

BMI indicates body mass index; CRP, C-reactive protein; HOMA-IR, homeostasis model assessment of insulin resistance; SD, standard deviation; TG, triglycerides.

* $P < 0.01$ for change between baseline and follow-up in log-transformed analysis.

** $P < 0.001$ for change between baseline and follow-up in log-transformed analysis.

[†]Homeostasis model for insulin resistance. Insulin sensitive is < 3 ; insulin resistant is ≥ 3 .

terol, LDL-C, triglycerides, glucose, and insulin (Table 4). Overall, a significant number of women entered the weight loss intervention with elevated cardiometabolic risk biomarkers including 200 with CRP > 3.0 mg/L, 178 with total cholesterol > 200 mg/dL, 139 with elevated LDL, and 252 with low HDL. In a sensitivity analysis that excluded women (n=10) who began a lipid-lowering agent during the trial, results of lipid changes were essentially unchanged (data not shown). At baseline, all women had a WHtR above the recommended cutoff point of 0.5; mean waist circumference was significantly reduced after 12 and 24 months. Elevated fasting insulin (n=227) and glucose (n=96) were also common. Regression to the mean was observed in some of the strata. In sensitivity analyses limited to postmenopausal women, outcomes were similar to those in the entire cohort.

Discussion

In this study of overweight women, participants in a longer term weight loss program lost an average of 7% of baseline body weight at the 2-year follow-up. Weight loss resulted in improvement in numerous cardiometabolic risk factors, including biochemical values and a functional test of cardiopulmonary fitness. Importantly, the improvements in most risk

factors were evident at both 12 and 24 months despite modest recidivism in weight loss over time. These results are supported by several shorter-term studies, including an 8-week study by Te Morenga et al in 83 adults, in which weight loss ranged from 3.3 to 4.5 kg and resulted in significant reductions in LDL-C, triglycerides, and glucose.⁸ Similarly, a weight loss intervention by Muzio et al resulted in significant reductions in weight, waist circumference, triglycerides, and LDL-C after 5 months.¹¹ Short-term studies have also demonstrated a reduction in CRP,²⁵ but seldom to the degree necessary to reduce clinical risk as observed in this trial. In a 12-week weight loss study by Clifton et al in 79 overweight/obese adult women, significant weight loss also was demonstrated, but importantly, this is 1 of only a few trials that reassessed the intervention efficacy longer term.¹⁰ As with the present study, long-term improvements in glucose, insulin, CRP, and LDL-C were demonstrated.¹⁰

There have been only a limited number of longer-term studies, but a few have evaluated changes in cardiometabolic parameters over time in relation to weight loss. Four-year data from the LookAHEAD trial, a study performed in individuals with type 2 diabetes, support our results and show that a lifestyle intervention aimed at weight reduction results in significant reductions in triglycerides and LDL-C.²⁶ In 1 of the larger samples studied, Foster et al showed an 11-kg average

Table 4. Cardiometabolic Risk Factors at Baseline and After 12 and 24 Months of Weight Loss Study Participation, Overall and Stratified by Clinical Risk/Cut Points

Risk Factor	Baseline (n=417)	n*	12 Months (n=407)	n*	24 Months (n=383)
CRP (mg/L), mean (SD)	4.2 (3.7)	407	3.4 (5.7)**	383	4.1 (9.4)**
CRP <1 (low risk)	0.6 (0.2)	38	0.8 (0.6)**	35	1.0 (0.7)**
CRP 1 to 2.99 (medium risk)	1.9 (0.6)	169	1.9 (3.9)	164	3.1 (12.8)
CRP ≥3 (high risk)	6.6 (3.9)	200	5.1 (6.8)**	184	5.4 (6.2)**
Total cholesterol (mg/dL), mean (SD)	196 (36)	406	189 (36)	383	184 (37)**
<200	171 (21)	228	172 (28)	215	168 (27)
≥200	229 (21)	178	211 (34)**	168	204 (37)**
LDL-cholesterol (mg/dL), mean (SD)	117 (33)	405	111 (34)*	382	113 (37)
<130	98 (20)	266	99 (27)	255	104 (33)
≥130	154 (17)	139	133 (35)**	127	132 (38)**
HDL-cholesterol (mg/dL), mean (SD)	57 (16)	405	57 (16)	383	53 (15)*
<60	47 (8)	252	52 (13)**	239	47 (10)
≥60	73 (12)	153	66 (15)**	144	63 (17)**
Non-HDL-cholesterol (mg/dL), mean (SD)	139 (36)	405	132 (35)*	383	131 (38)*
<160	122 (24)	294	121 (28)	280	120 (30)
≥160	185 (19)	111	162 (38)*	103	159 (42)*
Triglycerides (mg/dL), mean (SD)	90 (26)	406	106 (55)	383	117 (93)
<150	90 (26)	337	93 (34)	317	100 (43)**
≥150	204 (69)	69	168 (89)**	66	195 (186)*
Glucose (mg/dL), mean (SD)	94 (11)	406	95 (18)	383	94 (13)
<100	89 (7)	310	91 (9)	293	91 (10)
≥100	109 (9)	96	107 (32)	90	101 (18)**
Insulin (μU/mL), mean (SD)	17.8 (8.9)	407	15.6 (9.8)**	383	18.5 (11.2)
<15	11.1 (2.7)	180	11.0 (4.5)	174	13.4 (5.5)**
≥15	23.1 (8.5)	227	19.3 (11.2)**	209	22.7 (12.8)
HOMA-IR [‡]	4.21 (2.53)	405	3.72 (2.58)**	383	4.37 (3.10)
<3	2.26 (0.54)	138	2.39 (1.11)	135	2.95 (1.38)**
≥3	5.24 (2.57)	267	4.41 (2.84)**	248	5.15 (3.49)
Waist (cm), mean (SD)	108.3 (9.5)	409	99.0 (10.1)**	392	101.1 (10.7)**
<88	86.7 (1.2)	4	86.0 (11.3)	4	82.9 (6.5)
≥88	108.5 (9.3)	405	99.2 (10.0)**	388	101.3 (10.5)**
Waist/height ratio, mean (SD)					
<0.5	N/A				
≥0.5	0.66 (0.06)	409	0.60 (0.06)**	392	0.61 (0.07)**

CRP, C-reactive protein; HDL, high-density lipoprotein; HOMA-IR, homeostasis model assessment of insulin resistance; LDL, low-density lipoprotein; SD, standard deviation;

* $P < 0.01$, compared with baseline in log-transformed analysis.

** $P < 0.001$, compared with baseline in log-transformed analysis.

[†]Sample size represents the number within stratum at baseline and after 12 or 24 months and varies based on completion of 12- or 24-month clinic activities.

[‡]Homeostasis model for insulin resistance. Insulin sensitive is < 3 ; insulin resistant is ≥ 3 .

weight loss at year 1 ($n=307$; 68% female) and 7-kg loss at year 2 ($n=113$),²⁷ weight loss effects similar to what was achieved in our sample. Consistent with our findings, triglycerides improved at year 1 and were not significantly

different from baseline at year 2 in either study. Their results suggested no significant reduction in LDL-C at 24 months,²⁷ though participants in our trial who had elevated baseline values demonstrated significant reductions in LDL-C even

after 24 months. In a study of 176 overweight/obese adults of whom 87% were female, Burke et al showed significant improvements in total cholesterol and the LDL:HDL ratio as well as HOMA-IR, a measure of insulin resistance, after 12 months of intervention, changes that were generally sustained at the 6-month postintervention follow-up.²⁸ These results, taken into consideration with our results, suggest that efforts to favorably modulate cardiometabolic risk factors through modest weight loss in numerous short-term trials generally translate to longer-term follow-up, even with modest weight regain between 12 and 24 months.

Improvements in several CVD risk factors were demonstrated in response to participation in this weight loss study. WHtR is an adiposity index that more accurately predicts cardiometabolic risk than BMI or waist circumference.^{23,29} In this study, WHtR was reduced regardless of the degree of weight lost, although this effect was more pronounced in women who lost $\geq 10\%$ of their initial body weight, as evidenced by the significant interaction between WHtR and weight. Recent evidence also indicates that non-HDL-cholesterol more accurately predicts risk of coronary heart disease than LDL-C, as it encompasses all cholesterol-containing atherogenic particles including LDL-C and very-low-density lipoprotein-cholesterol.³⁰ On average, women with elevated baseline non-HDL-cholesterol (≥ 160 mg/dL) demonstrated reductions to within a normal range after 24 months, and non-HDL cholesterol values were reduced at 12 and 24 months, regardless of the degree of weight loss. The literature regarding lifestyle interventions for weight loss and non-HDL-cholesterol response is extremely limited, although our findings suggest that any amount of weight loss may reduce non-HDL levels.

Previous work has demonstrated that reductions in insulin resistance are observed with increases in physical activity, independent of weight loss.³¹ Other intervention studies for weight reduction agreed that exercise is a necessary component, in addition to diet modification, to reduce insulin resistance.³² In this study, diet was the major lifestyle modification addressed, but regular physical activity was also promoted through counseling. In response, insulin levels and, perhaps more importantly, HOMA-IR values were significantly reduced at 12 months, but only in women who demonstrated weight loss $\geq 10\%$ of their initial body weight. We cannot ascertain if this effect was due solely to changes in diet and/or physical activity.

This study also had the benefit of evaluating cardiopulmonary fitness, an indicator of health also known to improve with physical activity.³³ Performance improved over time regardless of percent weight loss but was not associated with cardiometabolic factors. These results suggest that efforts to expand outcome measures in weight loss studies targeting improved cardiometabolic risk should consider biochemical

biomarkers as well as functional measures. Improvements in cardiopulmonary fitness have been reported in other weight loss studies,^{32,34} although not consistently.³⁵

Although control for baseline values of the respective outcome (ie, weight, biomarker) when conducting regression analysis is sometimes applied, estimation of the predictive nature of the baseline value is seldom evaluated.³⁶ Our results suggest that 1 important predictor of change in cardiometabolic biomarkers is the baseline level of the individual biomarker. These data suggest that lifestyle interventions targeting weight loss are effective for improving cardiometabolic risk in individuals who present with unfavorable measures. Our data suggest that these measures show significant and sustained improvement if any degree of weight loss is achieved and maintained.

Limitations to this study include a lack of assessment of lipoprotein particle size, known to be associated with cardiovascular risk³⁷ and responsive to dietary intervention.⁹ Further, we did not conduct hemodynamic measures such as blood pressure during the 3-minute step test. These measures likely would have provided more robust data regarding cardiopulmonary fitness.³³ However, cardiopulmonary fitness was not a primary end point of this study, and although less accurate than measuring maximal oxygen uptake (VO_{2max}) and hemodynamics during test administration, the 3-minute step test has high reliability, is sensitive to change, and has been used broadly to assess change in cardiopulmonary fitness in clinical trials.²⁴ We did not collect specific information on diet or physical activity that can influence these biomarkers; however, it is apparent that women were generally in negative energy balance given the substantial weight loss demonstrated by so many study participants. It is well recognized that negative energy balance can modulate many of these targeted risk factors whether that occurs through dietary intervention or in combination with physical activity.^{11,25,26,30,35,36,38,39} Last, the results of this study may not represent effects of weight loss on cardiometabolic risk factors in a clinical population under direct care, including medications, to treat their disease.

Strengths of the trial include the 24-month intervention and follow-up period in a relatively large sample of women with robust and repeated measures of cardiometabolic risk factors. The sample size and variance in weight loss also afforded an opportunity to ascertain differences in risk factors over time in women who were more ($\geq 10\%$ body weight loss) versus less ($< 10\%$ body weight loss) responsive to the weight loss interventions.

In conclusion, participation in a weight loss study resulted in significant reductions in body weight, BMI, and waist circumference after 12 and 24 months. This weight loss was associated with significant improvements in several of the

cardiometabolic risk factors that are commonly assessed in the clinical setting. Moreover, weight loss of $\geq 10\%$ after 12 months produced significant reductions in these risk factors, often lowering the risk factor from a category of high risk to that of medium or low risk (ie, CRP and insulin). These data also show that baseline cardiometabolic risk factors (cholesterol, fasting glucose, insulin, triglycerides, and CRP) have significant predictive value in determining favorable risk reduction response to weight loss, suggesting that these risk factors could be used to identify women more likely to respond favorably to weight loss interventions.

Acknowledgments

The authors thank the data and safety monitoring committee (Ken Fujioka, MD, Karen Messer, PhD, and Kevin Patrick, MD). We also acknowledge Dennis Heath, MS, Mila Pruitt, Paul Mills, PhD, and Jeff Chang, MD (UCSD), for conducting the laboratory assays; Christine Zoumas-Morse, MS, RD (UCSD), for managing the project; Annie Hotop, MA, MS (University of Minnesota), Lucy Fulton, DTR (Kaiser Permanente Center for Health Research), Jennifer Ravia, MS, and Emily Nardi, MPH, RD (University of Arizona), and Elizabeth Quintana, MS, RD (UCSD), for coordinating activities at the clinical sites; and Lea Jacinto, BS (UCSD), for assistance with administrative support and manuscript preparation. Shirley Flatt, MS, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Source of Funding

This study was supported by Jenny Craig, Inc. By contractual agreement, scientists at UCSD and the other participating institutions have responsibility and independence regarding data management, analysis, and publication.

Disclosures

None.

References

- Gregg EW, Cheng YJ, Cadwell BL, Imperatore G, Williams DE, Flegal KM, Narayan KM, Williamson DF. Secular trends in cardiovascular disease risk factors according to body mass index in US adults. *JAMA*. 2005;293:1868–1874.
- Lumeng CN, Saltiel AR. Inflammatory links between obesity and metabolic disease. *J Clin Invest*. 2011;121:2111–2117.
- Taylor R. Pathogenesis of type 2 diabetes: tracing the reverse route from cure to cause. *Diabetologia*. 2008;51:1781–1789.
- Layman DK, Evans EM, Erickson D, Seyler J, Weber J, Bagshaw D, Griel A, Psota T, Kris-Etherton P. A moderate-protein diet produces sustained weight loss and long-term changes in body composition and blood lipids in obese adults. *J Nutr*. 2009;139:514–521.
- Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, Cuzick J, Jebb SA, Martin B, Cutler RG, Son TG, Maudsley S, Carlson OD, Egan JM, Flyvbjerg A, Howell A. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes (Lond)*. 2011;35:714–727.
- Laddu D, Dow C, Hingle M, Thomson C, Going S. A review of evidence-based strategies to treat obesity in adults. *Nutr Clin Pract*. 2011;26:512–525.
- Johnson WD, Brashear MM, Gupta AK, Rood JC, Ryan DH. Incremental weight loss improves cardiometabolic risk in extremely obese adults. *Am J Med*. 2011;124:931–938.
- Morenga LT, Williams S, Brown R, Mann J. Effect of a relatively high-protein, high-fiber diet on body composition and metabolic risk factors in overweight women. *Eur J Clin Nutr*. 2010;64:1323–1331.
- Varady KA, Bhutani S, Klempel MC, Kroeger CM. Comparison of effects of diet versus exercise weight loss regimens on LDL and HDL particle size in obese adults. *Lipids Health Dis*. 2011;10:119.
- Clifton PM, Keogh JB, Noakes M. Long-term effects of a high-protein weight-loss diet. *Am J Clin Nutr*. 2008;87:23–29.
- Muzio F, Mondazzi L, Harris WS, Sommariva D, Branchi A. Effects of moderate variations in the macronutrient content of the diet on cardiovascular disease risk factors in obese patients with the metabolic syndrome. *Am J Clin Nutr*. 2007;86:946–951.
- Rock CL, Flatt SW, Sherwood NE, Karanja N, Pakiz B, Thomson CA. Effect of a free prepared meal and incentivized weight loss program on weight loss and weight loss maintenance in obese and overweight women: a randomized controlled trial. *JAMA*. 2010;304:1803–1810.
- 1983 metropolitan height and weight tables. *Stat Bull Metrop Life Found*. 1983;64:3–9.
- Services UDoHaH. *2008 Physical Activity Guidelines for Americans*. Centers for Disease Control and Prevention: Atlanta, GA; 2008.
- Fukuyama N, Homma K, Wakana N, Kudo K, Suyama A, Ohazama H, Tsuji C, Ishiwata K, Eguchi Y, Nakazawa H, Tanaka E. Validation of the Friedewald equation for evaluation of plasma LDL-cholesterol. *J Clin Biochem Nutr*. 2008;43:1–5.
- Kirk E, Reeds DN, Finck BN, Mayurranjan SM, Patterson BW, Klein S. Dietary fat and carbohydrates differentially alter insulin sensitivity during caloric restriction. *Gastroenterology*. 2009;136:1552–1560.
- Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, Franklin B, Kris-Etherton P, Harris WS, Howard B, Karanja N, Lefevre M, Rudel L, Sacks F, Van Horn L, Winston M, Wylie-Rosett J. Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation*. 2006;114:82–96.
- Cornier MA, Donahoo WT, Pereira R, Gurevich I, Westergren R, Enerback S, Eckel PJ, Goalstone ML, Hill JO, Eckel RH, Draznin B. Insulin sensitivity determines the effectiveness of dietary macronutrient composition on weight loss in obese women. *Obes Res*. 2005;13:703–709.
- Ridker PM. High-sensitivity C-reactive protein and cardiovascular risk: rationale for screening and primary prevention. *Am J Cardiol*. 2003;92:17K–22K.
- Wilson PW, D'Agostino RB, Levy D, Belanger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation*. 1998;97:1837–1847.
- Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486–2497.
- Kirk EP, Klein S. Pathogenesis and pathophysiology of the cardiometabolic syndrome. *J Clin Hypertens (Greenwich)*. 2009;11:761–765.
- Ashwell M, Gunn P, Gibson S. Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes Rev*. 2012;13:275–286.
- McArdle WD, Katch FI, Pechar GS, Jacobson L, Ruck S. Reliability and interrelationships between maximal oxygen intake, physical work capacity and step-test scores in college women. *Med Sci Sports*. 1972;4:182–186.
- Camhi SM, Stefanick ML, Ridker PM, Young DR. Changes in C-reactive protein from low-fat diet and/or physical activity in men and women with and without metabolic syndrome. *Metabolism*. 2010;59:54–61.
- Wing RR. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med*. 2010;170:1566–1575.
- Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, Stein RI, Mohammed BS, Miller B, Rader DJ, Zemel B, Wadden TA, Tenhave T, Newcomb CW, Klein S. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet: a randomized trial. *Ann Intern Med*. 2010;153:147–157.
- Burke LE, Hudson AG, Warziski MT, Styn MA, Music E, Elci OU, Sereika SM. Effects of a vegetarian diet and treatment preference on biochemical and dietary variables in overweight and obese adults: a randomized clinical trial. *Am J Clin Nutr*. 2007;86:588–596.
- Ashwell M, Hsieh SD. Six reasons why the waist-to-height ratio is a rapid and effective global indicator for health risks of obesity and how its use could

- simplify the international public health message on obesity. *Int J Food Sci Nutr*. 2005;56:303–307.
30. Rana JS, Boekholdt SM, Kastelein JJ, Shah PK. The role of non-HDL cholesterol in risk stratification for coronary artery disease. *Curr Atheroscler Rep*. 2012;14:130–134.
 31. Earnest CP, Poirier P, Carnethon MR, Blair SN, Church TS. Autonomic function and change in insulin for exercising postmenopausal women. *Maturitas*. 2010;65:284–291.
 32. Blumenthal JA, Babyak MA, Sherwood A, Craighead L, Lin PH, Johnson J, Watkins LL, Wang JT, Kuhn C, Feinglos M, Hinderliter A. Effects of the dietary approaches to stop hypertension diet alone and in combination with exercise and caloric restriction on insulin sensitivity and lipids. *Hypertension*. 2010;55:1199–1205.
 33. Church TS, Earnest CP, Skinner JS, Blair SN. Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure: a randomized controlled trial. *JAMA*. 2007;297:2081–2091.
 34. Borel AL, Nazare JA, Smith J, Almeras N, Tremblay A, Bergeron J, Poirier P, Despres JP. Visceral and not subcutaneous abdominal adiposity reduction drives the benefits of a 1-year lifestyle modification program. *Obesity (Silver Spring)*. 2012;20:1223–1233.
 35. Christiansen T, Paulsen SK, Bruun JM, Pedersen SB, Richelsen B. Exercise training versus diet-induced weight-loss on metabolic risk factors and inflammatory markers in obese subjects: a 12-week randomized intervention study. *Am J Physiol Endocrinol Metab*. 2010;298:E824–E831.
 36. Albu JB, Heilbronn LK, Kelley DE, Smith SR, Azuma K, Berk ES, Pi-Sunyer FX, Ravussin E. Metabolic changes following a 1-year diet and exercise intervention in patients with type 2 diabetes. *Diabetes*. 2010;59:627–633.
 37. Krauss RM. Lipoprotein subfractions and cardiovascular disease risk. *Curr Opin Lipidol*. 2010;21:305–311.
 38. Due A, Larsen TM, Mu H, Hermansen K, Stender S, Astrup A. Comparison of 3 ad libitum diets for weight-loss maintenance, risk of cardiovascular disease, and diabetes: a 6-mo randomized, controlled trial. *Am J Clin Nutr*. 2008;88:1232–1241.
 39. Lavoie ME, Faraj M, Strychar I, Doucet E, Brochu M, Lavoie JM, Rabasa-Lhoret R. Synergistic associations of physical activity and diet quality on cardiometabolic risk factors in overweight and obese postmenopausal women. *Br J Nutr*. 2012; [Epub ahead of print]: doi:10.1017/S0007114512001699.