

Minireview

The Mechanism of Phloem Loading in Rice (*Oryza sativa*)

Joon-Seob Eom, Sang-Bong Choi¹, John M. Ward², and Jong-Seong Jeon*

Carbohydrates, mainly sucrose, that are synthesized in source organs are transported to sink organs to support growth and development. Phloem loading of sucrose is a crucial step that drives long-distance transport by elevating hydrostatic pressure in the phloem. Three phloem loading strategies have been identified, two active mechanisms, apoplastic loading via sucrose transporters and symplastic polymer trapping, and one passive mechanism. The first two active loading mechanisms require metabolic energy, carbohydrate is loaded into the phloem against a concentration gradient. The passive process, diffusion, involves equilibration of sucrose and other metabolites between cells through plasmodesmata. Many higher plant species including *Arabidopsis* utilize the active loading mechanisms to increase carbohydrate in the phloem to higher concentrations than that in mesophyll cells. In contrast, recent data revealed that a large number of plants, especially woody species, load sucrose passively by maintaining a high concentration in mesophyll cells. However, it still remains to be determined how the worldwide important cereal crop, rice, loads sucrose into the phloem in source organs. Based on the literature and our results, we propose a potential strategy of phloem loading in rice. Elucidation of the phloem loading mechanism should improve our understanding of rice development and facilitate its manipulation towards the increase of crop productivity.

INTRODUCTION

Sucrose (Suc) is the main carbohydrate product of photosynthesis in higher plants and is transported long distances from source to various sink organs such as flowers, seeds and roots, thus supporting plant growth and development. Long-distance transport of Suc is initiated by Suc uptake into collective phloem of minor veins of source leaves. This process is referred to as phloem loading and is the first important step of the long-distance transport of photoassimilate to sink organs. Suc is then translocated via transport phloem and unloaded into sink organs via release phloem (van Bel, 2003). The phloem contains two main cell types, sieve elements (SEs) and companion cells (CCs). This long-distance transport occurs in SEs, which

lack a nucleus and have very few organelles and thus rely on associated CCs for most metabolic requirements (van Bel and Knoblauch, 2000).

High concentrations of photoassimilate loaded into source collective phloem generates the driving force, a hydrostatic pressure gradient, that enables the mass flow of Suc to sink organs via transport and release phloem (Turgeon, 2010; van Bel, 1996). That is, a high concentration of Suc inside the phloem cells in source organs creates an osmotic potential gradient that draws water into the cells. Phloem sap thus moves from source to sink by means of turgor pressure gradient.

In collective phloem, the loading strategies can be divided into three major pathways: apoplastic loading, polymer trapping, and diffusion. The apoplastic loading strategy (Fig. 1A) uses the proton motive force as metabolic energy to load Suc from the apoplast (cell wall space) actively into the phloem by Suc transporters (SUTs). In the second mechanism, polymer trapping (Fig. 1B), Suc diffuses from mesophyll cells into the specialized CCs called intermediary cells, symplastically through the cell-to-cell connections called plasmodesmata. Suc then serves as a substrate to synthesize raffinose family oligosaccharides (RFOs) such as raffinose and stachyose to increase the concentration of these sugars in the phloem in this thermodynamically active process (McCaskill and Turgeon, 2007; Zhang and Turgeon, 2009). Therefore, these two strategies are metabolic energy-dependent pathways. In contrast, the diffusion strategy (Fig. 1C), a passive loading mechanism, is an energetically downhill process because Suc levels are higher in mesophyll cells than in collective phloem, and no energy is used to collect Suc in SEs.

Rice is an important crop worldwide and is the principal food source for a large proportion of the global population. In general, the yield potential of rice crop plants is dependent upon whole plant carbohydrate partitioning mediated by Suc translocation from source to sink organs. Therefore, understanding the entire carbohydrate partitioning process whereby Suc is assimilated in source organs and translocated through the phloem to sink organs should provide new approaches to improve plant growth and crop yield (Lim et al., 2006). In this review, we summarize putative phloem loading strategies by integrating recent reports on SUTs, and propose a potential phloem loading strategy that may be especially important in rice.

Graduate School of Biotechnology and Crop Biotech Institute, Kyung Hee University, Yongin 446-701, Korea, ¹Division of Bioscience and Bioinformatics, Myongji University, Yongin 449-728, Korea, ²Plant Biology Department, University of Minnesota, St. Paul, MN 55108, USA
*Correspondence: jjeon@khu.ac.kr




Fig. 1. Schematic representation of phloem loading mechanism. (A) Apoplastic phloem loading mechanism. Suc synthesized in mesophyll cells diffuses to vascular parenchyma cells, is exported to apoplast, then transported by a SUT into the phloem. (B) Symplastic polymer trapping mechanism. Suc is mobilized symplastically from mesophyll cells to the specialized CCs, intermediary cells of the collective minor veins. RFOs are then synthesized in intermediary cells. This generates high sugar concentrations in the phloem. (C) Symplastic passive loading (diffusion) mechanism. Suc is translocated passively from mesophyll cells to the phloem of minor veins. This passive loading mechanism requires high Suc concentrations in mesophyll cells to drive the diffusion via plasmodesmata into the phloem. (D) Revised symplastic passive loading mechanism mediated by vacuolar Suc trapping in rice. In this model, the tonoplast SUT is essential to regulate Suc flux to the phloem. CC, companion cell; M, mesophyll cell; SE, sieve element, IC, intermediary cell.

PUTATIVE PHLOEM LOADING MECHANISMS

Apoplastic loading

While a portion of fixed carbon is stored transiently as starch in the chloroplasts and Suc in the vacuole, a large amount of photoassimilate fixed during the day is exported directly through the phloem. Suc is synthesized in mesophyll cells, diffuses symplastically through plasmodesmata into bundle sheath cells and subsequently vascular parenchyma cells. Suc is then exported into the apoplast. In the apoplastic loading strategy, the phloem is symplastically isolated or has rare plasmodesmata connecting to surrounding cells. Subsequently, Suc must be transported across the plasma membrane of CCs and/or SEs from the apoplast (Gottwald et al., 2000). In this apoplastic loading mechanism, the function of SUTs localized in the plasma membrane is essential for pumping Suc into the phloem. The proton motive force generated by H^+ -ATPases is the form of energy used to drive Suc into the phloem by SUTs against a concentration gradient (Braun and Slewinski, 2009; Lalonde et al, 2004; Sauer, 2007; Turgeon, 2010). It is also

noteworthy that active phloem loading is assumed to allow plants to maintain low photoassimilate concentrations in source leaves and thus to avoid feedback inhibition of photosynthesis, as well as to elevate sufficient pressure in the phloem to enable long-distance transport (Fig. 1A; Turgeon, 2010).

A number of plant species including the model plant *Arabidopsis* utilize the apoplastic loading pathway. SUTs are the best characterized components involved in apoplastic phloem loading. The first SUT genes, *SoSUT1* and *StSUT1*, were isolated from spinach (*Spinacia oleracea*) and potato (*Solanum tuberosum*), respectively (Riesmeier et al., 1992; 1993). Antisense suppression of *StSUT1* in potato causes local breaching and curling of leaves and dramatic reduction in root development and tuber yield. The plant leaves contain over 20-fold higher soluble sugars and 5-fold higher starch compared with wild type plants. This work revealed an essential role of SUTs, localized in the phloem plasma membrane, in the primary step in the apoplastic loading pathway (Riesmeier et al., 1994). Subsequently, SUTs have been identified throughout the plant kingdom (Reinders et al., 2012).

In *Arabidopsis*, AtSUC2 is a potential apoplastic phloem loader, which acts as Suc/H⁺ symporter to uptake Suc into the phloem. *atsuc2* mutants display a severe growth retardation phenotype that is considered typical for an apoplastic phloem loader mutant (Gottwald et al., 2000). In the monocot species maize (*Zea mays*), *zmsut1* mutant plants develop chlorotic leaves that accumulate high concentrations of sugars and prematurely senesce. The feeding of [¹⁴C]Suc demonstrated that Suc export is diminished in the mutants compared with wild type. In addition, the *zmsut1* plants display reduced plant height, fewer leaves, delayed flowering, and stunted tassel development (Slewiniski et al., 2009). Therefore, in maize, the activity of ZmSUT1 is essential for phloem loading of Suc.

In the apoplastic loading strategy, the mechanism for Suc export from the vascular parenchyma cells to the apoplast as a prerequisite for phloem loading by SUTs has been elusive. However, a transporter responsible for this Suc efflux process was recently discovered to be SWEET proteins (Chen et al., 2012) that were previously known as sugar, glucose and fructose, efflux transporters in *Arabidopsis* and rice (Chen et al., 2010). Expression analysis of *Arabidopsis* AtSWEET11 and AtSWEET12 promoter:: β -glucuronidase (*GUS*) transgenic plants indicated that these genes are preferentially expressed in vascular parenchyma cells. In addition, *atsweet11/atsweet12* double mutants were defective in phloem loading (Chen et al., 2012), thus revealing an important function of SWEET-mediated uniport of Suc from the parenchyma cells feeding H⁺-coupled import into the phloem SE-CC complex.

It is noteworthy that while mutation of the Suc carriers such as *AtSUC2* and *ZmSUT1*, caused a typical phenotype of apoplastic phloem loading mutants, these plants can complete their life cycle and make viable seeds (Slewiniski et al., 2009; Srivastava et al., 2009). Therefore, the apoplastic loading plant species largely follow apoplastic loading pathway, but are capable to use an alternative loading mechanism such as the symplastic loading strategy that can partially compensate. In addition, apoplastic loading was found to be highly correlated with the herbaceous habit (Davidson et al., 2011).

Polymer trapping

In the polymer trapping mechanism, Suc is mobilized symplastically from mesophyll cells to intermediary cells of the collective minor veins. Polymer trapping is thus a specialized form of the symplastic phloem loading mechanism. Although polymer trapping does not involve active transport of Suc across the plasma membrane, it is considered a thermodynamically active process because Suc is converted to RFOs, a process that requires metabolic energy (Turgeon, 1996; 2010; Zhang and Turgeon, 2009). RFOs are assumed to be too large to diffuse back to mesophyll through the plasmodesmata but can proceed through wider plasmodesmata into SEs. Therefore, RFOs synthesized in intermediary cells are essential to maintain high sugar concentrations in the phloem, which is similar to that in apoplastic loading species. This allows mesophyll cells to maintain low Suc levels (Fig. 1B; Slewiniski and Braun, 2010; Turgeon, 2010).

The presence of intermediary cells is always correlated with the translocation of considerable RFOs. Plasmodesmata are especially numerous between bundle sheath cells and intermediary cells in the minor veins of species such as *Verbascum phoeniceum*, *Rehmannia glutinosa*, *Catalpa speciosa*, *Buddleja davidii*, *Citrullus lanatus*, and *Coleus blumei* that transport RFOs (Gamalei, 1989; Turgeon and Gowan, 1990; Rennie and Turgeon, 2009; Zhang and Turgeon, 2009).

The polymer trapping hypothesis is supported by recent mo-

lecular and genetic evidence. In *V. phoeniceum*, RNAi suppression of a plasma membrane-localized SUT did not produce a typical phenotype of apoplastic phloem loader mutants (Zhang and Turgeon, 2009). Galactinol synthase (GAS) catalyzes the first committed step of RFO synthesis, which produces galactinol from *myo*-inositol and UDP-galactose. Galactinol then serves as the galactosyl donor in the RFO synthesis. RNAi suppression of the two *VpGAS* genes resulted in pronounced inhibition of RFO synthesis. As a result, transgenic plants showed carbohydrate accumulation, reduced sugar export, leaf chlorosis and severe growth retardation (McCaskill and Turgeon, 2007). This indicates that the mechanism of polymer trapping is most likely dependent on RFO synthesis in intermediary cells but does not require active Suc transport from the apoplast (Zhang and Turgeon, 2009). Notably, a floristic analysis indicated that plants with intermediary cells are over-represented in the tropics and subtropics (Davidson et al., 2011).

Diffusion, a passive symplastic loading

A third proposed mechanism for phloem loading is a passive symplastic pathway called diffusion (Reidel et al., 2009; Slewiniski and Braun, 2010; Turgeon, 2010; Turgeon and Medville, 1998). This is a passive flux of Suc from mesophyll cells to the phloem of minor veins. Plant species that utilize this passive loading mechanism maintain high levels of Suc in mesophyll cells to drive diffusion via plasmodesmata into the phloem, which is an energetically downhill process (Fig. 1C; Reidel et al., 2009; Rennie and Turgeon, 2009; Turgeon and Medville, 1998).

Plasmodesmatal frequencies in the phloem of leaf minor veins vary considerably (Davidson et al., 2011; Gamalei, 1989; van Bel and Gamalei, 1992). Plants with high, intermediate, and low plasmodesmatal frequencies in the phloem of minor veins are classified to type 1, 1-2a, and 2, respectively (Gamalei, 1989). Analysis of minor vein ultrastructure and plasmodesmatal frequencies of CCs indicates that most of the symplastic loaders are type 1, having numerous plasmodesmata to maintain the passive flux (Rennie and Turgeon, 2009; Schulz, 2005; Turgeon and Ayre, 2005). In contrast, *Arabidopsis* and maize belong to type 1-2a and type 2, respectively, by Gamalei's definition (Evert et al., 1978; Gamalei, 1989; Haritatos et al., 2000). In the analysis of 45 herbaceous and woody species, 14 exhibited the characteristics of passive loading (Rennie and Turgeon, 2009). This number included 11 of 19 woody plants analyzed. This indicates strong association between passive symplastic loading and the tree growth form (Davidson et al., 2011).

Since mesophyll cells and the phloem are symplastically linked, these plants do not accumulate radiolabeled Suc in the minor veins when their leaf discs are incubated in [¹⁴C]Suc. In symplastic loaders, Suc taken up into the symplast diffuses through plasmodesmata into mesophyll cells as well as the phloem. In contrast, apoplastic loaders and polymer trapping species showed clear vein images (Rennie and Turgeon, 2009; Turgeon, 2010).

PHLOEM LOADING STRATEGY IN RICE

From observation of the ultrastructure of vascular bundles, it was suggested that both apoplastic and symplastic phloem loading pathways are possible in rice (Chonan et al., 1984; Kaneko et al., 1980), but the exact phloem loading mechanism in rice has not been clearly demonstrated.

Functions of rice SUTs

In apoplastic loaders, function of the plasma membrane-localized SUTs is essential for translocating Suc from apoplast into




Fig. 2. Isolation and characterization of the *ossut1* mutant. (A) Schematic diagram of the rice *OsSUT1* gene and the Tos17 insertion position of line NF8036. The 14 exons are indicated by boxes. (B) RT-PCR analysis of wild type (+/+) and homozygous (-/-) mutant. *OsSUT1* transcripts are not detectable in the homozygous mutant. (C) Growth phenotype of wild type (left) and homozygous *ossut1* mutant (right) at heading stage.

the phloem against concentration gradient (Braun and Slewinski, 2009; Lalonde et al., 2004; Sauer, 2007; Turgeon, 2010). Based on comparison of their deduced amino acid sequences, SUTs are categorized into three major groups: type I (specific to eudicots, plasma membrane localized), type II (present in all plants, plasma membrane localized), and type III (present in all plants, vacuolar membrane localized). Members of type I are responsible for phloem loading or Suc import into sink organs (Gottwald et al., 2000; Riesmeier et al., 1994). Monocot species utilize type II SUTs for phloem loading (Aoki et al., 2003; Kühn, 2003; Lim et al., 2006; Slewinski et al., 2009). Type III SUTs are localized at the vacuolar membrane (tonoplast) and function in Suc transport into the cytosol from the vacuole lumen (Endler et al., 2006; Eom et al., 2011; Reinders et al., 2008). Of five *OsSUTs* in rice, four, *OsSUT1*, *OsSUT3*, *OsSUT4*, and *OsSUT5*, are categorized into type II, and *OsSUT2* belongs to the type III family of tonoplast SUT members. Of type II *OsSUTs*, *OsSUT4* is closely related to dicot type SUTs of type II family, sharing a common feature of N-terminal extension and long central loop, whose function has not been clearly demonstrated (Aoki et al., 2003; Hirose et al., 1997). In addition, *OsSUT5* was further classified to a subgroup of monocot SUTs (Kühn and Grof, 2010), and its transport activity is less sensitive to pH, than other SUTs in the type II family (Sun et al., 2010). Therefore, it is hypothesized that the remaining members of type II, *OsSUT1* and/or *OsSUT3*, are candidates to function in apoplastic phloem loading.

OsSUT1 is most likely an ortholog of the phloem loader, *ZmSUT1*, of maize (Slewinski et al., 2009). In gene expression pattern analysis, *OsSUT1* mRNA was highly expressed in leaves as well as in stems and filling grains (Aoki et al., 2003;

Hirose et al., 1997). A detailed expression pattern of *OsSUT1* was conducted by immunolocalization using an *OsSUT1* peptide-specific antibody (Scofield et al., 2007a; 2007b). The results showed that during germination and early seedling growth *OsSUT1* is confined to the phloem of coleoptiles and first and second leaf blades (Scofield et al., 2007a). Also, in mature rice plants, *OsSUT1* is present in the mature phloem of all the vegetative tissues, from flag leaf blade to the base of the filling grain, involved in long-distance transport pathway during grain filling (Scofield et al., 2007b). The results obtained from expression and localization experiments suggest that *OsSUT1* may function in apoplastic phloem loading in leaves.

However, transgenic rice plants with antisense suppression of *OsSUT1* do not accumulate Suc or show significant effects on photosynthetic rates in source leaves. Also, their vegetative growth does not show any visible symptom (Ishimaru et al., 2001; Scofield et al., 2002), raising the possibility that apoplastic phloem loading is a minor pathway or not largely contributed by *OsSUT1* in rice vegetative tissues. Although *OsSUT1* antisense rice plants under the control of CaMV35S (Ishimaru et al., 2001) or maize ubiquitin (Scofield et al., 2002) promoter were assessed, it would be valuable to examine null mutants because the antisense lines have still low expression of *OsSUT1* that can be sufficient to maintain phloem loading (Ishimaru et al., 2001; Scofield et al., 2002). Recent analysis of *Tos17* insertional mutant of *OsSUT1* revealed that germination of mutant pollen is impaired (Hirose et al., 2010). Therefore, homozygous mutant plants of *OsSUT1* have not been produced by normal self-pollination. In our present study, we generated homozygous *ossut1* mutants from the *Tos17* insertional heterozygous line of *OsSUT1* by using anther culture (Fig. 2A).

RT-PCR analysis indicated that the *ossut1* mutant plants are null lacking any expression of *OsSUT1* (Fig. 2B). During their vegetative growth stage, the *ossut1* mutants are indistinguishable from wild type plants (Fig. 2C), which is therefore consistent with those of *OsSUT1* antisense transgenic rice plants (Ishimaru et al., 2001; Scofield et al., 2002). This indicates that *OsSUT1*-mediated pathway may not be a primary route for phloem loading in source leaves of rice. It is therefore now assumed that *OsSUT1* may function in retrieval by phloem reloading of Suc leaked from the phloem (Scofield et al., 2007b).

It is noteworthy that, by *in situ* localization, *OsSUT1* is expressed in the maternal nucellar projection and nucellar epidermis, and the filial aleurone tissues but not in the starchy endosperm of mid-developing seeds (Furbank et al., 2001). This is consistent with the severe defects in grain filling in *OsSUT1* antisense plants as well as retarded development at the early stage of vegetative growth (Scofield et al., 2002). In addition, in tissue slices of filling grains, treatment of *p*-chloromercuriben-zene sulfonate, a SUT inhibitor, interferes with Suc transport (Furbank et al., 2001). These results suggest that *OsSUT1* may play a role in Suc transport from maternal to aleurone tissues via an apoplastic route.

OsSUT3 is also a type II SUT. *In situ* localization was used to show that *OsSUT3* is highly expressed in developing pollen (Ngampanya et al., 2002; Takeda et al., 2001), suggesting a role in Suc translocation into pollen. Our preliminary analysis of *OsSUT3* promoter::GUS transgenic rice plants confirms that it is preferentially expressed in pollen. This expression suggests that *OsSUT3* may function in developing pollen rather than in the phloem loading of source organs. In summary, apoplastic loading mechanism may not be predominant for phloem loading in source leaves of rice.

Although phylogenetic analysis of SUTs above indicated that *OsSUT4* and *OsSUT5* are classified within different subgroups than *OsSUT1* (Aoki et al., 2003; Kühn and Grof, 2010; Lim et al., 2006), we cannot completely rule out the possibility that *OsSUT4* and/or *OsSUT5* may have redundant function in apoplastic loading. Cellular localization assay in detail on *OsSUT4* and *OsSUT5* may aid understanding of the *in planta* roles of these rice SUTs. In addition, at present the mutant lines of *OsSUT3*, *OsSUT4*, and *OsSUT5* are available from our rice T-DNA mutant population and their single, double, and multiple mutants are being analyzed in our laboratory.

Involvement of tonoplast SUT in phloem loading

In photoautotrophic cells, vacuoles store excess Suc during the day and export it at night (Ayre, 2011; Linka and Weber, 2010; Martinoia et al., 2007; Neuhaus, 2007). Recently, through proteomic and/or GFP fusion analyses, tonoplast-localized SUTs have been identified (Endler et al., 2006; Eom et al., 2011; Okhubo-Kurihara et al., 2011; Payyavula et al., 2011; Reinders et al., 2008; Schneider et al., 2012; Schulz et al., 2011). Among them, expression of *PtaSUT4* was highest in mature source leaves. By *in situ* localization, *PtaSUT4* expression was found in epidermal cells, spongy mesophyll, the lower layer of the palisade mesophyll and minor phloem traces (Payyavula et al., 2011). In rice, *OsSUT2* expression was mainly found in mesophyll and bundle sheath cells of leaf blades and sheaths, rarely in CCs and SEs of the phloem (Eom et al., 2011). Thus, these have been suggested to function in the transport and vacuolar storage of photosynthetically derived Suc.

Functional characterization of tonoplast SUTs have been performed in *Populus*, rice and *Arabidopsis* (Eom et al., 2011; Payyavula et al., 2011; Schneider et al., 2012). In poplar, *PtaSUT4* RNAi transgenic plants exhibit 1.5 to 2-fold higher Suc content

in source leaves. An increased ratio of leaf-to-stem biomass indicates a link between vacuolar transport of Suc and biomass partitioning (Payyavula et al., 2011). In rice, the T-DNA insertional mutant, *ossut2* has increased Suc content both at the end of day (2-fold) and at the end of night (4-fold). The *ossut2* mutants also have diminished plant growth, tiller number, plant height, root dry weight and 1,000 grain weight. Decreased translocation of Suc from source leaves to sink parts of the *ossut2* mutant was also demonstrated (Eom et al., 2011). These data suggest that the tonoplast SUTs influence Suc export from source to sink organs.

Suc transport into *Xenopus* oocytes by LjSUT4, a *Lotus japonicus* tonoplast SUT, was found to be inhibited by the protonophore carbonyl cyanide *m*-chlorophenylhydrazone (CCCP) and the extracellular application of Suc induced membrane depolarization in LjSUT4-expressing *Xenopus* oocytes (Reinders et al., 2008). Suc transport by *OsSUT2* expressed in yeast was inhibited by CCCP (Eom et al., 2011). These findings indicate that tonoplast SUTs function as Suc/H⁺ symporters. Considering the normal pH gradient across the tonoplast, they function in Suc transport across the tonoplast from the vacuole lumen to the cytosol. It is probable that the Suc transport activity of *OsSUT2* in yeast is due to mis-targeting of the overexpressed protein to the plasma membrane in the heterologous system. In summary, tonoplast SUTs are involved in Suc transport from the vacuole lumen to the cytosol, playing an essential role in Suc export from source leaves to sink organs. It is noteworthy that poplar utilizes passive symplastic export (Russin and Evert, 1985; Turgeon and Medville, 1998). In contrast, in the apoplastic phloem loading species, *Arabidopsis*, mutation of the tonoplast *AtSUC4* showed no visible phenotype in normal growth conditions (Schneider et al., 2012), suggesting a relatively a minor role for the tonoplast SUT during Suc translocation.

Proposed phloem loading mechanism in rice

Based on the concentration and form of transport sugars, function of SUTs, and vein structures in source leaves of various plant species, a potential phloem loading mechanism is considered (Turgeon, 2010). Although the phloem loading mechanism in rice is still uncertain and need more consideration, previous and our present data provide some evidence that rice may utilize a passive loading strategy in source leaves as a primary phloem loading pathway. It has been shown that rice stores relatively high ratios of Suc to transitory starch in leaves, which differs from other plant species including *Arabidopsis* that primarily stores starch (Lee et al., 2008; Murchie et al., 2002; Nakano et al., 1995; Trevanion, 2002; Winder et al., 1998). It is noteworthy that concentrations of transport sugars in leaves correlate with a qualitative assessment of plasmodesmatal frequencies. Most of the type 1 and type 1-2a species with high and intermediate numbers of plasmodesmata have high concentrations of transport sugars, while many of the type 2 species have low concentrations of transport sugars (Rennie and Turgeon, 2009). It is known that the former group utilizes passive loading while the latter adopted an apoplastic loading strategy. In poplar the osmolalities between the phloem and mesophyll cells are approximately the same, as they are in the passive loading species willow (*Salix babylonica*) (Russin and Evert, 1985; Turgeon and Medville, 1998). High concentration of Suc in leaves is a diagnostic feature of plants that load passively (Rennie and Turgeon, 2009).

Numerous plasmodesmatal connections between parenchyma cells and CCs were observed in rice leaves in ultrastructural studies of small and large vascular bundles, suggesting

that Suc loading could occur via symplastic pathway (Botha et al., 2008; Chonan et al., 1981; Kaneko et al., 1980). In a dye-feeding experiment, the low molecular weight dye, 5,6-carboxyfluorescein diacetate was observed to move freely out of the phloem into the surrounding tissues including the mesophyll cells in the flag leaf blade (Scofield et al., 2007b), further supporting the possibility of a symplastic pathway in source leaves.

Antisense suppression of *OsSUT1* (Ishimaru et al., 2001; Schofield et al., 2002) and *Tos17* transposon insertional homozygous plant (Fig. 2) did not show a visible abnormal phenotype of apoplastic phloem loader mutants during vegetative growth. Therefore, *OsSUT1* might not function in the main phloem loading pathway in source leaves. There is no current evidence that another Suc transport activity compensates for the loss of *OsSUT1*. A closely related type II SUT in rice, *OsSUT3*, appeared to be preferentially expressed in pollen (Ngampanya et al., 2002; Takeda et al., 2001). Unless the expression pattern of *OsSUT3* is altered in the *OsSUT1* mutant and antisense lines, the possibility that *OsSUT3* functions in apoplastic phloem loading in source leaves can be excluded. *OsSUT4* and *OsSUT5* are also type II SUTs but are less related to *OsSUT1* and the phloem loader *ZmSUT1*. Although the functions of *OsSUT4* and *OsSUT5* in plants have not been determined (Aoki et al., 2003; Kühn and Grof, 2010; Lim et al., 2006), it is not likely that they have redundant function in apoplastic loading in source organs. Therefore, we are now proposing that rice does not utilize SUT-mediated apoplastic phloem loading pathway as a primary route.

Considering that Suc is temporarily stored in the vacuoles of photosynthetic assimilatory tissues (Linka and Weber, 2010; Martinoia et al., 2007; Neuhaus, 2007; Riens et al., 1991; Winter et al., 1993), this raises the question of the involvement and function of tonoplast-localized SUT in plant growth and development in rice. In this regard, it is noteworthy that the *ossut2* phenotype resembles those of apoplastic loaders (Eom et al., 2011; Gottwald et al., 2000; Slewinski et al., 2009). Rice plants lacking *OsSUT2* display severe growth defects compared with wild type and *OsSUT2*-complemented lines (Eom et al., 2011). In the *ossut2* mutant, Suc was accumulated in source leaves. Thus, it is probable that the accumulation of sugar in *ossut2*, due to the decreased transport of Suc from the vacuole lumen to the cytosol, interferes with Suc translocation to sink organs, and thus affects plant growth.

In summary, in rice, Suc most likely diffuses through plasmodesmata into the phloem of the minor vein via passive process. Suc concentrations of mesophyll cells and thus in the entire leaves are relatively higher than those in the minor veins. In this diffusion process, the tonoplast SUT, *OsSUT2* controls mesophyll cytosolic Suc concentration and therefore functions as a valve to regulate Suc flux into the phloem (Fig. 1D).

FUTURE PROSPECTS

Elucidating the phloem loading mechanism in source leaves is essential for our understanding of whole plant carbon partitioning via long-distance phloem translocation of photoassimilate. Here, we summarized current putative phloem loading mechanism of Suc in plant species and hypothesized the potential primary phloem loading pathway in rice. Several lines of evidence now support our proposal that rice may utilize a passive symplastic pathway to translocate Suc through plasmodesmata into the phloem of the minor vein. In particular, we hypothesized that "vacuolar trapping" regulated by the tonoplast *OsSUT2* is an essential process for Suc translocation into the phloem. In poplar and rice, tonoplast SUTs were found to strongly influ-

ence plant growth and development. Now both species are believed to utilize passive phloem loading pathway as primary Suc translocation in source leaves. Therefore, it would be interesting to determine whether in most of passive loading species tonoplast SUTs have the similar function that have been found in poplar and rice.

Expression and phylogenetic analysis suggested that *OsSUT3*, *OsSUT4*, and *OsSUT5* have distinct functions from apoplastic loaders. Nevertheless, future characterization of the single and multiple mutants of *OsSUT3*, *OsSUT4*, and *OsSUT5* should help us understand their function in rice. In preliminary experiments, we did not see visible abnormal plant growth of any of single, double, and triple homozygous mutant plants of *OsSUT3*, *OsSUT4*, and *OsSUT5*. Detailed analysis of these resources is further needed to determine their function.

Recently, SWEET proteins in *Arabidopsis* were found to export Suc from vascular parenchyma cells to apoplast for phloem loading by SUTs (Chen et al., 2012). Rice also has the SWEET homologs including *OsSWEET11/Xa13* and *OsSWEET14*. Interestingly, *OsSWEET11* functions as a rice susceptibility gene for specific pathovars of *Xanthomonas oryzae* pv. *oryzae* (Chen et al., 2012). Rice SWEETs may have evolved to regulate the efflux from parenchyma cells and highly localized transfer of Suc into the phloem, and thus reduce Suc release to the apoplast in order to prevent pathogen infections. There is the possibility that *OsSWEETs* may function in apoplastic loading in concert with *OsSUT1* under particular circumstances that require more Suc translocation into the phloem. In this regard, in rice that appears to be mainly dependent on passive symplastic loading, it would be interesting to see whether double or multiple mutations of these functionally redundant *OsSWEETs* may cause growth a defect phenotype like *atsweet11/atsweet12*.

ACKNOWLEDGMENTS

This work was supported by grants from the Next-Generation BioGreen 21 Program (PJ008114022011 to J.-S.J.), Rural Development Administration of the Korean Ministry of Food, Agriculture, Forestry, and Fisheries, and from the World Class University program (R33-2008-000-10168-0 to J.-S.J.) and the Mid-Career Researcher Program (2010-0026679 to J.-S.J.) from the Korean Ministry of Education, Science and Technology. The Division of Chemical Sciences, Geosciences, and Biosciences, Office of Basic Energy Sciences of the U.S. Department of Energy (grant DE-FG02-10ER15886) is gratefully acknowledged for support to J.M.W.

REFERENCES

- Aoki, N., Hirose, T., Scofield, G.N., Whitfield, P.R., and Furbank, R.T. (2003). The sucrose transporter gene family in rice. *Plant Cell Physiol.* **44**, 223-232.
- Ayre, B.G. (2011). Membrane-transport systems for sucrose in relation to whole-plant carbon partitioning. *Mol. Plant* **4**, 377-394.
- Botha, C.E., Aoki, N., Scofield, G.N., Liu, L., Furbank, R.T., and White, R.G. (2008). A xylem sap retrieval pathway in rice leaf blades: evidence of a role for endocytosis? *J. Exp. Bot.* **59**, 2945-2954.
- Braun, D.M., and Slewinski, T.L. (2009). Genetic control of carbon partitioning in grasses: roles of sucrose transporters and tied-ye loci in phloem loading. *Plant Physiol.* **149**, 71-81.
- Chonan, N., Kaneko, M., Kawahara, H., and Matsuda, T. (1981). Ultrastructure of the large vascular bundles in the leaves of rice plants. *Jpn. J. Crop Sci.* **50**, 323-331.
- Chonan, N., Kawahara, H., and Matsuda, T. (1984). Ultrastructure of vascular bundles and fundamental parenchyma in relation to movement of photosynthate in leaf sheath of rice. *Jpn. J. Crop Sci.* **53**, 435-444.

- Chen, L.Q., Hou, B.H., Lalonde, S., Takanaga, H., Hartung, M.L., Qu, X.Q., Guo, W.J., Kim, J.G., Underwood, W., Chaudhuri, B., et al. (2010). Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* **468**, 527-532.
- Chen, L.Q., Qu, X.Q., Hou, B.H., Sosso, D., Osorio, S., Fernie, A.R., and Frommer, W.B. (2012). Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. *Science* **335**, 207-211.
- Davidson, A., Keller, F., and Turgeon, R. (2011). Phloem loading, plant growth form, and climate. *Protoplasma* **248**, 153-163.
- Endler, A., Meyer, S., Schelbert, S., Schneider, T., Weschke, W., Peters, S.W., Keller, F., Baginsky, S., Martinoia, E., and Schmidt, U.G. (2006). Identification of a vacuolar sucrose transporter in barley and *Arabidopsis* mesophyll cells by a tonoplast proteomic approach. *Plant Physiol.* **141**, 196-207.
- Eom, J.S., Cho, J.I., Reinders, A., Lee, S.W., Yoo, Y., Tuan, P.Q., Choi, S.B., Bang, G., Park, Y.I., Cho, M.H., et al. (2011). Impaired function of the tonoplast-localized sucrose transporter in rice, *OsSUT2*, limits the transport of vacuolar reserve sucrose and affects plant growth. *Plant Physiol.* **157**, 109-119.
- Evert, R.F., Eschrich, W., and Heyse, W. (1978). Leaf structure in relation to solute transport and phloem loading in *Zea mays* L. *Planta* **138**, 279-294.
- Furbank, R.T., Scofield, G.N., Hirose, T., Wang, X-D., Patrick, J.W., and Offler, C.E. (2001). Cellular localization and function of a sucrose transporter *OsSUT1* in developing rice grains. *Aust. J. Plant Physiol.* **28**, 1187-1196.
- Gamalei, Y. (1989). Structure and function of leaf minor veins in trees and herbs. *Trees* **3**, 96-110.
- Gottwald, J.R., Krysan, P.J., Young, J.C., Evert, R.F., and Sussman, M.R. (2000). Genetic evidence for the in planta role of phloem-specific plasma membrane sucrose transporters. *Proc. Natl. Acad. Sci. USA* **97**, 13979-13984.
- Haritatos, E., Medville, R., and Turgeon, R. (2000). Minor vein structure and sugar transport in *Arabidopsis thaliana*. *Planta* **211**, 105-111.
- Hirose, T., Imaizumi, N., Scofield, G.N., Furbank, R.T., and Ohsugi, R. (1997). cDNA cloning and tissue specific expression of a gene for sucrose transporter from rice (*Oryza sativa* L.). *Plant Cell Physiol.* **38**, 1389-1396.
- Hirose, T., Zhang, Z., Miyao, A., Hirochika, H., Ohsugi, R., and Terao, T. (2010). Disruption of a gene for rice sucrose transporter, *OsSUT1*, impairs pollen function but pollen maturation is unaffected. *J. Exp. Bot.* **61**, 3639-3646.
- Ishimaru, K., Hirose, T., Aoki, N., Takahashi, S., Ono, K., Yamamoto, S., Wu, J., Saji, S., Baba, T., Ugaki, M., et al., (2001). Antisense expression of a rice sucrose transporter *OsSUT1* in rice (*Oryza sativa* L.). *Plant Cell Physiol.* **42**, 1181-1185.
- Kaneko, M., Chonan, N., and Matsuda, T. (1980). Ultrastructure of the small vascular bundles and transfer pathways for photosynthate in the leaves of rice plant. *Jpn. J. Crop Sci.* **49**, 42-50.
- Kühn, C. (2003). A comparison of the sucrose transporter systems of different plant species. *Plant Biol.* **5**, 215-232.
- Kühn, C., and Grof, C.P.L. (2010). Sucrose transporters of higher plants. *Curr. Opin. Plant Biol.* **13**, 288-298.
- Lalonde, S., Wipf, D., and Frommer, W.B. (2004). Transport mechanisms for organic forms of carbon and nitrogen between source and sink. *Annu. Rev. Plant Biol.* **55**, 341-372.
- Lee, S.K., Jeon, J.S., Börmke, F., Voll, L., Cho, J.I., Goh, C.H., Jeong, S.W., Park, Y.I., Kim, S.J., Choi, S.B., et al., (2008). Loss of cytosolic fructose-1,6-bisphosphatase limits photosynthetic sucrose synthesis and causes severe growth retardation in rice (*Oryza sativa*). *Plant Cell Environ.* **31**, 1851-1863.
- Lim, J.D., Cho, J.I., Park, Y.I., Hahn, T.R., Choi, S.B., and Jeon, J.S. (2006). Sucrose transport from source to sink seeds in rice. *Physiol. Planta.* **126**, 572-584.
- Linka, N., and Weber, A.P. (2010). Intracellular metabolite transporters in plants. *Mol. Plant* **3**, 21-53.
- Martinoia, E., Maeshima, M., and Neuhaus, H.E. (2007). Vacuolar transporters and their essential role in plant metabolism. *J. Exp. Bot.* **58**, 83-102.
- McCaskill, A., and Turgeon, R. (2007). Phloem loading in *Verbascum phoeniceum* L. depends on the synthesis of raffinose-family oligosaccharides. *Proc. Natl. Acad. Sci. USA* **104**, 19619-19624.
- Murchie, E.H., Yang, J., Hubbart, S., Horton, P., and Peng, S. (2002). Are there associations between grain-filling rate and photosynthesis in the flag leaves of field-grown rice? *J. Exp. Bot.* **53**, 2217-2224.
- Nakano, H., Makino, A., and Mae, T. (1995). Effects of panicle removal on the photosynthetic characteristics of the flag leaf of rice plants during the ripening stage. *Plant Cell Physiol.* **36**, 653-659.
- Neuhaus, H.E. (2007). Transport of primary metabolites across the plant vacuolar membrane. *FEBS Lett.* **581**, 2223-2226.
- Ngampanya, B., Takeda, T., Sonoda, Y., Narangajavana, J., and Yamaguchi, J. (2002). Characterization of *OsSUT2* cDNA expressed before flowering stage. *Rice Genet. Newsl.* **19**, 49-51.
- Okubo-Kurihara, E., Higaki, T., Kurihara, Y., Kutsuna, N., Yamaguchi, J., and Hasezawa, S. (2011). Sucrose transporter NtSUT4 from tobacco BY-2 involved in plant cell shape during miniprotoplast culture. *J. Plant Res.* **124**, 395-403.
- Payyavula, R.S., Tay, K.H., Tsai, C.J., and Harding, S.A. (2011). The sucrose transporter family in *Populus*: the importance of a tonoplast *PtaSUT4* to biomass and carbon partitioning. *Plant J.* **65**, 757-770.
- Reinders, A., Sivitz, A.B., Starker, C.G., Ganitt, J.S., and Ward, J.M. (2008). Functional analysis of *LjSUT4*, a vacuolar sucrose transporter from *Lotus japonicas*. *Plant Mol. Biol.* **68**, 289-299.
- Reinders, A., Sivitz, A.B., and Ward, J.M. (2012). Evolution of plant sucrose uptake transporters (SUTs). *Front. Plant Sci.* **3**, 00022.
- Rennie, E.A., and Turgeon, R. (2009). A comprehensive picture of phloem loading strategies. *Proc. Natl. Acad. Sci. USA* **106**, 14162-14167.
- Reidel, E.J., Rennie, E.A., Amiard V., Cheng, L., and Turgeon, R. (2009). Phloem loading strategies in three plant species that transport sugar alcohols. *Plant Physiol.* **149**, 1601-1608.
- Riens, B., Lohaus, G., Heineke, D., and Heldt, H.W. (1991). Amino acid and sucrose content determined in the cytosolic, chloroplastic, and vacuolar compartments and in the phloem sap of spinach leaves. *Plant Physiol.* **97**, 227-233.
- Riesmeier, J.W., Willmitzer, L., and Frommer, W.B. (1992). Isolation and characterization of a sucrose carrier cDNA from spinach by functional expression in yeast. *EMBO J.* **11**, 4705-4713.
- Riesmeier, J.W., Hirner, B., and Frommer, W.B. (1993). Potato sucrose transporter expression in minor veins indicates a role in phloem loading. *Plant Cell* **5**, 1591-1598.
- Riesmeier, J.W., Willmitzer, L., and Frommer, W.B. (1994). Evidence for an essential role of the sucrose transporter in phloem loading and assimilate partition. *EMBO J.* **13**, 1-7.
- Russin, W.A., and Evert, R.F. (1985). Studies on the leaf of *Populus deltoids* (Salicaceae): ultrastructure, plasmodesmata frequency, and solute concentrations. *Am. J. Bot.* **72**, 1232-1247.
- Sauer, N. (2007). Molecular physiology of higher plant sucrose transporters. *FEBS Lett.* **581**, 2309-2317.
- Schneider, S., Hulpke, S., Schulz, A., Yaron, I., Höll, J., Imlau, A., Schmitt, B., Batz, S., Wolf, S., Hedrich, R., et al., (2012). Vacuoles release sucrose via tonoplast-localized SUC4-type transporters. *Plant Biol. (Stuttg)* **14**, 325-336.
- Schulz, A. (2005). Role of plasmodesmata in solute loading and unloading. In *Plasmodesmata*, K.J. Oparka, ed. Vol. Annual Plant Reviews, Vol. 18. Blackwell, Oxford, pp. 135-161.
- Schulz, A., Beyhl, D., Marten, I., Wormit, A., Neuhaus, E., Poschet, G., Buttner, M., Schneider, S., Sauer, N., and Hedrich, R. (2011). Proton-driven sucrose symport and antiport are provided by the vacuolar transporters SUC4 and TMT1/2. *Plant J.* **68**, 129-136.
- Scofield, G.N., Hirose, T., Gaudron, J.A., Upadhyaya, N.M., Ohsugi, R., and Furbank, R.T. (2002). Antisense suppression of the rice sucrose transporter gene, *OsSUT1*, leads to impaired grain filling and germination but does not affect photosynthesis. *Funct. Plant Biol.* **29**, 815-826.
- Scofield, G.N., Aoki, N., Hirose, T., Takano, M., Jenkins C.L., and Furbank, R.T. (2007a). The role of the sucrose transporter, *OsSUT1*, in germination and early seedling growth and development of rice plants. *J. Exp. Bot.* **58**, 483-495.
- Scofield, G.N., Hirose, T., Aoki, N., and Furbank, R.T. (2007b). Involvement of the sucrose transporter, *OsSUT1*, in the long-distance pathway for assimilate transport in rice. *J. Exp. Bot.* **58**, 3155-3169.
- Slewisinski, T.L., and Braun, D.M. (2010). Current perspectives on the regulation of whole-plant carbohydrate partitioning. *Plant Sci.* **178**, 341-349.
- Slewisinski, T.L., Meeley, R., and Braun, D.M. (2009). Sucrose transporter1 functions in phloem loading in maize leaves. *J. Exp. Bot.* **60**, 881-892.

- Srivastava, A.C., Dasgupta, K., Aijeren, E., Costilla, G., McGarry, R.C., and Ayre, B.G. (2009). Arabidopsis plants harboring a mutation in *AtSUC2*, encoding the predominant sucrose/proton symporter necessary for efficient phloem transport, are able to complete their life cycle and produce viable seed. *Ann. Bot.* *104*, 1121-1128.
- Sun, Y., Reinders, A., LaFleur, K.R., Mori, T., and Ward, J.M. (2010). Transport activity of rice sucrose transporters OsSUT1 and OsSUT5. *Plant Cell Physiol.* *51*, 114-122.
- Takeda, T., Toyofuku, K., Matsukura, C., and Yamaguchi, J. (2001). Sugar transporters involved in flowering and grain development of rice. *J. Plant Physiol.* *158*, 465-470.
- Trevanion, S.J. (2002). Regulation of sucrose and starch synthesis in wheat (*Triticum aestivum* L.) leaves: role of fructose 2,6-bisphosphate. *Planta* *215*, 653-665.
- Turgeon, R. (1996). Phloem loading and plasmodesmata. *Trends Plant Sci.* *1*, 415-423.
- Turgeon, R. (2010). The role of phloem loading reconsidered. *Plant Physiol.* *152*, 1817-1823.
- Turgeon, R., and Ayre, B.G. (2005). Pathways and mechanisms of phloem loading. In *Vascular Transport in Plants*. Elsevier/Academic Press, N.M. Holbrook, and M.A. Zwieniecki, eds. (Oxford, UK: Elsevier/Academic Press), pp. 45-67.
- Turgeon, R., and Gowan, E. (1990). Phloem loading in *Coleus blumei* in the absence of carrier-mediated uptake of export sugar from the apoplast. *Plant Physiol.* *94*, 1244-1249.
- Turgeon, R., and Medville, R. (1998). The absence of phloem loading in willow leaves. *Proc. Natl. Acad. Sci. USA* *95*, 12055-12060.
- van Bel, A.J.E. (1996). Interaction between sieve element and companion cell and the consequences for photoassimilate distribution. Two structural hardware frames with associated physiological software packages in dicotyledons? *J. Exp. Bot.* *47*, 1129-1140.
- van Bel, A.J.E. (2003). Phloem, a miracle of ingenuity. *Plant Cell Environ.* *26*, 125-149.
- van Bel, A.J.E., and Gamalei, Y. (1992). Ecophysiology of phloem loading in source leaves. *Plant Cell Environ.* *15*, 265-270.
- van Bel, A.J.E., and Knoblauch, M. (2000). Sieve element and companion cell: the story of the comatose patient and the hyperactive nurse. *Funct. Plant Biol.* *27*, 477-487.
- Winder, T.L., Sun, J., Okita, T.W., and Edwards, G.E. (1998). Evidence for the occurrence of feedback inhibition of photosynthesis in rice. *Plant Cell Physiol.* *154*, 665-677.
- Winter, H., Robinson, D.G., and Heldt, H.W. (1993). Subcellular volumes and metabolite concentrations in barley leaves. *Planta* *191*, 180-190.
- Zhang, C., and Turgeon, R. (2009). Downregulating the sucrose transporter *VpSUT1* in *Verbascum phoeniceum* does not inhibit phloem loading. *Proc. Natl. Acad. Sci. USA* *106*, 18849-18854.