

Protein Kinase C α Protects Against Multidrug Resistance in Human Colon Cancer Cells

Se-Kyoung Lee, Adeb Shehzad, Jae-Chang Jung¹, Jong-Kyung Sonn¹, Jae-Tae Lee², Jeen-Woo Park, and Young-Sup Lee*

Multidrug resistance is the phenomenon by which, after exposure to a single chemotherapeutic agent, cancer cells evade the agent's cytotoxic effects as well as become resistant to several classes of diverse drugs. ATP-binding cassette (ABC) transporters are a family of transporter proteins that contribute to drug resistance via an ATP-dependent drug efflux pump. P-glycoprotein (P-gp) is a prominent ABC superfamily protein encoded by the *mdr* gene which has the ability to mediate the cellular extrusion of xenobiotics and anticancer drugs from tumor cells. Exclusively expressed P-gp cells from the human colon cancer HCT15/DOX line showed resistance to doxorubicin while parental HCT15 cells treated with doxorubicin displayed typical signs of apoptosis. In order to verify the hypothesis that expression of MDR is controlled in part, by protein kinase C (PKC), expression patterns of different PKC isoforms were examined in both cell lines. Of the PKC isoforms evaluated, the membrane translocation and expression levels of PKC α were strikingly increased in HCT15/DOX cells. PKC α reversed doxorubicin-induced apoptosis through the scavenging of ROS as well as inhibition of PARP cleavage. In addition, inhibition of PKC α with Go6976, a specific inhibitor of classical PKC, led to reduced MDR expression and increased doxorubicin-induced apoptosis. Knockdown of PKC α by siRNA diminished the protective effects of PKC α for doxorubicin-induced apoptosis. These results suggested that over-expression and activity of PKC α is closely associated with the regulation of the MDR phenotype in human colon cancer HCT15 cells and provided insight into a new strategy for inhibiting doxorubicin resistance in human cancers.

INTRODUCTION

Membrane transporter families play a pivotal role in the circulation and excretion of many clinically essential therapeutic drugs. P-glycoprotein (P-gp) is a 170-kDa ATP-dependent transmembrane glycoprotein which is the best characterized drug efflux

pump and known to confer resistance to a variety of structurally and functionally unrelated anti-cancer drugs. This phenomenon is known as multidrug resistance (MDR) and represents a serious problem in the chemotherapeutic management of cancers. Manifestation of the MDR phenotype is most often due to the over-expression of drug efflux pumps in the plasma membrane of cancer cells (Endicott and Ling, 1989; Gottesman et al., 2002). It has been reported that P-gp drug efflux activity can be modulated through phosphorylation by different families of protein kinases (Gottesman and Pastan, 1993). The protein kinase responsible for the phosphorylation of P-gp is believed to be protein kinase C (PKC) (Ling, 1997). PKC is a serine/threonine kinase involved in the signal transduction required for cellular proliferation and differentiation (Clemens et al., 1992; Nishizuka and Kikkawa, 2003). Classical and novel PKC groups respond to activation by phorbol esters (Gill et al., 2001). Conventional PKCs are, like the originally described kinase activity, dependent on Ca²⁺ and phospholipids for their activation. This group includes PKC α , PKC β I, PKC β II and PKC γ . A second group, including the novel PKCs, does not require Ca²⁺ but are phospholipid dependent and known PKC δ , PKC ϵ , PKC η and PKC θ . A third, more recently identified group includes PKC ζ and PKC λ/τ . These are known as atypical PKCs, which require neither phospholipids nor Ca²⁺ for their activation (Mellor and Parker, 1998).

A wide array of research has postulated that specific isoforms of PKCs may be either pro-apoptotic or anti-apoptotic depending on the condition and stimulus of the cell type (Leszczynski, 1995; Lucas and Sánchez-Margalet, 1995). Of the PKC isoforms, PKC α has been reported to be of particular determinant in the development and prolongation of the MDR phenotype. Studies have shown that PKC α is expressed abundantly in surgical specimens of human colon cancer and contributes to drug resistance as observed in colon cancer (Gravitt et al., 1994). Breast carcinoma cells also develop resistance against doxorubicin through elevated levels of P-gp and PKC α , and display the MDR phenotype (Budworth et al., 1997). Additionally, doxorubicin was screened against a panel of human tumor cell lines derived from cancer patients. This study re-

School of Life Sciences, Kyungpook National University, Daegu 702-701, Korea, ¹Department of Biology, College of Natural Sciences, Kyungpook National University, Daegu 702-701, Korea, ²Department of Nuclear Medicine, Kyungpook National University Hospital, Daegu 700-721, Korea
*Correspondence: yselee@knu.ac.kr

Received March 21, 2012; revised April 26, 2012; accepted April 27, 2012; published online May 23, 2012

Keywords: apoptosis, doxorubicin, multidrug resistance, PKC α

vealed that increased PKC α levels were associated with reduced drug efficacy, which means that one or more substrates of this kinase may be related to the MDR phenotype (Gariboldi et al., 2003).

Doxorubicin is commonly used for the treatment of different types of cancer. However, its efficacy is often reduced because it is a substrate for P-gp. Therefore, in order to investigate the individual roles of various PKC isoforms in MDR, we examined the expression and activity of the major isoforms of PKC in the parent HCT15 cell line and its drug-resistant subline, HCT15/DOX. It was observed that the expression level and membrane translocation of PKC α alone markedly increased in HCT15/DOX cells and functioned as an antagonist to doxorubicin-induced apoptosis. Additionally, inhibition of PKC α by a specific inhibitor, Go6967, and by using PKC α siRNA, both lead to reduced MDR activity and enhanced doxorubicin-induced apoptosis. The results of this study demonstrated that over-expression of an endogenous PKC α conferred increased MDR through the modulation of reactive oxygen species (ROS) generation in human colon cancer HCT15 cells.

MATERIALS AND METHODS

Materials

Doxorubicin (adriamycin), 4,6-Diamidino-2-phenylindole (DAPI) and propidium iodide were purchased from Sigma-Aldrich (USA). Dichlorofluorescein diacetate (DCFHDA), dihydrohodamine 123 (DHR 123), and rhodamine 123 were obtained from Molecular Probes (USA). The electrophoresis reagents and Bio-Rad protein assay kit were purchased from Bio-Rad Laboratories (USA). The anti-PKC isotypes antibodies were obtained from Cell Signaling Technology Inc. (USA). These chemicals were used according to the manufacturer's instructions.

Cell culture

The human colorectal carcinoma (HCT15) cells were obtained from ATCC, CCL-225. An doxorubicin-resistant subline of HCT15 cells, named HCT15/doxorubicin (HCT15/DOX), were established by continuous exposure to doxorubicin as described previously (Choi et al., 1996). Cells were cultured at a density of 5×10^5 in RPMI-1640 medium containing 10% heat inactivated fetal bovine serum (FBS), L-glutamine and supplemented with 1% (v/v) antibiotic-penicillin streptomycin (Gibco, Invitrogen Corporation) at 37°C in a 5% CO₂-humidified atmosphere. Exponentially growing cells were seeded at 1×10^6 cells per dish and then the cells were exposed to various agents.

Preparation of the cytosolic and membrane protein fractions

Cytosolic and particulate membrane fractions of cells were prepared as described previously but with slight modification (Tentori et al., 2001). Briefly, the cells were sonicated in buffer A (20 mM Tris, pH 7.5, 250 mM sucrose, 10 mM ethyleneglycol tetra-acetic acid (EGTA), 2 mM ethylenediamine tetra-acetic acid (EDTA), 1 mM sodium fluoride (NaF), 1 mM sodium orthovanadate (Na₃VO₄), 1 mM phenylmethylsulfonyl fluoride (PMSF), 1 μg/ml aprotinin, 1 μg/ml leupeptin, and 1 μg/ml pepstatin) and centrifuged for 10 min at $1,000 \times g$ to remove cell debris. The supernatants were then centrifuged at $100,000 \times g$ for 30 min and the resulting supernatants were saved as the cytosolic fractions. Proteins in the pellets were extracted with buffer B [20 mM Tris, pH 7.5, 1% sodium dodecyl sulfate (SDS), 150 mM NaCl, 1 mM EGTA, 1 mM NaF, 1 mM Na₃VO₄ and various protease inhibitors]. Following centrifugation at $100,000 \times g$ for

30 min, the supernatants were referred to as the particulate membrane fractions.

Determination of apoptosis

Cell viability was determined using a trypan blue exclusion test. Morphological analysis of apoptosis was performed after staining using Hoechst 33342 dye. The cells were fixed in 4% paraformaldehyde and permeabilized with PBS/0.5% Triton X-100, and the nuclei were stained for 20 min using Hoechst dye. The coverslips were then washed, mounted onto slides and viewed with a fluorescence microscope. Apoptotic DNA fragments were isolated according to a previously described method (Ma et al., 2000). The DNA was separated by electrophoresis on 1% agarose gels, stained with ethidium bromide and visualized under ultraviolet light.

Cell cycle analysis

The cells were trypsinized and collected by centrifugation at $1,000 \times g$ for 10 min. For fixation, 70% ethanol was added and the cell suspension was kept overnight at 4°C. The cells were then stained with propidium iodide solution (50 μg/ml propidium iodide, 0.1% Triton X-100, 0.1 mM EDTA and 50 μg/ml RNase) for 20 min at 4°C. The stained DNA was analyzed by flow cytometer (Becton Dickinson).

Measurement of ROS

Intracellular ROS concentration was measured using the oxidant-sensitive fluorescent probe, DCFHDA, with inverted microscope. Cells were grown at a density of 1×10^6 cells per 35 mm culture dish and maintained in the growth medium for 24 h. Cells were exposed to 5 μM DCFHDA for 20 min and were then washed with PBS. DCF fluorescence (excitation, 480 nm; emission, 520 nm) was imaged using an inverted microscope (Zeiss Axiovert 200). For FACS analyses, cellular DCF fluorescence measurement included at least 10,000 events/test using a flow cytometer with a fluorescein isothiocyanate filter.

Dihydrohodamine 123 accumulation assay

The HCT15 cells were incubated with 10 μM of DHR123 in the culture medium for 20 min and then washed three times with ice-cold PBS. The cells were imaged on an inverted microscope using FITC fluorescence intensity.

Protein kinase assay

The PKC activity was determined by measuring the transfer of ³²P-ATP to a peptide substrate that was captured on P81 phosphocellulose papers. Aliquots of cell lysates were incubated with 50 mM Tris at pH 7.5 with 200 μM PKC α substrate AAKIQASFRGHMARKK, 40 mM of MgCl₂ and 400 μM of ATP. After 5 min at 30°C, the reactions were terminated by adding EDTA to a final concentration of 20 mM. Relative PKC activity was then quantified with a liquid scintillation counter.

Western blotting

In brief, aliquots of protein extracts (30 μg) from cells of different treatment groups were suspended in 0.1 M Tris-HCl buffer (pH 7.4) containing 1% SDS, 0.05% β-mercaptoethanol, 2.5% glycerol and 0.001% bromophenol blue, and subsequently fractionated by 10% SDS-polyacrylamide gel electrophoresis. The proteins were transferred electrophoretically onto nitrocellulose membranes (0.2 μm, Schleicher and Schuell). The membranes were blocked using 5% non-fat dry milk and 0.1% Tween 20 in Tris buffered saline (TBS). The membranes were subsequently probed with primary antibody in TBS containing 3% non-fat dry




Fig. 1. Doxorubicin induced oxidative-stress apoptosis in HCT15 and HCT15/DOX cells. HCT15 and HCT15/DOX cells were incubated in the absence (con) or presence of doxorubicin for 24 h. (A) Cleavages of PARP and lamin B were analyzed by Western blot. (B) Analysis of doxorubicin-induced DNA fragmentation of cells by agarose gel electrophoresis. (C) At the indicated time, the cells were harvested and stained with propidium iodide and their DNA content was analyzed by flow cytometry. (D) Doxorubicin induces ROS generation. HCT15 and HCT15/DOX cells were treated with doxorubicin for the indicated time and further incubated with DCFHDA for 20 min. Accumulation of the probe in the cells was measured by an increased emission at 520 nm when the sample was excited at 480 nm. Fluorescence images of DCF-loaded cells were obtained under a microscope.

milk and 0.1% Tween 20. The antibody-antigen complexes were detected using goat anti-mouse IgG or goat anti-rabbit IgG peroxidase conjugates followed by the use of an enhanced chemiluminescence (ECL) detection kit (Amersham Corp.).

Reverse-transcription polymerase chain reaction (RT-PCR)

In order to determine if HCT15 cells expressed transcriptional levels of *mdr*, RT-PCR was performed. Total RNA was isolated from the doxorubicin-resistant HCT15/DOX and parental HCT15 cells using TRI reagent (Molecular Research Center, Inc.). RT-PCR was performed using an access RT-PCR kit (Promega). The following PCR conditions were employed: 1 cycle at 95°C for 5 min, 30 cycles at 95°C for 1 min, 55°C for 1 min, 72°C for 1 min, and an additional extension step of 5 min at 72°C. The amplified PCR products were analyzed by 1 % agarose gel electrophoresis and ethidium bromide staining. The following primer sequences for *mdr1* gene sequence were utilized as follows, 5'-GCCTGGCAGCTGGAAGACAAATACACAAAATT-3' (sense) and 5'-CAGACAGCAGCTGACAGTCCAAGAACAGGACT-3' (antisense), corresponding to the previously published cDNA sequence (Limthrakul et al., 2004), and 5'-GTCCDAGTCAACGGATTGG-3' (sense) and 5'-GGGTGGAATCATATTTGGAAGCTTG-3' (antisense) for GAPDH as an

internal loading control (Oligo, Macrogen).

Generation of the siRNA-PKC α HCT15 clones

The siRNA-PKC α pool was obtained from Santa Cruz Biotechnology (SC-36243) and utilized according to the manufacturer's instruction. HCT15 cells were transfected using Lipofectamine RNAiMAX (Invitrogen, USA) according to the manufacturer's recommendations. Cells were cultured for 24 h at densities of 1×10^6 cells per dish to 40-50% confluence. The cells were then transfected with 1 nmol of siRNA. At the indicated times after transfection, the cells were treated with 5 μ M doxorubicin for 24 h. The effects of scrambled siRNA were also evaluated according to the modified protocol as described previously (Nakura et al., 2011).

Statistical analysis

All data are presented as the mean values \pm SD (standard deviation) and originate from 3 separate experiments. The data was evaluated using SPSS for the student *t*-test and subjected to one-way analysis of variance (ANOVA). The significance of the difference between the means was determined by the Tukey range test and considered significant at $P \leq 0.05$.




Fig. 2. Over-expression of MDR in doxorubicin resistant HCT15/DOX cells. (A) RT-PCR analysis of P-glycoprotein (MDR). Total RNA was isolated from HCT15 parental and resistant cells, and RT-PCR was performed. PCR products were separated on 1.2% agarose gel and photographed. GAPDH was also examined as a reference. (B) Western blot identification of MDR expression in HCT15 and HCT15/DOX cells. Whole-cell lysates were extracted from HCT15 cells and separated by SDS-PAGE and the resulting proteins were detected by immunoblotting. (C) Mitochondria and membrane potential were revealed by rhodamine 123. Rhodamine 123 retention was evaluated after accumulation for 30 min and 30 min of efflux in dye-free medium.

RESULTS

Doxorubicin induced oxidative-stress apoptosis

The quinone moiety of the anthracyclines are known to act as catalysts in the formation of ROS including superoxide anion and hydrogen peroxide (Doroshov et al., 1986). At relevant physiological concentrations, a consequence of treatment with doxorubicin is the induction of apoptosis associated with DNA fragmentation and cell shrinkage (Bose et al., 1995; Skladanowski and Konopa, 1993). HCT15 and HCT15/DOX cells were treated with 2 and 4 μ M of doxorubicin for 12, 24 and 36 h, and the cell lysates from both cell lines were analyzed by Western blot. Doxorubicin induced apoptosis through poly (ADP-ribose) polymerase (PARP) and lamin B cleavage in the HCT15 cells only, and failed to induce apoptosis in the HCT15/DOX cells (Fig. 1A). Additionally, doxorubicin induced DNA fragmentation with rapid kinetics in the parental cells (Fig. 1B). Lamin B and PARP were cleaved with similar kinetics as the DNA fragmentation, yielding cleavage bands (Fig. 1B). The use of doxorubicin in cancer therapy has been hampered by the ability of cancer cells to develop doxorubicin resistance (Dolfini et al., 1997). In order to evaluate the degree of apoptosis induced by doxorubicin, both cell types were treated with doxorubicin for 12 and 24 h. As shown in Fig. 1C, doxorubicin induced 13% apoptosis in parental cells, but no apoptosis was observed in the resistant cell line. These results indicated that doxorubicin is hindered in inducing apoptosis in HCT15/DOX cell lines.

In order to investigate the role of oxidative stress in doxorubicin-induced apoptosis, we used a cell-permeable fluorescent dye, DCFHDA, to examine ROS generation in both types of cells in response to doxorubicin stimulation. A considerable increase in oxidant-induced 2',7'-dichlorofluorescein fluorescence was observed in the HCT15 cells but not in the HCT15/DOX cells (Fig. 1D). It is assumed that doxorubicin-induced apoptosis occurs through increase in ROS generation, but only in parental cells. Under resistance conditions, it does not induce intracellular oxidative stress.

MDR expression in HCT15 and HCT15/DOX cells

Expression of the *mdr1* gene was evaluated in the parental and

HCT15/DOX cells using RT-PCR. After RNA isolation, the expression of *mdr1* was determined using *mdr1*-specific primers producing a 243 bp RT-PCR product. Control RT-PCR was carried out in parallel with GAPDH. Little *mdr* expression was revealed in parental cells, while robust expression was observed in the HCT15/DOX cells (Fig. 2A). Gel electrophoresis was performed in order to detect the expression of MDR1 in both cell lines. The cell lysates were analyzed by Western blot with anti-MDR1. The expression of MDR was very high in the HCT15/DOX cells while remaining very little or undetected in the parental cell lines (Fig. 2B).

In order to gain further insight into the multiple resistance mediated by HCT15/DOX cells, rhodamine 123 was used as a molecular probe and functional assay because this fluorescent dye is considered to be a relatively specific substrate for P-gp, reduced retention of rhodamine 123 in the cells indicates the functioning of P-gp (Altenberg et al., 1994; Gravitt et al., 1994). Rhodamine 123 accumulation was evaluated for 30 min along with 30 min efflux in dye-free medium. As shown in Fig. 2C, doxorubicin-resistant subline HCT15/DOX significantly reduced the retention of rhodamine 123 as compared to the parental cells.

Differential expressions of PKC isoforms

It was realized very early that the mechanism of action for PKC induced reversal of MDR would require a specific substrate. Therefore, the anti-apoptotic activity of PKC and the effects of doxorubicin on PKC activity were determined. PKC was measured in the parental and resistant cells with a radioisotope assay that utilizes the phosphorylation of a synthetic substrate AAKIQASFRGHMARPCK. The PKC activity increased almost six-fold after treatment with doxorubicin in the HCT15/DOX cell lines (Fig. 3A). This result suggested that the activity of PKC is implicated in the doxorubicin resistance observed in the HCT15/DOX cells.

PKC isotypes α , β I and ζ were evaluated by Western blot in the investigation into their role in drug resistance. It was noted that the PKC α , β I and ζ -isoforms were detectable in the cytosol and in membrane portion of both cell lines, although some differences in their expressions were apparent. β I- and ζ -PKC




Fig. 3. PKC α is specifically activated in doxorubicin resistant cell lines. (A) Measurement of PKC activity in HCT15 and HCT15/DOX cells. Lysates from HCT15 and HCT15/DOX cells were used for the kinase assay using the specific PKC substrate peptide (AAKIQA SFRGHMARKK) and the 32 P-incorporation was measured by scintillation counter. Notably, specific up-regulation of PKC α was detected in the doxorubicin-resistant HCT15/DOX cells. (B) HCT15 and HCT15/DOX cells were treated with doxorubicin. The cells were fractionated into cytosolic and membrane fractions and equal amounts of lysates were used for immunoblotting with PKC isoform (α , β and ζ)-specific antibodies.

Fig. 4. Western blot analysis of PKC isoforms in HCT15 and HCT15/DOX cells. HCT15 and HCT15/DOX cells were treated with doxorubicin for the indicated time. The cells were fractionated into cytosolic and membrane fractions and equal amounts of lysates were used for immunoblotting with PKC isoform-specific antibodies.

protein levels were similar in the parental and resistant cell lines, but an obvious difference was observed in PKC α expression. As shown in Fig. 3B, a higher level of PKC α was found in the HCT15/DOX cell membranes than in the cytosol, as compared to the results with the HCT15 cells. Furthermore, total expression of PKC α was markedly decreased in the parental cell line and highly expressed in the HCT15/DOX cell line.

Protein level of PKC α was reduced by doxorubicin in HCT15 cells

The conventional PKC family contains many isoforms including PKC α and PKC β I. In order to scrutinize the effects of doxorubicin on the expression of PKC α and PKC β I in parental and resistant cell lines, we performed immunoblot experiments on their separated cytosolic and membrane fractions. As shown in Fig. 4, doxorubicin treatment inhibited PKC α expression in the membrane fractions whereas the PKC β I isoform remained the same at both fractions in the HCT15 cells. Furthermore, the transient expression of PKC α in the cytosol suggests the translocation from cytosol to the membrane. In contrast, PKC α in the HCT15/DOX cells seemed to predominate in regards to doxorubicin treatment in the cytosol and membrane fractions (Fig. 4). The relative expression of PKC β I in the cytosolic and membrane fractions was not significantly altered by doxorubicin. This data suggests that PKC β I has no major role, whereas PKC α mediates, at least in part, multidrug resistance.

PKC α activity blocks ROS production in HCT15 and HCT15/DOX cells

Increasing evidence suggests that the mitochondria are the main site of ROS generation and high levels of ROS may mediate the induction of apoptosis (Fleury et al., 2002). In order to investigate if doxorubicin killed cancer cells through ROS generation, we evaluated the redox status in the HCT15 and HCT15/DOX cells using oxidation-sensitive fluorescent dye, DCFHDA. Doxorubicin treatment increased oxidant-induced 2',7'-dichlorofluorescein fluorescence, only in the parental cells, but with treatment of the cells with Go6976, a rapid accumulation of fluorescence was observed in both cell types (Fig. 5). The occurrence of ROS generation were displayed by the mitochondrial-specific redox-sensitive fluorophore dihydrorhodamine (DHR) 123 (Fig. 5) when compared to both types of cell lines, either with or without inhibiting PKC α .

Go6976 reversed the PKC α inhibition of doxorubicin - induced apoptosis in HCT15/DOX cells

Much compelling evidence has illuminated the anti-apoptotic roles of Bcl $_2$ and Bad proteins. In fact, PKC α can phosphorylate Bcl $_2$ at Ser 70 , a site which is critical for its anti-apoptotic effects. In order to determine the relative contribution of the PKC α isoform to decreased activity of doxorubicin in HCT15/DOX cells, we treated both cell types with classical PKC inhibitor, Go6976. After treatment, a significant decrease was observed in Bcl $_2$




Fig. 5. Inhibition of PKC α by Go6976 induces ROS generation in doxorubicin treated HCT15/DOX cells. Both cell types were challenged with Go6976 and then incubated with doxorubicin. ROS generation was exhibited by DCFHDA. Dihydrorhodamine 123 results indicated mitochondrial ROS generation.




Fig. 6. Go6976 reversed the PKC α inhibition of doxorubicin-induced apoptosis in HCT15/DOX cells. Cells were pretreated, either with or without 5 μ M Go6976, for 2 h and then with 5 μ M of doxorubicin. (A) Lysates of HCT15/DOX cells were used for Western blotting to detect PARP, Bcl $_2$ and p-Bad, and actin was used as a control. (B) The cells were harvested and stained with propidium iodide and their DNA content was analyzed by flow cytometry.

and p-Bad in the parental fraction as well as in HCT15/DOX cells lines (Fig. 6A). Upon the inhibition of PKC α with inhibitor Go6976, doxorubicin-induced apoptosis increased 45% in the parental cells and 17% in the HCT15/DOX cells relative to the untreated cells (Fig. 6B). These results are in agreement with previously published work (La Porta et al., 1998) that indicated PKC α plays a pivotal role in drug resistance and mediation of anti-apoptotic events in doxorubicin treated cells. Furthermore, inhibition of PKC α led to increased cleavage of PARP in both types of cells (Fig. 6A).

Knockdown of PKC α by siRNA induced apoptosis in HCT15/DOX cells

To gain further insight into the anti-apoptotic effects of PKC α ,

targeted siRNA was used against PKC α mRNA in order to knockdown PKC α in the resistant cell line. The result was analyzed by FACS and Western blot. As shown in Fig. 7A, apoptotic cell death increased in transfected PKC α siRNA cells as compared to control siRNA transfected cells. Additionally, knockdown of PKC α down-regulated Bcl $_2$ and p-Bad as well as enhanced PARP cleavage and induced apoptosis in the resistant cell line (Fig. 7B). We also examined the dose dependent effects of siRNA by exposing HCT15/DOX cells to 5 μ M doxorubicin along with different concentrations of siRNA (Fig. 7B). These results clearly indicated that depletion of PKC α was associated with doxorubicin-induced apoptosis in the resistant cell line.

Conventional evidence has indicated that doxorubicin-indu-




Fig. 7. RNA interference-mediated inhibition of PKC α increased doxorubicin-induced apoptosis. (A) FACS analysis of the cell cycle. Cells were pre-incubated with either siRNA, scrambled siRNA, or vehicle alone for 24 h followed by stimulation with doxorubicin. Cell cycle analysis was done by FACS. (B) PKC α protein levels of siRNA-PKC α cells. HCT15/DOX cells were transfected with various concentrations of siRNA and then incubated with either doxorubicin or carrier for 24 h. Total cell lysates were prepared and used for Western blotting. (C) PKC α inhibition induces ROS generation. HCT15/DOX cells were transfected with siRNA and then incubated with doxorubicin or carrier for 24 h. PKC α siRNA diminished the protective effect of PKC α and induced ROS generation as shown through DCFHDA fluorescence and translocation of DHR.

ced apoptosis is a complex multifunctional process in which ROS generation production plays a vital role in the induction of apoptosis (Konorev et al., 1999). In order to investigate whether or not doxorubicin-induced apoptosis in HCT15/DOX cells is associated with the generation of cellular oxidative stress, the levels of intracellular peroxides were evaluated with DCFHDA. It was assumed that PKC α reversed doxorubicin-induced apoptosis in resistant cells through decrease in DCF-detectable ROS. As shown in Fig. 7C, transfection of cells with PKC α siRNA, followed by doxorubicin treatment, diminished the protective effect of PKC α and induced ROS generation. Therefore, it is likely that ROS production is the agent of doxorubicin-induced apoptosis in HCT15/DOX cells. Additionally, the HCT15/DOX cells transfected with scramble siRNA mediated or induced the same effect as the control cells without transfection (Fig. 7C). These results suggested that PKC α provides protection from the apoptotic actions of doxorubicin and enhanced MDR by decreasing the steady-state level of intracellular oxidants.

DISCUSSION

The mechanisms of MDR relative to the treatment failure of chemotherapeutic drugs has been studied extensively for more than two decades, but the exact molecular mechanism mediating the chemotherapy resistance and the role of drug resistance genes involved in the transport of anticancer agents remains unclear. There are a number of different mechanisms by which cancer cells develop MDR (Takara et al., 2006). The most often reported mechanism with known clinical significance is activation of ATP-dependent efflux pumps resulting in reduced intracellular drug concentrations, in which P-gp is the best known efflux pump gene alteration. MDR can also be caused by proteins involved in the control of apoptosis, such as Bcl₂ (Han et al., 2002). The current study postulated a role for PKC in the

regulation of the MDR phenotype. Other studies have shown that PKC produces prominent activity in MDR cell lines and that PKC inhibitors could be of therapeutic value in reversing MDR and inhibiting P-gp phosphorylation (Fine et al., 1996; O'Brian, 2001). It has been shown that P-gp is phosphorylated by PKC and that this phosphorylation entirely modulates P-gp efflux function (Ling, 1997). At present, the PKC family consists of at least 12 isoforms whose expression and regulation varies between cell types. In particular, the over-expression and activity of the PKC α isoform is closely associated with the development of the MDR phenotype, but in some cancers the expression patterns of PKC β I, PKC γ and PKC η mediate MDR. Similarly, the PKC θ isoform is also activated by diverse stimuli and participates in MDR, but this contribution is minor and occurs only in concert with PKC α (Aquino et al., 1999; Gill et al., 2001). In addition, the pro- and anti-apoptotic roles of PKCs isoforms has been well documented (Leszczynski, 1995; Lucas and Sánchez-Margalet, 1995). Previously, it has been shown that over-expression of PKC α resulted in increased mitochondrial PKC α localization and Bcl₂ phosphorylation, which are required for their anti-apoptotic function (Ruvolo et al., 1998). Therefore, it is pertinent to understand the mechanism by which PKC α is activated in the apoptotic process.

HCT15/DOX is an established doxorubicin resistant cell line selected by the stepwise exposure of parental HCT15 cells to increasing concentrations of doxorubicin. Studies have shown that HCT15/DOX cells possess characteristics of classical MDR including over-expression of P-gp as well as high resistance to the cytotoxicity of doxorubicin (Choi et al., 1996). The above mentioned results imply that doxorubicin is hindered by some protein residues which may alter the resistant cell efflux rate. The over-expression of PKC α has been radically elevated yet no significant difference was observed in the expression of PKC β I, PKC β II and PKC γ in colon cancer cells (Gravitt et al., 1994). Furthermore, the expression patterns of PKC α and

PKC β 1 in the resistant cell line remained constant upon the treatment of doxorubicin, but in HCT15 cells the membrane translocation of PKC α was reduced. After treatment with doxorubicin, similar expression levels of PKC β 1 in the cytosol and membrane revealed that it shared no major contribution to the effect, suggesting that only PKC α has a role in MDR. This finding means that PKC α is capable of enhancing *mdr* gene expression and inducing MDR because the total protein expression of PKC α was also increased in the HCT15/DOX cells (Kuranami et al., 1995). The over-expression of PKC α in HCT15/DOX cells, not only induced MDR, but also reversed apoptosis through the down-regulation of ROS generation in doxorubicin treated cells. Doxorubicin has been reported to induce apoptosis in various cancer cell lines through ROS generation (Gouazé et al., 2001), but the exact role of ROS in the resistance of chemotherapy remains to be elucidated. Early evidence has shown that ROS production is responsible for G α q-mediated activation of phospholipase C (PLC)- β , which is followed by the activation of a complex network of signaling proteins governing cell-cycle transition and redox balance (Bae et al., 2011). Although PKC stimulated the induction of ROS generation, this effect seemed to be cell and tissue dependent (Inoguchi et al., 2003).

The current study proved that PKC α was over-expressed in resistant cells as compared to parental cells and attenuated doxorubicin induced oxidative-stress apoptosis. In fact, PKC α phosphorylated Bcl $_2$, p-Bad, decreased PARP cleavage and prevented apoptosis in doxorubicin treated resistant cell lines. But incubation with PKC α inhibitor Go6976, doxorubicin induced apoptosis in the parental fraction and the HCT15/DOX cells line. Inhibition of PKC α activity has been shown to efficiently restore the drug-sensitive phenotype of cancer cells (Wang et al., 2010). To gain further insight into the possible effects of PKC α on MDR in human colon cancer cells, we transfected the cells with PKC α -siRNA because it has been shown that transfection of exogenous PKC α into cells over-expressing the MDR gene conferred an increased MDR phenotype into these cells (Yu et al., 1991). The knockdown of PKC α , through a PKC α -siRNA dose, dependently down-regulated Bcl $_2$ and p-Bad, enhanced PARP cleavage and induced apoptosis in the resistant cell line. These results were in agreement with a previously published report that suggested that transfection of PKC α antisense into cells reversed the MDR phenotype (Osborn et al., 1999).

In conclusion, the results presented here support the notion that PKC α activation phosphorylates P-gp and over-expression of MDR, and ultimately leads to drug resistance. PKC α reversed doxorubicin induced apoptosis through the scavenging of ROS and expression of anti-apoptotic proteins such as Bcl $_2$ as well as inhibited PARP cleavage. Using PKC α inhibitor Go6976 and PKC α siRNA knockdown revealed that PKC α provides a major contribution to drug resistance. We hope that the outcome of this study will make further insights into doxorubicin-induced apoptosis as well as therapeutic strategies directed against PKC α as a tool for restructuring the resistance of tumors against cytotoxic agents.

ACKNOWLEDGMENTS

This work was supported by the Nuclear Research and Development Program of the National Research Foundation of Korea funded by the Korean Ministry of Education, Science and Technology (grant code: 2011-0006331).

REFERENCES

- Altenberg, G.A., Vanoye, C.G., Han, E.S., Deitmer, J.W., and Reuss, L. (1994). Relationships between rhodamine 123 transport, cell volume, and ion-channel function of P-glycoprotein. *J. Biol. Chem.* *269*, 7145-7149.
- Aquino, A., Warren, B.S., Omichinski, J., Hartman, K.D., and Glazer, R.I. (1990). Protein kinase C-gamma is present in doxorubicin resistant HL-60 leukemia cells. *Biochem. Biophys. Res. Commun.* *166*, 723-728.
- Bae, Y.S., Oh, H., Rhee, S.G., and Yoo, Y.D. (2011). Regulation of reactive oxygen species generation in cell signaling. *Mol. Cells* *32*, 491-509.
- Bose, R., Verheij, M., Haimovitz-Friedman, A., Scotto, A., Fuks, Z., and Kolesnick, R.N. (1995). Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell* *82*, 405-414.
- Budworth, J., Gant, T.W., and Gescher, A. (1997). Co-ordinate loss of protein kinase C and multidrug resistance gene expression in revertant MCF-7/ADR breast carcinoma cells. *Br. J. Cancer* *75*, 1330-1335.
- Clemens, M.J., Trayner, I., and Menaya, J. (1992). The role of protein kinase C isoenzymes in the regulation of cell proliferation and differentiation. *J. Cell Sci.* *103*, 881-888.
- Choi, S.U., Kim, N.Y., Choi, E.J., Kim, K.H., and Lee, C.O. (1996). Establishment of doxorubicin-resistant subline derived from HCT15 human colorectal cancer cells. *Arch. Pharm. Res.* *19*, 342-347.
- Dolfini, E., Dasdial, T., Arancia, G., Molinari, A., Calcabrin, A., Scheper, R.J., Flens, M.J., Gariboldi, M.B., and Monti, E. (1997). Characterization of a clonal human colon adenocarcinoma line intrinsically resistant to doxorubicin. *Br. J. Cancer* *76*, 67-76.
- Doroshov, J.H., Locker, G.Y., and Myers, C.E. (1980). Enzymatic defenses of the mouse heart against reactive oxygen metabolites: alter alterations produced by doxorubicin. *J. Clin. Invest.* *65*, 128-135.
- Endicott, J.A., and Ling, V. (1989). The biochemistry of P-glycoprotein mediated multidrug resistance. *Ann. Rev. Biochem.* *58*, 137-171.
- Fine, R.L., Chambers, T.C., and Sachs, C.W. (1996). P-glycoprotein, multidrug resistance and protein kinase C. *Stem Cells.* *14*, 47-55.
- Fleury, C., Mignotte, B., and Vayssiere, J.L. (2002). Mitochondrial reactive oxygen species in cell death signaling. *Biochimie* *84*, 131-141.
- Gariboldi, M.B., Ravizza, R., Riganti, L., Meschini, S., Calcabrin, A., Marra, M., Arancia, G., Dolfini, E., and Monti, E. (2003). Molecular determinants of intrinsic resistance to doxorubicin in human cancer cell lines. *Int. J. Oncol.* *22*, 1057-1064.
- Gill, P.K., Gescher, A., and Gant, T.W. (2001). Regulation of MDR1 promoter activity in human breast carcinoma cells by protein kinase C isozymes alpha and theta. *Eur. J. Biochem.* *268*, 4151-4157.
- Gottesman, M.M., and Pastan, I. (1993). Biochemistry of multidrug resistance mediated by the multidrug transporter. *Annu. Rev. Biochem.* *62*, 385-427.
- Gottesman, M., Fojo, T., and Bates, S. (2002). Multidrug resistance in cancer: role of ATP-dependent transporters. *Nat. Rev. Cancer* *2*, 48-58.
- Gouazé, V., Mirault, M.E., Carpentier, S., Salvayre, R., Levade, T., and Andrieu-Abadie, N. (2001). Glutathione peroxidase-1 over-expression prevents ceramide production and partially inhibits apoptosis in doxorubicin-treated human breast carcinoma cells. *Mol. Pharmacol.* *60*, 488-496.
- Gravitt, K.R., Ward, N.E., Fan, D., Skibber, J.M., Levin, B., and O'Brian, C.A. (1994). Evidence that protein kinase C-alpha activation is a critical event in phorbol ester-induced multiple drug resistance in human colon cancer cells. *Biochem. Pharmacol.* *48*, 375-381.
- Han, Y., Han, Z.Y., Zhou, X.M., Shi, R., Zheng, Y., Shi, Y.Q., Miao, J.Y., Pan, B.R., and Fan, D.M. (2002). Expression and function of classical protein kinase C isoenzymes in gastric cancer cell line and its drug-resistant sublines. *World J. Gastroenterol.* *8*, 441-445.
- Inoguchi, T., Sonta, T., Tsubouchi, H., Etoh, T., Kakimoto, M., Sonoda, N., Sato, N., Sekiguchi, N., Kobayashi, K., Sumimoto, M.

- H., et al. (2003). Protein kinase C-dependent increase in reactive oxygen species (ROS) production in vascular tissues of diabetes: role of vascular NAD(P)H oxidase. *J. Am. Soc. Nephrol.* **14**, S227-232.
- Konorev, E.A., Kennedy, M.C., and Kalyanaraman, B. (1999). Cell-permeable superoxide dismutase and glutathione peroxidase mimetics afford superior protection against doxorubicin-induced cardiotoxicity: The role of reactive oxygen and nitrogen intermediates. *Arch. Biochem. Biophys.* **368**, 421-428.
- Kuranami, M., Powell, C.T., Hug, H., Zeng, Z., Cohen, A.M., and Guillem, J.G. (1995). Differential expression of protein kinase C isoforms in human colorectal cancers. *J. Surg. Res.* **58**, 233-239.
- La Porta, C.A., Dolfini, E., and Comolli, R. (1998). Inhibition of protein kinase C-alpha isoform enhances the P-glycoprotein expression and the survival of LoVo human colon adenocarcinoma cells to doxorubicin exposure. *Br. J. Cancer* **78**, 1283-1287.
- Leszczynski, D. (1995). Regulation of cell cycle and apoptosis by protein kinase C in rat myeloid leukemia cell line. *Oncol. Res.* **7**, 471-480.
- Limtrakul, P., Anuchapreeda, S., and Buddhasukh, D. (2004). Modulation of human multidrug-resistance MDR-1 gene by natural curcuminoids. *BMC Cancer* **4**, 13.
- Ling, V. (1997). Multidrug resistance: molecular mechanisms and clinical relevance. *Cancer Chemother. Pharmacol.* **40**, S3-S8.
- Lucas, M., and Sanchez-Margalet, V. (1995). Protein kinase C involvement in apoptosis. *Gen. Pharmacol.* **26**, 886-887.
- Ma, Y.Y., Wei, S.J., Lin, Y.C., Lung, J.C., Chang, T.C., Whang-Peng, J., Liu, J.M., Yan, D.M., Yang, W.K., and Shen, C.Y. (2000). PIK3CA as an oncogene in cervical cancer. *Oncogene* **19**, 2739-2744.
- Mellor, H., and Parker, P.J. (1998). The extended protein kinase C superfamily. *Biochem. J.* **332**, 281-292.
- Nakura, A., Higuchi, C., Yoshida, K., and Yoshikawa, H. (2011). PKC α suppresses osteoblastic differentiation. *Bone* **48**, 476-484.
- Nishizuka, Y., and Kikkawa, U. (2003). Early studies of protein kinase C: a historical perspective. *Methods Mol. Biol.* **233**, 9-18.
- O'Brian, C.A., Ward, N.E., Stewart, J.R., and Chu, F. (2001). Prospects for targeting protein kinase C isozymes in the therapy of drug resistant cancer--an evolving story. *Cancer Met. Rev.* **20**, 95-100.
- Osborn, M.T., Berry, A., Ruberu, M.S., Ning, B.T., Bell, L.M., and Chambers, T.C. (1999). Phorbol ester induced MDR1 expression in K562 cells occurs independently of mitogen-activated protein kinase signaling pathways. *Oncogene* **18**, 5756-5764.
- Ruvolo, P.P., Deng, X., Carr, B.K., and May, W.S. (1998). A functional role for mitochondrial protein kinase Calpha in Bcl2 phosphorylation and suppression of apoptosis. *J. Biol. Chem.* **273**, 25436-25442.
- Skladanowski, A., and Konopa, J. (1993). Doxorubicin and daunomycin induce programmed cell death (apoptosis) in tumour cells. *Biochem. Pharmacol.* **46**, 375-382.
- Takara, K., Sakaeda, T., and Okumura, K. (2006). An update on overcoming MDR1-mediated multidrug resistance in cancer chemotherapy. *Curr. Pharm.* **12**, 273-228.
- Tentori, L., Portarena, I., Bonmassar, E., and Graziani, G. (2001). Combined effects of adenovirus-mediated wild-type p53 transduction, temozolomide and poly (ADP-ribose) polymerase inhibitor in mismatch repair deficient and non-proliferating tumor cells. *Cell Death Differ.* **8**, 457-469.
- Wang, N., Li, Z., Tian, F., Feng, Y., Huang, J., Li, C., and Xie, F. (2010). PKC α inhibited apoptosis by decreasing the activity of JNK in MCF-7/ADR cells. *Exp. Toxicol. Pathol.* **64**, 459-464.
- Yu, G., Ahmad, S., Aquino, A., Fairchild, C.R., Tepel, J.B., Ohno, S., Suzuki, K., Tsuru, T., Cowan, K.H., and Glazer, R.I. (1991). Transfection with protein kinase C α confers increased multidrug resistance to MCF-7 cells expressing P-glycoprotein. *Cancer Commun.* **3**, 181-189.

