



Published in final edited form as:

Hypertension. 2014 February ; 63(2): 404–412. doi:10.1161/HYPERTENSIONAHA.113.02043.

Negative impact of β arrestin1 on post-myocardial infarction heart failure via cardiac and adrenal-dependent neurohormonal mechanisms

Ashley Bathgate-Siryk^{*}, Samalia Dabul^{*}, Krunal Pandya^{*}, Karlee Walklett^{*}, Giuseppe Rengo[^], Alessandro Cannavo[^], Claudio De Lucia[^], Daniela Liccardo[^], Erhe Gao[#], Dario Leosco[^], Walter J. Koch[#], and Anastasios Lymperopoulos^{*}

^{*}Laboratory for the Study of Neurohormonal Control of the Circulation, Department of Pharmaceutical Sciences, Nova Southeastern University College of Pharmacy, Fort Lauderdale, FL 33328, USA

[^]Department of Translational Medical Sciences, University “Federico II”, Naples, Italy

[#]Center for Translational Medicine, Temple University, Philadelphia, PA 19140, USA

Abstract

β arrestin (β arr)-1 and -2 (β arrs) are universal G protein-coupled receptor adapter proteins that negatively regulate cardiac β -adrenergic receptor (β AR) function via β AR desensitization & downregulation. In addition, they mediate G protein-independent β AR signaling, which might be beneficial, e.g. antiapoptotic, for the heart. However, the specific role(s) of each β arr isoform in cardiac β AR dysfunction, the molecular hallmark of chronic heart failure (HF), remain unknown. Furthermore, adrenal β arr1 exacerbates HF by chronically enhancing adrenal production, and hence circulating levels of aldosterone and catecholamines. Herein, we sought to delineate specific roles of β arr1 in post-myocardial infarction (MI) HF by testing the effects of β arr1 genetic deletion on normal and post-MI cardiac function and morphology. We studied β arr1 knockout (β arr1KO) mice alongside wild type (WT) controls under normal conditions and after surgical MI. Normal (sham-operated) β arr1KO's display enhanced β AR-dependent contractility and post-MI β arr1KO's enhanced overall cardiac function (and β AR-dependent contractility) compared to WT's. Post-MI β arr1KO's also show increased survival, and decreased cardiac infarct size, apoptosis, and adverse remodeling, as well as circulating catecholamines & aldosterone, compared to post-MI WT's. The underlying mechanisms are on one hand improved cardiac β AR signaling and function, as evidenced by increased β AR density and pro-contractile signaling, via reduced cardiac β AR desensitization due to cardiac β arr1 absence, and on the other hand decreased production leading to lower circulating levels of catecholamines & aldosterone due to adrenal β arr1 absence. Thus, β arr1, via both cardiac and adrenal effects, is detrimental for cardiac structure and function and significantly exacerbates post-MI HF.

Address correspondence to: Anastasios Lymperopoulos, PhD, FAHA, Assistant Professor, Department of Pharmaceutical Sciences, Nova Southeastern University College of Pharmacy, 3200 S. University Dr., HPD (Terry) Bldg/Room 1338, Fort Lauderdale, FL 33328, USA, Tel.: 954-262-1338, FAX: 954-262-2278, al806@nova.edu.

Disclosures

NONE.

The authors declare no relationships with industry or any other conflict of interest.

Keywords

β arrestin1; β -adrenergic receptor desensitization; knockout; aldosterone; catecholamine; post-myocardial infarction heart failure

Despite recent advances in its prevention and management, heart disease, such as hypertension and post-myocardial infarction (MI) heart failure (HF), remains the leading cause of death in the western world and new treatments are needed.¹ The molecular hallmark in chronic HF is cardiac β -adrenergic receptor (β AR) dysfunction due to increased receptor desensitization and downregulation.²⁻⁶ β ARs belong to the superfamily of G protein-coupled receptors (GPCRs), which, upon agonist binding, activate the G_s protein-adenylyl cyclase (AC)-cyclic adenosine monophosphate (cAMP)-protein kinase A (PKA) signaling pathway leading to increased inotropy and chronotropy in the heart.² In chronic HF though, there is a selective downregulation of cardiac β_1 ARs and significant functional desensitization (G protein uncoupling) of the remaining membrane β_1 ARs and β_2 ARs, which dramatically diminish cardiac adrenergic and inotropic reserves.³⁻⁶ These processes are mediated at the molecular level by the β arrestins (β arrestins), which exist in two distinct isoforms (β arr1 and -2, also known as arrestin-2 and -3, respectively) in mammals, and are abundantly expressed in the heart.⁷⁻⁹ β arrestins bind agonist-activated cardiac β ARs, terminating their G protein-mediated signaling and targeting them for internalization, following receptor phosphorylation by GPCR kinase (GRK)-2 (mainly), the main cardiac GRK that is significantly elevated in HF.^{2,8-16} Consistent with their crucial role in cardiac β AR desensitization, β arr1 knockout (KO) mice display enhanced cardiac contractile responses to isoproterenol (a standard β AR full agonist) stimulation in vivo.¹⁷

In addition, the β AR-bound β arrestins have been shown to elicit a second wave of G protein-independent signals, some of which might be beneficial for the failing heart, e.g. extracellular signal-regulated kinase (ERK) activation and epidermal growth factor receptor (EGFR) transactivation, both of which can promote cardiac cell survival, proliferation, etc.¹⁸⁻²¹ However, the actions of the two β arrestins in vivo are rarely complementary; in fact, they might even oppose each other's effects in certain tissues/organs, including in the cardiovascular system (e.g. vascular smooth muscle cells and cardiomyocytes).^{22,23} With regard to the heart, the exact role of each β arr in cardiac β AR desensitization or G protein-independent signaling is not presently known. The physiological hallmark of chronic HF is the sustained hyperactivity of several neurohormonal systems, in particular the sympathetic nervous (SNS) and the renin-angiotensin-aldosterone (RAAS) systems, accompanied by elevated circulating levels of the catecholamines (CAs) norepinephrine (NE) and epinephrine (Epi), and of aldosterone, respectively.^{9,14,15} All these circulating hormones exert detrimental effects on the failing heart, leading both to diminished function and to accelerated and aggravated adverse remodeling in chronic HF.^{24,25} The main source of these circulating hormones is the adrenal gland and we have shown in previous studies that β arr1, in particular, is a crucial mediator of adrenocortical aldosterone production,^{26,27} as well as of adrenomedullary CA production (acting in concert with adrenal GRK2),²⁸⁻³⁰ both normally and in post-MI HF in vivo.

Given that the specific roles of cardiac β arr1 in β AR (dys)function and signaling in HF have never been investigated and also the crucial role this protein plays in the adrenals in elevating the neurohormonal burden of the failing heart, we sought to investigate, in the present study, the effects of its genetic deletion on cardiac function, both normally and in post-MI HF. In order to study the (patho)physiological role(s) specifically of this β arr isoform in vivo, we took advantage of the available β arr1KO mouse model¹⁷ and studied these mice alongside age-matched wild type controls (of note, the double β arr1/2 KO mice

are embryonic-lethal).⁹ We found that the absence of β arr1 from the heart and the adrenals leads to significantly elevated cardiac function in post-MI HF, accompanied by improved adverse remodeling, reduced neurohormonal activation and better cardiac β AR signaling and function.

Materials and Methods

Experimental animals and surgical procedures

The animals in this study were handled according to animal welfare regulations and protocols approved by the authors' Institutional Review Boards. An expanded "Materials and Methods" section is available in the Supplementary Material online, which includes detailed methods on the following: animals and surgical procedures, echocardiography and hemodynamics, ELISA measurements of hormones and cytokines, infarct size measurements, TUNEL and Masson-Trichrome staining, real-time PCR, β AR density, cAMP and SERCA activity measurements, and western blotting.

Statistical analyses

Data are generally expressed as mean \pm SEM. Unpaired 2-tailed Student's *t* test and one- or two-way ANOVA with Bonferroni test were generally performed for statistical comparisons, unless otherwise indicated. For most 3-group statistical comparisons Dunnett's test using SAS version 8.2 software was used, as well. For all tests, a *p* value of <0.05 was generally considered to be significant.

Results

Cardiac function of normal and post-MI β arr1KO mice

In order to investigate the impact on cardiac function of the genetic deletion exclusively of β arr1, we utilized the available global β arr1KO mouse model (Figure S1). These mice breed normally, without any gross abnormalities and present no overt cardiovascular or other phenotype.¹⁷ To induce HF, three-month-old male mice underwent surgical MI and were studied alongside age-matched male WT mice. We first examined the cardiac function parameters of these mice, both in sham and post-MI groups. Echocardiography revealed that, although similar between the sham groups (a finding consistent with the results of a previous study on these mice¹⁷), (basal) ejection fraction was severely diminished in WT mice at 4 weeks post-MI, as expected, but significantly higher in the post-MI β arr1KO's at the same time point (Figure 1A & Table S1). Consistent with these findings, upon *in vivo* catheterization for hemodynamic measurements, post-MI β arr1KO mice also show significantly enhanced cardiac contractility, both basally and in response to isoproterenol stimulation, compared to post-MI WT mice (Figure 1B & Table S1), even though left ventricular end-systolic and end-diastolic pressures are similar between the two post-MI groups upon maximal isoproterenol challenge (Table S1). Notably, isoproterenol-induced contractility is elevated also in sham (normal) β arr1KO's compared to sham WT's (Figure 1B & Table S1), which is again consistent with the results of a previous study on these mice.¹⁷ Taken together, these findings strongly indicate that cardiac β arr1 is a major negative regulator of β AR-dependent contractility *in vivo* and that its absence leads to significant attenuation of cardiac dysfunction post-MI.

Survival and neurohormonal status of post-MI β arr1KO mice

Next, we sought to further examine the phenotype of the post-MI β arr1KO mice. Kaplan-Meier survival curves indicated a markedly lower (overall) mortality of the post-MI β arr1KO's compared to post-MI WT controls ($p=0.012$, Figure 2A). This was accompanied

by significantly reduced elevations in the plasma circulating levels of the CAs NE and Epi in post-MI β arr1KO's compared to post-MI WT's (Figure 2B), indicative of reduced overall sympathetic activation in post-MI HF β arr1KO mice. As for the other major cardiotoxic hormone, aldosterone, post-MI β arr1KO's, remarkably, failed to exhibit any hyperaldosteronism (i.e. elevation of circulating aldosterone levels) whatsoever, in marked contrast to post-MI WT's, which, as expected, display severe hyperaldosteronism (Figure 2C). This finding strongly corroborates the essential role of adrenal β arr1 in post-MI HF-associated hyperaldosteronism we have previously reported.^{26,27} Finally, and consistent with these effects on circulating levels of catecholamines and aldosterone, post-MI β arr1KO's also display significantly lower mean arterial blood pressure (125 ± 4 mm Hg) than post-MI WT's (141 ± 5 mm Hg) at 4 weeks post-MI ($p < 0.05$, $n = 5$).

Cardiac apoptosis, dilatation and infarct size in post-MI β arr1KO mice

The increased survival of the post-MI β arr1KO's prompted us to investigate the cardiac apoptosis in these mice. Cardiac cellular apoptosis, as measured with the TUNEL assay in the hearts of the post-MI mice, was found significantly reduced in β arr1KO's compared to WT mice at 24 hrs post-MI, which is consistent with the Kaplan-Meier survival results (Figure 3A). In addition, cardiac dilatation was also decreased in the β arr1KO's compared to WT controls at 4 weeks post-MI (Figure 3B & Table S1), and, importantly, infarct size was also significantly reduced in the β arr1KO hearts compared to control WT hearts at the same post-MI time point (4 weeks) (Figure 3C & 3D). To exclude the possibility that differences in the extent of the initial injury inflicted by our surgical MI method upon the two lines might have been responsible for this result, infarct size at 24 hrs post-MI was also measured and found indistinguishable between the two groups (Figure S2A & S2B). Taken together, these results indicate that post-MI β arr1KO hearts have decreased apoptosis, dilatation, and infarct size, which might underlie their favorable overall survival phenotype.

Cardiac inflammation, fibrosis and adverse remodeling markers in post-MI β arr1KO mice

Next, we examined markers of post-MI cardiac inflammation and adverse remodeling in the two animal groups. Levels of all three major pro-inflammatory cytokines, tumor necrosis factor (TNF)- α (Figure 4A), interleukin (IL)-6 (Figure 4B), and IL-1 β (Figure 4C), were found significantly decreased in post-MI β arr1KO hearts compared to control WT hearts at 4 weeks post-MI, indicating decreased overall cardiac inflammation in the post-MI β arr1KO mice. Moreover, cardiac fibrosis, as measured with Masson-Trichrome staining, was also markedly decreased in the post-MI β arr1KO mice compared to post-MI WT controls (Figure 4D & 4E). Finally, mRNA expression of all major adverse remodeling-associated biomarkers, i.e. collagen synthesis (Figure 4F), B-type natriuretic peptide (BNP) (Figure 4G), atrial natriuretic factor (ANF) (Figure S3A), and transforming growth factor (TGF)- β (Figure S3B), was significantly reduced in the post-MI β arr1KO hearts compared to post-MI WT hearts, strongly suggesting that adverse remodeling is markedly attenuated in post-MI β arr1KO hearts.

Cardiac β AR density, signaling and function in post-MI β arr1KO mice

In an effort to dissect the molecular mechanisms underlying the dramatic effect of the absence of β arr1 on cardiac contractile function, we also investigated several aspects of cardiac β AR signaling and function in the two post-MI groups. Consistent with the phenotypic and functional data, total β AR density in the post-MI β arr1KO hearts was significantly higher compared to post-MI WT hearts, which display marked β AR downregulation post-MI, as expected (Figure 5A). This attenuated β AR downregulation in post-MI β arr1KO hearts was accompanied by preserved total cAMP levels, again in sharp contrast with the WT hearts whose cAMP content is dramatically diminished post-MI (Figure 5B). Given that cAMP is the most critical mediator of cardiac β AR pro-contractile

signaling,^{2,15} this result strongly indicates that the absence of cardiac β arr1 is sufficient to restore β AR-dependent contractility in post-MI HF. Another important component of the cardiac contractile machinery regulated by β ARs is sarco(endo)plasmic reticulum Ca^{2+} -ATPase (SERCA)-2a.³¹ SERCA activity was also found significantly increased in post-MI β arr1KO hearts compared to post-MI WT hearts (Figure 5C), thus providing another line of evidence for the increased contractile function of post-MI β arr1KO hearts. Of note though, cardiac SERCA activity of β arr1KO's appears elevated also in the normal (sham) groups (Figure 5C), indicating that cardiac β arr1 might affect the cardiac contractile machinery even under normal conditions (no HF). Finally, since cardiac β arrs, and in particular β arr2, have been shown to mediate EGFR transactivation from the β_1 AR upon the latter's phosphorylation by GRK5, an effect that is considered anti-apoptotic in the heart,¹⁸ we also examined levels of EGFR transactivation in the post-MI hearts of β arr1KO's and WT's after acute stimulation with isoproterenol in the presence of ICI-118,551 (a β_2 AR-selective antagonist) in vivo (in order to activate the β_1 ARs only). Western blotting for phospho-Tyr845-EGFR (active EGFR) in cardiac extracts from these mice revealed significantly elevated EGFR transactivation in β arr1KO hearts compared to WT hearts at 24 hrs post-MI (and also in sham hearts) (Figure 5D & 5E), suggesting that cardiac β arr1 inhibits (rather than promotes) this β_1 AR-induced EGFR transactivation in the heart. Finally, in an effort to better dissect the relative contributions of cardiac vs. adrenal β arr1 on the observed phenotypes of the mice of the study, we overexpressed the known β arr1 inhibitor mini-gene β arr1ct²⁷ specifically in the hearts of post-MI WT mice, in order to inhibit cardiac (only) β arr1 in vivo, and studied these hearts alongside control post-MI WT hearts receiving adenovirus encoding green fluorescent protein (GFP) at the time of MI. As shown in Figure S4, cardiac β arr1 inhibition with β arr1ct led to significant improvements in cardiac β AR density (Figure S4A) and cAMP accumulation (Figure S4B) at 4 weeks post-MI, indicating improved cardiac adrenergic and inotropic reserves, as expected; however, no significant changes in circulating catecholamines of the post-MI WT mice were observed with cardiac β arr1ct at 4 weeks post-MI (Figure S4C), indicating that catecholamine production/secretion is mediated by adrenal (rather than cardiac) β arr1.

Discussion

β arr1 (and to a lesser extent β arr2) are abundantly expressed throughout the cardiovascular system⁷⁻⁹ and regulate the vast majority of cardiovascular GPCRs, including cardiac β ARs, cardiac and adrenocortical angiotensin II (AngII) type 1 receptors (AT_1Rs), adrenal and central SNS $\alpha_2\text{ARs}$, etc.^{9,32,33} Their importance in cardiovascular (and other systems') biology is evidenced by the fact that the double β arr1/2 KO mouse is embryonic-lethal,^{32,33} Their actions on the heart have been attributed to both their functions as receptor desensitizers/internalizers and as G protein-independent signal transducers.^{9,32} However, and especially for their latter function, the exact in vivo roles of each cardiac β arr isoform in GPCR-dependent signal transduction are currently unknown. In the present study, we were able to delineate the cardiovascular phenotype of β arr1KO mice in post-MI HF, along with its associated underlying molecular signaling mechanisms. Specifically, we found that the absence of β arr1 from the heart leads to significantly better overall cardiac function in post-MI HF, dramatically increased cardiac β AR-dependent cardiac function both physiologically and in post-MI HF, and significantly attenuated adverse remodeling, apoptosis, inflammation and infarct size, leading to improved overall survival post-MI. At the same time, absence of β arr1 from the adrenal gland leads to a dramatically improved neurohormonal profile of the failing post-MI heart, with reduced circulating catecholamine and aldosterone levels. Finally, cardiac β AR signaling and function are dramatically elevated in the post-MI β arr1KO mice, translating into markedly improved cardiac adrenergic and inotropic reserves.

Thus, β arr1 appears to be the single most important negative regulator of β AR-dependent pro-contractile signaling and function, via the classical processes of desensitization and downregulation of the cardiomyocyte β AR content, and its absence results in dramatically elevated β AR-dependent contractility, both normally and after MI, and in almost restored functional β AR number and cAMP-mediated signaling post-MI, i.e. in dramatically improved β -adrenergic and inotropic reserves of the failing heart. Consistent with this, cardiac SERCA activity is also markedly elevated in post-MI β arr1KO's compared to control WT mice. Of note however, SERCA activity is also significantly elevated in normal (sham) β arr1KO mice, suggesting that the effects of cardiac β arr1 on SERCA activity extend beyond (merely) β AR desensitization. Indeed, current investigations suggest that cardiac β arr2 directly interacts with SERCA2a, promoting its SUMOylation and increasing its activity,³⁴ and that this is actually opposed by β arr1 (Lymeropoulos, A., unpublished data). On the other hand, β arr2 appears unable to compensate for the absence of β arr1 for any of these effects in vivo (and it is not compensatively upregulated, either; data not shown & Ref. 17). Taken together, these phenotypic findings strongly suggest that the β arr1 isoform is the salient β arr responsible for β AR desensitization and downregulation in the heart (Figure 6), processes whose dramatic elevation constitutes a molecular hallmark of chronic HF^{2,3,5}. Therefore, prevention of the interaction of β arr1 with cardiac β ARs, which is achieved (indirectly) via cardiac GRK2 inhibition with β ARKct for instance,^{13,15,35} might be beneficial as a positive inotropic therapy for HF (Figure 6).

It is now well known and documented that β arrs, in addition to the original biological role ascribed to them as terminators of G protein-dependent signaling by GPCRs, can also serve as signal transducers in their own right independently of G proteins.^{32,33} Indeed, specifically in the heart, β arrs have been shown to mediate beneficial, anti-apoptotic and proliferation-promoting signals from the β_1 AR (upon its phosphorylation by GRK5 and not GRK2 though) and from the mechanical stretch-activated AT₁R via EGFR transactivation and subsequent ERK phosphorylation/activation.^{18,36} In addition, they might even mediate pro-contractile signaling from the AT₁R in cardiac myocytes, an effect ascribed to β arr2 and presumably inhibited by GRK2 and β arr1.²³ We found that β_1 AR-induced EGFR transactivation is significantly elevated in β arr1KO hearts in vivo, both in normal (sham) conditions and (even more so) in post-MI HF, consistent with the apoptotic phenotype of the β arr1KO hearts in vivo. This strongly suggests that cardiac β arr2 normally mediates this beneficial, anti-apoptotic β_1 AR-dependent EGFR transactivation, whereas cardiac β arr1 opposes this signaling mechanism (Figure 6). Of course, studies in β arr2KO mice are warranted to further validate this scenario. Another important aspect of HF pathophysiology, in which the two β arr isoforms might exert differential or even opposing effects, is peri-infarct area neoangiogenesis;³⁷ in fact, given the difference observed in infarct size at 4 weeks post-MI (Figure 3C–D), this is quite likely and certainly warrants further investigation in the future.

Finally, the neurohormonal profile of the post-MI β arr1KO mice confirms the essential roles of adrenal β arr1 in: a) chronically elevated CA secretion in post-MI HF (via chronic desensitization/downregulation of the sympatho-inhibitory α_2 ARs in the chromaffin cells of the adrenal medulla) (Figure 6), and b) post-MI HF-associated hyperaldosteronism (via increased AngII-dependent aldosterone synthesis and secretion in adrenocortical zona glomerulosa cells) (Figure 6), which we have uncovered over the past several years.^{26–30}

Admittedly, the biggest limitation of the present study is the use of a global knockout model which precludes any safe conclusions on the relative contributions of adrenal vs. cardiac β arr1 on the observed phenotypic effects of the post-MI β arr1KO mice. Only development of tissue-specific KO mice can provide definitive answers to this important question. However, our data on post-MI WT mice having cardiac-specific blockade of β arr1 courtesy

of cardiac-specific β arr1ct overexpression strongly suggest that cardiac β arr1 absence is responsible for the observed improvements in the adrenergic and inotropic reserves of the post-MI β arr1KO mice, while their vastly improved neurohormonal profile (i.e. lower circulating catecholamine levels and complete absence of hyperaldosteronism) is (mainly) due to β arr1 absence from the adrenal glands. Given the extensive cross-talk between the two organs (heart-adrenals) and that aldosterone and catecholamines can be produced also in the heart per se however,^{38,39} drawing of these conclusions warrants extreme caution. Nevertheless, all of the above findings are clinically important: we report for the first time that β arr1 is the β arr isoform mainly (if not exclusively) responsible for cardiac β AR desensitization and downregulation in vivo, a hallmark abnormality in chronic HF. In addition, β arr1 promotes elevation of the neurohormonal burden of the failing heart through its actions in the adrenal gland, and seems to oppose several beneficial signaling effects of its counterpart, β arr2, in the myocardium, such as EGFR transactivation and ERK signaling, which lead to inhibition of apoptosis and inflammation and to promotion of cardiac survival. Thus, β arr1 removal from both the heart and the adrenals results in significantly improved cardiac function, structural remodeling, β -adrenergic and inotropic reserves, reduced neurohormonal burden and increased cardiac and overall survival in post-MI HF. These findings strongly suggest that inhibition of β arr1 activity towards β ARs in the heart and β arr1 blockade in the adrenals, either directly (with a pharmacological β arr1-specific inhibitor or via siRNA knockdown) or indirectly (e.g. via blockade of GRK2, which promotes association of β arr1 with cardiac β ARs) might be a viable therapeutic strategy for chronic post-MI HF treatment, with the potential also of complementing or enhancing the benefits of β -blocker therapy, which is hampered by the adverse effect of negative inotropy.

Perspectives

The present study reports for the first time that cardiac β arr1 is the β arr isoform responsible for cardiac β AR desensitization and downregulation in vivo, a hallmark molecular abnormality in chronic HF. In addition, adrenal β arr1 promotes elevation of the neurohormonal burden of the failing heart by mediating catecholamine secretion and aldosterone synthesis and secretion in the adrenal medulla and cortex, respectively. Finally, β arr1 seems to oppose several beneficial signaling effects of its counterpart, β arr2, in the myocardium, such as EGFR transactivation and ERK signaling, which lead to inhibition of apoptosis and inflammation and to promotion of cardiac survival. Thus, β arr1 removal from both the heart and the adrenals results in significantly improved cardiac function, halted adverse remodeling, elevated β -adrenergic and inotropic reserves, reduced neurohormonal burden and increased cardiac and overall survival in post-MI HF. These findings strongly suggest that inhibition of β arr1 activity towards β ARs in the heart and β arr1 blockade in the adrenals, either directly (with a pharmacological β arr1-specific inhibitor or via siRNA knockdown) or indirectly (e.g. via blockade of GRK2, which promotes association of β arr1 with receptors) might be a viable therapeutic strategy for chronic post-MI HF treatment.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. Yuhui Wen (Univ. of Miami, Miller School of Medicine) for excellent technical assistance with the in vivo experiments.

Sources of Funding

This work was supported in part by a Scientist Development Grant from the American Heart Association (AHA #09SDG2010138, National Center) and a Nova Southeastern University's Health Professions Division (HPD) Research Grant (to A.L.), and NIH grants R37 HL061690, R01 HL085503, P01 HL075443 (Project 2) and P01 HL091799 (to W.J.K.).

Non-standard abbreviations and acronyms

MI	myocardial infarction
HF	heart failure
AR	adrenergic receptor
GPCR	G protein-coupled receptor
AC	adenylyl cyclase
PKA	protein kinase A
GRK	GPCR kinase
βarr	beta-arrestin
KO	knockout
EGFR	epidermal growth factor receptor
ERK	extracellular signal-regulated kinase
SNS	sympathetic nervous system
CA	catecholamine
NE	norepinephrine
Epi	epinephrine
WT	wild type
TUNEL	terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling
TNFα	tumor necrosis factor alpha
IL	interleukin
cAMP	3'-5' adenosine monophosphate (cyclic adenosine monophosphate)
SERCA	sarco(endo)plasmic reticulum Ca ²⁺ -ATPase
AngII	angiotensin II
AT₁R	AngII type 1 receptor
siRNA	small interfering RNA

References

1. Tamargo J, López-Sendón J. Novel therapeutic targets for the treatment of heart failure. *Nat Rev Drug Disc.* 2011; 10:536–555.
2. Rockman HA, Koch WJ, Lefkowitz RJ. Seven-transmembrane-spanning receptors and heart function. *Nature.* 2002; 415:206–212. [PubMed: 11805844]
3. Brodde OE. Beta-adrenoceptors in cardiac disease. *Pharmacol Ther.* 1993; 60:405–430. [PubMed: 7915424]
4. Bristow MR, Ginsburg R, Umans V, Fowler M, Minobe W, Rasmussen R, Zera P, Menlove R, Shah P, Jamieson S. Beta 1- and beta 2-adrenergic-receptor subpopulations in nonfailing and failing human ventricular myocardium: coupling of both receptor subtypes to muscle contraction and

- selective beta 1-receptor down-regulation in heart failure. *Circ Res.* 1986; 59:297–309. [PubMed: 2876788]
5. Bristow MR, Ginsburg R, Minobe W, Cubicciotti RS, Sageman WS, Lurie K, Billingham ME, Harrison DC, Stinson EB. Decreased catecholamine sensitivity and β -adrenergic receptor density in failing human hearts. *N Engl J Med.* 1982; 307:205–211. [PubMed: 6283349]
 6. Port JD, Bristow MR. Altered beta-adrenergic receptor gene regulation and signaling in chronic heart failure. *J Mol Cell Cardiol.* 2001; 33:887–905. [PubMed: 11343413]
 7. Sterne-Marr R, Gurevich VV, Goldsmith P, Bodine RC, Sanders C, Donoso LA, Benovic JL. Polypeptide variants of β -arrestin and arrestin3. *J Biol Chem.* 1993; 268:15640–15648. [PubMed: 8340388]
 8. Lymperopoulos A, Bathgate A. Pharmacogenomics of the heptahelical receptor regulators G-protein-coupled receptor kinases and arrestins: the known and the unknown. *Pharmacogenomics.* 2012; 13:323–341. [PubMed: 22304582]
 9. Lymperopoulos A. Beta-arrestin biased agonism/antagonism at cardiovascular seven transmembrane-spanning receptors. *Curr Pharm Des.* 2012; 18:192–198. [PubMed: 22229558]
 10. Ungerer M, Böhm M, Elce JS, Erdmann E, Lohse MJ. Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. *Circulation.* 1993; 87:454–463. [PubMed: 8381058]
 11. Rengo G, Lymperopoulos A, Koch WJ. Future G protein-coupled receptor targets for treatment of heart failure. *Curr Treat Options Cardiovasc Med.* 2009; 11:328–338. [PubMed: 19627665]
 12. Belmonte SL, Blaxall BC. G protein coupled receptor kinases as therapeutic targets in cardiovascular disease. *Circ Res.* 2011; 109:309–319. [PubMed: 21778435]
 13. Rengo G, Lymperopoulos A, Leosco D, Koch WJ. GRK2 as a novel gene therapy target in heart failure. *J Mol Cell Cardiol.* 2011; 50:785–792. [PubMed: 20800067]
 14. Lymperopoulos A, Rengo G, Koch WJ. Adrenal adrenoceptors in heart failure: fine-tuning cardiac stimulation. *Trends Mol Med.* 2007; 13:503–511. [PubMed: 17981507]
 15. Lymperopoulos A, Rengo G, Koch WJ. GRK2 inhibition in heart failure: something old, something new. *Curr Pharm Des.* 2012; 18:186–191. [PubMed: 22229578]
 16. Penn RB, Pronin AN, Benovic JL. Regulation of G protein-coupled receptor kinases. *Trends Cardiovasc Med.* 2000; 10:81–89. [PubMed: 11150735]
 17. Conner DA, Mathier MA, Mortensen RM, Christe M, Vatner SF, Seidman CE, Seidman JG. beta-Arrestin1 knockout mice appear normal but demonstrate altered cardiac responses to beta-adrenergic stimulation. *Circ Res.* 1997; 81:1021–1026. [PubMed: 9400383]
 18. Noma T, Lemaire A, Naga Prasad SV, Barki-Harrington L, Tilley DG, Chen J, Le Corvoisier P, Violin JD, Wei H, Lefkowitz RJ, Rockman HA. Beta-arrestin-mediated beta1-adrenergic receptor transactivation of the EGFR confers cardioprotection. *J Clin Invest.* 2007; 117:2445–2458. [PubMed: 17786238]
 19. Tilley DG. G protein-dependent and G protein-independent signaling pathways and their impact on cardiac function. *Circ Res.* 2011; 109:217–230. [PubMed: 21737817]
 20. Noor N, Patel CB, Rockman HA. β arrestin: a signaling molecule and potential therapeutic target for heart failure. *J Mol Cell Cardiol.* 2011; 51:534–541. [PubMed: 21074538]
 21. Luttrell LM, Roudabush FL, Choy EW, Miller WE, Field ME, Pierce KL, Lefkowitz RJ. Activation and targeting of extracellular signal-regulated kinases by beta-arrestin scaffolds. *Proc Natl Acad Sci USA.* 2001; 98:2449–2454. [PubMed: 11226259]
 22. Kim J, Zhang L, Poppel K, Wu JH, Zidar DA, Brian L, DeWire SM, Exum ST, Lefkowitz RJ, Freedman NJ. Beta-arrestins regulate atherosclerosis and neointimal hyperplasia by controlling smooth muscle cell proliferation and migration. *Circ Res.* 2008; 103:70–79. [PubMed: 18519945]
 23. Rajagopal K, Whalen EJ, Violin JD, Stiber JA, Rosenberg PB, Premont RT, Coffman TM, Rockman HA, Lefkowitz RJ. Beta-arrestin2-mediated inotropic effects of the angiotensin II type 1A receptor in isolated cardiac myocytes. *Proc Natl Acad Sci USA.* 2006; 103:16284–16289. [PubMed: 17060617]
 24. Swedberg K, Eneroth P, Kjekshus J, Wilhelmsen L. Hormones regulating cardiovascular function in patients with severe congestive heart failure and their relation to mortality. CONSENSUS Trial Study Group. *Circulation.* 1990; 82:1730–1736. [PubMed: 2225374]

25. Weber KT. Aldosterone in congestive heart failure. *N Engl J Med.* 2001; 345:1689–1697. [PubMed: 11759649]
26. Lympopoulos A, Rengo G, Zincarelli C, Kim J, Soltys S, Koch WJ. An adrenal beta-arrestin 1-mediated signaling pathway underlies angiotensin II-induced aldosterone production in vitro and in vivo. *Proc Natl Acad Sci USA.* 2009; 106:5825–5830. [PubMed: 19289825]
27. Lympopoulos A, Rengo G, Zincarelli C, Kim J, Koch WJ. Adrenal beta-arrestin 1 inhibition in vivo attenuates post-myocardial infarction progression to heart failure and adverse remodeling via reduction of circulating aldosterone levels. *J Am Coll Cardiol.* 2011; 57:356–365. [PubMed: 21232674]
28. Lympopoulos A, Rengo G, Funakoshi H, Eckhart AD, Koch WJ. Adrenal GRK2 upregulation mediates sympathetic overdrive in heart failure. *Nat Med.* 2007; 13:315–323. [PubMed: 17322894]
29. Lympopoulos A, Rengo G, Zincarelli C, Soltys S, Koch WJ. Modulation of adrenal catecholamine secretion by in vivo gene transfer and manipulation of G protein-coupled receptor kinase-2 activity. *Mol Ther.* 2008; 16:302–307. [PubMed: 18223549]
30. Lympopoulos A, Rengo G, Gao E, Ebert SN, Dorn GW 2nd, Koch WJ. Reduction of sympathetic activity via adrenal-targeted GRK2 gene deletion attenuates heart failure progression and improves cardiac function after myocardial infarction. *J Biol Chem.* 2010; 285:16378–16386. [PubMed: 20351116]
31. Bers DM. Calcium cycling and signaling in cardiac myocytes. *Annu Rev Physiol.* 2008; 70:23–49. [PubMed: 17988210]
32. Luttrell LM, Gesty-Palmer D. Beyond desensitization: physiological relevance of arrestin-dependent signaling. *Pharmacol Rev.* 2010; 62:305–330. [PubMed: 20427692]
33. DeWire SM, Ahn S, Lefkowitz RJ, Shenoy SK. Beta-arrestins and cell signaling. *Annu Rev Physiol.* 2007; 69:483–510. [PubMed: 17305471]
34. Kho C, Lee A, Jeong D, Oh JG, Chaanine AH, Kizana E, Park WJ, Hajjar RJ. SUMO1-dependent modulation of SERCA2a in heart failure. *Nature.* 2011; 477:601–605. [PubMed: 21900893]
35. Rengo G, Lympopoulos A, Zincarelli C, Donniacuo M, Soltys S, Rabinowitz JE, Koch WJ. Myocardial adeno-associated virus serotype 6-betaARKct gene therapy improves cardiac function and normalizes the neurohormonal axis in chronic heart failure. *Circulation.* 2009; 119:89–98. [PubMed: 19103992]
36. Rakesh K, Yoo B, Kim IM, Salazar N, Kim KS, Rockman HA. Beta-arrestin-biased agonism of the angiotensin receptor induced by mechanical stress. *Sci Signal.* 2010; 3:ra46. [PubMed: 20530803]
37. Santulli G, Cipolletta E, Sorriento D, Del Giudice C, Anastasio A, Monaco S, Maione AS, Condorelli G, Puca A, Trimarco B, Illario M, Iaccarino G. CaMK4 Gene Deletion Induces Hypertension. *J Am Heart Assoc.* 2012; 1:e001081. [PubMed: 23130158]
38. Silvestre JS, Heymes C, Oubénaïssa A, Robert V, Aupetit-Faisant B, Carayon A, Swynghedauw B, Delcayre C. Activation of cardiac aldosterone production in rat myocardial infarction: effect of angiotensin II receptor blockade and role in cardiac fibrosis. *Circulation.* 1999; 99:2694–2701. [PubMed: 10338465]
39. Kuroko Y, Yamazaki T, Tokunaga N, Akiyama T, Kitagawa H, Ishino K, Sano S, Mori H. Cardiac epinephrine synthesis and ischemia-induced myocardial epinephrine release. *Cardiovasc Res.* 2007; 74:438–444. [PubMed: 17448453]

Novelty and Significance

What Is New?


- β_{arr1} (rather than β_{arr2}) is the β_{arr} isoform responsible for β_{AR} desensitization and downregulation in vivo in the heart, a hallmark molecular abnormality in chronic HF.
- β_{arr1} in the adrenals is absolutely essential for post-MI hyperaldosteronism and catecholamine elevation in vivo, thereby heightening the neurohormonal burden of the failing heart.
- Cardiac β_{arr1} also counters some beneficial effects of β_{arr2} in the myocardium, e.g. anti-apoptosis and suppression of inflammation.

What Is Relevant?

- Adrenal β_{arr1} can promote hyperaldosteronism and sympathetic hyperactivity, two neurohormonal mechanisms that contribute significantly to the development of hypertension and HF.
- Pharmacological or genetic β_{arr1} inhibition in vivo, at least in the adrenals and in the heart, might be a valid anti-hypertensive and anti-HF treatment strategy.

Summary

Blockade of β arr1 activity in both the heart and the adrenal gland might be of therapeutic value for halting or even reversing post-MI HF progression.

**Figure 1.**

(A) Ejection fraction (EF) % of sham-operated (sham) or of 4-week post-MI (MI) β arr1KO and WT mice. *, $p < 0.05$, vs. either Sham; **, $p < 0.05$, vs. WT MI; $n = 7$ mice/group. (B) Basal and maximal dose (333 ng/kg) of isoproterenol (Max. Iso)-stimulated $+dP/dt_{max}$ responses of these mice. *, $p < 0.05$, vs. WT Sham-Max. Iso; #, $p < 0.05$, vs. either Sham; **, $p < 0.05$, vs. WT MI-Max. Iso; ^, $p < 0.05$, vs. WT MI-Basal; $n = 5$ mice/group.




Figure 2.

(A) Kaplan-Meier survival curves of the 4 groups of the study: sham-operated (Sham) and post-MI (MI) β arr1KO and WT mice. $p=0.012$ between MI WT and MI β arr1KO; $n=15$ mice/group for sham, 37 mice/group for MI mice. (B) Plasma circulating NE and Epi levels in sham-operated (Sham) or in 4-week post-MI (MI) β arr1KO (KO) and WT mice. *, $p<0.05$, vs. Sham-either genotype; ^, $p<0.05$, vs. WT MI; $n=6$ mice/group. (C) Serum aldosterone levels in these mice. *, $p<0.05$, vs. all other groups; $n=6$ mice/group.





Figure 3.

(A) Apoptotic cell death at 24 hrs post-MI in β arr1KO (KO) and WT mice, as measured by TUNEL performed in the border zone of the infarct. Representative images of TUNEL-positive nuclei are shown (left), along with their quantitation (right). No difference in rate of apoptosis in the remote zone was found (data not shown). *, $p < 0.05$, vs. either Sham; #, $p < 0.05$, vs. WT MI; $n = 6$ hearts/group. (B) Left Ventricular End Diastolic Diameter (LVEDD) of these mice at 4 weeks post-MI or post-sham operation. #, $p < 0.05$, vs. either Sham; *, $p < 0.05$, vs. WT MI; $n = 7$ mice/group. (C–D) Infarct size at 4 weeks post-MI. Representative triphenyltetrazolium chloride (TTC)-stained cardiac cross-sections are shown in (C), and average left ventricular (LV) infarct size, expressed as % LV infarction and measured from 6 mice/group, is shown in (D). **, $p < 0.05$, vs. WT MI.

**Figure 4.**

(A–C) Levels of pro-inflammatory cytokines TNF α (A), IL-6 (B), and IL-1 β (C), measured in serum of intra-cardiac blood from β arr1KO (KO) and WT mice at 4 weeks post-MI (MI) or post-sham operation (Sham). *, $p < 0.05$, vs. all other groups; **, $p < 0.05$, vs. WT Sham; $n = 5$ mice/group. (D–E) Cardiac fibrosis staining in the 4-week post-MI mice. Blue denotes collagen fibers, red denotes muscle fibers, and black represents cell nuclei. Representative images are shown in (D) and quantification of the % fibrotic area in (E). *, $p < 0.05$; $n = 5$ hearts/group. (F–G) Heart mRNA levels of Collagen-1 α 1 (Col-1 α 1) (F) and of brain

natriuretic peptide (BNP) (**G**) in these mice. [#], $p < 0.05$, vs. either Sham; ^{*}, $p < 0.05$, vs. WT MI; $n = 6$ hearts/group.




Figure 5.

(A) β AR density in cardiac plasma membranes of sham-operated (Sham) or of 4-week post-MI (MI) β arr1KO (KO) and WT mice. *, $p < 0.05$, vs. all other groups; $n = 5$ hearts/group. (B) Steady-state total cAMP levels in cardiac homogenates purified from these mice. *, $p < 0.05$ vs. all other groups; $n = 5$ hearts/group. (C) Cardiac SERCA activity measured in membrane homogenates from the hearts of these mice. *, $p < 0.05$, vs. WT Sham; **, $p < 0.05$, vs. WT MI; $n = 5$ hearts/group. (D–E) Cardiac EGFR transactivation in these mice. Representative western blots in total cardiac protein extracts for phospho-Tyr845-EGFR or total EGFR (loading control) are shown in (D) and their densitometric quantitation in (E). Relative EGFR transactivation was calculated by comparing the ratios of phospho-EGFR to total EGFR for each sample. *, $p < 0.05$, vs. either WT; **, $p < 0.05$, vs. KO Sham; $n = 5$ hearts/group.




Figure 6.

Schematic illustration of the cardiac (left) and adrenal (right) signaling mechanisms underlying the effects of β arr1 on cardiovascular function. G_s : stimulatory G protein; P: Phosphorylation; α_2 AR: alpha2-adrenergic receptor; AT_1R : Angiotensin II type 1 receptor; CA: Catecholamine. See text for details and all other molecular acronym descriptions.