

# Nuclear pore proteins regulate chromatin structure and transcriptional memory by a conserved mechanism

William H Light<sup>†</sup> and Jason H Brickner\*

Department of Molecular Biosciences; Northwestern University; Evanston, IL USA

<sup>†</sup>Current affiliation: Molecular and Cell Biology Laboratory; Salk Institute for Biological Studies; San Diego, CA USA

**P**revious experience alters the rate of transcriptional induction of many genes in yeast, and this phenomenon persists through several cell division cycles. This phenomenon is called epigenetic transcriptional memory. For the yeast gene *INO1*, transcriptional memory requires a physical interaction with the nuclear pore complex (NPC) and changes in the chromatin structure of the promoter. These changes lead to binding of a preinitiation form of RNA Polymerase II (RNAPII) to the *INO1* promoter, bypassing the need to recruit RNAPII to the promoter during reactivation. In our recent study, we found that in human cells, hundreds of interferon- $\gamma$  responsive genes exhibit a mechanistically similar form of transcriptional memory. Transcriptional memory requires a homologous nuclear pore protein in yeast and humans, which interacts with the promoters of genes that exhibit transcriptional memory and promotes both alteration of chromatin structure and binding of RNAPII. Whereas the interaction of yeast genes with nuclear pore proteins occurs at the NPC, the interaction of human genes with nuclear pore proteins occurs in the nucleoplasm. Thus, the interaction of nuclear pore proteins with genes plays an important and conserved role in affecting long-term epigenetic changes in transcriptional regulation.

## Introduction

The nuclear pore complex (NPC) is a large (~60 megadalton), universally

conserved 8-fold symmetric complex made up of ~30 different nuclear pore proteins (Nups).<sup>1-3</sup> This structure plays an essential role in the transit of RNA and protein between the nucleus and the cytoplasm and has been implicated in many diverse processes aside from transport.<sup>4-11</sup> Despite early evidence in yeast indicating that the NPC interacts with silenced or repressed portions of the genome,<sup>12-14</sup> more recent work has indicated that many active genes in yeast interact with the NPC.<sup>15,16</sup> Furthermore, several yeast genes remain associated with the NPC after being repressed and, in the case of the model gene *INO1*, this interaction leads to an altered chromatin structure and binding of a poised RNA Polymerase II (RNAPII) to the recently repressed promoter.<sup>1,17,18</sup> This phenomenon is called transcriptional memory and can be inherited through several generations, indicating that it is epigenetic. The *GAL* genes and a class of stress-induced genes exhibit a similar behavior.<sup>19</sup> The mechanistic details for these genes are similar but not identical to *INO1*.<sup>19</sup> Furthermore, although there is a clear role for the interaction of Nups in promoting transcription in flies and humans,<sup>20-23</sup> it was unclear if metazoan Nups were also involved in transcriptional memory. A study published in 2010 suggested that the induction of the human *HLA-DRA* gene in response to interferon gamma (IFN- $\gamma$ ) was faster in cells that had previously been exposed to IFN- $\gamma$ .<sup>24</sup> This led us to ask if IFN- $\gamma$  induced genes are regulated by a mechanistically similar form of epigenetic transcriptional memory. In our recent paper, we showed

**Keywords:** nuclear pore complex, nuclear porins, chromatin, epigenetics, transcriptional memory

\*Correspondence to: Jason H Brickner;  
Email: j-brickner@northwestern.edu

Submitted: 07/31/2013

Revised: 08/16/2013

Accepted: 08/19/2013

<http://dx.doi.org/10.4161/nucl.26209>

Extra View to: Light WH, Freaney J, Sood V, Thompson A, D'Urso A, Horvath CM, Brickner JH. A conserved role for human Nup98 in altering chromatin structure and promoting epigenetic transcriptional memory. PLoS Biol 2013; 11:e1001524; PMID:23555195; <http://dx.doi.org/10.1371/journal.pbio.1001524>

©2013 Landes Bioscience. Do not distribute

that hundreds of such genes exhibit transcriptional memory and that this form of memory is mechanistically very similar to *INO1* memory in yeast.<sup>25</sup> Here we discuss these findings in a broader context.

### Transcriptional Memory Regulates RNAPII Recruitment

We previously discovered a persistent interaction of a poised, unphosphorylated RNAPII at the *INO1* promoter for generations after repression.<sup>1</sup> Yeast strains that lack the histone variant H2A.Z or the nuclear pore protein Nup100 do not exhibit transcriptional memory; *INO1* relocalizes to the nucleoplasm after repression, RNAPII binding to the promoter is lost and the rate of reactivation of *INO1* is significantly slowed (while the rate of the initial activation is unaffected).<sup>1</sup> This suggests that transcriptional memory promotes faster reactivation by bypassing the rate-limiting step of RNAPII recruitment. To better understand this, we compared the binding of general transcription factors (GTFs) to the promoter of active and recently repressed *INO1* by chromatin immunoprecipitation (ChIP). Every GTF tested (TFIIA, TFIIB, TFIID, TFIIE, TFIIF, TFIIH, TFIK, TFIIS, and Mediator) was bound to the active *INO1* promoter. Most of the GTFs (TFIIA, TFIIB, TFIID, TFIIE, TFIIF, and TFIIH) were bound to the recently repressed *INO1* promoter, but others (TFIIK, TFIIS, and Mediator) were not. This result was consistent with our observation that the RNAPII that was bound to the recently repressed *INO1* promoter was not phosphorylated on Ser2 and Ser5 of the carboxyl terminal domain of RNAPII, indicating that this is a preinitiation form of RNAPII.<sup>26</sup> The mechanism of *INO1* transcriptional memory suggests that transcription can be regulated at the level of RNAPII recruitment, at the level of initiation and at the level of elongation.<sup>27,28</sup> Consistent with this conclusion, in both stationary phase yeast cells and G<sub>0</sub> lymphocytes, thousands of genes are poised for future expression through binding of preinitiation complex to their promoters.<sup>29,30</sup>

*INO1* transcriptional memory requires an 11 base pair *cis* acting promoter element

called the memory recruitment sequence (MRS). The promoters of stress-induced yeast genes that display transcriptional memory are enriched for a very similar element.<sup>19</sup> Furthermore, the MRS is sufficient to recapitulate the chromatin structure associated with transcriptional memory when inserted at ectopic loci.<sup>1</sup> However, the MRS is not sufficient to promote PIC association with an ectopic locus, suggesting that PIC association is a context-dependent (and presumably promoter-specific) output of transcriptional memory. Therefore, peripheral localization and chromatin structure probably function to regulate PIC formation and can be uncoupled from these downstream events.

The human gene *HLA-DRA* is induced by IFN- $\gamma$  and has much faster induction kinetics if cells have previously been treated with IFN- $\gamma$ .<sup>31,32</sup> This effect persists for up to four cell divisions, suggesting that it is epigenetically inherited through mitosis.<sup>24,25</sup> We found that several hundred human genes displayed significantly faster induction kinetics in cells that had been exposed to IFN- $\gamma$  compared with their initial activation. Transcriptional memory does not require full transcription of *HLA-DRA*; treatment of cells with IFN- $\gamma$  for 2 h, which was not sufficient to lead to measurable increase in *HLA-DRA*, was sufficient to lead to transcriptional memory. After removing IFN- $\gamma$ , a preinitiation form of RNAPII remained bound to the promoters of genes that exhibited transcriptional memory, suggesting that human genes exhibit a form of transcriptional memory that is mechanistically similar to *INO1* memory in yeast.

### Non-NPC Bound Roles for Nups

Nups interact with a large number of genes in eukaryotic genomes, leading to both positive and negative impacts on gene expression.<sup>15,16,33,34</sup> *INO1* transcriptional memory requires interactions with several yeast Nups (Fig. 1).<sup>1</sup> The active *INO1* promoter interacts with different Nups than the recently repressed *INO1* promoter.<sup>1</sup> Based on ChIP, active *INO1* interacts with Nup2, one of several Nups having Phe-X-Phe-Gly repeats, but does not interact

with Nup100, a Gly-Leu-Phe-Gly repeat protein. In contrast, recently repressed *INO1* interacts weakly with Nup2 and strongly with Nup100.<sup>1</sup> Using antibodies against Phe-X-Phe-Gly or against Nup98, a human homolog of yeast Nup100, we performed ChIP and measured the interaction of each of these antigens with the promoters of human genes that exhibit transcriptional memory. In humans, Phe-X-Phe-Gly Nups include Nup62, Nup153, Nup214, Nup358/RanBP2, and Pom121. Similar to yeast *INO1*, we observed an interaction of Phe-X-Phe-Gly proteins but not Nup98, with the active promoters and a weaker interaction with Phe-X-Phe-Gly proteins but a strong interaction of Nup98 with previously expressed promoters for up to 96h after removing IFN- $\gamma$ . Therefore, different human Nups bind to active and recently expressed genes that exhibit transcriptional memory.

Although all of the interactions between yeast genes and Nups occur at the NPC, many Nup-gene interactions in metazoan cells occur in the nucleoplasm, away from the NPC. We localized the *HLA-DRA* gene with respect to the nuclear periphery using DNA FISH before IFN- $\gamma$  treatment, during IFN- $\gamma$  treatment and 48 h after IFN- $\gamma$  treatment. Consistent with a role for non-NPC-bound Nups in transcriptional memory, *HLA-DRA* did not co-localize with the nuclear periphery under any condition. This suggests that in HeLa cells, nucleoplasmic Nups interact with *HLA-DRA*.<sup>21,23</sup> Although these interactions occur at a different subnuclear location in yeast and humans, it seems likely that they represent a similar, highly conserved, biochemical process.

### Chromatin Structure and Transcriptional Memory

Changes in the chromatin state of gene promoters have been tied to transcriptional memory in both yeast and humans.<sup>1,24</sup> In yeast, the histone variant H2A.Z was required for *INO1* transcriptional memory and the *INO1* MRS was both necessary and sufficient to promote the deposition of H2A.Z.<sup>1,17</sup> *HLA-DRA* transcriptional memory is associated with persistent dimethylation of histone H3 lysine 4 at the promoter (H3K4me2).<sup>24</sup> This led us to ask

if this chromatin mark was also associated with yeast *INO1* transcriptional memory. Indeed, H3K4me2 persists at the recently repressed *INO1* promoter. Mutation of the MRS led to loss of H3K4me2 after repression. The MRS was also sufficient to induce dimethylation of H3K4 at an ectopic locus. Moreover, mutants lacking the enzymes responsible for H3K4 methylation blocked *INO1* transcriptional memory, suggesting that H3K4 methylation is required for memory. Finally, H3K4me2 deposition occurred in the absence of H2A.Z, suggesting that H3K4me2 incorporation occurred upstream, or in parallel to, H2A.Z deposition.

### hNup98 is Required for Transcriptional Memory

Yeast *INO1* transcriptional memory specifically requires Nup100.<sup>1</sup> Using siRNAs to deplete Nup98 in HeLa cells, we found that loss of Nup98 led to loss of all measures of transcriptional memory: RNAPII and H3K4me2 were lost from the promoters of genes that exhibit IFN- $\gamma$  memory and the rate of reactivation of these genes was much slower. This effect was specific to Nup98 depletion; Nup107 depletion did not affect transcriptional memory. We also found that mutants lacking yeast Nup100 failed to retain H3K4me2 at the recently repressed *INO1* promoter, further supporting the conservation of this mechanism.

### Conclusions

Some Nups can interact with promoters to modulate chromatin structure and gene expression in both yeast and humans,


#### References

1. Light WH, Brickner DG, Brand VR, Brickner JH. Interaction of a DNA zip code with the nuclear pore complex promotes H2A.Z incorporation and *INO1* transcriptional memory. *Mol Cell* 2010; 40:112-25; PMID:20932479; <http://dx.doi.org/10.1016/j.molcel.2010.09.007>
2. Suntharalingam M, Wente SR. Peering through the pore: nuclear pore complex structure, assembly, and function. *Dev Cell* 2003; 4:775-89; PMID:12791264; [http://dx.doi.org/10.1016/S1534-5807\(03\)00162-X](http://dx.doi.org/10.1016/S1534-5807(03)00162-X)
3. Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, et al. The molecular architecture of the nuclear pore complex. *Nature* 2007; 450:695-701; PMID:18046406; <http://dx.doi.org/10.1038/nature06405>

suggesting that Nups have functions independent of their role in intranuclear transport. Our recent work has further defined the molecular mechanism of yeast transcriptional memory and found strong evidence that facets of this mechanism are conserved from yeast to humans. Based on this, we predict that many genes are regulated by transcriptional memory in different signaling contexts. Consistent with this notion, the promoters of stress-responsive yeast genes that exhibit a form of epigenetic transcriptional memory are enriched for an element that is very similar to the MRS element.<sup>19</sup> Also, these genes require Nup42 for this behavior.<sup>19</sup> If these genes are regulated by a mechanism similar to that used by *INO1* and *HLA-DRA*, their promoters may be marked with H3K4me2 and bound to RNAPII following the primary stress.

Transcriptional memory reflects both the cell's current physiology and previous experiences. In the context of the body, the response of cells and tissues to the same stimulus might differ dramatically based on recent (or not so recent) experience. A complete understanding of gene expression patterns must also consider how these patterns are related to a cell's or a tissue's history. Transcriptional memory may allow qualitatively different transcriptional responses to a particular stimulus, by integrating signals with recent experience. If so, this mechanism might provide better responses to infection. However, it is

4. Ishii K, Arib G, Lin C, Van Houwe G, Laemmli UK. Chromatin boundaries in budding yeast: the nuclear pore connection. *Cell* 2002; 109:551-62; PMID:12062099; [http://dx.doi.org/10.1016/S0092-8674\(02\)00756-0](http://dx.doi.org/10.1016/S0092-8674(02)00756-0)
5. Fischer T, Strässer K, Rácz A, Rodríguez-Navarro S, Oppizzi M, Ihrig P, Lechner J, Hurt E. The mRNA export machinery requires the novel Sac3p-Thp1p complex to dock at the nucleoplasmic entrance of the nuclear pores. *EMBO J* 2002; 21:5843-52; PMID:12411502; <http://dx.doi.org/10.1093/emboj/cdf590>
6. Siniosoglou S, Wimmer C, Rieger M, Doye V, Tekotte H, Weise C, Emig S, Segref A, Hurt EC. A novel complex of nucleoporins, which includes Sec13p and a Sec13p homolog, is essential for normal nuclear pores. *Cell* 1996; 84:265-75; PMID:8565072; [http://dx.doi.org/10.1016/S0092-8674\(00\)80981-2](http://dx.doi.org/10.1016/S0092-8674(00)80981-2)



**Figure 1.** Nup-dependent transcriptional memory in yeast and humans. Left, *INO1* transcriptional memory in yeast. After repression, the *INO1* gene remains bound to the NPC. RNAPII clears the coding region but remains associated with the promoter in a poised, preinitiation form. Promoter nucleosomes are altered by incorporation of H2A.Z and dimethylation of H3K4 (blue nucleosome). Transcriptional memory persists at the NPC for approximately four generations. Right, IFN- $\gamma$  transcriptional memory in humans. After the removal of IFN- $\gamma$ , a poised preinitiation form of RNAPII remains associated with the promoter. Promoter nucleosomes are marked with H3K4me2 (blue) and bind to nucleoplasmic Nups such as Nup98. This epigenetic state is inherited for at least four generations.

also possible that this mechanism might contribute to pathological outcomes such as chronic inflammation. If so, transcriptional memory may represent a novel pathway for therapeutic intervention for the treatment of inflammatory disorders, which often involve mis-regulation of IFN- $\gamma$  responsive genes.

#### Disclosure of Potential Conflicts of Interest

No potential conflict of interest was disclosed.

7. Mendjan S, Taipale M, Kind J, Holz H, Gebhardt P, Schelder M, Vermeulen M, Buscaino A, Duncan K, Mueller J, et al. Nuclear pore components are involved in the transcriptional regulation of dosage compensation in *Drosophila*. *Mol Cell* 2006; 21:811-23; PMID:16543150; <http://dx.doi.org/10.1016/j.molcel.2006.02.007>
8. Menon BB, Sarma NJ, Pasula S, Deminoff SJ, Willis KA, Barbara KE, Andrews B, Santangelo GM. Reverse recruitment: the Nup84 nuclear pore sub-complex mediates Rap1/Gcr1/Gcr2 transcriptional activation. *Proc Natl Acad Sci U S A* 2005; 102:5749-54; PMID:15817685; <http://dx.doi.org/10.1073/pnas.0501768102>
9. Lei EP, Krebber H, Silver PA. Messenger RNAs are recruited for nuclear export during transcription. *Genes Dev* 2001; 15:1771-82; PMID:11459827; <http://dx.doi.org/10.1101/gad.892401>

10. Dilworth DJ, Suprpto A, Padovan JC, Chait BT, Wozniak RW, Rout MP, Aitchison JD. Nup2p dynamically associates with the distal regions of the yeast nuclear pore complex. *J Cell Biol* 2001; 153:1465-78; PMID:11425876; <http://dx.doi.org/10.1083/jcb.153.7.1465>
11. Dilworth DJ, Tackett AJ, Rogers RS, Yi EC, Christmas RH, Smith JJ, Siegel AF, Chait BT, Wozniak RW, Aitchison JD. The mobile nucleoporin Nup2p and chromatin-bound Prp20p function in endogenous NPC-mediated transcriptional control. *J Cell Biol* 2005; 171:955-65; PMID:16365162; <http://dx.doi.org/10.1083/jcb.200509061>
12. Galy V, Olivo-Marin JC, Scherthan H, Doye V, Rascalou N, Nehrbass U. Nuclear pore complexes in the organization of silent telomeric chromatin. *Nature* 2000; 403:108-12; PMID:10638763; <http://dx.doi.org/10.1038/47528>
13. Hediger F, Dubrana K, Gasser SM. Myosin-like proteins 1 and 2 are not required for silencing or telomere anchoring, but act in the Tel1 pathway of telomere length control. *J Struct Biol* 2002; 140:79-91; PMID:12490156; [http://dx.doi.org/10.1016/S1047-8477\(02\)00533-6](http://dx.doi.org/10.1016/S1047-8477(02)00533-6)
14. Hediger F, Neumann FR, Van Houwe G, Dubrana K, Gasser SM. Live imaging of telomeres: yKu and Sir proteins define redundant telomere-anchoring pathways in yeast. *Curr Biol* 2002; 12:2076-89; PMID:12498682; [http://dx.doi.org/10.1016/S0960-9822\(02\)01338-6](http://dx.doi.org/10.1016/S0960-9822(02)01338-6)
15. Casolari JM, Brown CR, Drubin DA, Rando OJ, Silver PA. Developmentally induced changes in transcriptional program alter spatial organization across chromosomes. *Genes Dev* 2005; 19:1188-98; PMID:15905407; <http://dx.doi.org/10.1101/gad.1307205>
16. Casolari JM, Brown CR, Komili S, West J, Hieronymus H, Silver PA. Genome-wide localization of the nuclear transport machinery couples transcriptional status and nuclear organization. *Cell* 2004; 117:427-39; PMID:15137937; [http://dx.doi.org/10.1016/S0092-8674\(04\)00448-9](http://dx.doi.org/10.1016/S0092-8674(04)00448-9)
17. Brickner DG, Cajigas I, Fondufe-Mittendorf Y, Ahmed S, Lee PC, Widom J, Brickner JH. H2A.Z-mediated localization of genes at the nuclear periphery confers epigenetic memory of previous transcriptional state. *PLoS Biol* 2007; 5:e81; PMID:17373856; <http://dx.doi.org/10.1371/journal.pbio.0050081>
18. Kundu S, Horn PJ, Peterson CL. SWI/SNF is required for transcriptional memory at the yeast GAL gene cluster. *Genes Dev* 2007; 21:997-1004; PMID:17438002; <http://dx.doi.org/10.1101/gad.1506607>
19. Guan Q, Haroon S, Bravo DG, Will JL, Gasch AP. Cellular memory of acquired stress resistance in *Saccharomyces cerevisiae*. *Genetics* 2012; 192:495-505; PMID:22851651; <http://dx.doi.org/10.1534/genetics.112.143016>
20. Faria AM, Levay A, Wang Y, Kamphorst AO, Rosa ML, Nussenzeig DR, Balkan W, Chook YM, Levy DE, Fontoura BM. The nucleoporin Nup96 is required for proper expression of interferon-regulated proteins and functions. *Immunity* 2006; 24:295-304; PMID:16546098; <http://dx.doi.org/10.1016/j.immuni.2006.01.014>
21. Kalverda B, Pickersgill H, Shloma VV, Fornerod M, Silver PA. Nucleoporins directly stimulate expression of developmental and cell-cycle genes inside the nucleoplasm. *Cell* 2010; 140:360-71; PMID:20144760; <http://dx.doi.org/10.1016/j.cell.2010.01.011>
22. Brown CR, Kennedy CJ, Delmar VA, Forbes DJ, Silver PA. Global histone acetylation induces functional genomic reorganization at mammalian nuclear pore complexes. *Genes Dev* 2008; 22:627-39; PMID:18316479; <http://dx.doi.org/10.1101/gad.1632708>
23. Capelson M, Liang Y, Schulte R, Mair W, Wagner U, Hetzer MW. Chromatin-bound nuclear pore components regulate gene expression in higher eukaryotes. *Cell* 2010; 140:372-83; PMID:20144761; <http://dx.doi.org/10.1016/j.cell.2009.12.054>
24. Gialitakis M, Arampatzi P, Makatounakis T, Papamatheakis J. Gamma interferon-dependent transcriptional memory via relocalization of a gene locus to PML nuclear bodies. *Mol Cell Biol* 2010; 30:2046-56; PMID:20123968; <http://dx.doi.org/10.1128/MCB.00906-09>
25. Light WH, Freaney J, Sood V, Thompson A, D'Urso A, Horvath CM, Brickner JH. A conserved role for human Nup98 in altering chromatin structure and promoting epigenetic transcriptional memory. *PLoS Biol* 2013; 11:e1001524; PMID:23555195; <http://dx.doi.org/10.1371/journal.pbio.1001524>
26. Fuda NJ, Ardehali MB, Lis JT. Defining mechanisms that regulate RNA polymerase II transcription in vivo. *Nature* 2009; 461:186-92; PMID:19741698; <http://dx.doi.org/10.1038/nature08449>
27. Peterlin BM, Price DH. Controlling the elongation phase of transcription with P-TEFb. *Mol Cell* 2006; 23:297-305; PMID:16885020; <http://dx.doi.org/10.1016/j.molcel.2006.06.014>
28. Sims RJ 3<sup>rd</sup>, Belotserkovskaya R, Reinberg D. Elongation by RNA polymerase II: the short and long of it. *Genes Dev* 2004; 18:2437-68; PMID:15489290; <http://dx.doi.org/10.1101/gad.1235904>
29. Radonjic M, Andrau JC, Lijnzaad P, Kemmeren P, Kockelkorn TT, van Leenen D, van Berkum NL, Holstege FC. Genome-wide analyses reveal RNA polymerase II located upstream of genes poised for rapid response upon *S. cerevisiae* stationary phase exit. *Mol Cell* 2005; 18:171-83; PMID:15837421; <http://dx.doi.org/10.1016/j.molcel.2005.03.010>
30. Kouzine F, Wojtowicz D, Yamane A, Resch W, Kieffer-Kwon KR, Bandle R, Nelson S, Nakahashi H, Awasthi P, Feigenbaum L, et al. Global regulation of promoter melting in naive lymphocytes. *Cell* 2013; 153:988-99; PMID:23706737; <http://dx.doi.org/10.1016/j.cell.2013.04.033>
31. Schroder K, Hertzog PJ, Ravasi T, Hume DA. Interferon-gamma: an overview of signals, mechanisms and functions. *J Leukoc Biol* 2004; 75:163-89; PMID:14525967; <http://dx.doi.org/10.1189/jlb.0603252>
32. Boehm U, Klamp T, Groot M, Howard JC. Cellular responses to interferon-gamma. *Annu Rev Immunol* 1997; 15:749-95; PMID:9143706; <http://dx.doi.org/10.1146/annurev.immunol.15.1.749>
33. Green EM, Jiang Y, Joyner R, Weis K. A negative feedback loop at the nuclear periphery regulates GAL gene expression. *Mol Biol Cell* 2012; 23:1367-75; PMID:22323286; <http://dx.doi.org/10.1091/mbc.E11-06-0547>
34. Ahmed S, Brickner DG, Light WH, Cajigas I, McDonough M, Froysheter AB, Volpe T, Brickner JH. DNA zip codes control an ancient mechanism for gene targeting to the nuclear periphery. *Nat Cell Biol* 2010; 12:111-8; PMID:20098417; <http://dx.doi.org/10.1038/ncb2011>