

Published in final edited form as:

Biochem Biophys Res Commun. 2014 January 3; 443(1): 115–119. doi:10.1016/j.bbrc.2013.11.060.

Synthetic embryonic lethality upon deletion of the ER cochaperone p58^{IPK} and the ER stress sensor ATF6 α

Javier A. Gomez^a, Heather M. Tyra^a, Diane DeZwaan McCabe^a, Alicia K. Olivier^b, and D. Thomas Rutkowski^{*,a,c}

^aDepartment of Anatomy and Cell Biology, 1-570 BSB, 51 Newton Rd., University of Iowa Carver College of Medicine, Iowa City, IA, 52242

^bDepartment of Pathology, 1-570 BSB, 51 Newton Rd., University of Iowa Carver College of Medicine, Iowa City, IA, 52242

^cDepartment of Internal Medicine, 1-570 BSB, 51 Newton Rd., University of Iowa Carver College of Medicine, Iowa City, IA, 52242

Abstract

The unfolded protein response (UPR) is activated as a consequence of alterations to ER homeostasis. It upregulates a group of ER chaperones and cochaperones, as well as other genes that improve protein processing within the secretory pathway. The UPR effector ATF6 α augments—but is not essential for—maximal induction of ER chaperones during stress, yet its role, if any, in protecting cellular function during normal development and physiology is unknown. A systematic analysis of multiple tissues from *Atf6 α -/-* mice revealed that all tissues examined were grossly insensitive to loss of ATF6 α . However, combined deletion of ATF6 α and the ER cochaperone p58^{IPK} resulted in synthetic embryonic lethality. These findings reveal for the first time that an intact UPR can compensate for the genetic impairment of protein folding in the ER *in vivo*. The also expose a role for p58^{IPK} in normal embryonic development.

Keywords

ER stress; unfolded protein response; chaperones; protein folding

Introduction

Disruption of protein folding in the endoplasmic reticulum (ER) is sensed by ER-resident transmembrane proteins which initiate signaling cascades collectively known as the unfolded protein response (UPR) [1]. Three pathways, initiated by ATF6 α , IRE1 α , and PERK, characterize the canonical UPR, and each culminates in the production of a transcriptional activator [2]. IRE1 α is conserved amongst eukaryotes and catalyzes splicing of an mRNA encoding the transcription factor XBP1 in vertebrates [3,4,5]. PERK is metazoan-specific, and phosphorylates the translation factor eIF2 α to transiently depress general protein synthesis but also to stimulate translation of a subset of mRNAs, including

© 2013 Elsevier Inc. All rights reserved.

*Corresponding author thomas-rutkowski@uiowa.edu Tel: (319) 335-7739.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

that of the transcription factor ATF4 [6,7,8]. ATF6 α is likewise metazoan-specific, and is cleaved by regulated intramembrane proteolysis to liberate an active transcription factor [9,10].

The IRE1 α and PERK pathways have roles in embryonic or postnatal development. Deficiency of IRE1 α or XBP1 compromises fetal liver development [11,12], and liver-specific rescue of XBP1 results in postnatal exocrine pancreas insufficiency [13]. PERK ablation leads to postnatal loss of insulin-producing pancreatic β cells [14]. While the IRE1 α and PERK pathways converge upon transcriptional regulation of genes that improve ER protein folding and processing [3,5,6], both can also regulate genes involved in process unrelated to ER folding such as membrane biosynthesis and lipid metabolism [15,16,17,18].

In contrast to ablation of the IRE1 α or PERK pathways, deletion of ATF6 α does not result in prenatal or neonatal lethality and *Atf6 α -/-* mice are fertile and grossly normal [19,20]. Microarray experiments have suggested that the direct targets of ATF6 α are largely restricted to genes encoding ER chaperones and cochaperones, ER-associated degradation factors, and other genes with direct roles in protein folding and processing [19,21]. Its function is partially compensated by its paralog ATF6 β , and chaperone insufficiency appears to underlie the embryonic lethality of *Atf6 α -/-*; *Atf6 β -/-* animals [20,22]. Perhaps because of this redundancy, to date *Atf6 α -/-* mice have only yielded phenotypes in the presence of exogenous superphysiological challenges such as inhibition of N-linked glycosylation, exposure to neurotoxin, exercise, experimental colitis, or maintenance on a high fat diet [23,24,25,26,27]. Thus, the role of ATF6 α , if any, in maintaining homeostasis during normal development and physiology is not clear.

The sensitivity of mice lacking ATF6 α to inhibition of N-linked glycosylation or experimental colitis is phenocopied in animals lacking the ER-resident DnaJ protein p58^{IPK}/ERDJ6 [23,24]. While this protein was originally identified as a negative regulator of the cytosolic eIF2 α kinase PKR [28], and thereafter as a negative regulator of PERK [29,30], it has since been found to contain an N-terminal ER targeting sequence, and p58^{IPK} is localized essentially quantitatively in the ER lumen; no detectable amounts of the protein were found in the cytosol under either basal or ER stress conditions [31]. In the ER lumen, p58^{IPK} interacts with the HSP70 chaperone BiP/GRP78 [31,32,33], as would be expected for a DnaJ-family protein [34]. Mice lacking p58^{IPK} are smaller than wild-type littermates, and develop diabetes attributed to compromised pancreatic β cell function [35]. The similar sensitivity to ER stress-inducing agents in animals lacking p58^{IPK} or ATF6 α suggests that a major function of p58^{IPK} is to protect the protein folding capacity of the ER, and raises the possibility that p58^{IPK} might serve in this capacity in tissues in addition to the endocrine pancreas, but that loss of p58^{IPK} function in these tissues is masked by compensatory activation of the UPR.

Materials and Methods

All protocols for animal use were reviewed and approved by the University Committee on Use and Care of Animals at the University of Iowa. Animals were fed standard rodent chow and housed in a controlled environment with 12 hr light and dark cycles. *Atf6 α -/-* [19] and *p58^{IPK}-/-* [35] animals have been backcrossed into the *C57BL/6J* strain for >10 generations. For histological analysis of wild-type and *Atf6 α -/-* mice, tissues were fixed in formalin, embedded in paraffin, sectioned, stained with hematoxylin and eosin, and examined in groups matched by age and gender for histological signs of pathology by a veterinary pathologist (A.K.O.) who was blinded to genotype. Multiple regions of each organ were examined. Numbers of mice examined were as follows: female wild-type, aged 7-17 months: 6; female *Atf6 α -/-*, aged 7-17 months: 8; male wild-type, aged 17 months: 3;

and male *Atf6a*^{-/-}, aged 17 months: 5. Timed intercrosses were dated by detection of a mucus plug, which was considered as E0.5; staging was confirmed by embryo size and developmental features. ADRP immunostaining, injections with tunicamycin, and RNA analysis from liver were performed as described [36].

Results

Atf6a^{-/-} mice are histologically normal

To date, the phenotypes reported for *Atf6a*^{-/-} mice or fish have been associated with exogenous challenges that are likely to compromise ER function and elicit *bona fide* ER stress [23,24,25,26,27]. While *Atf6a*^{-/-} mice are viable, fertile, and grossly normal, it had not previously been reported whether deletion of ATF6 α compromises the integrity of any tissue. Thus, we collected multiple tissues from wild-type and age-matched *Atf6a*^{-/-} mice of both genders and of varying ages with an emphasis on older (>1 year) animals, in which a modest phenotype might be most likely to be manifested. These tissues—skeletal muscle, brown and white adipose, brain, kidney, lung, pancreas, heart, spleen, and liver—were fixed, paraffin-embedded, stained, and analyzed by a veterinary pathologist (A.K.O.) who was blinded to genotype. In this analysis, no systematic genotype-dependent abnormalities of any sort were observed in any tissue examined. Although multiple regions of each tissue were analyzed, here we show representative images demonstrating that the structural organization of each of these tissues is intact in *Atf6a*^{-/-} mice (Figure 1). While these findings do not exclude the possibility of phenotypes for *Atf6a*^{-/-} mice in tissues not examined, or by non-histological criteria, they are in clear contrast to the effects of manipulation of other UPR pathways on secretory tissues such as pancreas and liver [11,13,37]. Taken together with the susceptibility of *Atf6a*^{-/-} animals to various exogenous challenges, this result suggests that ATF6 α is necessary to protect against superphysiological stressors but is dispensable for normal physiological maintenance. Therefore, to whatever extent normal physiological processes elicit ER stress, the remaining limbs of the UPR are sufficient to protect tissue structure and function.

Embryonic lethality upon deletion of ATF6 α and p58^{IPK}

Given that the most dramatic metabolic phenotypes in otherwise viable UPR-compromised animals are observed only upon challenge, we reasoned that a phenotype might be elicited in *Atf6a*^{-/-} mice by intercrossing them with mice lacking the ER cochaperone p58^{IPK}/ERDJ6. We predicted that deletion of p58^{IPK} would be a sufficient stimulus for eliciting ER stress and hence a phenotype in *Atf6a*^{-/-} animals in whatever tissues p58^{IPK} is important during normal physiology.

To our surprise, homozygous deletion of both genes resulted in synthetic embryonic lethality prior to embryonic day 14.5. Intercrosses among animals heterozygous for both genes yielded no *Atf6a*^{-/-}; *p58*^{-/-} animals among 255 weaned progeny (Figure 2A). Intercrosses among animals homozygous for deletion of one gene and heterozygous for the other similarly produced no *Atf6a*^{-/-}; *p58*^{-/-} animals at weaning (Figure 2B). To exclude the possibility that double knockout animals were born but then died prior to weaning, timed intercrosses were established using various combinations of heterozygous and homozygous animals (Table S1), with pups analyzed at embryonic day 14.5. We obtained a sufficient number of embryos to conclude that embryonic lethality of *Atf6a*^{-/-}; *p58*^{-/-} animals occurs prior to E14.5 (Figure 2C). We did recover one double-nullizygous pup from among two litters of E9.5 embryos (Table S1), which was in the process of being reabsorbed and which provided only enough embryonic material for genotyping.

We next tested whether heterozygosity for $p58^{IPK}$ would be sufficient to elicit a phenotype in $Atf6\alpha^{-/-}$ animals. We found that $Atf6\alpha^{-/-}; p58^{+/+}$ animals were grossly normal, were comparably fertile to wild-type animals, and displayed no overt tissue pathology. Histological examination of the pancreas, kidney, and liver—tissues in which a basal or stress-induced phenotype in $p58^{-/-}$ animals has been reported [24,35,38]—of these animals was also normal (data not shown).

We then examined the consequence of $p58^{IPK}$ heterozygosity in an $Atf6\alpha^{-/-}$ background in the liver. We chose the liver because $p58^{IPK}$ is expressed at the highest levels in that tissue [31,39] and because of our interest in the relationship between ER stress and fatty liver—also known as hepatic steatosis. As exogenous ER stress in the liver elicits dysregulation in hepatic lipid metabolism and consequent fatty liver that is dramatically exacerbated in $Atf6\alpha^{-/-}$ mice [24,40], we hypothesized that loss of one allele of $p58^{IPK}$ might be a sufficient ER stress—in the sensitizing context of loss of ATF6 α —to lead to hepatic lipid accumulation. However, in contrast to the stark steatosis seen upon challenge of even wild-type animals with the ER stress-inducing agent tunicamycin (TM), animals lacking both alleles of $Atf6\alpha$ and one allele of $p58^{IPK}$ did not show evidence for accumulated hepatic lipid droplets, as evidenced by immunostaining for the droplet marker protein ADRP (Figure 3A). Similarly, while exogenous ER stress causes a pronounced suppression of lipid metabolic genes in $Atf6\alpha^{-/-}$ animals [24,40,41], $Atf6\alpha^{-/-}; p58^{+/+}$ animals showed no significant changes in the basal expression of such genes compared to $Atf6\alpha^{-/-}; p58^{+/+}$ animals, nor in expression of the UPR target genes *Bip* and *Chop* (Figure 3B). Loss of one allele of $p58^{IPK}$ also did not further sensitize $Atf6\alpha^{-/-}$ animals to ER stress in the liver; these animals did not evince altered *Xbp1* mRNA splicing nor enhanced upregulation of *Bip* and *Chop* upon challenge with TM (Figure 3C, D).

Discussion

The embryonic lethality of $Atf6\alpha^{-/-}; p58^{-/-}$ animals suggests that $p58^{IPK}$ plays a role in embryonic development, the loss of which can be compensated by ATF6 α . Several possibilities could account for this lethality: $p58^{IPK}$ might facilitate the folding of one or a small number of developmentally important proteins, and if its deletion induces ER stress, then ATF6 α might compensate by upregulating other chaperoning pathways, including those mediated by other ER DnaJ proteins such as ERDJ3 and ERDJ4 [42,43]. Alternatively, $p58^{IPK}$ might contribute in a less specific way to the overall efficiency of protein folding in one or more tissues; and in the absence of ATF6 α , loss of this contribution might result in unameliorated ER stress and either tissue malfunction (e.g., failure to secrete important developmental morphogens) or insurmountable levels of ER stress-induced apoptosis, which would be enhanced when the UPR is compromised. It is also possible that loss of ATF6 α produces ER dysfunction and increases the load on the ER, and that an ER lacking $p58^{IPK}$ is ill-equipped to handle such increased load—i.e., that ATF6 α acts upstream of $p58^{IPK}$ in inducing ER dysfunction rather than the other way around. However, given that there is as yet no evidence that loss of ATF6 α alone induces ER dysfunction in any tissue or cell type yet studied, here or elsewhere, we think this possibility is unlikely. We also cannot formally exclude that the developmental role of $p58^{IPK}$ is exerted through putative regulation of PERK or PKR, but the absence of evidence for detectable cytosolic $p58^{IPK}$ militates against this possibility.

The site of $p58^{IPK}$ embryonic function might be in the nascent pancreas, the development of which takes place from E8.5-E14.5 [44]. However, the pattern of expression of $p58^{IPK}$ in the embryo is unknown, and in the adult it is ubiquitously expressed, suggesting that it might also function broadly in the embryo. Ultimately, a systematic series of timed intercrosses

will be necessary to better understand the timing of the lethality, the tissue or tissues affected, and the molecular mechanisms likely at work.

This work is the first to our knowledge to combine deletion of an ER chaperone or cochaperone with that of a UPR signaling molecule in a vertebrate. A key conclusion from the synthetic embryonic lethality of *Atf6a*^{-/-}; *p58*^{-/-} animals is that the UPR in general, and the ATF6 α pathway in particular, can compensate for ER chaperone insufficiency and reverse a protein folding defect. Although it is possible that ATF6 α accomplishes this by regulating other processes that indirectly improve ER function and so reduce the requirement for p58-dependent folding, the fact that p58^{IPK} functions as a cochaperone and that ATF6 α is known to regulate chaperone and cochaperone expression makes ATF6 α -dependent upregulation of compensatory folding pathways the most likely explanation for survival of *Atf6a*^{+/-}; *p58*^{-/-} animals. While this conclusion at first glance seems self-evident, both pharmacological (e.g., tunicamycin, thapsigargin...) and physiological (obesity, exercise...) inducers of ER stress perturb cellular function in many ways, and it is conceivable that the UPR could protect cellular function by other means—for example through regulation of protein synthesis rates, calcium homeostasis, glycosylation efficiency, etc. rather than by a UPR pathway that predominantly regulates the expression of chaperones. Thus, while *Atf6a*^{-/-} animals have no apparent basal phenotype, compounding their deletion with that of p58^{IPK} elucidates physiological roles for UPR-regulated protein folding pathways.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank K. E. Berquam-Vrieze for assistance with timed mating experiments and the Tootle and Cornell labs in the Department of Anatomy and Cell Biology at the University of Iowa for sharing equipment. This work was funded by grant R01 DK084058 from the National Institute of Diabetes and Digestive and Kidney Disorders to D.T.R.; D.D.M. was supported by an Institutional Postdoctoral Training Grant from the University of Iowa Cardiovascular Center, and J.A.G. was supported by an Institutional Predoctoral T32 Training Grant from the University of Iowa Department of Pharmacology. Funders had no role in study design, data collection or analysis, manuscript preparation, or decision to submit. J.A.G. and D.T.R. conceived and designed the experiments. J.A.G., H.M.T., and D.D.M. performed the experiments. A.K.O. analyzed tissue specimens for pathology. J.A.G. and D.T.R. analyzed data. D.T.R. wrote the paper.

References

- [1]. Rutkowski DT, Kaufman RJ. A trip to the ER: coping with stress. *Trends Cell Biol.* 2004; 14:20–28. [PubMed: 14729177]
- [2]. Ron D, Walter P. Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol.* 2007; 8:519–529. [PubMed: 17565364]
- [3]. Lee K, Tirasophon W, Shen X, Michalak M, Prywes R, Okada T, Yoshida H, Mori K, Kaufman RJ. IRE1-mediated unconventional mRNA splicing and S2P-mediated ATF6 cleavage merge to regulate XBP1 in signaling the unfolded protein response. *Genes Dev.* 2002; 16:452–466. [PubMed: 11850408]
- [4]. Shen X, Ellis RE, Lee K, Liu CY, Yang K, Solomon A, Yoshida H, Morimoto R, Kurnit DM, Mori K, Kaufman RJ. Complementary signaling pathways regulate the unfolded protein response and are required for *C. elegans* development. *Cell.* 2001; 107:893–903. [PubMed: 11779465]
- [5]. Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell.* 2001; 107:881–891. [PubMed: 11779464]

- [6]. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D. Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol Cell*. 2000; 6:1099–1108. [PubMed: 11106749]
- [7]. Harding HP, Zhang Y, Ron D. Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature*. 1999; 397:271–274. [PubMed: 9930704]
- [8]. Shi Y, Vattem KM, Sood R, An J, Liang J, Stramm L, Wek RC. Identification and characterization of pancreatic eukaryotic initiation factor 2 alpha-subunit kinase, PEK, involved in translational control. *Mol Cell Biol*. 1998; 18:7499–7509. [PubMed: 9819435]
- [9]. Ye J, Rawson RB, Komuro R, Chen X, Dave UP, Prywes R, Brown MS, Goldstein JL. ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Mol Cell*. 2000; 6:1355–1364. [PubMed: 11163209]
- [10]. Yoshida H, Okada T, Haze K, Yanagi H, Yura T, Negishi M, Mori K. ATF6 activated by proteolysis binds in the presence of NF-Y (CBF) directly to the cisacting element responsible for the mammalian unfolded protein response. *Mol Cell Biol*. 2000; 20:6755–6767. [PubMed: 10958673]
- [11]. Reimold AM, Etkin A, Clauss I, Perkins A, Friend DS, Zhang J, Horton HF, Scott A, Orkin SH, Byrne MC, Grusby MJ, Glimcher LH. An essential role in liver development for transcription factor XBP-1. *Genes Dev*. 2000; 14:152–157. [PubMed: 10652269]
- [12]. Zhang K, Wong HN, Song B, Miller CN, Scheuner D, Kaufman RJ. The unfolded protein response sensor IRE1alpha is required at 2 distinct steps in B cell lymphopoiesis. *J Clin Invest*. 2005; 115:268–281. [PubMed: 15690081]
- [13]. Lee AH, Chu GC, Iwakoshi NN, Glimcher LH. XBP-1 is required for biogenesis of cellular secretory machinery of exocrine glands. *Embo J*. 2005; 24:4368–4380. [PubMed: 16362047]
- [14]. Harding HP, Zeng H, Zhang Y, Jungries R, Chung P, Plesken H, Sabatini DD, Ron D. Diabetes mellitus and exocrine pancreatic dysfunction in perk^{-/-} mice reveals a role for translational control in secretory cell survival. *Mol Cell*. 2001; 7:1153–1163. [PubMed: 11430819]
- [15]. Lee AH, Scapa EF, Cohen DE, Glimcher LH. Regulation of hepatic lipogenesis by the transcription factor XBP1. *Science*. 2008; 320:1492–1496. [PubMed: 18556558]
- [16]. Oyadomari S, Harding HP, Zhang Y, Oyadomari M, Ron D. Dephosphorylation of translation initiation factor 2alpha enhances glucose tolerance and attenuates hepatosteatosis in mice. *Cell Metab*. 2008; 7:520–532. [PubMed: 18522833]
- [17]. So JS, Hur KY, Tarrío M, Ruda V, Frank-Kamenetsky M, Fitzgerald K, Koteliensky V, Lichtman AH, Iwakoshi T, Glimcher LH, Lee AH. Silencing of Lipid Metabolism Genes through IRE1alpha-Mediated mRNA Decay Lowers Plasma Lipids in Mice. *Cell Metab*. 2012; 16:487–499. [PubMed: 23040070]
- [18]. Wang S, Chen Z, Lam V, Han J, Hassler J, Finck BN, Davidson NO, Kaufman RJ. IRE1alpha-XBP1s Induces PDI Expression to Increase MTP Activity for Hepatic VLDL Assembly and Lipid Homeostasis. *Cell Metab*. 2012; 16:473–486. [PubMed: 23040069]
- [19]. Wu J, Rutkowski DT, Dubois M, Swathirajan J, Saunders T, Wang J, Song B, Yau GD, Kaufman RJ. ATF6alpha optimizes long-term endoplasmic reticulum function to protect cells from chronic stress. *Dev Cell*. 2007; 13:351–364. [PubMed: 17765679]
- [20]. Yamamoto K, Sato T, Matsui T, Sato M, Okada T, Yoshida H, Harada A, Mori K. Transcriptional induction of mammalian ER quality control proteins is mediated by single or combined action of ATF6alpha and XBP1. *Dev Cell*. 2007; 13:365–376. [PubMed: 17765680]
- [21]. Adachi Y, Yamamoto K, Okada T, Yoshida H, Harada A, Mori K. ATF6 Is a transcription factor specializing in the regulation of quality control proteins in the endoplasmic reticulum. *Cell Struct Funct*. 2008; 33:75–89. [PubMed: 18360008]
- [22]. Ishikawa T, Okada T, Ishikawa-Fujiwara T, Todo T, Kamei Y, Shigenobu S, Tanaka M, Saito TL, Yoshimura J, Morishita S, Toyoda A, Sakaki Y, Taniguchi Y, Takeda S, Mori K. ATF6alpha/beta-mediated adjustment of ER chaperone levels is essential for development of the notochord in medaka fish. *Mol Biol Cell*. 2012; 24:1387–1395. [PubMed: 23447699]
- [23]. Cao SS, Zimmermann EM, Chuang BM, Song B, Nwokoye A, Wilkinson JE, Eaton KA, Kaufman RJ. The Unfolded Protein Response and Chemical Chaperones Reduce Protein Misfolding and Colitis in Mice. *Gastroenterology*. 2013

- [24]. Rutkowski DT, Wu J, Back SH, Callaghan MU, Ferris SP, Iqbal J, Clark R, Miao H, Hassler JR, Fornek J, Katze MG, Hussain MM, Song B, Swathirajan J, Wang J, Yau GD, Kaufman RJ. UPR pathways combine to prevent hepatic steatosis caused by ER stress-mediated suppression of transcriptional master regulators. *Dev Cell*. 2008; 15:829–840. [PubMed: 19081072]
- [25]. Wu J, Ruas JL, Estall JL, Rasbach KA, Choi JH, Ye L, Bostrom P, Tyra HM, Crawford RW, Campbell KP, Rutkowski DT, Kaufman RJ, Spiegelman BM. The unfolded protein response mediates adaptation to exercise in skeletal muscle through a PGC-1alpha/ATF6alpha complex. *Cell Metab*. 2011; 13:160–169. [PubMed: 21284983]
- [26]. Egawa N, Yamamoto K, Inoue H, Hikawa R, Nishi K, Mori K, Takahashi R. The endoplasmic reticulum stress sensor, ATF6alpha, protects against neurotoxin-induced dopaminergic neuronal death. *J Biol Chem*. 2011; 286:7947–7957. [PubMed: 21131360]
- [27]. Usui M, Yamaguchi S, Tanji Y, Tominaga R, Ishigaki Y, Fukumoto M, Katagiri H, Mori K, Oka Y, Ishihara H. Atf6alpha-null mice are glucose intolerant due to pancreatic beta-cell failure on a high-fat diet but partially resistant to diet-induced insulin resistance. *Metabolism*. 2012; 61:1118–1128. [PubMed: 22386934]
- [28]. Lee TG, Tang N, Thompson S, Miller J, Katze MG. The 58,000-dalton cellular inhibitor of the interferon-induced double-stranded RNA-activated protein kinase (PKR) is a member of the tetratricopeptide repeat family of proteins. *Mol Cell Biol*. 1994; 14:2331–2342. [PubMed: 7511204]
- [29]. van Huizen R, Martindale JL, Gorospe M, Holbrook NJ. P58IPK, a novel endoplasmic reticulum stress-inducible protein and potential negative regulator of eIF2alpha signaling. *J Biol Chem*. 2003; 278:15558–15564. [PubMed: 12601012]
- [30]. Yan W, Frank CL, Korth MJ, Sopher BL, Novoa I, Ron D, Katze MG. Control of PERK eIF2alpha kinase activity by the endoplasmic reticulum stress-induced molecular chaperone P58IPK. *Proc Natl Acad Sci U S A*. 2002; 99:15920–15925. [PubMed: 12446838]
- [31]. Rutkowski DT, Kang SW, Goodman AG, Garrison JL, Taunton J, Katze MG, Kaufman RJ, Hegde RS. The role of p58IPK in protecting the stressed endoplasmic reticulum. *Mol Biol Cell*. 2007; 18:3681–3691. [PubMed: 17567950]
- [32]. Petrova K, Oyadomari S, Hendershot LM, Ron D. Regulated association of misfolded endoplasmic reticulum luminal proteins with P58/DNAJc3. *Embo J*. 2008; 27:2862–2872. [PubMed: 18923430]
- [33]. Tao J, Petrova K, Ron D, Sha B. Crystal structure of P58(IPK) TPR fragment reveals the mechanism for its molecular chaperone activity in UPR. *J Mol Biol*. 2010; 397:1307–1315. [PubMed: 20184891]
- [34]. Qiu XB, Shao YM, Miao S, Wang L. The diversity of the DnaJ/Hsp40 family, the crucial partners for Hsp70 chaperones. *Cell Mol Life Sci*. 2006; 63:2560–2570. [PubMed: 16952052]
- [35]. Ladiges WC, Knoblaugh SE, Morton JF, Korth MJ, Sopher BL, Baskin CR, Macauley A, Goodman AG, Leboeuf RC, Katze MG. Pancreatic {beta}-Cell Failure and Diabetes in Mice With a Deletion Mutation of the Endoplasmic Reticulum Molecular Chaperone Gene P58IPK. *Diabetes*. 2005; 54:1074–1081. [PubMed: 15793246]
- [36]. Chikka MR, McCabe DD, Tyra HM, Rutkowski DT. C/EBP homologous protein (CHOP) contributes to suppression of metabolic genes during endoplasmic reticulum stress in the liver. *J Biol Chem*. 2013; 288:4405–4415. [PubMed: 23281479]
- [37]. Harding HP, Zhang Y, Bertolotti A, Zeng H, Ron D. Perk is essential for translational regulation and cell survival during the unfolded protein response. *Mol Cell*. 2000; 5:897–904. [PubMed: 10882126]
- [38]. Datta R, Shah GN, Rubbelke TS, Waheed A, Rauchman M, Goodman AG, Katze MG, Sly WS. Progressive renal injury from transgenic expression of human carbonic anhydrase IV folding mutants is enhanced by deficiency of p58IPK. *Proc Natl Acad Sci U S A*. 2010; 107:6448–6452. [PubMed: 20308551]
- [39]. Korth MJ, Lyons CN, Wambach M, Katze MG. Cloning, expression, and cellular localization of the oncogenic 58-kDa inhibitor of the RNA-activated human and mouse protein kinase. *Gene*. 1996; 170:181–188. [PubMed: 8666242]

- [40]. Arensdorf AM, Dezwaan McCabe D, Kaufman RJ, Rutkowski DT. Temporal clustering of gene expression links the metabolic transcription factor HNF4alpha to the ER stress-dependent gene regulatory network. *Front Genet.* 2013; 4:188. [PubMed: 24069029]
- [41]. Yamamoto K, Takahara K, Oyadomari S, Okada T, Sato T, Harada A, Mori K. Induction of liver steatosis and lipid droplet formation in ATF6alpha-knockout mice burdened with pharmacological endoplasmic reticulum stress. *Mol Biol Cell.* 2010; 21:2975–2986. [PubMed: 20631254]
- [42]. Shen Y, Hendershot L. ERdj3, a Stress-inducible Endoplasmic Reticulum DnaJ Homologue, Serves as a CoFactor for BiP's Interactions with Unfolded Substrates. *Mol Biol Cell.* 2005; 16:40–50. [PubMed: 15525676]
- [43]. Lai CW, Otero JH, Hendershot LM, Snapp E. ERdj4 protein is a soluble endoplasmic reticulum (ER) DnaJ family protein that interacts with ER-associated degradation machinery. *J Biol Chem.* 2012; 287:7969–7978. [PubMed: 22267725]
- [44]. Jorgensen MC, Ahnfelt-Ronne J, Hald J, Madsen OD, Serup P, Hecksher-Sorensen J. An illustrated review of early pancreas development in the mouse. *Endocr Rev.* 2007; 28:685–705. [PubMed: 17881611]

Highlights

- The ER stress sensing molecule ATF6 α is dispensable for normal tissue organization
- Heterozygosity for p58^{IPK} does not sensitize *Atf6 α* ^{-/-} animals to hepatic ER stress
- Combined homozygous deletion of ATF6 α and the ER cochaperone p58^{IPK} leads to embryonic lethality
- An intact UPR can compensate for genetic disruption of ER protein folding *in vivo*




Figure 1.
 No histological abnormalities in *Atf6α*^{-/-} mice.
 The indicated tissues were stained with hematoxylin and eosin and examined for histological signs of pathology by a veterinary pathologist who was blinded to genotype. Multiple regions of each organ were examined. Representative images of the indicated tissues are shown, all from 17 month-old female mice. Results from male mice and from younger female mice were similar. Scale bar = 50 μm.

A *double het x double dhet*

		Observed			Expected			
		<i>p58</i>			<i>p58</i>			
<i>Atf6</i>		<i>W</i>	<i>H</i>	<i>K</i>		<i>W</i>	<i>H</i>	<i>K</i>
	<i>W</i>	12	46	14	<i>W</i>	16	32	16
	<i>H</i>	30	82	27	<i>H</i>	32	64	32
	<i>K</i>	14	30	0	<i>K</i>	16	32	16

$p < 0.01$

B *Atf6 α +/-; p58-/- x Atf6 α -/-; p58+/-*

		Observed			Expected			
		<i>p58</i>			<i>p58</i>			
<i>Atf6</i>		<i>W</i>	<i>H</i>	<i>K</i>		<i>W</i>	<i>H</i>	<i>K</i>
	<i>W</i>	0	0	0	<i>W</i>	0	0	0
	<i>H</i>	0	34	40	<i>H</i>	0	26	26
	<i>K</i>	0	30	0	<i>K</i>	0	26	26

$p < 0.001$

C *Timed intercrosses E14.5*

		Observed			Expected			
		<i>p58</i>			<i>p58</i>			
<i>Atf6</i>		<i>W</i>	<i>H</i>	<i>K</i>		<i>W</i>	<i>H</i>	<i>K</i>
	<i>W</i>	2	7	2	<i>W</i>	1	3	2
	<i>H</i>	6	15	11	<i>H</i>	3	13	10
	<i>K</i>	0	8	0	<i>K</i>	2	10	7

$p = 0.0301$

Figure 2.

Combined deletion of ATF6 α and p58^{IPK} results in embryonic lethality.

(A) *Atf6 α +/-; p58^{IPK}+/-* animals were intercrossed, and the genotypes of the progeny from multiple litters were recorded. W = wild-type; H = heterozygous; K = knockout. The observed genotype frequencies are given in the table on the left, while the expected frequencies are in the table on the right. P-value was determined by Chi-square analysis. Total animal numbers do not correspond because of rounding for the expected group.

(B) Same as (A), except *Atf6 α +/-; p58-/-* males were crossed to *Atf6 α -/-; p58+/-* females and vice-versa.

(C) Animals of various combinations of *Atf6 α* and *p58* genotypes were subjected to timed intercross, and embryos were harvested and genotyped at embryonic day 14.5. The observed and expected preponderances of each genotype from all crosses are shown. The outcome of each cross is given in Table S1.




Figure 3.

Heterozygosity for $p58^{IPK}$ does not sensitize *Atf6α-/-* animals to ER stress in the liver.

(A) Liver sections from animals of the indicated genotypes were examined for lipid droplet accumulation by immunostaining for ADRP. For comparative purposes, the increase in ADRP immunostaining seen upon challenge of wild-type animals with 1 mg/kg TM for 18h from a separate experiment is shown.

(B) Littermate *Atf6α-/-; p58+/+* and *Atf6α-/-; p58+/-* animals of both genders were sacrificed and the expression of UPR target genes and metabolic genes in the liver was determined by qRT-PCR. None of the genes showed statistically significant differences in expression between genotypes.

(C) *Atf6α-/-; p58+/+* and *Atf6α-/-; p58+/-* animals were challenged with either vehicle or 250 μg/kg TM. 16 hours after treatment RNA was prepared from liver samples and *Xbp1* mRNA splicing was assessed by conventional RT-PCR. Unspliced (us) and spliced (spl) forms are indicated. Image is shown in black-to-white inverted form for greater visual clarity.

(D) Same as (C), except the expression of *Bip* and *Chop* mRNA was determined by qRT-PCR from liver samples. Differences in expression between genotypes upon TM treatment were not statistically significant.