



Botanical ethnoveterinary therapies in three districts of the Lesser Himalayas of Pakistan

Abbasi *et al.*



RESEARCH

Open Access

Botanical ethnoveterinary therapies in three districts of the Lesser Himalayas of Pakistan

Arshad Mehmood Abbasi^{1*}, Shujaul Mulk Khan², Mushtaq Ahmad³, Mir Ajab Khan³, Cassandra Leah Quave^{4,5} and Andrea Pieroni⁶

Abstract

Background: Ethnoveterinary knowledge is highly significant for persistence of traditional community-based approaches to veterinary care. This is of particular importance in the context of developing and emerging countries, where animal health (that of livestock, especially) is crucial to local economies and food security. The current survey documents the traditional veterinary uses of medicinal plants in the Lesser Himalayas-Pakistan.

Methods: Data were collected through interviews, focus groups, participant observation, and by administering questionnaires. A total of 105 informants aged between 20–75 years old who were familiar with livestock health issues (i.e. farmers, shepherds, housewives and herbalists) participated in the study.

Results: A total of 89 botanical taxa, belonging to 46 families, were reported to have ethnoveterinary applications. The most quoted families were Poaceae (6 taxa), Fabaceae (6), Asteraceae (5), and Polygonaceae (5). *Adhatoda vasica* was the most cited species (43%), followed by *Trachyspermum ammi* (37%), and *Zanthoxylum armatum* var. *armatum* (36%). About 126 medications were recorded against more than 50 veterinary conditions grouped into seven categories. The highest cultural index values were recorded for *Trachyspermum ammi*, *Curcuma longa*, *Melia azedarach*, *Zanthoxylum armatum* var. *armatum* and *Adhatoda vasica*. The highest informant consensus factor was found for pathologies related to respiratory and reproductive disorders. Comparison with the local plant-based remedies used in human folk medicine revealed that many of remedies were used in similar ways in local human phytotherapy. Comparison with other field surveys conducted in surrounding areas demonstrated that approximately one-half of the recorded plants uses are novel to the ethnoveterinary literature of the Himalayas.

Conclusion: The current survey shows a remarkable resilience of ethnoveterinary botanical knowledge in the study area. Most of the species reported for ethnoveterinary applications are wild and under threat. Thus, not only is it imperative to conserve traditional local knowledge of folk veterinary therapies for bio-cultural conservation motives, but also to assist with *in-situ* and *ex-situ* environmental conservation initiatives, which are urgently needed. Future studies that focus on the validation of efficacy of these ethnoveterinary remedies can help to substantiate emic concepts regarding the management of animal health care and for rural development programs.

Keywords: Medicinal plants, Ethnobotany, Ethnoveterinary, Lesser Himalayas, Pakistan

Introduction

Ethnoveterinary medicine is a broad field encompassing people's beliefs, skills, knowledge and practices related to veterinary health care [1]. Medicinal plants traditionally used in the treatment of animal diseases play a crucial role in local health modalities. Specifically, phytotherapeutics

often represent the primary form of therapy in rural veterinary care as allopathic modalities remain inaccessible, especially in the developing world [2]. Therefore, local knowledge of ecological resources for veterinary care is of particular importance to pastoral and agro-pastoral communities that rely heavily on livestock for their livelihood and food security. However, traditional ethnoveterinary knowledge is still mainly orally transmitted from generation to generation (i.e., in the form of traditional remedies, poems, drawings stories, folk myths, proverbs and

* Correspondence: amabbasi@ciit.net.pk

¹Department of Environmental Sciences, COMSATS Institute of Information Technology, Abbottabad 22060, Pakistan

Full list of author information is available at the end of the article

songs). Due to the nature of oral transmission, this form of local knowledge remains fragile and threatened, and presents an urgent need for being recorded and documented.

An increasing number of studies have very recently focused on the documentation of local ethnoveterinary practices in South Asia [3-24]. These studies hold potential for having a tremendous impact on the Himalayan region, in particular, where efforts for sustaining endogenous development and ultimately improving the health and well-being of both animals and humans is still largely neglected. Pakistan has a very large livestock population composed of a number of local breeds that are well adapted to local conditions. In particular, there are an estimated 27 million buffaloes, 30 million cattle, 27 million sheep, 54 million goats, one million camels, 0.3 million horses, 4 million asses, 0.2 million mules and 74 million poultry in Pakistan [25].

The objectives of this field study were multifold: 1. to record the local knowledge related to medicinal plants used for treating animal diseases in the Lesser Himalayan region in Pakistan; 2. to compare the collected data with the traditional medical knowledge devoted to humans in the same region; 3. to compare the collected data with those of other ethnoveterinary studies conducted in the Himalayan region over the last decades; 4. to assess their cultural importance and the consensus among the informants regarding cited veterinary pathologies; and 5. to examine local perceptions of factors that threaten wild medicinal plant resources.

Materials and methods

Study site

An ethnobotanical study was conducted from March 2010 to April 2013 in different locations of the Lesser Himalayas, which is a hotspot for plant biodiversity in Pakistan. Fifty-five localities in three districts (Haripur, Abbottabad and Mansehra) within the Khyber Pakhtunkhwa (KPK) province were selected for inclusion in the study (Figure 1). The Lesser Himalayan range in Pakistan lies between 33°-44' and 35°-35' north latitude and between 72°-33' and 74°-05' east longitude, comprising an area of 23,295 km². The climate of the area is subtropical in the lowland plains and foot-hills zone and subtropical-sub alpine in middle Himalayas, Siwalik, Murree and entire Hazara hills. The average rainfall varies from 70–90 mm in southern and 100–130 mm in the northern parts. The vegetation of the Lesser Himalayas falls within the subtropical, temperate, sub-alpine and alpine zones. The region is divided into six vegetation zones, namely: the subtropical sub-humid zone, the subtropical humid zone, the temperate humid zone, the sub alpine zone and the zone of the glaciers/snowfields. This area is populated by several ethnic groups (Syed, Abbasi, Karaal, Jadoon, Tanoli, Ghakar, Gujar, and Awan), all speaking the

Hindko dialect of the Western Punjabi, and belonging in turn to the Indo-Aryan (Indic) language family spoken in Northern Pakistan.

Ethnobotanical data collection

Ethnobotanical surveys were conducted in all four seasons. Participatory rural appraisal (PRA) approaches were adopted during fieldwork and prior informed consent was obtained before conducting interviews. Information regarding ethnoveterinary practices was collected through semi-structured interviews and guided fieldtrips with the help of traditional healers. A total of 105 informants (75 males and 30 females), ranging from 20–75 years old and including farmers, shepherds, housewives and herbalists familiar with livestock problems and use of conventional recipes, were interviewed and their responses recorded in detail.

Information regarding the vernacular plant names, part (s) used, methods of preparation, mode/route of application and treated diseases were documented during each interview. Taxonomic identification of the collected plant samples was carried out with the help of Flora of Pakistan [26], The Plant List [27] and by one of the authors (MAK, plant taxonomist). Family nomenclature follows the Angiosperm Phylogeny Group III designations [28]. Additionally, 15 key informants were selected at four locations within three study districts (Haripur, Abbottabad and Mansehra) and specific information regarding the perceived threats for the local medicinal flora was obtained. Following identification of 5 key perceived threats (agricultural land expansion, overharvesting, overgrazing, fuel, fire), we employed pair-wise ranking techniques in which respondents were presented with two threats and chose one from the two threats at a time [29]. Respondent scores were then summed up and ranks for each threat determined by region.

Data analysis

Cultural importance index (CI) values for each species and mean cultural importance values for each family (mCI_f) were calculated as described in a previous quantitative ethnobotanical work [30]. Briefly, CI values of species were calculated based on previously described methods [31] and express the sum of the proportion of informants that mention each species used. The CI values for each species were calculated using the following formula, with UR_i: use reports in each use-category and N: total number of survey participants:

$$CI = \sum \frac{UR_i}{N}$$

Moreover, we calculated the mean cultural importance (mCI) index of plant species as measured in three study districts (Haripur, Abbottabad, Mansehra) within the Khyber Pkahtunkhaw Province of Pakistan, on the basis



of their cultural importance index (CI) calculated for each single district. To calculate the mCif, CI values of all reported species within a family were added. Regression analysis was performed upon comparison of mCif with the number of species in each respective family.

Informant consensus on the reported cures for a given group of ailments was calculated as an informant consensus factor (ICF) [32]. All of the quoted veterinary diseases were grouped into seven categories, which included: gastrointestinal

disorders, skin infections, parasites/worms, fever/cold/respiratory diseases, reproductive disorders, musculoskeletal disorders and galactagogue remedies. As previously reported [32], we used the following formula, with n_{ur} : number of use citations in each category and n_i : number of species used:

$$ICF = \frac{n_{ur} - n_i}{n_{ur} - 1}$$

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan

Botanical name and voucher number	Local name	PU ^a	Preparation and application	Veterinary condition	AT ^b	C ^c	mCl ^d	FR ^e	SU ^f
ACANTHACEAE									
<i>Adhatoda vasica</i> Nees CIITH-2	Bhekker	L	1 kg fresh leaves each of <i>Adhatoda vasica</i> and <i>Rhazya stricta</i> are soaked in water over night and liquid is orally administered for 3–4 days.	Stomach disorder, fever, dehydration	B, C	45	1.975	-	+
			1 kg leaves are mixed with grass or husk and fed to animal for 2–3 days.	Dysentery	B, C, G, Sh				
			½ kg each of leaves of <i>Adhatoda vasica</i> and bulbs of <i>Allium cepa</i> are crushed and paste is fed to animal for 3–4 days.	Indigestion	B, C, G, Sh, H				
			½ kg fresh leaves are crushed along with table salt and resulting paste is fed to animal for 5–6 days.	Diarrhea, dysentery, gas trouble	A, B, C, G, Sh, Cam				
AMARANTHACEAE									
<i>Aerva javanica</i> (Burm. f.) Juss. CIITH-4	Chittibui	Wp	1 kg roots are boiled in water and decoction is given orally twice a day for 7–8 days.	Skin infection	B, C, G, Sh	6	0.276	+	-
			Whole plant boiled in water and decoction is orally administered at night for 2–3 days.	Intestinal worms	A, B, C, H, M				
<i>Amaranthus viridis</i> L. CIITH-10	Chulai	Wp	1 kg fresh plant is crushed and mixed in wheat husk; paste is fed to animal twice a day for 10–15 days.	Against weakness	A, C, Cam, G, M	3	0.648	-	+
<i>Chenopodium album</i> L. CIITH-25	Bathu	L	Leaves are boiled in mustard oil, and then crushed and topically applied to wounds.	Wound healing	B, C, G, H, Sh	3	0.014	-	+
AMARYLLIDACEAE									
<i>Allium cepa</i> L. CIITH-7	Piaz	B	Bulbs are crushed with sugar. The resulting paste is mixed with milk and orally administered at night and early morning up to 1 week.	Galactagogue	B, C	35	1.839	-	+
			½ kg of each leaves of <i>Adhatoda vasica</i> and bulbs of <i>Allium cepa</i> are crushed and this paste is orally administered for 3–4 days.	Indigestion	B, C, G, H, Cam, Sh,				
			½ kg bulbs are crushed along with salt and mixed in flour. This paste is administered orally for 4–5 days.	Stomach disorder, fever	B, C, G,				
<i>Allium jacquemontii</i> Kunth CIITH-8	Jangli-Thoom	B	100 g fresh bulbs are ground and mixed with wheat flour. This paste is applied topically for 10–15 days.	Unequal mammary glands	C, G	2	0.131	+	-
ANACARDIACEAE									
<i>Mangifera indica</i> L. CIITH-51	Aam	F	Pickled fruit is fed to animal for 4–5 days.	Mouth infection	A, B, C, G	3	0.011	-	-

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

APIACEAE									
<i>Anethum sowa</i> Roxb. ex Fleming CIITH-11	Soay	S	100 g seeds are mixed in wheat flour and orally administered for up to 15–20 days.	Galactagogue	B, C	5	1.028	+	-
			100 g each of <i>Anethum sowa</i> , <i>Trachyspermum ammi</i> and <i>Foeniculum vulgare</i> are ground together and paste is orally administered for 2–3 days.	Abdominal pain, swelling	A, C, Cam, G				
<i>Foeniculum vulgare</i> Mill. CIITH-46	Souf	Ap	200 g of aerial parts are boiled in water. The decoction is mixed with <i>Trachyspermum ammi</i> , <i>Camelia sinensis</i> , brown sugar and ghee. This paste is fed to the animal for 5–6 days.	Indigestion	A, B, C, Cam, H, M	22	1.793	-	+
			200 g each of <i>Foeniculum vulgare</i> aerial parts and <i>Punica granatum</i> rind are ground together. The resulting powder is orally administered for 4–5 days.	Diarrhea	B, C, G, H Sh				
<i>Trachyspermum ammi</i> (L.) Sprague CIITH-79	Ajwain	S	¼ kg each of <i>Trachyspermum ammi</i> seeds, <i>Anethum sowa</i> , <i>Allium cepa</i> and <i>Foeniculum vulgare</i> are mixed in flour; paste is orally administered to animals for 10–15 days.	Appetite stimulant, galactagogue	A, B, B, C, Cam, G, H, M, Sh	39	2.317	-	-
APOCYNACEAE									
<i>Calotropis procera</i> (Aiton) W.T. Aiton CIITH-19	Ak	L	Fresh leaves and black salt are fed to animals for 1–2 days.	Mouth and eye watering	C, Cam, G	15	0.125	-	+
				Colic, indigestion	A, C, H, M				
				Pain, inflammation	C, Cam, H, M				
<i>Carissa opaca</i> Stapf ex Haines CIITH-23	Granda	L, S	½ kg fresh leaves and ripened seeds are ground and the resulting powder is mixed in water and orally administered to animals for 2–3 days.	Throat infection	G, Sh	3	0.013	+	+
			100 g dried roots are ground into a powder and sprinkled onto wounds for 2–3 days.	Infected sores, wound healing	B, C, G, Sh				
<i>Periploca aphylla</i> Decne. CIITH-57	Kathi	Ltx	Latex is topically applied for 4–5 days.	Skin infection	A, C, Cam, D, H, Sh	5	0.213	+	+
<i>Rhazya stricta</i> Decne. CIITH-65	Veran	L	Fresh leaves are soaked in water and the resulting liquid is orally administered to animals for 8–10 days.	Skin infection and blood purification	B, C, G, H, Sh	7	0.193	+	-
			A decoction of fresh leaves is orally administered for 2–3 days.	Abdominal pain	A, B, C, G, M				

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

ARACEAE									
<i>Arisaema flavum</i> (Forssk.) Schott CIITH-12	Adbais	S	8-10 ripened seeds are orally administered.	New Castle disease	P	2	0.103	-	+
ARALIACEAE									
<i>Hedera nepalensis</i> K. Koch CIITH-48	Hurr Bumbal	L	200 g fresh leaves are crushed and soaked in water. The resulting liquid is instilled in the nose twice a day for 1–2 days.	To remove leeches	B, C	4	0.131	+	+
ASTERACEAE									
<i>Erigeron</i> sp. CIITH-40	Taku Booti	Wp	500 g of fresh plant material is crushed and mixed with flour; the paste is orally administered for up to a week.	Fever, stomach collapse	B, C, D, G	14	0.031	+	-
<i>Launaea procumbens</i> (Roxb.) Ramayya & Rajagopal CIITH-49	Doodh Pathar	L	Paste of fresh leaves is topically applied for 3–4 days.	External worms (skin infection)	A, B, C, D, M	2	0.314	-	+
<i>Saussurea heteromalla</i> (D. Don) Hand.-Mazz. CIITH-72	Kali Zeri	Wp	50 g seeds are wrapped in paper and fed to animals along with ghee for up to 1 week.	Edema	B, C, H	11	0.874	-	-
<i>Senecio chrysanthemoides</i> DC. CIITH-73	Chitta Hola	Wp	200 g dried plant material is ground and the resulting powder is mixed with kneaded flour; paste is orally administered for 8–10 days. Root decoction is orally administered for 5–6 days.	Blood purification Arthritis	B, C, G, Sh A, C Cam, G, H	4	1.153	-	-
<i>Tagetes minuta</i> L. CIITH-77	Saat Barga	L	Paste of fresh plant is topically applied. Fresh leaves are soaked in water and the resulting liquid is instilled into the ear for 2–3 days.	Sore joints Earache	A, C Cam, G, H A, C, D, G, M	2	0.675	+	+
BERBERIDACEAE									
<i>Berberis lycium</i> Royle CIITH-14	Sumbol	Rt	A decoction of ¼ kg bark is prepared and orally administered for 10–12 days. 100 g bark is ground and powder is sprinkled on wounds up to a week.	Bone fracture Wound healing	A, B, C, G, H, M, Sh, A, B, C, G, H, M	25	1.753	-	+
BORAGINACEAE									
<i>Cordia obliqua</i> Willd. CIITH-32	Lasoor	F	2-3 fruits are mixed in fodder and fed for 4–5 days.	Throat infection, common cold	C, G, Sh	8	0.561	+	-
<i>Trichodesma indicum</i> (L.) Lehm. CIITH-81	Hadusi	Wp	Decoction of seeds is administered orally for 10–15 days. Paste of fresh plant is administered orally for a week.	Stomach ulcer Stomach disorder, intestinal worms	C, G, Sh B, C, G, Sh	22	1.769	-	+

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

BRASSICACEAE									
<i>Brassica campestris</i> L. CIITH-17	Sarain	S	200 g seeds are ground with 50 g of sulfur and mixed with mustard oil. This paste is topically applied for a week.	Skin infection	A, B, C, Cam, D, H	34	1.359	-	+
			100 g seeds are ground, and then powder is mixed with eggs and orally administered for 2–3 days.	Stomach disorder/ infection	B				
			2-3 peppers are soaked in mustard oil for a few days and cooked along with bread, which is fed to animals for 8–10 days.	Eye disease (cornea opacity)	B, C				
<i>Eruca sativa</i> Mill. CIITH-41	Tara Mira	S	200 ml seed oil is mixed with 200 g of sugar orally administered for 4–5 days.	Dysentery	B, C, G, H	12	1.176	+	-
CANNABACEAE									
<i>Cannabis sativa</i> L. CIITH-21	Bhang	L	100 g of fresh leaves are crushed and paste is applied topically.	Leeches, lice	B, C, G, Sh	5	1.348	-	+
			½ kg dried leaves are ground and the resulting powder is mixed with kneaded flour; the paste is fed to animals twice a day for a week.	Appetite stimulant; abdominal swelling	B, C, D, G, H, M				
			L,Fb, S	1 kg of dried leaves, floral buds and seeds are made into a powder and mixed with wheat flour, salt and water. This paste orally administered for 10–15 days.	Indigestion	B, C, D, G, H, M			
CAPPARACEAE									
<i>Capparis decidua</i> (Forssk.) Edgew. CIITH-22	Kirir	Ap	½ kg of fresh aerial parts is boiled in water; 1 glass of resulting decoction is orally administered for 5–6 days.	Stomach gripe, indigestion	C, Cam, H, M	4	0.011	+	-
CONVOLVULACEAE									
<i>Convolvulus arvensis</i> L. CIITH-31	Leli	Wp	1 kg fresh plant is crushed along with sugar and water; this juice is given orally for 3–4 days.	Constipation	B, C, G, Sh	9	0.419	+	+
<i>Cuscuta reflexa</i> Roxb. CIITH-35	NilaTahri	Wp	½ kg plant material is crushed and mixed with flour; paste is applied topically for 8–10 days.	Galactagogue	B, C, G, Sh	5	0.151	-	+
			Paste of fresh plant is fed to goats and sheep for 4–5 days.	Indigestion	G, Sh				
CRASSULACEAE									
<i>Bryophyllum pinnatum</i> (Lam.) Oken CIITH-18	Zakham Hayat	L	Fresh leaves are wormed in mustard oil and bandage on topically on wounds.	Bleeding wounds	C, G, H, M	9	0.086	-	-

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

CUCURBITACEAE									
<i>Citrullus colocynthis</i> (L.) Schrad. CIITH-27	Tumba	F	Juice is extracted by heating the fruit of <i>Citrullus colocynthis</i> and <i>Calotropis procera</i> in a mud pot on a garbage fire for 2–3 weeks; it is orally administered for up to a week.	Indigestion, gas trouble, abdominal worms	B, C, Cam, D, G, H, M, Sh	17	0.514	+	-
			100 g fruit is crushed and mixed in <i>Aloe vera</i> pulp. This paste is orally administered for 2–3 days.	Constipation	B, C, Cam, D, Sh				
<i>Citrullus vulgaris</i> Schrad. CIITH-28	Rainda	F	1 kg fresh fruit coat is ground with salt and orally administered for 10–12 days.	Appetite stimulant, galactagogue	A, C, Cam, G	13	0.417	+	-
<i>Cucumis melo</i> L. CIITH-33	Chibber	L, F	Paste from fresh leaves and fruits are fed to animals for up to 1 week.	Indigestion	B, C, G, Sh	5	0.463	-	-
EUPHORBIACEAE									
<i>Euphorbia wallichii</i> Hook. f. CIITH-43	Hervi	Ltx	50 ml stem latex is topically applied for 2–3 days.	Rashes, wound healing	A, C, Cam, D, G, M	3	0.041	+	+
<i>Mallotus philippensis</i> (Lam.) Müll. Arg. CIITH-50	Kamila	F	¼ kg dried fruits are ground and the resulting powder is fed to animals along with wheat flour for 2–3 days.	Intestinal worms	C, G, Sh	21	1.103	-	+
<i>Ricinus communis</i> L. CIITH-66	Hernoli	S	½ cup of seed oil is orally administered for up to a week.	Constipation	B, C, Cam, G, H, Sh	8	0.052	-	+
FABACEAE									
<i>Acacia nilotica</i> (L.) Willd. ex Delile CIITH-1	Kiker	Bk	Decoction of ½ kg bark is orally administered twice a day for 5–6 days.	Stomach disorder	B, C, H	10	0.686	-	+
<i>Cassia fistula</i> L. CIITH-24	Kinjal	S	4–6 seeds are mixed with chicken feed and fed to hens.	Newcastle disease	P	3	0.413	-	+
<i>Cicer arietinum</i> L. CIITH-26	Kalay	S	200 g seeds are ground and resulting powder is mixed with yogurt; this paste is orally administered for 10–15 days.	Piles	C	2	0.139	-	-
<i>Dalbergia sissoo</i> Roxb. ex DC. CIITH-37	Tahli	L	½ kg fresh leaves of <i>Dalbergia sissoo</i> and 200 g linseeds are boiled in water. This decoction is administered orally for 8–10 days.	Constipation	C, G	3	0.117	-	+
<i>Phyllodium pulchellum</i> (L.) Desv. CIITH-39	Ladan	Rt	Root decoction is administered orally for 10–15 days.	Fever, weakness	A, B, C, H, M	12	0.075	+	+
<i>Trigonella foenum-graecum</i> L. CIITH-82	Mathray	S	200 g seeds are ground and the resulting powder is used orally after washing urethra with a sugar and potash alum (potassium alum) mixture for 4–5 days.	Urethra prolapse	B, C	13	1.541	-	-
			50 g seeds are mixed with fodder and fed to animal for 3–4 days.	Diarrhea	B, C				

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

LAMIACEAE							
<i>Ajuga bracteosa</i> Wall. ex Benth. CIITH-6	Ratti Booti	Wp	125 g shade-dried plant is ground and resulting powder is mixed with flour and orally administered for 2–3 days.	Abdominal pain	B, C, G	3	0.433 - +
LYTHRACEAE							
<i>Punica granatum</i> L. CIITH-63	Druni	Fr	¼ kg dried rind is ground and the resulting powder is fed to animals along with flour up to 1 week.	Dysentery	C, G, Sh	9	1.769 - +
MELIACEAE							
<i>Melia azedarach</i> L. CIITH-52	Dhrek	L	200 g fresh leaves are crushed along with sugar and water; the mixture is administered orally to animals for 2–3 days.	Foot, mouth infection	G, Sh	29	2.101 - +
		Tw	100 g fresh twigs are crushed and soaked in water; the resulting liquid is given orally for 2–3 days.	Skin infections	B, C, G, H		
		Fr	200 g fruit rinds are soaked in water and the resulting juice is given orally for 4–5 days.	Rashes	B, C, G, H		
		L,Tw, F	200 g of fresh leaves, twigs and fruits are crushed. This paste is fed to animals up to a week.	Gas trouble, indigestion	B, C, G, H, Sh		
MORACEAE							
<i>Ficus palmata</i> Forssk. CIITH-45	Phagwar	L, F	½ kg dried leaves and fruits are ground and the resulting powder is administered orally with water for 5–6 days.	Indigestion	C,G	2	0.316 - +
MYRTACEAE							
<i>Eucalyptus camaldulensis</i> Dehnh. CIITH-42	Safada	L	Fresh leaves are fed to animals for 4–5 days.	Common cold	B, C, G	4	0.010 + -
<i>Syzygium cumini</i> (L.) Skeels CIITH-76	Jaman	L	Fresh leaves are fed to animals.	Diarrhea	B, C, G	3	0.010 + -
NITRARIACEAE							
<i>Peganum harmala</i> L. CIITH-56	Hremal	L,Br	Smoke of leaves and branches is used for 4–5 days.	Mastitis	B, C, H	17	1.736 - -
		L	½ kg fresh leaves ground with salt; paste is orally administered for 5–6 days.	Gastric problems	B, C, Cam, H, M		
		S	200 g dried seeds are burnt and mixed in mustard oil. This infusion is applied topically 2–3 days.	Ticks and mites	B, G, Sh		

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

OLEACEAE									
<i>Olea ferruginea</i> Royle. CIITH-54	Kahu	F	Extract of fruits is given orally for 5–6 days.	Indigestion	C, G	3	0.017	+	+
PAPAVERACEAE									
<i>Fumaria indica</i> (Hauskn.) Pugsley CIITH-47	Papra	Wp	Fresh plant material is fed to animals for 2–3 days.	Diarrhea	C, G	6	1.511	-	+
PLANTAGINACEAE									
<i>Plantago lanceolata</i> L. CIITH-58	Batti	L	Leaf paste is topically applied.	Neck rashes	C, H	2	0.173	+	+
<i>Plantago major</i> L. CIITH-59	ChimchipAtra	Wp	200 g dried plant is ground and the resulting powder is sprinkled on infected hooves for 6–7 days.	Infected hooves	C, Cam, G, H, M	3	0.185	-	+
POACEAE									
<i>Cynodon dactylon</i> (L.) Pers. CIITH-36	Khabul	Wp	100 g of fresh plant material is ground and the paste is topically applied for 2–3 days.	Wound healing	A, B, C, D, H, M	7	0.351	-	+
<i>Oryza sativa</i> L. CIITH-55	Chawal	S	1 kg rice is boiled in water along with yoghurt and <i>Eruca sativa</i> oil; paste is fed to animals for 15–20 days.	Weakness, lung infection	B, C, Cam, H, M	6	0.075	-	-
<i>Saccharum bengalense</i> Retz. CIITH-71	Kana	Rt	½ kg roots are boiled along with ½ kg <i>Solanum surattense</i> ; decoction is orally administered for 8–10 days.	Intestinal worms, appetite stimulant	C, G	3	0.010	+	+
<i>Sorghum halepense</i> (L.) Pers. CIITH-75	Baru	Rt	Fresh roots are crushed and soaked in water; the resulting liquid is orally administered for 2–3 days.	Indigestion	D	3	0.011	-	-
<i>Triticum aestivum</i> L. CIITH-83	Kank	S	200 g seed porridge is given orally to animals for a week.	Galactagogue, dysentery	B, C, G	27	1.753	-	+
			Hot bread is fed to cattle for a week.	Mouth sores	C, G, Sh				
			¼ kg seeds are ground with brown sugar and the resulting paste is fed to animal for 8–10 days.	Galactagogue	B, C, G				
<i>Zea mays</i> L. CIITH-88	Maki	Stg	A decoction of the female inflorescence is orally administered for 4–5 days.	Urinary inflammation	B, C, G	15	0.015	-	+
POLYGONACEAE									
<i>Polygonum amplexicaule</i> D. Don. CIITH-60	Mosloon	Rt	Root decoction is orally administered to animals for 1 week.	Dehydration, fever	B, C, G, Sh	2	0.138	+	+
		L	Fresh leaves are fed to animals for 3–4 days.	Indigestion	B, C, G				

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

<i>Polygonum plebeium</i> R. Br. CIITH-61	Sarwar Booti	Wp	Paste of the fresh plant is applied topically for 2–3 days.	Scorpion bite	B, C, D, H	3	0.109	+	-
<i>Rumex dentatus</i> L. CIITH-68	Jngli Palak	Rt	Root decoction is orally administered for up to a week.	Foot, mouth infection	C, G, Cam, H, M	4	0.014	+	+
<i>Rumex hastatus</i> D. Don CIITH-69	Khitiml	Ap	Arial parts are used as a brush for a week.	Scabies	B, C	3	0.915	-	+
		Rt	1 kg of each of <i>Rumex hastatus</i> roots and <i>Quercus incana</i> bark are boiled; decoction is mixed in sugar and flour. This sweet meal is fed to animals for 8–10 days.	Cough, fever, weakness	B, C, G				
<i>Rumex nepalensis</i> Spreng. CIITH-70	Hoola	L	Extract of fresh leaves is topically applied to infected parts for 4–5 days.	Antiseptic and anti-inflammatory	C, G, M, Sh	2	0.013	-	+
			Leaf paste is applied topically for 2–3 days.	Hemostatic	A, B, C, D				
		Rt	½ kg fresh roots are crushed with salt and the resulting paste is administered orally for 5–6 days.	Diarrhea, dysentery, and intestinal worms	G, Sh				
PRIMULACEAE									
<i>Myrsine africana</i> L. CIITH-53	Khukan	L	Fresh leaves are fed to animals for up to a week.	Indigestion, worms	G,Sh	17	0.735	+	+
PTDERIDACEAE									
<i>Adiantum incisum</i> Forssk. CIITH-3	Sarhaaj	L	Paste of crushed leaves is made by mixing with wheat flour and orally administered for 2–3 days.	Abdominal pain	B, C	5	0.336	+	+
RANUNCULACEAE									
<i>Clematis grata</i> Wall. CIITH-30	Dhand	L	Paste of fresh leaves is applied topically on infection sites.	To kill external worms in wounds	A, C, Cam, D, M	17	0.432	+	+
RHAMNACEAE									
<i>Ziziphus nummularia</i> (Burm. f.) Wight & Arn. CIITH-89	Beri	L	100 g leaves are boiled and decoction is orally administered for 1–2 days.	To discharge placenta following birth	B, C	7	0.018	-	+
ROSACEAE									
<i>Prunus persica</i> (L.) Batsch CIITH-62	Aru	L	Juice of fresh leaves is applied topically for 4–5 days.	To kill germs/worms	A, B, C, D, H, M	21	1.783	-	+
			Leaf decoction is orally administered for 2–3 days.	Dysentery	G, Sh				
<i>Pyrus pashia</i> Buch.-Ham. ex D. Don CIITH-64	Batangi	F	Powder of dried fruits is given orally for 5–6 days.	Dysentery, diarrhea	B, C	4	0.241	-	+

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

<i>Rosa cymosa</i> Tratt. CIITH-67	Ghulab	Fl	100 g flowers are soaked in a sugar solution and mixed with milk; mixture is orally administered for 2–3 days at 10–20 days post-delivery.	To clean uterus	B, C	3	0.101	-	+
RUTACEAE									
<i>Citrus limon</i> (L.) Osbeck CIITH-29	Nimbu	F	Fruit juice is mixed with sugar and this paste is fed to animals and applied topically (to the mammary glands) for 10–15 days.	Mastitis	B, C, G	8	0.131	-	-
<i>Zanthoxylum armatum</i> var. <i>armatum</i> CIITH-87	Timer	L	100 g leaves are crushed and mixed with kneaded flour; the resulting paste fed to animals for 3–4 days.	Vomiting, indigestion	B, C, G, Sh	38	2.101	-	+
		S,F	50 g seeds/fruit are orally administered with flour.	Indigestion, mouth infection	B, C, G, Sh				
SAPINDACEAE									
<i>Aesculus indica</i> (Wall. ex Cambess.) Hook. CIITH-5	Bankhor	F	¼ kg fruits are ground and the powder mixed with husk and fed to animals for 5–6 days.	Cough, fever, abdominal pain	A, B, C, Cam, G, H, M	18	1.576	-	+
			Fruit juice is applied topically up to a week.	External wounds	B, C				
SAXIFRAGACEAE									
<i>Bergenia ciliata</i> (Haw.) Sternb. CIITH-15	Batpia	Rh	Fresh leaves are slightly crushed and applied as bandages on bleeding wounds.	Hemostatic (for bleeding wounds)	B, C	15	1.341	-	+
			Dried rhizome powder is sprinkled onto wounds for 8–10 days.	Wound healing	A, B, C, G, D, M				
<i>Bergenia stracheyi</i> (Hook. f. & Thomson) Engl. CIITH-16	Batpia	Rh	Dry root powder is sprinkled onto wounds for 8–10 days.	Wound healing	A, B, C, G, D, M	12	0.785	+	-
SCROPHULARIACEAE									
<i>Verbascum thapsus</i> s. L. CIITH-85	Gidhar Tumaku	Wp	100 g fresh plant is crushed and paste is fed to animal for a week.	Diarrhea	C, D	2	1.135	-	+
SOLANACEAE									
<i>Datura innoxia</i> Mill. CIITH-38	Datura	L	Leaf extract is applied topically for 1–2 days.	Anti-lice	B, C	9	0.543	-	+
<i>Solanum surattense</i> Burm. f. CIITH-74	Mhokri	Wp	Fresh fruit paste is applied topically for 2–3 days.	Wound healing	B, C, G, D, H, M	18	0.984	-	+
			½ kg fresh plant is cooked with salt, peppers and yoghurt. This paste is orally administered for 10–15 days.	Tonic	B, C, G, H, Sh				

Table 1 Botanical ethnoveterinary therapies for treating livestock in communities of the Lesser Himalayas in Pakistan (Continued)

			200 g fresh plant is boiled along with black pepper and salt; decoction is given orally for 8–10 days.	Fever, indigestion, cough	C, G				
<i>Withania somnifera</i> (L.) Dunal CIITH-86	Aksan	Rt	200 g fresh roots are crushed and paste is applied topically up to a week.	Mastitis	B, C, G	8	0.736	-	+
TAMARICACEAE									
<i>Tamarix aphylla</i> (L.) H. Karst. CIITH-78	Rokh	L	An infusion of the dried, burnt leaves is applied topically onto the skin, nose and ear in livestock.	To kill external worms in wounds	A, C, D, H	5	0.011	+	-
THEACEAE									
<i>Camellia sinensis</i> (L.) Kuntze CIITH-20	Chay	L	½ kg fresh leaves are boiled in water along with sugar; 1 glass of this decoction is given orally for 5–6 days.	Fever	C, G	4	0.101	+	+
VITACEAE									
<i>Vitis vinifera</i> L. CIITH-84	Dakh	L, Wd	Ash prepared from wood and leaves of is orally administered with milk for 8–10 days.	Hemoglobinuria	C, G	3	0.191	+	+
XANTHORRHOACEAE									
<i>Aloe vera</i> (L.) Burm. f. CIITH-9	Kwar Gandal	L	½ kg of leaf pulp, salt and <i>Trachyspermum ammi</i> are mixed and paste is administered orally for up to 1 week.	Digestive problems	C, G, H	8	0.175	-	+
<i>Asphodelus tenuifolius</i> Cav. CIITH-13	Bokhat	L	Fresh leaves are crushed and fed to horses for 8–10 days.	Constipation	A, C, Cam, H	3	0.037	+	-
ZINGIBERACEAE									
<i>Curcuma longa</i> L. CIITH-34	Haldi	Rt	Powder from dried roots is applied and wrapped or sprinkled onto wounds for 2–3 days.	Wound healing	A, C, Cam, H	4	2.181	-	-
ZYGOPHYLACEAE									
<i>Fagonia indica</i> Burm. f. CIITH-44	Dhamian	Wp	Fresh leaves are fed to animals for 8–10 days.	Appetite stimulant, indigestion	B, C, Cam, G, H	2	0.015	+	-
<i>Tribulus terrestris</i> L. CIITH-89	Bhakra	L	200 g of each of <i>Tribulus terrestris</i> dried leaves, <i>Curcuma domestica</i> and <i>Foeniculum vulgare</i> are ground together and the resulting powder is orally administered to cattle for 10–15 days.	Appetizer, joint pain	B, C, Cam, G, H	7	0.472	+	+
			Paste of fresh plant is orally administered for up to a week.	Gastric problems	B, C				

^aPart(s) of the plant used: Ap: Aerial parts; Bk: Bark; Br: Branches; B: Bulb; L: Leaves; Fb: Flower buds; F: Fruit; Fl: Flower; Fr: Fruit rind; Rh: Rhizome; S: Seed; Rt: Root; O: Oil; St: Stem; Stg: Stigma; Twg: Twigs; Ltx: Latex; Wd: Wood; Wp: Whole plant.

^bAT: Animals treated. A: ass; B: buffalo; C: cow; Cam: camel; D: dog; Dk: donkey; G: goat; H: horse; M: mule; P: poultry, Sh: sheep; Sw: swine.

^cC: Number of citations.

^dmCI: mean cultural importance index value.

^eFR: First report of this Ethnoveterinary practice in the Himalayan region. +: yes; -: no.

^fSU: Similar use/application in local folk medical practice for humans.+: yes; -: no.

geographic areas, it appears that nearly half of the quoted plants have never been described before as useful in folk veterinary practices. The other half has already been reported in the literature, but in some cases, for different ethnoveterinary purposes. In this section, we explore some reports on other ethnoveterinary applications of these species in the literature. This discussion is organized by plant family.

Acanthaceae

Regarding *Adhatoda vasica*, the leaves are used to treat stomach pains, fever, dehydration, diarrhea, dysentery, indigestion and gas troubles. The leaf paste of this plant has been reported for uses in the treatment of hoof rot in the literature [46]. Interestingly, aqueous extracts from the leaves have shown significant activity against *Bacillus* bacteria [47,48].

Amaranthaceae

Paste prepared from whole plant of *Amaranthus viridis* is used here against weakness in cattle. The leaves of the same plant were reported as emollient in amenorrhea, scorpion sting and snake bite in a study conducted in Islamabad, Pakistan [49].

Amaryllidaceae

The crushed bulbs of *Allium cepa* are administered to treat indigestion, stomach gripe, fever and for lactation in the study area, whereas in Italy, they are used to prevent pestilence [50]. The leaves, flowers and bulb extracts of *A. cepa* have demonstrated activity against pathogens such as *Staphylococcus aureus*, *Salmonella typhi*, *Candida albicans*, and nematodes [48,51].

Anacardiaceae

The fruit pickle of *Mangifera indica* is used for mouth infections. Others have reported that the leaves of same plant are fed to livestock to treat retained fetal membrane [20]. Chloroform, ethanolic, water and petroleum ether extracts of *M. indica* were found to have antibacterial, anti-viral anti-fungal activities, as well as anti-inflammatory properties [52,53].

Apiaceae

Aerial parts of *Foeniculum vulgare* were used to treat indigestion and diarrhea. Flowers and fruit of the same species have been reported as galactagogues and ruminative [50]. Seeds of *Trachyspermum ammi* are given to cattle as appetite stimulant and to increase milk production. In the Sargodha district of Pakistan, seed powder and decoctions of the same plant were reported for treatments against genital prolapse and to treat retained fetal membrane [20]. Alcoholic and aqueous extracts of this plant species have shown antibacterial activity [54].

Apocynaceae

Leaves, stems and twigs of *Calotropis procera* are applied to cure mouth and eye watering, colic, indigestion, pain and inflammation. Other reports regarding use of this plant include crushed leaves for the relief of flatulence, latex to increase lactation and bark decoction for hoof rot [46]. The leaves and seeds are also reported to be useful for silent estrus and delayed puberty [20]. Alcoholic and aqueous extracts of *C. procera* have shown antibacterial activities [54]. Powder prepared from the roots and leaves of *Carissa opaca* is given to cattle to treat infected or sore throats and to heal wounds. In Uttar Pradesh, India, aerial parts of *C. opaca* were reported to be administered orally to kill pest in cattle [46].

Araliaceae

An aqueous extract of *Hedera nepalensis* is applied to remove leech in cattle. In Italy, the use of fresh leaves and plant decoctions for abortive and anti-inflammatory purposes have been reported [50].

Asteraceae

The seeds and paste made from the whole plant of *Saussurea heteromalla* are used to treat edema and to purify the blood. In Islamabad, the seeds were reported as carminatives and used also in tonics for horses and camels [49]. In the present study, we found that decoctions and pastes of *Senecio chrysanthemoides* are used for the treatment of sore joints and arthritis, whereas other work has reported the use of roots and leaves for treating blackleg disease and Evil-eye [55].

Boraginaceae

The leaf paste of *Trichodesma indicum* is used to treat stomach disorders and intestinal worms in cattle in the study area, whereas others have reported the use of this paste in the treatment of mastitis and for uterine prolapse [46].

Brassicaceae

Brassica campestris seed oil is used for skin, eye and stomach infections. Other studies in Pakistan and India [20,46] have reported the use of this oil in topical applications for sores and the treatment of genital prolapses. *Eruca sativa* seed oil is used to treat dysentery in the study area. *E. sativa* seed powder has been reported for diarrhea in other work [46].

Cannabaceae

Paste from the leaves, seeds and floral buds of *Cannabis sativa* are applied as an appetite stimulant, anti-leech, anti-lice, and for abdominal swelling and indigestion. Other studies have reported the use of decoctions and

infusions for measles and East coast fever [56] and leaves for genital prolapse [20].

Convolvulaceae

Paste prepared from *Cuscuta reflexa* is fed to cattle for treatment of swelling (rumination problems), indigestion and short mammary glands. Other studies have documented its use as a galactagogue food (after being fried) [57].

Euphorbiaceae

Seed oil of *Ricinus communis* is administered to treat constipation. Other studies have documented the use of *R. communis* for intestinal obstruction, hoof problems, digestive problems, wounds, abscesses, to expel retained placenta and for silent estrus/delayed puberty in cattle [20,46,58]. The stem/leaf hexane extract of *R. communis* was suggested to be active against *Escherichia coli*, *Enterococcus faecalis*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* [59].

Fabaceae

Acacia nilotica bark decoctions are used for the treatment of stomach pains in livestock. The bark of this same plant has been reported to be used in the case of hoof rot and genital prolapse in cattle [20,46]. The seeds of *Trigonella foenum-graecum* are reported to treat diarrhea here, whereas in other areas of Pakistan they are used for treatment of genital prolapse, silent estrus and delayed puberty [20].

Lythraceae

The fruit rind of *Punica granatum* is used to cure dysentery. Other work reports the use of leaf paste for enteritis, bark powder for helminthic infection, flowers as a tonic and the rind as an astringent and to treat diarrhea [60]. Antibacterial studies on the alcoholic and aqueous extracts of this plant have demonstrated activity against *Bacillus subtilis*, *Escherichia coli*, *Proteus vulgaris*, *Salmonella typhimurium*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* [51].

Meliaceae

The leaves and fruit of *Melia azedarach* are used against foot, mouth, skin infections, gas trouble and indigestion in the study areas while according to other studies; it is used as a cooling agent and for genital prolapse [20,46].

Pteridaceae

Adiantum incisum leaf paste is used for abdominal pain in the study area, whereas in Italy, a decoction of the plant is used to expel the placenta following delivery [50].

Poaceae

Paste prepared from the seeds of *Oryza sativa* is used to treat weakness and respiratory infection. It was reported [20,50] that seeds of the same plant are also used against diarrhea and to treat retained fetal membrane. *Triticum aestivum* seeds are used against dysentery, sore mouth and to increase milk production in livestock. Other studies have reported its use as a ruminative, laxative, for dermatitis, delayed puberty, silent estrus and to treat retained fetal membrane [20,50]. *Zea mays* inflorescences are given to cure urinary inflammation in cattle. *Z. mays* has been reported for applications in wound healing and treating genital prolapse in other studies [20,50].

Polygonaceae

Local people use the roots and leaves of *Rumex nepalensis* for treating diarrhea, dysentery, intestinal worms, allergies and to stop bleeding in cattle. Crushed roots of this plant have been reported for treatment of blackleg disease (an infectious disease attributed to *Clostridium* spp.) [55].

Rutaceae

Citrus limon juice is used in the treatment of mastitis. Others have reported the use of citrus juice for uterine prolapse in cattle [20].

Sapindaceae

The powder and juice of *Aesculus indica* fruit and seeds is used against cough, fever, abdominal pain and to heal wounds in animals in the study area. However, in other regions of Pakistan, the seed endocarp is given to horses to relieve stomach pain, colic and swelling [61-63].

Saxifragaceae

Fresh leaves and powder derived from the rhizomes of *Bergenia ciliata* are topically applied for use in wound healing. Dried and fresh leaves of the same plant have been used to treat diarrhea in animals [64]. Alcoholic and aqueous extracts of *B. ciliata* rhizome has shown antibacterial and antifungal activities [65].

Scrophulariaceae

The fresh leaf paste of *Verbascum thapsus* is used to treat diarrhea. Others report the use of a leaf ointment for the treatment of rectal prolapse [50].

Solanaceae

Solanum surattense is used for healing wounds, fever, indigestion, cough and as a tonic. Others have reported the use of the leaves for genital prolapse [20]. A leaf extracts of *S. surattense* was found to be active against *Staphylococcus aureus*, *Salmonella typhi*, *Candida albicans* and nematodes [48,51]. The root paste of *Withania somnifera* is topically applied to treat bovine mastitis in

Table 2 Informant consensus values based on categories of veterinary conditions

Category	Botanical taxa used	Plant reports	ICF value
Gastrointestinal disorders (incl. tympany, colics)	47	70	0.33
Skin diseases (incl. wounds and diseases affecting eye, ear, and throat)	30	54	0.45
Diseases related to milk production	4	7	0.50
Rheumatoid disorders and inflammations	3	5	0.50
Worms and other parasitic diseases	14	35	0.62
Reproductive disorders	9	24	0.63
Respiratory disorders (cold, cough) and fever	10	29	0.68

this study area, whereas the crushed roots of this same species are used against an evil spirit (Wan laffa) in animals in Ethiopia [55]. Alcoholic and aqueous extracts of *W. somnifera* have shown antibacterial activity against *Bacillus subtilis*, *Escherichia coli*, *Proteus vulgaris*, *Salmonella typhimurium*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*, as well diuretic and anti-hypercholesterolemic activities [48,54].

Theaceae


Decoctions of *Camellia sinensis* leaves are used to cure fever in cattle in this region, while another study in Sargodha district (Pakistan) reported the use of this decoction for treating retained fetal membrane in cows [20]. Fermented tea has been shown to be hypolipidemic and to reduce high blood pressure [49].

Xanthorrhoeaceae

The leaf pulp of *Aloe vera* is administered orally as ruminative. The pulp of this same species has also been reported for similar use in the treatment of digestive problems [58]. Alcoholic and aqueous extracts of this plant have shown significant activity against *Bacillus subtilis*, *Escherichia coli*, *Proteus vulgaris*, *Salmonella typhimurium*, *Pseudomonas aeruginosa* and *Staphylococcus aureus* [54,66,67]. Leaves of *Asphodelus tenuifolius* were used to cure weakness in horses in our study, while others have reported that root paste of this plant is applied to wounds in cattle [46].

Zingiberaceae

Turmeric powder (from *Curcuma longa* rhizomes) is topically applied for wound healing in cattle in the study area, while a study on equine medicines has mentioned that roots of this plant are used for hoof problems and



sore joints [58]. Alcoholic and aqueous extracts of *C. longa* have shown antibacterial activity [54]. Chloroform, ethanol, water and petroleum ether extracts of *C. longa* rhizome were also found to be active against bacteria, viruses, and fungi, and have shown anti-inflammatory activities [52,53].

Cultural importance of the species

The Cultural Importance index (CI) of species is useful for estimating the significance of certain plants to a given culture [68] and takes into account not only the spread of the use (number of informants) for each species, but also its versatility, i.e. the diversity of its uses [68].

Based on medicinal applications, *Trachyspermum ammi* was found to be the most cited species followed by *Curcuma longa*, *Melia azedarach*, *Zanthoxylum armatum* var. *armatum*, *Adhatoda vasica*, *Allium cepa*, *Foeniculum vulgare*, *Prunus persica*, *Punica granatum*, *Trichodesma indicum*, *Berberis lycium*, *Triticum aestivum* and *Peganum harmala* (Table 1). It is notable that the top ten species of medicinal plants used to treat various livestock conditions were cited in all three major study sites (Haripur, Abbottabad, and Mansehra).

Cultural importance of the families

With regards to the diversity of species used, Fabaceae and Poaceae were the most important, with 6 species cited. Like the study by Pardo-de-Santayana et al. [30], we also elected to add the sum of CI of species in each family in order to measure the mean cultural importance of the families (mCI_f). Unlike the aforementioned study, however, the number of species reported here did not strongly correlate with the number of species ($R^2 = 0.211$). This could be explained, perhaps, by the greater diversity of families (with a more limited number of species per family, average of 1.9 species/family) quoted for ethnoveterinary applications. Of the families reported, Apiaceae had the highest mCI_f value, despite having only three species in this group (Figure 2).

Informant consensus

Perceived efficacy of medicinal plants can be assessed by ICF values, with those plants that are supposed to be effective in curing diseases having elevated ICF levels [69]. We identified seven major disease categories and the highest ICF values were recorded for respiratory disorders and fever (0.68), followed by reproductive disorders (0.63), worms and other parasitic diseases (0.63) (Table 2).

Comparison with human medicine

A large number of the veterinary plant reports share commonalities with the folk medical practices used in traditional ethnomedicine for humans in surrounding sites (last column in Table 1). This overlap may be a reflection

on how folk veterinary remedies may be the diachronic result of a deep observation of the efficacy of certain plants used in animal diseases or at least of intense transfers of local knowledge between the folk veterinary and the ethnomedical domains.

Bio-conservation concerns

Various human activities may be implicated in placing some of the local medicinal flora under a state of threat within their natural habitat. The perceptions that local people share regarding this phenomenon of threats to local ecological resources – medicinal plants, in particular, was examined based on interviews with 15 key respondents in study districts. We examined these perceived threats using pair-wise ranking [29] of five central factors: agricultural land expansion, over-harvesting, over-grazing, uncontrolled fire setting and fuel wood collection. It was observed that agricultural land expansion was perceived as the dominant threat to medicinal plants used in ethnoveterinary medicine, followed by over-harvesting, over-grazing, fire and fuel wood collection (Figure 3). Current conservation efforts concerning medicinal plants in this region are very limited, and as a result, the majority of them have no protection. This a major issue to be considered in future research and in local rural development initiatives.

Conclusions

A remarkable heritage of folk veterinary knowledge has been preserved within the framework of local knowledge and practices in the Pakistani communities of the Lesser Himalayas. However, like many other studies in this discipline have found, local knowledge is fragile and susceptible to rapid erosion with the expansion of biomedical paradigms and replacement of traditional resources with modern allopathic medicines. This is increasingly the case in both human and veterinary medicine. Nevertheless, as the majority of the reported species are wild and sometimes rare or under threat, much heed must be taken not to diminish these plant populations.

It is more urgent now than ever to record this rich body of knowledge not only for the purpose of bio-cultural conservation, but also to provide insights to scientists engaged in the search for new herbal veterinary therapies and especially to local stakeholders, who work on fostering endogenous trajectories of community-based rural development projects in mountainous areas. The latter perspective is of crucial importance in the possible implementation of ethnobiological studies in disadvantaged areas, such as the mountain regions of Pakistan [70-73] as it may have a tremendous impact in sustaining and/or revitalizing communal forms of natural resource management [74]. Moreover, *emic* visions of environmental protection and provision of health and dietary care both for humans and animals may represent

the key to environmental and social sustainability of social-ecological systems [75]. The validation and eventual application of this knowledge into concrete, comprehensive and culturally appropriate participatory initiatives aimed at fostering the sustainable use of local natural resources would promote the well-being of both animals and local communities.

Competing interest

Authors declare that they have no competing interest.

Authors' contributions

AMA conducted the ethnobotanical survey and drafted the manuscript; SMK helped in the data compilation; MA supported the field data collection; MAK supervised the project and helped in plant identification; CLQ analyzed the data and reviewed the manuscript; AP critically reviewed the manuscript and wrote the discussion and the conclusions. All authors read and approved the final manuscript.

Acknowledgement

We are grateful to all the study participants and the local communities for having shared their valuable traditional knowledge.

Author details

¹Department of Environmental Sciences, COMSATS Institute of Information Technology, Abbottabad 22060, Pakistan. ²Department of Botany, Hazara University Mansehra, Mansehra 21300, Pakistan. ³Department of Plant Sciences, Quaid-i-Azam University, Islamabad 45320, Pakistan. ⁴Department of Dermatology, Emory University School of Medicine, 1518 Clifton Rd NE, CNR Bldg. 5000, Atlanta, GA 30322, USA. ⁵Center for the Study of Human Health, Emory College of Arts and Sciences, 550 Asbury Circle, Candler Library 107, Atlanta, GA 30322, USA. ⁶University of Gastronomic Sciences, Piazza Vittorio Emanuele 9, Pollenzo I-12042 Bra/Pollenzo, Italy.

Received: 21 March 2013 Accepted: 5 December 2013

Published: 20 December 2013

References

- McCorkle CM: An introduction to ethnoveterinary research and development. *J Ethnobiol* 1986, **6**:129–149.
- Katerere DR, Luseba D: *Ethnoveterinary botanical medicine. Herbal medicines for animal health*. Boca Raton: CRC Press; 2010.
- Eswaran S, Boomibalan P, Rathinavel S: Ethnoveterinary medicinal practices of the villagers of Usilampatti Taluk of Madurai district, India. *Int J Bot* 2013, **9**:37–43.
- Galav P, Jain A, Katewa SS: Ethnoveterinary medicines used by tribals of Tadgarh-Raoli wildlife sanctuary, Rajasthan, India. *Indian J Tradit Knowl* 2013, **12**:56–61.
- Mishra D: Cattle wounds and ethnoveterinary medicine: a study in Polasara block, Ganjam district, Orissa, India. *Indian J Tradit Knowl* 2013, **12**:62–65.
- Mallik BK, Panda T, Paddy RN: Ethnoveterinary practices of aborigine tribes in Odisha, India. *Asian Pacific Journal of Tropical Biomedicine* 2012, **2**:S1520–S1525.
- Pragada PM, Rao GMN: Ethnoveterinary medicinal practices in tribal regions of Andhra Pradesh, India. *Bangladesh Journal of Plant Taxonomy* 2012, **19**:7–16.
- Bharati KA, Sharma BL: Plants used as ethnoveterinary medicines in Sikkim Himalayas. *Ethnobot Res Appl* 2012, **10**:339–356.
- Sharma R, Manhas RK, Magotra R: Ethnoveterinary remedies of diseases among milk yielding animals in Kathua, Jammu and Kashmir, India. *J Ethnopharmacol* 2012, **141**:265–272.
- Naik RM, Venugopalan V, Kumaravelayutham P, Krishnamurthy YL: Ethnoveterinary uses of medicinal plants among the Lambani community in Chitradurga district, Karnataka, India. *Asian Pacific Journal of Tropical Biomedicine* 2012, **2**:S470–S476.
- Mishra D: Ethnoveterinary practices and use of herbal medicines for treatment of skin diseases in cattle: a study in Polasara Block, Ganjam District, Orissa, India. *Veterinary World* 2011, **4**:250–253.
- Phondani PC, Maikhuri RK, Kala CP: Ethnoveterinary uses of medicinal plants among traditional herbal healers in Alaknanda catchment of Uttarakhand, India. *Afr J Tradit Complement Altern Med* 2010, **7**:195–206.
- Rajput DS, Tripathi H, Bhanja SK: Ethnoveterinary practices of Raika pastoralists for camel health management in Bikaner District of Rajasthan. *Asian Agri-History* 2005, **9**:243–252.
- Sikanwar RLS, Kumar V: Ethnoveterinary knowledge and practices prevalent among the tribals of central India. *Journal of Natural Remedies* 2005, **5**:147–152.
- Sharma UK, Sapkota D: Studies on some ethnoveterinary practices on Dhemaji district of Assam. *Journal of Natural Remedies* 2003, **3**:73–77.
- Maiti S, Chakravarty P, Garai S, Bandyopadhyay S, Chouhan VS: Ethno-veterinary practices for ephemeral fever of Yak: a participatory assessment by the Monpa tribe of Arunachal Pradesh. *Indian J Tradit Knowl* 2013, **12**:36–39.
- Selvaraju A, Ayyanar M, Rathinakumar SS, Sekar T: Plants used in ethno-veterinary medicine by malayali tribals in Salem district of Tamil Nadu, India. *Medicinal Plants* 2011, **3**:209–215.
- Raziq A, de Verdier K, Younas M: Ethnoveterinary treatments by dromedary camel herders in the Suleiman mountainous region in Pakistan: an observation and questionnaire study. *J Ethnobiol Ethnomed* 2010, **6**:16.
- Hussain A, Khan MN, Iqbal Z, Sajid MS: An account of the botanical anthelmintics used in traditional veterinary practices in Sahiwal district of Punjab, Pakistan. *J Ethnopharmacol* 2008, **119**:185–190.
- Dilshad SMR, Rehman N, Iqbal Z, Muhammad G, Iqbal A, Ahmed N: An inventory of the ethnoveterinary practices for reproductive disorders in cattle and buffaloes, Sargodha district of Pakistan. *J Ethnopharmacol* 2008, **117**:393–402.
- Dilshad SMR, Rehman NU, Ahmad N, Iqbal A: Documentation of ethnoveterinary practices for mastitis in dairy animals in Pakistan. *Pakistan Vet J* 2010, **30**:167–171.
- Farooq Z, Iqbal Z, Mushtaq S, Muhammad G, Iqbal MZ, Arshad M: Ethnoveterinary practices for the treatment of parasitic diseases in livestock in Cholistan desert (Pakistan). *J Ethnopharmacol* 2008, **118**:213–219.
- Sindhu ZUD, Iqbal Z, Khan MN, Jonsson NN, Siddique M: Documentation of ethnoveterinary practices used for treatment of different ailments in a selected hilly area of Pakistan. *Int J Agric Biol* 2010, **12**:353–358.
- Muhammad G, Khan MZ, Hussain MH, Iqbal Z, Iqbal M, Athar M: Ethnoveterinary practices of owners of pneumatic-cart pulling camels in Faisalabad City (Pakistan). *J Ethnopharmacol* 2005, **97**:241–246.
- Statistical Division, Government of Pakistan: *Livestock Population in Pakistan*. Lahore: Agricultural Census Organization; 2006.
- Ali SI, Qaiser M: *Flora of Pakistan*. Karachi-Pakistan: Botany Department, University of Karachi; 1995–2005.
- The Plant List. <http://www.theplantlist.org>.
- Stevens PF: *Angiosperm Phylogeny Website. Version 12. 2*. <http://www.mobot.org/MOBOT/research/APweb>.
- Balemie K, Kebebew F: Ethnobotanical study of wild edible plants in Derashe and Kucha Districts, South Ethiopia. *J Ethnobiol Ethnomed* 2006, **2**:53.
- Pardo-de-Santayana M, Tardío J, Blanco E, Carvalho AM, Lastra JJ, San Miguel E, Morales R: Traditional knowledge of wild edible plants used in the northwest of the Iberian Peninsula (Spain and Portugal): a comparative study. *J Ethnobiol Ethnomed* 2007, **3**:27.
- Tardío J, Pardo-de-Santayana M: Cultural importance indices: a comparative analysis based on the useful wild plants of Southern Cantabria (Northern Spain). *Economic Botany* 2008, **62**:24–39.
- Heinrich M, Ankli A, Frei B, Weimann C, Sticher O: Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc Sci Med* 1998, **47**:1859–1871.
- Ashok A, Reddy PG: Some reports on traditional ethno-veterinary practices from Savargaon areas of Ashti Taluka in Beed district (M.S.) India. *International Journal of Advanced Biological Research* 2012, **2**:115–119.
- Gaur RD, Sharma J, Painuli RM: Plants used in traditional health care of livestock by Gujjar community of Sub-Himalayan Tract, Uttarakhand, India. *Indian Journal of Traditional Knowledge* 2010, **1**:243–248.
- Islam M, Anwar Z, Tabassum S, Khan SA, Zeb AC, Abrar M, Khattak KU, Khattak JZK: Plants of Ethno-Veterinary Uses of Tuglaj Mountain Baffa Mansehra, Pakistan. *International Journal of Animal and Veterinary Advances* 2012, **4**:221–224.

36. Khan MA, Khan MA, Hussain M: **Ethnoveterinary medicinal uses of plants of Poonch valley Azad Kashmir.** *Pakistan Journal of Weed Sciences* 2012, **18**:495–507.
37. Mondal T: **An investigation on ethno-veterinary medicinal plants of Siliguri Subdivision of Darjeeling district, West Bengal, India.** *Journal of Today's Biological Sciences* 2012, **1**:45–50.
38. Pande PC, Tiwari L, Pande HC: **Ethnoveterinary plants of Utranchal - a review.** *Indian Journal of Traditional Knowledge* 2007, **6**:444–458.
39. Pandit PK: **Inventory of ethno veterinary medicinal plants of Ghargram division, West Bengal, India.** *Indian Forester* 2010, **136**:1183–1194.
40. Phondani PC, Maikhuri K, Kala P: **Ethno veterinary uses of medicinal plants among traditional herbal Healers in Alaknanda catchment of Uttarakhand, India.** *African Journal of Traditional and Complementary Medicines* 2010, **7**:195–206.
41. Shah GM, Ahmad M, Arshad M, Khan MA, Zafar M, Sultana S: **Ethno-phyto-veterinary medicines in northern Pakistan.** *The Journal of Animal and Plant Sciences* 2012, **22**:791–797.
42. Shah R, Pande PC, Tiwari L: **Traditional veterinary herbals medicines of western part of Almora district, Uttarakhand, Himalayas.** *Indian Journal of Traditional Knowledge* 2008, **7**:355–359.
43. Singh P, Singh BK, Joshi GC, Lalit M: **Veterinary Ethno-Medicinal Plants in Uttarakhand Himalayan Region Tewari.** *Nature and Science* 2009, **7**:44–52.
44. Singh PK, Singh S, Kumar V, Krishna B: **Ethno veterinary healthcare practices in Marihan sub-division of district Mirzapur, Uttar Pradesh, India.** *Life Sciences Leaflets* 2011, **16**:561–569.
45. Tiwari L, Pande PC: **Ethno veterinary medicines in Indian perspective: reference to Uttarkhand, Himalaya.** *Indian Journal of Traditional Knowledge* 2010, **9**:611–617.
46. Ali ZA: **Folk veterinary medicine in Moradabad district Uttar Pradesh, India.** *Fitoterapia* 1999, **70**:340–347.
47. Parsad G, Panday S, Gupta KK: **Antimicrobial activity of certain indigenous drugs and drug plants against certain bacterial species.** *Hamdard Medicus* 1999, **27**:53–57.
48. Srinivasan D, Nathan S, Suresh T, Perumalsamy PL: **Antimicrobial activity of certain Indian medicinal plants used in folklore medicine.** *J Ethnopharmacol* 2001, **74**:217–220.
49. Shinwari MI, Khan MA: **Folk use of medicinal herbs of Margalla hills national park, Islamabad.** *J Ethnopharmacol* 2000, **69**:45–56.
50. Uncini Manganelli RE, Camangi F, Tomei PE: **Curing animals with plants, traditional usage in Tuscany, Italy.** *J Ethnopharmacol* 2001, **78**:171–191.
51. Qarar F, Kalhora MA, Badar Y: **Anthelmintic properties of some indigenous plants.** *Hamdard Medicus* 1998, **21**:115–117.
52. Grewal RC: *Medicinal Plants.* New Delhi: Campus Books International; 2000.
53. Khattak S, Rehman S, Shah HU, Ahmed W, Ahmed M: **Biological effects of indigenous medicinal plants *Curcuma longa* and *Alpinia galangal*.** *Fitoterapia* 2005, **76**:254–257.
54. Ahmad I, Mehmood Z, Mohammad F: **Screening of some Indian medicinal plants for their antimicrobial properties.** *J Ethnopharmacol* 1998, **62**:183–193.
55. Yineger H, Kelbessa E, Bekele T, Lulekal E: **Ethno veterinary medicinal plants at Bale Mountains National Park, Ethiopia.** *Journal of Ethnopharmacology* 2007, **112**:55–70.
56. Tabuti JR, Dhillion SS, Lye KA: **Ethno veterinary medicine for cattle (*Bos indicus*) in Bulamogi County, Uganda.** *Journal of Ethnopharmacology* 2003, **88**:279–286.
57. Bandyopadhyay S, Mukherjee SK: **Ethno veterinary medicine from Koch Bihar district, West Bengal.** *Indian Journal of Traditional Knowledge* 2005, **4**(4):456–461.
58. Lans C, Turner N, Khan T, Brauer G, Lourenco G, Georges K: **Ethnoveterinary medicines used for horses in Trinidad and in British Columbia, Canada.** *J Ethnobiol Ethnomed* 2006, **2**:31.
59. McGaw LJ, Merwe DVD, Eloff JN: **In vitro anthelmintic, antibacterial and cytotoxic effects of extracts from plants used in South African ethno veterinary medicine.** *The Veterinary Journal* 2007, **173**:366–372.
60. Karthickayan SMK, Gajendran K: **Indigenous technical know-how in the health care of domestic animals.** *Indian Journal of Traditional Knowledge* 2005, **4**(4):462–463.
61. Abbasi AM, Khan MA, Ahmad M, Zafar M: *Medicinal plant biodiversity of Lesser Himalayas, Pakistan.* New York: Springer; 2012.
62. Shah GM, Khan MA: **Common medicinal folk recipes of Siran Valley, Mansehra, Pakistan.** *Ethnobotanical Leaflet* 2006, **4**:110–130.
63. Matin A, Khan MA, Ashraf M, Qureshi RA: **Traditional uses of shrubs and trees of Himalayan Region, Shogran valley, District Mansehra, Pakistan.** *Hamdard Medicus* 2002, **45**:50–56.
64. Shinwari ZK, Gilani SS: **Sustainable harvest of medicinal plants at Bulashbar Nullah, Astore (Northern Pakistan).** *J Ethnopharmacol* 2003, **84**:289–298.
65. Gorski MS: **Antimicrobial activity of some medicinal plants of Muzaffarabad.** *Hamdard Medicus* 2005, **48**:27–41.
66. Sweelam ME: **The potential use of some ornamental plants for nematodal control in Egypt.** *Bulletin Agriculture University Cairo* 1989, **40**:391–393.
67. Trivedi PC: *Medicinal plants.* Jodhpur (India) Agrobios: Ethnobotanical Approaches; 2006.
68. Albuquerque UP, Andrade LHC, Silva ACO: **Use of plant resources in a seasonal dry forest (Northeastern Brazil).** *Acta Botanica Brasiliica* 2005, **19**:27–38.
69. Teklehaymanot T, Giday M: **Ethnobotanical study of medicinal plants used by people in Zegie Peninsula. Northwestern Ethiopia.** *J Ethnobiol Ethnomed* 2007, **3**:12.
70. Akhtar N, Rashid A, Murad W, Bergmeier E: **Diversity and use of ethno-medicinal plants in the region of Swat, North Pakistan.** *J Ethnobiol Ethnomed* 2013, **9**:25.
71. Khan SM, Page S, Ahmad H, Shaheen H, Ullah Z, Ahmad M, Harper DM: **Medicinal flora and ethnoecological knowledge in the Naran Valley, Western Himalaya, Pakistan.** *J Ethnobiol Ethnomed* 2013, **9**:4.
72. Abbasi AM, Khan MA, Shah MH, Shah MM, Pervez A, Ahmad M: **Ethnobotanical appraisal and cultural values of medicinally important wild edible vegetables of Lesser Himalayas-Pakistan.** *J Ethnobiol Ethnomed* 2013, **9**:66.
73. Rotherham ID: *Cultural Severance and the Environment. The Ending of Traditional and Customary Practice on Commons and Landscapes Managed in Common.* Dordrecht/Heidelberg/New York/London: Springer; 2013.
74. Pieroni A, Nedelcheva A, Hajdari A, Mustafa B, Scaltriti B, Cianfaglione K, Quave CL: **Local Knowledge on Plants and Domestic Remedies in the Mountain Villages of Peshkopia (Eastern Albania).** *Journal of Mountain Science* 2014, **11**:180–194.
75. Berkes F, Colding J, Folke C: *Navigating Social-Ecological Systems. Building Resilience for Complexity and Change.* Cambridge: Cambridge University Press; 2003.

doi:10.1186/1746-4269-9-84

Cite this article as: Abbasi et al.: Botanical ethnoveterinary therapies in three districts of the Lesser Himalayas of Pakistan. *Journal of Ethnobiology and Ethnomedicine* 2013 **9**:84.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

