

Published in final edited form as:

Toxicol Appl Pharmacol. 2009 September 1; 239(2): 184–192. doi:10.1016/j.taap.2009.01.010.

Arsenic Exposure at Low-to-Moderate Levels and Skin Lesions, Arsenic Metabolism, Neurological Functions, and Biomarkers for Respiratory and Cardiovascular Diseases: Review of Recent Findings from the Health Effects of Arsenic Longitudinal Study (HEALS) in Bangladesh

Yu Chen¹, Faruque Parvez², Mary Gamble², Tariqul Islam³, Alauddin Ahmed³, Maria Argos⁴, Joseph H. Graziano², and Habibul Ahsan⁴

¹Departments of Environmental Medicine and Medicine and New York University Cancer Institute, New York University School of Medicine, New York, NY, USA

²Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York City, NY, USA

³Columbia University Arsenic Research Project, Dhaka, Bangladesh

⁴Departments of Health Studies, Medicine and Human Genetics and Cancer Research Center, The University of Chicago, Chicago, IL, USA

Abstract

The contamination of groundwater by arsenic in Bangladesh is a major public health concern affecting 35–75 million people. Although it is evident that high levels (> 300 µg/L) of arsenic exposure from drinking water are related to adverse health outcomes, health effects of arsenic exposure at low-to-moderate levels (10–300 µg/L) are not well understood. We established the Health Effects of Arsenic Longitudinal Study (HEALS) with more than 20,000 men and women in Araihaazar, Bangladesh, to prospectively investigate the health effects of arsenic predominately at low-to-moderate levels (0.1 to 864 µg/L, mean 99 µg/L) of arsenic exposure. Findings to date suggest adverse effects of low-to-moderate levels of arsenic exposure on the risk of pre-malignant skin lesions, high blood pressure, neurological dysfunctions, and all-cause and chronic disease mortality. In addition, the data also indicate that the risk of skin lesion due to arsenic exposure is modifiable by nutritional factors, such as folate and selenium status, lifestyle factors, including cigarette smoking and body mass index, and genetic polymorphisms in genes related to arsenic metabolism. The analyses of biomarkers for respiratory and cardiovascular functions support that there may be adverse effects of arsenic on these outcomes and call for confirmation in large studies. A unique strength of the HEALS is the availability of outcome data collected prospectively and data on detailed individual-level arsenic exposure estimated using water, blood

© 2009 Elsevier Inc. All rights reserved.

Address for correspondence and reprints: Reprint requests and correspondence should be addressed to: Dr. Habibul Ahsan, Center for Cancer Epidemiology and Prevention, The University of Chicago, 5841 South Maryland Avenue, Suite N102, Chicago, IL 60637, Phone: 773-834-9956, Fax: 773-834-0139, habib@uchicago.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest:

The authors declare that there are no conflicts of interest.

and repeated urine samples. Future prospective analyses of clinical endpoints and related host susceptibility will enhance our knowledge on the health effects of low-to-moderate levels of arsenic exposure, elucidate disease mechanisms, and give directions for prevention.

INTRODUCTION

Arsenic is abundant in the earth's crust and can be released into groundwater under certain conditions. In many parts of the world where groundwater is an important source of drinking water, long-term exposure to arsenic from drinking water has been considered a public health hazard. In addition to be classified as a Class I human carcinogen by the International Agency for Research on Cancer (IARC) for its association with an increase in skin cancer risk, arsenic exposure has also been linked to increased risks of internal cancers, diabetes, cardiovascular disease, adverse pregnancy outcomes, and a decrease in children's intellectual function.

However, epidemiologic evidence on many of these health effects of arsenic exposure has not been well-established, with uncertainties in latency, dose-response relationships and population differences. In particular, although it is evident that high levels of arsenic exposure ($> 300 \mu\text{g/L}$) are related to internal cancer and cardiovascular disease, epidemiologic evidence of the effects of arsenic exposure from drinking water at low-to-moderate levels (< 300 or $< 100 \mu\text{g/L}$) remains inconclusive. Most existing studies include limited sample size and/or unreliable long-term measures of arsenic exposure with exposure measured ecologically or only cross-sectionally. Ecologic exposure measurement that uses the mean or median values in a county or community as the exposure level for individuals could result in considerable measurement errors. Since the effects of low-to-moderate levels of arsenic on disease are likely to be modest in magnitude, these studies are particularly susceptible to measurement errors in exposure ascertainment, which, in most cases, would lead to bias towards the null, but could also generate spurious associations under certain conditions. In addition, the heterogeneity of drinking water resources, the limited exposure range, and the relatively high migration rates within the nation altogether pose a challenge for epidemiologic studies in the U.S. or other populations with surface water as the main source of drinking water that aim to investigate the impact of low-to-moderate levels of arsenic exposure to have a valid long-term arsenic measure at the individual level.

The contamination of groundwater by arsenic in Bangladesh is the largest poisoning of a population in history. As 95% of the country's 140 million population rely on well water, an estimated 57 million people have been chronically exposed to drinking water with arsenic levels exceeding the WHO standard of $10 \mu\text{g/L}$, and 35 million people were exposed to arsenic levels above the country's government standard of $50 \mu\text{g/L}$ (British Geological Survey, 1999). Given the known health consequences of arsenic exposure, there is an imminent need to develop arsenic mitigation programs. Although there have been several case-control and cross-sectional studies published on arsenic exposure in Bangladesh, large epidemiologic studies are needed to assess the health effects of arsenic exposure in the population.

In Bangladesh, where 95% of the population exclusively drinks groundwater through wells and the population is remarkable stable, long-term arsenic exposure can be measured and tracked in large cohort at the individual level. We have established the Health Effects of Arsenic Longitudinal Study (HEALS) in Araihaazar, Bangladesh, a prospective cohort with more than 20,000 participants, to investigate the health effects of arsenic exposure, avoiding many of the limitations in previous studies. In particular, the study population has been exposed to a wide range of arsenic exposure at low-to-moderate levels, providing us with a

unique opportunity to evaluate health effects of arsenic exposure at these levels and avoiding many of the limitations of previous studies. We summarize findings from the HEALS to date and discuss the implications of our findings and future directions.

Health Effects of Arsenic Longitudinal Study (HEALS)

The HEALS was established as part of the Columbia University's Superfund Basic Research Program (CU-SBRP). Details of the HEALS have been presented elsewhere. Briefly, prior to subject recruitment, water samples and geographic positional system data were collected for a set of 5,966 contiguous wells in a well-defined geographic area of 25 km² in Araihaazar. As a baseline measurement, well owners were interviewed to create an initial roster list of 65,876 regular well users and residents in the area. From October, 2000 to May, 2002, 12,050 potential participants meeting eligibility criteria were approached, and 11,746 of those eligible agreed to participate (97.5 % response rate). In 2007, an additional 8,000 participants were recruited using the same methodologies, with a roster established based on arsenic measurement of a different set of 5,000 well with a response rate of 95%. In order to update changes in arsenic exposure and the detection of a variety of health outcomes of interest, all cohort members were actively followed every two years through in-person home visits by trained physicians and interviewers.

Water samples from all the tube wells in the study area were collected and tested for total arsenic concentrations first by graphite furnace atomic-absorption spectrometry (GFAA) with a Hitachi Z-8200 system at the Lamont-Doherty Earth Observatory (LDEO) of Columbia University. Samples that fell below the detection limit of GFAA (5 µg/L) were subsequently analyzed by inductively coupled plasma mass spectrometry (ICP-MS), with a detection limit of 0.1 µg/L. The study population has been exposed to a wide range of arsenic exposure at low-to-moderate levels, ranging from 0.1 to 864 µg/L, with a mean of 99 µg/L.

At baseline recruitment, venous whole blood samples and a spot urine sample were collected for > 90% cohort participants. At follow-up visits, a spot urine sample is also collected. Both blood and urine samples were kept in portable coolers immediately after collection. All samples were kept frozen and shipped to USA on dry ice within 1–2 months. All urine samples collected at baseline and at follow-up visits were analyzed for total arsenic concentration by GFAA, using the Analyst 600 graphite furnace system at Columbia University, as previously described.

At baseline and in each in-person biennial follow-up visit, participants undergo a comprehensive clinical examination conducted by trained physicians. Information on patterns and history of well use, demographics and lifestyle characteristics are also collected at every visit. At baseline, the questionnaire also included a validated food frequency questionnaire (FFQ) designed specifically for the study population. In addition, a field clinic was established solely for cohort study participants and their family members to passively follow up participants between their biennial in-person visits to augment the detection of study outcomes through active follow-up.

Given its biological samples and its data on arsenic exposure, dietary factors and lifestyle habits, the HEALS provides a unique opportunity to study the health effects of arsenic exposure, host susceptibility and pre-clinical disease states measured using biomarkers. In addition, based on findings from observational studies, a series of intervention studies have been initiated, including those pertaining to health education and chemoprevention trials.

KEY FINDINGS FROM HEALS, TO DATE

Well Arsenic Concentration and Biochemical Measures of Arsenic Exposure

Relationships among Different Measures of Arsenic Exposure—Although total urinary and blood arsenic levels have been considered indicators of short-term internal dose of arsenic exposure, with chronic and continuing exposure, steady-state concentrations in blood and urine can be achieved. Repeated measurements of these indices may be used to depict long-term exposure level and changes in exposure over time. Among 849 individuals randomly selected from the HEALS participants, baseline urinary arsenic was highly correlated with baseline blood arsenic ($r = 0.85$, p -value < 0.01) and with water arsenic levels ($r = 0.76$, p -value < 0.01).

Using urinary arsenic data collected during the first three visits from 10,224 participants, we estimated that the pair-wise correlation between urinary arsenic measured at each visit was high (all > 0.60 , p -values < 0.01). The correlation of baseline urinary arsenic levels with urinary arsenic levels at follow-up visits somewhat decreased from 0.66 for Visit 2 to 0.60 for Visit 3, suggesting a change in arsenic exposure among some participants; however, the change in arsenic exposure is not substantial in the overall cohort. On average, urinary arsenic levels decreased by 62 $\mu\text{g/g}$ creatinine from baseline to the first follow-up visit ($\sim -20 \mu\text{g/L}$ in well arsenic), remained at the same level at Visit 3, with a high correlation (0.74, p -value < 0.01) between urinary arsenic at Visits 2 and 3.

These data suggest that the level of urinary arsenic does not fluctuate greatly overtime, and urinary arsenic can serve as a long-term biomarker of arsenic exposure to track arsenic exposure levels during the follow-up time in the cohort.

Measures and Determinants of Arsenic Metabolism—Arsenic in drinking water is present as inorganic arsenic (InAs), i.e., arsenite (As^{III}) and arsenate (As^{V}). In Bangladesh, As^{III} is the predominant form to which people are exposed from groundwater. Methylation of InAs^{III} , which is primarily hepatic, relies on folate-dependent, one-carbon metabolism; folate contributes methyl groups used in generating *S*-adenosylmethionine (SAM). Arsenic (+3 oxidation state) methyltransferase (AS3MT) transfers the methyl group from SAM to InAs^{III} to generate monomethylarsonic acid (MMA^{V}). After the reduction of MMA^{V} to monomethylarsonous acid (MMA^{III}), AS3MT can catalyze a second methylation, generating dimethylarsinic acid (DMA^{V}). The relative distribution of urinary arsenic metabolites varies from person to person and has been interpreted to reflect biologically effective doses of exposure to the various arsenic metabolites as well as arsenic methylation efficiency, with some evidence suggesting that increased proportions of MMA and InAs species in urine may be associated with a higher risk of cancer.

Among 98 HEALS cohort members participating in the placebo arm of a double-blind randomized folate supplementation trial, the intraclass correlations (ICCs) of three urine samples collected over a three-month period were all > 0.65 for urinary arsenic metabolites. Of particular note, the ICCs for %MMA, %DMA, and the ratio of MMA-to-DMA were 0.84 (95% CI, 0.78–0.89), 0.82 (95% CI, 0.74–0.87), and 0.82 (95% CI, 0.75–0.88), respectively. The data suggest that, in the absence of interventions, there is little within-subject variability of urinary arsenic metabolite profiles and thus justifies the use of single measurements in epidemiologic studies.

Factors that are related to the composition of urinary arsenic metabolites may contribute to the susceptibility to health effects of arsenic exposure. Table 1 summarizes several studies that utilized the resources of the HEALS with a focus on inter-individual variability in the distribution of urinary arsenic metabolites. In 1,041 individuals randomly selected from the

HEALS participants who were free of skin lesions at baseline, water arsenic concentration was positively associated with urinary %MMA and inversely associated with urinary %DMA, suggesting that either methylation of arsenic to DMA is saturable or, alternatively, that arsenic inhibits the arsenic methyltransferase enzyme. Body mass index (BMI) was positively related to urinary %MMA and inversely related to urinary %DMA, however the association was not statistically significant. Specific dietary food/nutrient intakes may be more relevant to arsenic metabolism. In the same study population, higher intakes of cysteine, methionine, calcium, protein and vitamin B-12 measured by the FFQ were associated with higher %MMA in urine. Additionally, higher intakes of niacin (beta = 0.22, p-value = 0.02) and choline (beta = 0.10, p-value = 0.02) were associated with higher DMA-to-MMA ratios.

In a sample of 1,650 HEALS cohort members who were recruited into the Nutritional Influences of Arsenic Toxicity (NIAT) study, an unusually high prevalence of hyperhomocysteinemia was found, particularly among males. In subsequent analyses of 300 randomly selected participants from the NIAT study, urinary %DMA was positively associated with plasma folate ($r = 0.14$, $p = 0.02$) and negatively associated with total homocysteine (tHcys; $r = -0.14$, $p = 0.01$). Conversely, urinary %MMA was negatively associated with folate ($r = -0.12$, $p = 0.04$) and positively associated with tHcys ($r = 0.21$, $p < 0.01$). An unanticipated finding was that urinary creatinine, a breakdown product of creatine, is negatively associated with %InAs and positively associated with %DMA in urine. Creatine is synthesized endogenously in a process that relies on one-carbon metabolism; it also consumes methyl groups, and is itself consumed in the diet, predominantly from meat. The finding warrants further investigation. Collectively, these findings indicate that nutritional factors involved in one-carbon metabolism contribute to the variability in arsenic methylation.

Recently, the CU-SBRP Trace Metals Laboratory has developed the technology to measure arsenic metabolites in blood. This has led to the observation that there is relatively more MMA and less DMA in blood than in urine both in adults and in children. This is not surprising, given that DMA has a shorter circulating half-life than InAs, and is consistent with the role of methylation in facilitating arsenic elimination. This finding has important implications with regard to risk assessment in epidemiologic studies that traditionally rely on measures of arsenic metabolites in urine.

Pre-malignant Skin Lesions

Cutaneous abnormalities are well-known early signs of chronic inorganic arsenic poisoning. Melanosis is considered as an early-stage skin lesions, while keratosis is the most frequent manifestation preceding the appearance of arsenic-related skin cancer. They give rise to the majority of arsenic-induced basal and squamous cell skin cancers. Unlike arsenic-related internal cancers that could have long latencies, these premalignant skin lesions may appear with shorter periods of arsenic exposure (Saha, 2003). We have conducted several cross-sectional, case-control, case-cohort, and nested case-control studies of skin lesions to evaluate the full dose-response relationship between arsenic exposure and the risk of skin lesions, as well as relevant nutritional and genetic susceptibility factors (Table 2).

Arsenic Exposure at Low-to-moderate Levels and Skin Lesions—At baseline, we identified 714 cases of premalignant skin lesions. In cross-sectional analyses of baseline data comparing cases and non-cases of skin lesions, we observed a dose-response effect of arsenic on the risk of skin lesions. In particular, arsenic exposure appears to increase the risk of skin lesions, even at the low end of exposure in this population. Compared with drinking water containing <8.1 $\mu\text{g/L}$ of arsenic, drinking water containing 8.1–40.0, 40.1–91.0, 91.1–

175.0 and 175.1–864.0 $\mu\text{g/L}$ of arsenic was associated with adjusted prevalence odds ratios of skin lesions of 1.91 (95% confidence interval (CI): 1.26, 2.89), 3.03 (95% CI: 2.05, 4.50), 3.71 (95% CI: 2.53, 5.44) and 5.39 (95% CI: 3.69, 7.86), respectively. A prospective case-cohort analysis of skin lesion cases diagnosed during two years of follow-up also showed a similar dose-response relationship.

Modifiable Determinants of Effects of Arsenic on Skin Lesions—In our analyses, we found that males and older participants were more likely to be affected by arsenic exposure. Additionally, a synergistic effect between the highest level of arsenic exposure ($> 113 \mu\text{g/L}$) and tobacco smoking on risk of skin lesions was observed in men. Furthermore, the risk of skin lesions associated with any given level of arsenic exposure was greater in men with excessive sun exposure. The effect of arsenic was also modified by land ownership on a multiplicative scale, with an increased risk among non-land owners associated with well water arsenic. Part of the modification effect due to socioeconomic status may be explained by nutritional status. In particular, participants with a comparatively high BMI or with comparatively high intake levels of riboflavin, pyridoxine, folate, and vitamins A, C and E were less likely to be affected by arsenic exposure. In a prospective case-cohort analysis of 303 incident skin lesion cases and 849 subcohort members, there was an inverse association between baseline blood selenium status and the incidence of skin lesions.

In another nested case-control study of 274 skin lesion cases identified two years after recruitment and 274 controls matched to cases for gender, age, and water arsenic, we found that folate deficiency, hyperhomocysteinemia, low urinary creatinine—each associated with decreased arsenic methylation—are risk factors for arsenic-induced skin lesions. The positive association between DNA methylation and arsenic exposure that we previously observed in the NIAT study was confirmed among the controls in this study. However, we believe that this may be an adaptive change, as hypomethylation of leukocyte DNA was found to be associated with an increased risk for skin lesions. These data suggest that lifestyle and nutritional factors may modify the risk of arsenic-related skin lesions.

Genetic Susceptibility to Effects of Arsenic on Skin Lesions—Interindividual variability in arsenic metabolism capacity may also contribute to the variation in susceptibility to the effect of arsenic. Glutathione S-transferase $\omega 1$ (*GSTO1*) and methylenetetrahydrofolate reductase (*MTHFR*) are enzymes involved in arsenic metabolism pathways. In a case-control study of 594 skin lesion cases and 1,041 controls, the dose-response relationship of skin lesion risk with urinary monomethylarsonous acid percentage (%MMA) was more apparent than those with other methylation indices. Individuals with the *MTHFR* 677TT/1298AA and 677CT/1298AA diplotypes were 1.66 (95% CI, 1.00–2.77) and 1.77 (95% CI, 0.61–5.14) times more likely to have skin lesions, compared with those carrying 677CC/1298CC diplotype. The OR for skin lesions in relation to the *GSTO1* diplotype containing all at-risk alleles was 3.91 (95% CI, 1.03–14.79). These findings reiterate that arsenic-induced health effects may be especially deleterious in subsets of the population carrying susceptible variants of genes relevant to arsenic metabolism. Based on the risk estimates observed in this study, the proportion of skin lesions in our study population attributable to the *MTHFR* 677TT/1298AA and 677CT/1298AA diplotypes was estimated to be 7.5%. The corresponding estimated attributable proportion for the *GSTO1* at-risk diplotype was 8.9%.

Peripheral Neuropathy and Children's Intellectual Function

Among 137 HEALS participants randomly selected from those visited the field clinic over eight weeks, peripheral neuropathy was assessed by a vibration sensitivity tester (Vibratron

II). Arsenic exposure was associated with elevated toe vibration threshold (TVT). Specifically, urinary arsenic was significantly associated with elevated TVT (p -value < 0.01) after adjustment for age and gender.

In cross-sectional analyses of 201 children at 10 years of age (whose parents participate in HEALS), children's intellectual function was assessed using tests drawn from the Wechsler Intelligence Scale for Children, version III (WISC-III). Children provided urine specimens for the measurement of urinary As and creatinine. Information on the primary source of drinking water was obtained from the child's mother. Exposure to arsenic from drinking water was associated with reduced intellectual function after adjustment for sociodemographic covariates in a dose-response manner, such that children exposed to water arsenic levels $> 50 \mu\text{g/L}$ achieved significantly lower Performance and Full-Scale scores than did children exposed to lower water arsenic levels. In a similar investigation of 301 randomly selected six-year-olds, water arsenic exposure was significantly negatively associated with both Performance and Processing Speed raw scores of the Wechsler Preschool and Primary Scale of Intelligence, version III (WPPSI-III). Although, these associations were less strong than in our previously studied 10-year-olds, this second cross-sectional study of arsenic exposure expands the concerns about arsenic neurotoxicity to a younger age group.

Arsenic Exposure at Low-to-Moderate Levels and Blood Pressure

Using baseline data in 10,910 participants, we assessed the association between arsenic exposure from drinking water and blood pressure. We observed a positive association between low-to-moderate levels of arsenic exposure from drinking water and high pulse pressure (pulse pressure $\geq 55 \text{ mmHg}$), an indicator of arterial stiffness, which is associated with an increased risk of atherosclerosis. In addition, among participants with lower-than-average dietary intake levels of B vitamins and folate, those with a higher well arsenic concentration were 1.83–1.89 times more likely to have a pulse pressure $\geq 55 \text{ mmHg}$, compared with those in the bottom quintile of well arsenic concentration ($< 8 \mu\text{g/L}$). These findings indicate that the effect of low-level arsenic exposure on blood pressure is nonlinear and may be more pronounced in persons with a lower intake of nutrients related to arsenic metabolism and cardiovascular health.

Intermediate Biomarkers of Early Biological Effect or Altered Structures and Functions of Target Organs

Arsenic and Pre-clinical Markers of Cardiovascular Diseases—Previous publications have documented dose-response relationships between Carotid Artery Intima-Medial Thickness (IMT), which is measured using ultrasound imaging, stroke, angina pectoris, myocardial infarction (MI), intermittent claudication and essential hypertension. These relationships indicate that IMT is a valid surrogate marker for clinical endpoints.

In a pilot cross-sectional study of 66 healthy, normotensive, relatively young cohort members, the ORs for carotid IMT $> 0.75 \text{ mm}$ were 1.61 (95% CI: 0.29–8.81) and 2.84 (95% CI: 0.39–20.86) comparing middle and highest levels with the lowest level of total urinary arsenic, respectively. Although the observed associations were not statistically significant, the trend in effect estimates suggests a possible role of low-to-moderate levels of arsenic exposure in atherosclerosis, which warrants future investigation in a larger study.

Adhesion of circulating leucocytes to the endothelial cell and subsequent transendothelial migration is an important step in the initiation of atherosclerosis. In part, this process is mediated by cellular adhesion molecules (CAMs), expressed on the endothelial membrane, in response to inflammatory stimuli. Circulating markers of systemic inflammation and

endothelial dysfunction, such as soluble intercellular adhesion molecule-1 (sICAM-1) and soluble vascular adhesion molecule-1 (sVCAM-1), have been shown to predict future cardiovascular disease (CVD). In a subgroup of 115 individuals with arsenic-related skin lesions, there was a positive association of urinary arsenic and well arsenic concentration with plasma levels of sICAM-1 and sVCAM-1. The positive associations of well arsenic with baseline and changes in plasma sVCAM-1 and sICAM-1 suggest a potential pathway underlying the effect of long-term arsenic exposure on CVD. Future larger studies are required to further examine the associations of low-level arsenic exposure with markers of vascular inflammation and endothelial dysfunction in healthy persons.

Arsenic and Biomarkers of Reno-vascular Diseases—Microalbuminuria, a marker of glomerular hyperfiltration, has been correlated with and may be a manifestation of impaired endothelial function. Impairment of endothelial function is recognized as one of the initial mechanisms that lead to atherosclerosis. In a study of arsenic and proteinuria (detected by dipstick analysis) in 11,121 persons in HEALS, we observed a dose-response relationship between well arsenic concentration and prevalence of proteinuria. In addition, cohort analysis with repeated measures of proteinuria and urinary arsenic concentrations revealed that a change in urinary arsenic was positively related to incidence of proteinuria during the four years of follow-up. The dose-response relationship between arsenic exposure and proteinuria provides evidence of the effect of low-to-moderate arsenic exposure levels on a common causal intermediate of CVD and kidney disease.

Arsenic and Biomarkers of Cancer—The development of skin lesions from arsenic exposure may be mediated by increases in the expression of various growth factors, including transforming growth factor- α (TGF α). To investigate this association in humans, levels of total urinary arsenic and urinary TGF α were determined in 41 individuals from HEALS, with and without arsenic-associated skin lesions; all individuals had chronic exposure to arsenic in their drinking water. Linear regression analyses suggest that total urinary arsenic explained a substantial variation of urinary TGF α (R-squared value of the model = 0.40; p-value <0.01), particularly in individuals with arsenic-associated skin lesions (R-squared value of the model = 0.70; p-value < 0.01). The R-squared values did not change appreciably with the adjustment for age and gender. There was also a trend of increasing odds ratios for the presence of arsenic-associated skin lesions with increasing urinary TGF α , although this was not significant (p-value = 0.15). However, the results were based on a small number of subjects.

In a follow-up study of 574 participants of the HEALS, the extracellular domain of the epidermal growth factor receptor (EGFR), to which TGF α binds to promote carcinogenesis, was measured by enzyme-linked immunosorbent assay in serum. Serum EGFR was found to be positively associated with three different measures of arsenic exposure (well water arsenic, urinary arsenic and a cumulative arsenic index) at statistically significant levels (p 0.034). In addition, the risk of skin lesions for a given level of arsenic increased with increasing levels of EGFR. These results indicate that TGF α -/EGFR-dependent mechanisms may play a role in the development of arsenic-related skin lesions and skin cancer.

Arsenic and Molecular and Clinical Biomarkers of Pulmonary Diseases—Serum level of Clara cell protein (CC16), one of the 20 proteins secreted by Clara cells in the lung's alveolar epithelium, has been indicated as a novel biomarker for respiratory illnesses, with a reduced CC16 concentration indicating damages in alveolar Clara cells. In cross-sectional analyses of 241 nonsmoking individuals randomly selected from the cohort, there was an inverse association between urinary arsenic and serum CC16 among persons with skin lesions ($\beta = -0.13$, p = 0.01). There was also a positive association between the

ratio of DMA to MMA in the urine and CC16 levels ($\beta = 0.12$, $p = 0.05$). In a subsample of study participants undergoing spirometric measures ($n = 31$), urinary arsenic was inversely associated with lung function measured by predictive FEV1 (forced expiratory volume measured in 1 sec) ($r = -0.37$).

Arsenic and Biomarkers of Gene Expressions in Peripheral Blood—In a microarray-based gene expression analysis comparing skin lesion cases with non-cases, Affymetrix HG-U133A GeneChip arrays were used to measure the expression of ~22,000 transcripts, using RNA from peripheral blood lymphocytes. Down-regulating of several genes, including *superoxide dismutase 2 (SOD2)* gene, *tumor necrosis factor (TNF)* gene and chemokine factor *CCL20* gene was observed in skin lesion cases. These findings suggest the involvement of inflammation, oxidative stress defense and chemokine response pathways in the development of skin lesions, or as a consequence of skin lesion manifestation.

In a separate comparison of samples collected from skin lesions cases before and after selenium supplementation, genes up-regulated by selenium supplementation included *TNF*, *IL1B*, *IL8*, *SOD2*, *CXCL2* and several other immunological and oxidative stress-related genes. These findings suggest that selenium supplementation may reverse some gene expression changes in individuals with pre-malignant skin lesions.

Prevention of Health Effects of Arsenic Exposure

Based on our findings of potential nutritional influence on arsenic toxicity (described above), we desired to assess whether dietary supplementation with specific micronutrients could prevent the occurrence of skin lesions, improve existing skin lesions and/or prevent skin and other cancers. In a pilot randomized, placebo-controlled, double-blind trial of 121 skin lesion cases, supplementation with vitamin E and selenium, either alone or in combination, slightly improved skin lesion status, although the improvement was not statistically significant. A large 2×2 factorial randomized placebo control trial is currently underway to evaluate whether supplementation with vitamin E or selenium can prevent the risk of skin cancer among 6,000 patients with skin lesions.

In a randomized, double-blind, placebo-controlled folic acid-supplementation trial of 200 folate-deficient cohort participants, we tested the hypothesis that folic acid supplementation could increase arsenic methylation and lower blood arsenic concentrations. In the trial, the increase of urinary %DMA in the folic acid group was significantly greater than that in the placebo group, as was the reduction in urinary %MMA. Furthermore, the concentrations of total blood arsenic and of MMA in the blood were significantly reduced, by 14% and 22%, respectively.

Arsenic Exposure Reduction by Mitigation Program

Since the baseline recruitment of the HEALS, we implemented interventions including: 1) person-to-person reporting of well test results and health education; 2) well labeling and village-level health education; and, 3) installations of 50 deep, low-arsenic community wells in villages with the highest levels of arsenic exposure. Two years after these interventions, 58% of the 6,512 participants with unsafe wells at baseline (arsenic > 50 $\mu\text{g/L}$) had responded by switching to other wells. Urinary arsenic levels in participants who switched to a well identified as safe (arsenic < 50 $\mu\text{g/L}$) dropped from an average of 375 to 200 $\mu\text{g arsenic/g creatinine}$, a 46% reduction towards the average urinary arsenic content of 136 $\mu\text{g arsenic/g creatinine}$ for participants that used safe wells throughout. These findings suggest that interventions such as these can effectively encourage switching to safe wells and lower arsenic exposure.

All-cause Mortality and Chronic Disease Mortality

A total of 219 participants have passed on from 2000 to 2006 in HEALS, with ages ranging from 30–70 years and a mean of 51 years. Major causes of chronic disease death within the cohort included diseases of the cardiovascular system (n=87), which primarily included hypertensive diseases, ischemic heart diseases, and cerebrovascular diseases; neoplasms (n=32); and diseases of the respiratory system (n=18), which primarily included asthma, chronic obstructive pulmonary disease, bronchiectasis, and respiratory infections. Other causes of death were related to the nervous system (n=3), the digestive system (n=14), the genitourinary system (n=8), pregnancy complications (n=5), diabetes (n=1), musculoskeletal disorders (n=1), or infectious disease (n=18).

We estimated hazard ratios for all-cause mortality and chronic disease mortality, the latter category included deaths due to cardiovascular disease, cancers, and non-infectious diseases, in relation to baseline well arsenic levels. The adjusted hazard ratios for all-cause mortality are 1.00 (ref), 1.23 (95% CI: 0.80, 1.88), 0.98 (95% CI: 0.65, 1.49) and 1.71 (95% CI: 1.16, 2.53) in increasing levels of well arsenic (<10, 10–50, 51–150, and 151–864 µg/L), respectively. The adjusted hazard ratios for chronic disease mortality are 1.00 (ref), 1.23 (95% CI: 0.80, 1.88), 0.98 (95% CI: 0.65, 1.49) and 1.71 (95% CI: 1.16, 2.53) in increasing levels of well arsenic, respectively. The hazard ratios for chronic disease mortality associated with a baseline well arsenic concentration >50 and >100 µg/L were 1.40 (0.93–2.10) and 1.61 (1.06–2.44), respectively.

DISCUSSION

Recent findings from the HEALS provide valuable information on the effects of arsenic exposure at low-to-moderate levels on arsenic-related skin lesions, children's intelligence function, blood pressure, all-cause and chronic disease mortality, and an array of biomarkers for early biological effects or altered structures and functions of target organs.

Consistent with studies of bladder and lung cancer, we also found a synergistic effect between the use of tobacco products and arsenic exposure on the risk of skin lesions. Cigarette smoking is likely to influence arsenic toxicity and should be taken into consideration in any studies on low-to-moderate levels of arsenic exposure. In addition, the data strongly suggest effect-modification roles of nutritional factors (i.e., folate and selenium) that are either involved in arsenic metabolism or have an antagonistic relationship with arsenic. While mounting evidence suggests that folate has an effect in arsenic metabolism, epidemiologic data on the mechanisms through which selenium may influence arsenic toxicity is limited. Similar to arsenic, methylation of selenium also uses SAM as the methyl donor. In addition to continue the investigation of gene expression changes due to selenium, future studies are needed to assess whether the effect of arsenic on DNA methylation is modifiable by selenium status. Intervention trials with clinical disease endpoints may be the next step in evaluating the potential of using some of these agents as treatments or preventive measures. In this regard, a large randomized clinical trial of 6,000 cases of skin lesions investigating the effects of selenium and/or vitamin E supplementation with five years of follow-up is ongoing. Future studies are needed to evaluate whether the associations of arsenic exposure with cancer, cardiovascular disease or other arsenic-related clinical and intermediate endpoints also differ by these nutritional factors.

Our observation that at a given level of As exposure, %MMA was most strongly associated with increased risk of As-induced skin lesions is consistent with studies of skin cancer, urothelial carcinoma, and bladder cancer. These data suggest a role of interindividual variability that may be regulated by arsenic methylation capacity. There is also evidence that genes coding the enzymes that catalyze the metabolism processes of arsenic can influence

the risk of arsenic-related skin lesions. Beside *MTHFR* and *GSTO1*, it is likely that there are many other candidate genes that may influence susceptibility to arsenic toxicity. Other GSTs may play a role in cellular antioxidant defense mechanisms by catalyzing the reduction of potentially harmful peroxides. Several studies have indicated that SNPs in genes for GST mu 1 (*GSTM1*) and theta 1 (*GSTT1*) were associated with urinary arsenic metabolite profiles. A case-control of 600 cases (with average exposure level at 174 µg/L) and 600 controls in Bangladesh, and a cross-sectional study of IMT with 605 subjects in Taiwan, have found that effects of arsenic on these outcomes were modifiable by polymorphisms in *glutathione S-transferase P1* gene. However, the studies did not have enough sample size to evaluate the effect-modification at low levels of exposure (< 50 or <100 µg/L). More recently, arsenic (+III) methyl-transferase (AS3MT) was characterized in rodents as an arsenic methyltransferase capable of methylating inorganic arsenic to its monomethyl form, and monomethylarsenic to its dimethylarsenic form. In a study of 147 individuals in northern Argentina, three intronic polymorphisms in the *AS3MT* gene were associated with a lower %MMA and a higher %DMA in urine. Although studies of arsenic-related skin lesions and urinary arsenic metabolites point to the role of genetic susceptibility, case-control studies of skin or bladder cancer generated mixed results, most probably due to the limited sample sizes (number of cases < 400). Taken together, additional studies with larger sample size and comprehensive genomic approach are needed to systemically evaluate gene-arsenic interactions, not only the risk of skin lesions but also on other arsenic-related disease conditions.

The studies of biomarkers for early biological effects or altered structures may aid in the recognition of early effects of arsenic exposure, as well as the elucidation of underlying pathogenic mechanisms. For instance, the study of serum CC16 shows promise as a biomarker for assessing early respiratory damage induced by arsenic. Our results confirm findings from several *in vivo* and *in vitro* studies. For instance, consistent with studies that have shown arsenic to be an inducer of TNF, a trigger of chemokine signal responses including chemokine factor CCL20, we observed down-regulation of *CCL20* and *TNF* genes in skin lesion cases. ICAM-1 and VCAM-1 expression in human umbilical vein endothelial cells (HUVEC) was higher after stimulation with arsenic. In mice tumors, treatment with As trioxide was associated with a clear increase in expression of ICAM-1 and VCAM-1. To our knowledge, our studies on EGFR and CC16 are the first published studies that link arsenic exposure with these biomarkers. However, aspects of the data are consistent with other reports in the literature. For example, individuals at risk for cancer from carcinogen exposures are shown to have increased levels of serum EGFR ECD years prior to their clinical diagnosis of malignancy. Serum CC16 concentrations are decreased in individuals with compromised lung condition induced by chronic environmental exposures such as cigarette smoking or ozone. Reliable biomarkers may also serve as intermediate endpoints for intervention trials, arsenic mitigation programs, screening and/or large observational epidemiologic studies. Analyses of participants with available data for both biomarkers and disease endpoints may be useful to further support biological plausibility of the putative causal pathways linking arsenic exposure to disease.

The observed positive associations between arsenic exposure and biomarkers relevant to cardiovascular disease and respiratory illness underscore the potentially adverse effects of arsenic exposure at low-to-moderate levels on these conditions. In a study of 463 subjects from the high-exposure area in southwestern Taiwan, a dose-response relationship was observed between arsenic exposure and carotid atherosclerosis assessed by duplex ultrasonography. The only epidemiologic evidence of low-to-moderate levels of arsenic exposure on cardiovascular disease came from the results of a cross-sectional study of > 8,000 subjects in northeastern coast of Taiwan that showed an elevated risk of cerebrovascular disease associated with arsenic exposure level at < 50 and 50–300 µg/L.

Because IMT is a reliable marker of atherosclerosis, which is a strong risk factor of both coronary heart disease and stroke and, larger studies of IMT are needed to assess the influence of the low-to-moderate levels of arsenic.

Using the hazard ratios of chronic disease mortality for a baseline well arsenic concentration >50 and >100 µg/L and the prevalence of these exposure levels, we estimated population attributable risk (PAR%); 18.1% and 13.1% of chronic disease mortality in the population can be attributable to arsenic exposure >50 and > 100 µg/L, respectively. Apparently, arsenic exposure from drinking water is associated with a significant proportion of chronic disease mortality in the overall population, especially in those with comparatively high exposure levels. In future, with a longer follow-up of the HEALS, we will be able to further assess the dose-response relationship between arsenic exposure and cause-specific mortality. The data will be valuable, given that existing studies of arsenic exposure at low-to-moderate levels and disease endpoints mostly include limited sample size and unreliable or group-level long-term measures of arsenic exposure.

Importantly, the use of multiple exposure measures, including well arsenic, urinary arsenic and blood arsenic, with repeated measurements in HEALS, improves the ascertainment of long-term exposure levels. The availability of detailed exposure data also provides a unique opportunity to evaluate the health effects of changes in exposure status over time and the joint status of difference exposure measures. In light of complex data, future analyses will need to employ modern epidemiology techniques, such as longitudinal analyses methods, survival analyses methods with time-dependent variables and/or marginal structural models to enhance causal inferences.

In conclusion, findings from the HEALS to date indicate that arsenic exposure has a profound influence on the health of the arsenic-exposed population in Bangladesh. Our findings on skin lesions, biomarkers and susceptibility altogether indicate that health effects of arsenic exposure at low-to-moderate levels may be cause for further investigation. Future analyses of clinical endpoints of cancer and cardiovascular disease are likely to provide useful knowledge on dose-response relationships. The HEALS will continue to be a valuable resource for the investigation of the health effects of arsenic exposure for years to come.

Acknowledgments

The research is supported by NIH grants: P42ES010349, R01CA102484, R01CA107431, R01ES011601, P30ES09089, CA016087, ES000260, and CA014599.

Reference List

- Adams DH, Shaw S. Leucocyte-endothelial interactions and regulation of leucocyte migration. *Lancet*. 1994; 343:831–836. [PubMed: 7908081]
- Ahsan H, Chen Y, Kibriya MG, Slavkovich V, Parvez F, Jasmine F, Gamble MV, Graziano JH. Arsenic metabolism, genetic susceptibility, and risk of premalignant skin lesions in Bangladesh. *Cancer Epidemiol Biomarkers Prev*. 2007; 16:1270–1278. [PubMed: 17548696]
- Ahsan H, Chen Y, Parvez F, Argos M, Hussain AI, Momotaj H, Levy D, van Geen A, Howe G, Graziano J. Health Effects of Arsenic Longitudinal Study (HEALS): description of a multidisciplinary epidemiologic investigation. *J. Expo. Sci. Environ. Epidemiol*. 2006a; 16:191–205. [PubMed: 16160703]
- Ahsan H, Chen Y, Parvez F, Zablotska L, Argos M, Hussain I, Momotaj H, Levy D, Cheng Z, Slavkovich V, van Geen A, Howe GR, Graziano JH. Arsenic exposure from drinking water and risk of premalignant skin lesions in Bangladesh: baseline results from the Health Effects of Arsenic Longitudinal Study. *Am J Epidemiol*. 2006b; 163:1138–1148. [PubMed: 16624965]

- Argos M, Kibriya MG, Parvez F, Jasmine F, Rakibuz-Zaman M, Ahsan H. Gene expression profiles in peripheral lymphocytes by arsenic exposure and skin lesion status in a Bangladeshi population. *Cancer Epidemiol Biomarkers Prev.* 2006; 15:1367–1375. [PubMed: 16835338]
- Argos M, Parvez F, Chen Y, Hussain AZ, Momotaj H, Howe GR, Graziano JH, Ahsan H. Socioeconomic status and risk for arsenic-related skin lesions in Bangladesh. *Am. J. Public Health.* 2007; 97:825–831. [PubMed: 17395836]
- Bernard AM, Roels HA, Buchet JP, Lauwerys RR. Serum Clara cell protein: an indicator of bronchial cell dysfunction caused by tobacco smoking. *Environ. Res.* 1994; 66:96–104. [PubMed: 8013441]
- Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE. Common carotid intima-media thickness and risk of stroke and myocardial infarction: the Rotterdam Study. *Circulation.* 1997; 96:1432–1437. [PubMed: 9315528]
- British Geological Survey, Natural Environment Research Council. Groundwater studies for arsenic contamination in Bangladesh—summary of phase 1 report. 1999. (<http://www.bgs.ac.uk/arsenic/>).
- Chen CJ, Hsu LI, Wang CH, Shih WL, Hsu YH, Tseng MP, Lin YC, Chou WL, Chen CY, Lee CY, Wang LH, Cheng YC, Chen CL, Chen SY, Wang YH, Hsueh YM, Chiou HY, Wu MM. Biomarkers of exposure, effect, and susceptibility of arsenic-induced health hazards in Taiwan. *Toxicol. Appl. Pharmacol.* 2005; 206:198–206. [PubMed: 15967209]
- Chen CL, Hsu LI, Chiou HY, Hsueh YM, Chen SY, Wu MM, Chen CJ. Ingested arsenic, cigarette smoking, and lung cancer risk: a follow-up study in arseniasis-endemic areas in Taiwan. *JAMA.* 2004b; 292:2984–2990. [PubMed: 15613666]
- Chen Y, Ahsan H, Parvez F, Howe GR. Validity of a food-frequency questionnaire for a large prospective cohort study in Bangladesh. *Br. J. Nutr.* 2004a; 92:851–859. [PubMed: 15533275]
- Chen Y, Factor-Litvak P, Howe GR, Graziano JH, Brandt-Rauf P, Parvez F, van Geen A, Ahsan H. Arsenic exposure from drinking water, dietary intakes of B vitamins and folate, and risk of high blood pressure in Bangladesh: a population-based, cross-sectional study. *Am. J. Epidemiol.* 2007b; 165:541–552. [PubMed: 17164464]
- Chen Y, Graziano JH, Parvez F, Hussain I, Momotaj H, van Geen A, Howe GR, Ahsan H. Modification of Risk of Arsenic-Induced Skin Lesions by Sunlight Exposure, Smoking, and Occupational Exposures in Bangladesh. *Epidemiology.* 2006a; 17:459–467. [PubMed: 16755266]
- Chen Y, Hakim ME, Parvez F, Islam T, Rahman AM, Ahsan H. Arsenic exposure from drinking-water and carotid artery intima-medial thickness in healthy young adults in Bangladesh. *J Health Popul. Nutr.* 2006b; 24:253–257. [PubMed: 17195567]
- Chen Y, Hall M, Graziano JH, Slavkovich V, van Geen A, Parvez F, Ahsan H. A prospective study of blood selenium levels and the risk of arsenic-related premalignant skin lesions. *Cancer Epidemiol Biomarkers Prev.* 2007a; 16:207–213. [PubMed: 17301251]
- Chen Y, Santella RM, Kibriya MG, Wang Q, Kappil M, Verret WJ, Graziano JH, Ahsan H. Association between Arsenic Exposure from Drinking Water and Plasma Levels of Soluble Cell Adhesion Molecules. *Environ. Health Perspect.* 2007c; 115:1415–1420. [PubMed: 17938729]
- Chen Y, van Geen A, Graziano JH, Pfaff A, Madajewicz M, Parvez F, Hussain AZ, Slavkovich V, Islam T, Ahsan H. Reduction in urinary arsenic levels in response to arsenic mitigation efforts in Araihasar, Bangladesh. *Environ. Health Perspect.* 2007d; 115:917–923. [PubMed: 17589600]
- Chen YC, Guo YL, Su HJ, Hsueh YM, Smith TJ, Ryan LM, Lee MS, Chao SC, Lee JY, Christiani DC. Arsenic methylation and skin cancer risk in southwestern Taiwan. *J Occup. Environ. Med.* 2003b; 45:241–248. [PubMed: 12661181]
- Chen YC, Guo YL, Su HJ, Hsueh YM, Smith TJ, Ryan LM, Lee MS, Chao SC, Lee JY, Christiani DC. Arsenic methylation and skin cancer risk in southwestern Taiwan. *J. Occup. Environ. Med.* 2003a; 45:241–248. [PubMed: 12661181]
- Chen YC, Su HJ, Guo YL, Hsueh YM, Smith TJ, Ryan LM, Lee MS, Christiani DC. Arsenic methylation and bladder cancer risk in Taiwan. *Cancer Causes Control.* 2003c; 14:303–310. [PubMed: 12846360]
- Chen YC, Xu L, Guo YL, Su HJ, Smith TJ, Ryan LM, Lee MS, Christiani DC. Polymorphisms in GSTT1 and p53 and urinary transitional cell carcinoma in south-western Taiwan: a preliminary study. *Biomarkers.* 2004c; 9:386–394. [PubMed: 15764300]

- Chiou HY, Chiou ST, Hsu YH, Chou YL, Tseng CH, Wei ML, Chen CJ. Incidence of transitional cell carcinoma and arsenic in drinking water: a follow-up study of 8,102 residents in an arseniasis-endemic area in northeastern Taiwan. *Am J Epidemiol*. 2001; 153:411–418. [PubMed: 11226969]
- Chiou HY, Huang WI, Su CL, Chang SF, Hsu YH, Chen CJ. Dose-response relationship between prevalence of cerebrovascular disease and ingested inorganic arsenic. *Stroke*. 1997; 28:1717–1723. [PubMed: 9303014]
- Cybulsky MI, Gimbrone MA Jr. Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Science*. 1991; 251:788–791. [PubMed: 1990440]
- Das S, Santra A, Lahiri S, Guha Mazumder DN. Implications of oxidative stress and hepatic cytokine (TNF-alpha and IL-6) response in the pathogenesis of hepatic collagenesis in chronic arsenic toxicity. *Toxicol. Appl. Pharmacol*. 2005; 204:18–26. [PubMed: 15781290]
- Do T, Ambelunghe A, Ahsan H, Graziano J, Perrin M, Slavkovic V, Parvez F, Milton AH, Brandt-Rauf P. Urinary transforming growth factor-alpha in individuals exposed to arsenic in drinking water in Bangladesh. *Biomarkers*. 2001; 6:127–132. [PubMed: 23886108]
- Engel RR, Smith AH. Arsenic in drinking water and mortality from vascular disease: an ecologic analysis in 30 counties in the United States. *Arch. Environ. Health*. 1994; 49:418–427. [PubMed: 7944575]
- Gamble MV, Ahsan H, Liu X, Factor-Litvak P, Ilievski V, Slavkovich V, Parvez F, Graziano JH. Folate and cobalamin deficiencies and hyperhomocysteinemia in Bangladesh. *Am. J. Clin. Nutr*. 2005a; 81:1372–1377. [PubMed: 15941889]
- Gamble MV, Liu X, Ahsan H, Pilsner JR, Ilievski V, Slavkovich V, Parvez F, Chen Y, Levy D, Factor-Litvak P, Graziano JH. Folate and arsenic metabolism: a double-blind, placebo-controlled folic acid-supplementation trial in Bangladesh. *Am J Clin Nutr*. 2006; 84:1093–1101. [PubMed: 17093162]
- Gamble MV, Liu X, Ahsan H, Pilsner R, Ilievski V, Slavkovich V, Parvez F, Levy D, Factor-Litvak P, Graziano JH. Folate, homocysteine, and arsenic metabolism in arsenic-exposed individuals in Bangladesh. *Environ. Health Perspect*. 2005b; 113:1683–1688. [PubMed: 16330347]
- Gamble MV, Liu X, Slavkovich V, Pilsner JR, Ilievski V, Factor-Litvak P, Levy D, Alam S, Islam M, Parvez F, Ahsan H, Graziano JH. Folic acid supplementation lowers blood arsenic. *Am J Clin Nutr*. 2007; 86:1202–1209. [PubMed: 17921403]
- Germolec DR, Spalding J, Boorman GA, Wilmer JL, Yoshida T, Simeonova PP, Bruccoleri A, Kayama F, Gaido K, Tennant R, Burleson F, Dong W, Lang RW, Luster MI. Arsenic can mediate skin neoplasia by chronic stimulation of keratinocyte-derived growth factors. *Mutat. Res*. 1997; 386:209–218. [PubMed: 9219559]
- Greenland S, Robins J. Invited commentary: ecologic studies--biases, misconceptions, and counterexamples. *Am. J. Epidemiol*. 1994; 139:747–760. [PubMed: 8178788]
- Griffin RJ, Lee SH, Rood KL, Stewart MJ, Lyons JC, Lew YS, Park H, Song CW. Use of arsenic trioxide as an antivasular and thermosensitizing agent in solid tumors. *Neoplasia*. 2000; 2:555–560. [PubMed: 11228548]
- Hafeman DM, Ahsan H, Louis ED, Siddique AB, Slavkovich V, Cheng Z, van Geen A, Graziano JH. Association between arsenic exposure and a measure of subclinical sensory neuropathy in Bangladesh. *J. Occup. Environ. Med*. 2005; 47:778–784. [PubMed: 16093927]
- Hall M, Chen Y, Ahsan H, Slavkovich V, van Geen A, Parvez F, Graziano J. Blood arsenic as a biomarker of arsenic exposure: results from a prospective study. *Toxicology*. 2006; 225:225–233. [PubMed: 16860454]
- Hall M, Gamble M, Slavkovich V, Liu X, Levy D, Cheng Z, van Geen A, Yunus M, Rahman M, Pilsner JR, Graziano J. Determinants of arsenic metabolism: blood arsenic metabolites, plasma folate, cobalamin, and homocysteine concentrations in maternal-newborn pairs. *Environ. Health Perspect*. 2007; 115:1503–1509. [PubMed: 17938743]
- Heck JE, Gamble MV, Chen Y, Graziano JH, Slavkovich V, Parvez F, Baron JA, Howe GR, Ahsan H. Consumption of folate-related nutrients and metabolism of arsenic in Bangladesh. *Am J Clin Nutr*. 2007; 85:1367–1374. [PubMed: 17490975]

- Hou YC, Hsu CS, Yeh CL, Chiu WC, Pai MH, Yeh SL. Effects of glutamine on adhesion molecule expression and leukocyte transmigration in endothelial cells exposed to arsenic. *J Nutr. Biochem.* 2005; 16:700–704. [PubMed: 16084078]
- Hsueh YM, Chiou HY, Huang YL, Wu WL, Huang CC, Yang MH, Lue LC, Chen GS, Chen CJ. Serum beta-carotene level, arsenic methylation capability, and incidence of skin cancer. *Cancer Epidemiol. Biomarkers Prev.* 1997; 6:589–596. [PubMed: 9264271]
- Huang YK, Pu YS, Chung CJ, Shiue HS, Yang MH, Chen CJ, Hsueh YM. Plasma folate level, urinary arsenic methylation profiles, and urothelial carcinoma susceptibility. *Food Chem. Toxicol.* 2008; 46:929–938. [PubMed: 18054417]
- Hwang SJ, Ballantyne CM, Sharrett AR, Smith LC, Davis CE, Gotto AM Jr, Boerwinkle E. Circulating adhesion molecules VCAM-1, ICAM-1, and E-selectin in carotid atherosclerosis and incident coronary heart disease cases: the Atherosclerosis Risk In Communities (ARIC) study. *Circulation.* 1997; 96:4219–4225. [PubMed: 9416885]
- Karagas MR, Park S, Nelson HH, Andrew AS, Mott L, Schned A, Kelsey KT. Methylene tetrahydrofolate reductase (MTHFR) variants and bladder cancer: a population-based case-control study. *Int. J. Hyg. Environ. Health.* 2005; 208:321–327. [PubMed: 16217917]
- Karagas MR, Tosteson TD, Morris JS, Demidenko E, Mott LA, Heaney J, Schned A. Incidence of transitional cell carcinoma of the bladder and arsenic exposure in New Hampshire. *Cancer Causes Control.* 2004; 15:465–472. [PubMed: 15286466]
- Kibriya MG, Jasmine F, Argos M, Verret WJ, Rakibuz-Zaman M, Ahmed A, Parvez F, Ahsan H. Changes in gene expression profiles in response to selenium supplementation among individuals with arsenic-induced pre-malignant skin lesions. *Toxicol. Lett.* 2007; 169:162–176. [PubMed: 17293063]
- Lagerkvist BJ, Bernard A, Blomberg A, Bergstrom E, Forsberg B, Holmstrom K, Karp K, Lundstrom NG, Segerstedt B, Svensson M, Nordberg G. Pulmonary epithelial integrity in children: relationship to ambient ozone exposure and swimming pool attendance. *Environ. Health Perspect.* 2004; 112:1768–1771. [PubMed: 15579425]
- Lewis DR, Southwick JW, Ouellet-Hellstrom R, Rench J, Calderon RL. Drinking water arsenic in Utah: A cohort mortality study. *Environ. Health Perspect.* 1999; 107:359–365.
- Li Y, Chen Y, Slavkovic V, Ahsan H, Parvez F, Graziano JH, Brandt-Rauf PW. Serum levels of the extracellular domain of the epidermal growth factor receptor in individuals exposed to arsenic in drinking water in Bangladesh. *Biomarkers.* 2007; 12:256–265. [PubMed: 17453740]
- McCarty KM, Chen YC, Quamruzzaman Q, Rahman M, Mahiuddin G, Hsueh YM, Su L, Smith T, Ryan L, Christiani DC. Arsenic methylation, GSTT1, GSTM1, GSTP1 polymorphisms, and skin lesions. *Environ. Health Perspect.* 2007; 115:341–345. [PubMed: 17431481]
- Meliker JR, Wahl RL, Cameron LL, Nriagu JO. Arsenic in drinking water and cerebrovascular disease, diabetes mellitus, and kidney disease in Michigan: a standardized mortality ratio analysis. *Environ. Health.* 2007; 6:4. [PubMed: 17274811]
- Milton AH, Smith W, Rahman B, Hasan Z, Kulsum U, Dear K, Rakibuddin M, Ali A. Chronic arsenic exposure and adverse pregnancy outcomes in bangladesh. *Epidemiology.* 2005; 16:82–86. [PubMed: 15613949]
- Mitra SR, Mazumder DN, Basu A, Block G, Haque R, Samanta S, Ghosh N, Smith MM, von Ehrenstein OS, Smith AH. Nutritional factors and susceptibility to arsenic-caused skin lesions in West Bengal, India. *Environ. Health Perspect.* 2004; 112:1104–1109. [PubMed: 15238285]
- Moore LE, Wiencke JK, Bates MN, Zheng S, Rey OA, Smith AH. Investigation of genetic polymorphisms and smoking in a bladder cancer case-control study in Argentina. *Cancer Lett.* 2004; 211:199–207. [PubMed: 15219943]
- Nixon DE, Mussmann GV, Eckdahl SJ, Moyer TP. Total arsenic in urine: palladium-persulfate vs nickel as a matrix modifier for graphite furnace atomic absorption spectrophotometry. *Clin. Chem.* 1991; 37:1575–1579. [PubMed: 1893592]
- Partanen R, Hemminki K, Koskinen H, Luo JC, Carney WP, Brandt-Rauf PW. The detection of increased amounts of the extracellular domain of the epidermal growth factor receptor in serum during carcinogenesis in asbestosis patients. *J Occup. Med.* 1994; 36:1324–1328. [PubMed: 7884573]

- Parvez F, Chen Y, Argos M, Hussain AZ, Momotaj H, Dhar R, van Geen A, Graziano JH, Ahsan H. Prevalence of arsenic exposure from drinking water and awareness of its health risks in a Bangladeshi population: results from a large population-based study. *Environ. Health Perspect.* 2006; 114:355–359. [PubMed: 16507457]
- Parvez F, Chen Y, Brandt-Rauf PW, Bernard A, Dumont X, Slavkovich V, Argos M, D'Armiento J, Foronjy R, Hasan MR, Eunos HE, Graziano JH, Ahsan H. Nonmalignant respiratory effects of chronic arsenic exposure from drinking water among never-smokers in Bangladesh. *Environ. Health Perspect.* 2008; 116:190–195. [PubMed: 18288317]
- Pilsner JR, Liu X, Ahsan H, Ilievski V, Slavkovich V, Levy D, Factor-Litvak P, Graziano J, Gamble M. Folate deficiency, hyperhomocysteinemia, low urinary creatinine and hypomethylation of leukocyte DNA are risk factors for arsenic-induced skin lesions. *Environ. Health Perspect.* 2008 In press.
- Pilsner JR, Liu X, Ahsan H, Ilievski V, Slavkovich V, Levy D, Factor-Litvak P, Graziano JH, Gamble MV. Genomic methylation of peripheral blood leukocyte DNA: influences of arsenic and folate in Bangladeshi adults. *Am. J. Clin. Nutr.* 2007; 86:1179–1186. [PubMed: 17921400]
- Pu YS, Yang SM, Huang YK, Chung CJ, Huang SK, Chiu AW, Yang MH, Chen CJ, Hsueh YM. Urinary arsenic profile affects the risk of urothelial carcinoma even at low arsenic exposure. *Toxicol. Appl. Pharmacol.* 2007; 218:99–106. [PubMed: 17196235]
- Rahman M, Tondel M, Ahmad SA, Axelson O. Diabetes mellitus associated with arsenic exposure in Bangladesh. *Am. J. Epidemiol.* 1998; 148:198–203. [PubMed: 9676702]
- Saha KC. Diagnosis of arsenicosis. *J Environ. Sci Health Part A Tox. Hazard. Subst. Environ. Eng.* 2003; 38:255–272.
- Schlawicke EK, Broberg K, Concha G, Nermell B, Warholm M, Vahter M. Genetic polymorphisms influencing arsenic metabolism: evidence from Argentina. *Environ. Health Perspect.* 2007; 115:599–605. [PubMed: 17450230]
- Smith AH, Lingas EO, Rahman M. Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. *Bull. World Health Organ.* 2000; 78:1093–1103. [PubMed: 11019458]
- Spiekstra SW, Toebak MJ, Sapat-Sardjoepersad S, van Beek PJ, Boorsma DM, Stoof TJ, von Blomberg BM, Scheper RJ, Bruynzeel DP, Rustemeyer T, Gibbs S. Induction of cytokine (interleukin-1 α and tumor necrosis factor- α) and chemokine (CCL20, CCL27, and CXCL8) alarm signals after allergen and irritant exposure. *Exp. Dermatol.* 2005; 14:109–116. [PubMed: 15679580]
- Stehouwer CD, Henry RM, Dekker JM, Nijpels G, Heine RJ, Bouter LM. Microalbuminuria is associated with impaired brachial artery, flow-mediated vasodilation in elderly individuals without and with diabetes: further evidence for a link between microalbuminuria and endothelial dysfunction—the Hoorn Study. *Kidney Int. Suppl.* 2004:S42–S44. [PubMed: 15485416]
- Steinmaus C, Moore LE, Shipp M, Kalman D, Rey OA, Biggs ML, Hopenhayn C, Bates MN, Zheng S, Wiencke JK, Smith AH. Genetic polymorphisms in MTHFR 677 and 1298, GSTM1 and T1, and metabolism of arsenic. *J Toxicol. Environ. Health A.* 2007; 70:159–170. [PubMed: 17365577]
- Steinmaus C, Yuan Y, Bates MN, Smith AH. Case-control study of bladder cancer and drinking water arsenic in the western United States. *Am. J. Epidemiol.* 2003; 158:1193–1201. [PubMed: 14652304]
- Tondel M, Rahman M, Magnuson A, Chowdhury IA, Faruquee MH, Ahmad SA. The relationship of arsenic levels in drinking water and the prevalence rate of skin lesions in Bangladesh. *Environ. Health Perspect.* 1999; 107:727–729. [PubMed: 10464073]
- Tseng WP, Chu HM, How SW, Fong JM, Lin CS, Yeh S. Prevalence of skin cancer in an endemic area of chronic arsenicism in Taiwan. *J Natl. Cancer Inst.* 1968; 40:453–463. [PubMed: 5644201]
- Verret WJ, Chen Y, Ahmed A, Islam T, Parvez F, Kibriya MG, Graziano JH, Ahsan H. A randomized, double-blind placebo-controlled trial evaluating the effects of vitamin E and selenium on arsenic-induced skin lesions in Bangladesh. *J Occup. Environ. Med.* 2005; 47:1026–1035. [PubMed: 16217243]
- Wang CH, Jeng JS, Yip PK, Chen CL, Hsu LI, Hsueh YM, Chiou HY, Wu MM, Chen CJ. Biological gradient between long-term arsenic exposure and carotid atherosclerosis. *Circulation.* 2002; 105:1804–1809. [PubMed: 11956123]

- Wang YH, Wu MM, Hong CT, Lien LM, Hsieh YC, Tseng HP, Chang SF, Su CL, Chiou HY, Chen CJ. Effects of arsenic exposure and genetic polymorphisms of p53, glutathione S-transferase M1, T1, and P1 on the risk of carotid atherosclerosis in Taiwan. *Atherosclerosis*. 2006;305–312. [PubMed: 16973168]
- Wasserman GA, Liu X, Parvez F, Ahsan H, Factor-Litvak P, Kline J, van Geen A, Slavkovich V, Loiacono NJ, Levy D, Cheng Z, Graziano JH. Water arsenic exposure and intellectual function in 6-year-old children in Arai hazar, Bangladesh. *Environ. Health Perspect*. 2007; 115:285–289. [PubMed: 17384779]
- Wasserman GA, Liu X, Parvez F, Ahsan H, Factor-Litvak P, van Geen A, Slavkovich V, Loiacono NJ, Cheng Z, Hussain I, Momotaj H, Graziano JH. Water arsenic exposure and children's intellectual function in Arai hazar, Bangladesh. *Environ. Health Perspect*. 2004; 112:1329–1333. [PubMed: 15345348]
- Yang L, Wang W, Hou S, Peterson PJ, Williams WP. Effects of Selenium Supplementation on Arsenism: An Intervention Trial in Inner Mongolia. *Environmental Geochemistry and Health*. 2002; 24:359–374.
- Yu RC, Hsu KH, Chen CJ, Froines JR. Arsenic methylation capacity and skin cancer. *Cancer Epidemiol Biomarkers Prev*. 2000; 9:1259–1262. [PubMed: 11097236]
- Zablotska LB, Chen Y, Graziano JH, Parvez F, van Geen A, Howe GR, Ahsan H. Protective effects of B vitamins and antioxidants on the risk of arsenic-related skin lesions in bangladesh. *Environ. Health Perspect*. 2008; 116:1056–1062. [PubMed: 18709164]
- Zierold KM, Knobeloch L, Anderson H. Prevalence of chronic diseases in adults exposed to arsenic-contaminated drinking water. *Am. J. Public Health*. 2004; 94:1936–1937. [PubMed: 15514231]

Table 1

Findings from HEALS on Determinants of Urinary or Blood Arsenic Metabolites

Reference	Study Design	Characteristics of Subjects	Main Findings
Heck et al, 2007 Ahsan et al, 2007	Cross-sectional	1,041 randomly selected subjects free of skin lesions	<ul style="list-style-type: none"> • Positive associations of intakes of cysteine, methionine, calcium, protein and vitamin B-12 with urinary %MMA • Positive associations of intakes of niacin, choline with urinary DMA-to-MMA ratios • Positive associations between water arsenic and urinary %MMA, and inverse association between water arsenic and urinary %DMA
Gamble et al, 2005b	Cross-sectional	300 randomly selected participants	<ul style="list-style-type: none"> • Positive correlation between plasma folate and urinary %DMA, between total homocysteine (tHcys) and urinary %MMA, and between urinary creatinine and urinary %DMA • Inverse correlation tHcys and urinary %DMA, and between plasma folate and urinary %MMA
Gamble et al, 2005 Gamble et al, 2007	Randomized, double-blind, placebo-controlled trial	200 folate-deficient cohort participants	<ul style="list-style-type: none"> • Greater increase of urinary %DMA and reduction in urinary %MMA in the folic acid group than in the placebo group • Significant reduction in total blood arsenic and of MMA in the blood in folic acid group.

Table 2

Findings from HEALS on the Risk of Arsenic-related Pre-malignant Skin Lesions

Reference	Study Design	Characteristics of Subjects	Arsenic Measurement	Main Findings
Ahsan et al, 2006b	Cross-sectional	714 cases of skin lesion and 10,724 non-skin lesion subjects	Water, urinary arsenic, time-weighted arsenic concentration	<ul style="list-style-type: none"> Dose-response relationship between arsenic exposure and risk of skin lesions Synergistic effect between high levels of arsenic exposure and male gender, low BMI, and old age.
Chen et al, 2006a	Cross-sectional	714 cases of skin lesion and 10,724 non-skin lesion subjects	Water, urinary arsenic, time-weighted arsenic concentration	<ul style="list-style-type: none"> Synergistic effect between the highest level of arsenic exposure and tobacco smoking in men. Additive effects of sun exposure and arsenic exposure in men
Argos et al, 2007	Cross-sectional	714 cases of skin lesion and 10,724 non-skin lesion subjects	Water, urinary arsenic	<ul style="list-style-type: none"> Effect-modification by land ownership on the multiplicative scale
Zablotska et al, 2008	Cross-sectional	714 cases of skin lesion and 10,724 non-skin lesion subjects	Water, urinary arsenic	<ul style="list-style-type: none"> Effect-modification by high intake of riboflavin, pyridoxine, folate, and vitamins A, C and E
Ahsan et al, 2007	Case-control	594 skin lesion cases and 1,041 controls	Water, urinary arsenic, urinary %MMA, %DMA, MMA/DMA	<ul style="list-style-type: none"> Stronger dose-response relationship with urinary %MMA compared with other methylation indices Effect differs by <i>MTHFR</i> and <i>GSTO1</i> diplotypes
Chen et al, 2007a	Prospective case-cohort	303 incident skin lesion cases and 849 subcohort members	Water, urinary arsenic	<ul style="list-style-type: none"> An inverse association between baseline blood selenium status and the incidence of skin lesions Additive effect of high arsenic exposure and low selenium status.
Hall et al, 2006	Prospective case-cohort	303 incident skin lesion cases and 849 subcohort members	Water, urinary, and blood arsenic	<ul style="list-style-type: none"> Dose-response relationship between baseline blood and well arsenic and incidence of skin lesions
Pilsner et al, 2007	Prospective nested case-control study	274 skin lesion cases 274 matched controls	Water, blood, urinary arsenic	<ul style="list-style-type: none"> Positive relationships of folate deficiency, hyperhomocysteinemia, low urinary creatinine, with skin lesion risk