

ORIGINAL ARTICLE

Plasma fibrinogen: now also an antidepressant response marker?

D Martins-de-Souza^{1,2,3}, G Maccarrone¹, M Ising¹, S Kloiber¹, S Lucae¹, F Holsboer¹ and CW Turck¹

Major depressive disorder (MDD) is one of the leading causes of global disability. It is a risk factor for noncompliance with medical treatment, with about 40% of patients not responding to currently used antidepressant drugs. The identification and clinical implementation of biomarkers that can indicate the likelihood of treatment response are needed in order to predict which patients will benefit from an antidepressant drug. While analyzing the blood plasma proteome collected from MDD patients before the initiation of antidepressant medication, we observed different fibrinogen alpha (FGA) levels between drug responders and nonresponders. These results were replicated in a second set of patients. Our findings lend further support to a recently identified association between MDD and fibrinogen levels from a large-scale study.

Translational Psychiatry (2014) 4, e352; doi:10.1038/tp.2013.129; published online 28 January 2014

Keywords: antidepressant; biomarkers; drug response; fibrinogen; proteomics

INTRODUCTION

Major depressive disorder (MDD) is a leading cause of global disability and a risk factor for noncompliance with medical treatment.¹ As a multivariate disorder presenting a wide range of symptoms, MDD treatment success varies among patients. About 40% of MDD patients do not respond to current treatments. Moreover, there is a high rate of relapse and treatment resistance,² and MDD patients have a high risk of attempting suicide.³

The identification and clinical implementation of biomarkers that can indicate the likelihood of treatment response are needed to improve this situation.⁴ In a recent article, Wium-Andersen *et al.*⁵ reported an association of elevated plasma fibrinogen levels and MDD. In the present study, we have asked the question whether fibrinogen levels can also serve as a predictive marker for the antidepressant treatment response. We used a mass spectrometry-based proteomics screen of blood plasma samples from MDD patients before antidepressant treatment. We found altered levels of fibrinogen alpha chain (FGA) in the plasma from drug responders compared to nonresponders. These differences were quantified and compared by western blot.

MATERIALS AND METHODS

Subjects

Initially, we analyzed the proteome of plasma specimens from 25 inpatients suffering from MDD, who participated in the Munich Antidepressant Response Signature (MARS) project.⁶ A first round of validation was performed in 17 patients—8 responders and 9 nonresponders, followed by a second round of validation in a separate set of 24 patients—16 responders and 8 nonresponders (Table 1). The two cohorts were built based on the time of sample collection in the clinic. While the first round of experiments was performed, samples from the second cohort were collected with no bias. Details of the study were explained to patients and a written informed consent was obtained. The study was approved by the ethics committee of the Medical Faculty at the *Ludwig-Maximilians*

University Munich. Patients were diagnosed by trained psychiatrists according to the DSM-IV criteria. Antidepressant treatment outcome was weekly evaluated with the 21 item version of the Hamilton Rating Scale for Depression (HAM-D). Response was defined as a 50% or larger reduction of the HAM-D score after 6 weeks of treatment compared with the HAM-D score on admission to the hospital of the *Max Planck Institute of Psychiatry*. Two patients were discharged with no HAM-D ratings at T6. Their weekly ratings at an earlier time point were used to classify them as responder or nonresponder. The mean age for responders and nonresponders was 45 and 47 years, respectively. No significant differences were observed in this regard between the two groups ($P=0.6398$, Mann–Whitney). Significant differences between the two groups were also not observed regarding body mass index (BMI: $P=0.2239$, Mann–Whitney), cholesterol levels ($P=0.6883$, Mann–Whitney) and gender ($P=1.000$, two-sided Fisher's exact test). None of the study subjects had diabetes. When admitted to the clinic, the majority of patients were not taking any anti-inflammatory or immunosuppressant medications. Only three patients were using nonsteroidal anti-inflammatory drugs (Supplementary Table 1).

Blood plasma samples

Blood samples were collected shortly after admission to the hospital in sample tubes containing potassium EDTA. Plasma was separated from blood using an Accuspin System HystopaqueTM-1077 (Sigma Aldrich, Taufkirchen, Germany) according to the manufacturer's protocol and protein concentrations determined by Bradford assay.

Western blot

Twenty micrograms of total protein extracts from each blood plasma sample was run individually on a 12% SDS minigel (BioRad, Hercules, CA, USA). Proteins were transferred to Immobilon PVDF membrane (Millipore, Bedford, MA, USA) at 100 V for 1 h using a cooling system. Membranes were treated with 5% Carnation Instant Nonfat Dry Milk in TBS-T for 4 h, rinsed in TBS-T three times for 20 min and incubated with an FGA antibody (Abcam, Cambridge, UK) at a 1:1000 dilution in TBS-T overnight at 4 °C. After the incubation, membranes were washed twice for 15 min with TBS-T. Next, membranes were incubated with an anti-c-MYC-peroxidase antibody

¹Max Planck Institute of Psychiatry, Proteomics and Biomarkers, Munich, Germany and ²Department of Psychiatry and Psychotherapy, Ludwig-Maximilians-University Munich, Munich, Germany. Correspondence: Professor CW Turck, Max Planck Institute of Psychiatry, Proteomics and Biomarkers, Kraepelinstr, 2-10, Munich 80804, Germany. E-mail: turck@mpipsykl.mpg.de

³Current address: Laboratory of Neuroproteomics, Department of Biochemistry, Institute of Biology, State University of Campinas (UNICAMP), Campinas, SP, Brazil.

Received 29 June 2013; revised 13 December 2013; accepted 14 December 2013

Table 1. Patients' clinical data including gender, age, BMI, plasma cholesterol levels and HAM-D

Patient ID	Gender	Age	BMI	Cholesterol	HAM-D T0	HAM-D T6	Western blot relative signal
<i>First experiment</i>							
<i>Nonresponders</i>							
1	M	48	25.15	207	29	22	67.37
2	M	51	25.39	187	27	22	49.30
3	M	28	28.28	187	26	14	58.03
4	M	58	27.58	215	36	35	60.41
5	F	33	20	203	24	18	64.59
6	F	53	24.49	211	28	22	59.27
7	F	54	19.61	192	27	17	68.12
8	F	39	26.76	219	35	28	60.20
<i>Responders</i>							
9	F	70	24.73	208	35	11	52.99
10	M	31	25	214	21	0	46.30
11	M	68	28.08	215	32	3	52.82
12	F	36	19.72	194	20	6	57.45
13	F	25	18.71	154	28	1	55.81
14	M	67	23.12	171	38	3	50.86
15	F	21	18.59	114	33	0	36.01
16	F	41	20	175	30	7	36.88
17	M	33	28.41	123	33	13	31.39
<i>Second experiment</i>							
<i>Nonresponders</i>							
18	M	75	23.80	172	28	16	65.07
19	F	36	24.80	135	21	14	66.15
20	M	53	28.04	162	30	20	53.03
21	F	22	20.25	188	33	23	63.51
22	M	22	25.54	190	22	25	56.01
23	F	43	24.10	190	18	NA	51.74
24	F	68	24.51	185	24	24	57.75
25	M	72	27.28	190	29	25	56.11
<i>Responders</i>							
26	F	49	24.98	211	21	2	32.83
27	F	25	20.20	208	34	10	59.28
28	M	43	24.89	181	29	10	51.37
29	M	34	20.05	132	22	0	55.71
30	M	87	26.07	155	27	4	63.12
31	F	39	22.28	136	21	9	64.87
32	F	24	18.75	209	39	14	44.64
33	M	68	24.88	157	21	10	55.68
34	M	58	28.09	210	34	8	50.45
35	M	31	23.80	207	41	20	49.50
36	M	44	26.94	211	25	0	37.13
37	M	49	24.30	200	25	9	28.55
38	M	62	24.38	207	30	10	32.66
39	F	56	24.54	185	28	NA	51.11
40	F	39	27.18	215	25	6	51.10
41	F	35	19.96	124	30	12	51.14

Abbreviations: BMI, body mass index; HAM-D, Hamilton Rating Scale for Depression; F, female; M, male; NA, not applicable.

(GE Healthcare, Uppsala, Sweden) for 40 min at RT, washed with water and TBS-T, and incubated with ECL solution (GE Healthcare) for 1 min. Membranes were subsequently scanned using a Gel Doc XR+ System (BioRad). The optical densities of the FGA band were assessed using Quantity One software (BioRad). Afterwards, the Immobilon PVDF membrane was stained with Coomassie Blue to ascertain equal protein loading in each gel lane. In addition, the Coomassie Blue-stained serum albumin protein band was used as an internal control for equal protein loading.

Statistics

All analyses were carried out using SPSS statistical software (IBM, Armonk, NY, USA). According to the D'Agostino & Pearson normality test, the two groups evaluated here presented a normal distribution. For the first group, the *P*-value for nonresponders is 0.6100 and for responders 0.3716; for the

second group, the *P*-value for nonresponders is 0.7213 and for responders 0.5509. Considering that 95% of the Gaussian distributed samples should be within two s.d. of the mean, values that differed more than two s.d. from the mean were excluded from the analysis (Chauvenet's Criterion). Differences between the groups were assessed using analysis of covariance (ANCOVA) considering BMI and age as covariates. *P* < 0.05 was considered significant.

RESULTS AND DISCUSSION

The plasma proteome analyses by mass spectrometry of antidepressant responders and nonresponders revealed altered FGA levels at baseline (data not shown). In the present study, we confirmed this result by western blot analysis of crude plasma samples from two separate patient groups (Table 1 and Figure 1).




Figure 1. Western blot FGA protein band densities of nonresponders and responders before treatment. (a) First patient validation set, with BMI ($P=0.149$) and age ($P=0.114$) as covariates; (b) second patient validation set with BMI ($P=0.174$) and age ($P=0.452$) as covariates. Insets show representative western blot images. $*P < 0.05$; $**P < 0.01$.

As our aim was to find a generic antidepressant response biomarker, we did not take into account the type of antidepressant used by each patient. Consistent with the results from our initial mass spectrometry analysis, we found higher FGA levels in nonresponders compared with responders in both patient groups (Figures 1a and b). BMI and age were considered as covariates and did not influence the results (Figure 1).

Our findings add further support to recently published data on an association between elevated plasma fibrinogen levels and depression in 73 367 subjects.^{5,7} Other studies have also shown an association of acute phase proteins including fibrinogen with MDD.^{8,9} A meta-analysis of plasma cytokine levels in MDD patients has found elevated TNF- α and IL-6 levels in depressed patients compared with healthy controls.¹⁰ Also, C-reactive protein, IL-1 and IL-6 have been associated with depression.¹¹ All these results suggest an activated inflammatory status in MDD¹² that may affect serotonergic neurotransmission.^{13,14} Other studies suggest an association of fibrinogen with MDD pathogenesis. Patients with coronary heart disease presenting depression symptoms tended to have elevated fibrinogen levels.¹⁵ Also, male MDD patients with hypertension had lower fibrinogen levels.¹⁶ Our data add one more piece to this puzzle, suggesting that baseline plasma fibrinogen levels can serve as a biomarker to gauge the success of antidepressant treatment response.

The expression of fibrinogen in hepatocytes depends on acute phase inflammatory stimuli.¹⁷ Our results show that high plasma fibrinogen levels are associated with a poor antidepressant response and indicate that nonresponders present an elevated inflammatory status compared with responders. Indeed, other studies also suggest that MDD patients with increased inflammatory protein levels tend to be treatment resistant.¹⁸ An increased production of acute phase proteins, such as fibrinogen, causes the activation of the indoleamine 2,3 dioxygenase, which catabolizes tryptophan, thus decreasing serotonin availability.¹⁹ High levels of circulating fibrinogen are also associated with and targeted in the treatment of cardiovascular disease.²⁰ Similarly, in the case of antidepressant resistance, MDD patients could be subjected to medication that modulates fibrinogen levels^{21,22} before drug treatment in order to increase the likelihood of response. The

administration of aspirin for example is currently used in the treatment of cardiovascular diseases. Aspirin promotes fibrinogen acetylation, which alters its clotting properties.²³ Acetylated fibrinogen is a less active acute phase protein²⁴, which might affect serotonin levels.

Although preliminary, our data obtained from a naturalistic set of samples imply that plasma FGA levels are not only of value as a biomarker for depression⁵ but can also serve as a predictive antidepressant response marker. This could have potential for clinical implementation following validation in larger sample cohorts.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

We thank Gertrud Ernst-Jansen, Karin Hofer, Elisabeth Kappelman, Maria Lebar and Christiane Rewerts for excellent technical assistance with clinical samples.

REFERENCES

- DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for non-compliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med* 2000; **160**: 2101–2107.
- Rosenzweig-Lipson S, Beyer CE, Hughes ZA, Khawaja X, Rajarao SJ, Malberg JE *et al*. Differentiating antidepressants of the future: efficacy and safety. *Pharmacol Ther* 2007; **113**: 134–153.
- Nrugham L, Hohen A, Sund AM. Suicide attempters and repeaters: depression and coping: a prospective study of early adolescents followed up as young adults. *J Nerv Ment Dis* 2012; **200**: 197–203.
- Martins-de-Souza D, Harris LW, Guest PC, Turck CW, Bahn S. The role of proteomics in depression research. *Eur Arch Psy Clin Neurosci* 2010; **260**: 499–506.
- Wium-Andersen MK, Orsted DD, Nordestgaard BG. Association between elevated plasma fibrinogen and psychological distress, and depression in 73 367 individuals from the general population. *Mol Psychiatry* 2012; **18**: 854–855.
- Hennings JM, Owashi T, Binder EB, Horstmann S, Menke A, Kloiber S *et al*. Clinical characteristics and treatment outcome in a representative sample of depressed inpatients—findings from the Munich Antidepressant Response Signature (MARS) project. *J Psychiatr Res* 2009; **43**: 215–229.
- Wium-Andersen MK, Orsted DD, Nordestgaard BG. Elevated plasma fibrinogen, psychological distress, antidepressant use, and hospitalization with depression: Two large population-based studies. *Psychoneuroendocrinology* 2012; **38**: 638–647.
- Panagiotakos DB, Pitsavos C, Chrysohoou C, Tsetsekou E, Papageorgiou C, Christodoulou G *et al*. Inflammation, coagulation, and depressive symptomatology in cardiovascular disease-free people; the ATTICA study. *Eur Heart J* 2004; **25**: 492–499.
- Maes M, Delange J, Ranjan R, Meltzer HY, Desnyder R, Cooremans W *et al*. Acute phase proteins in schizophrenia, mania and major depression: modulation by psychotropic drugs. *Psychiatry Res* 1997; **66**: 1–11.
- Dowlati Y, Herrmann N, Swardfager W, Liu H, Sham L, Reim EK *et al*. A meta-analysis of cytokines in major depression. *Biol Psychiatry* 2010; **67**: 446–457.
- Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom Med* 2009; **71**: 171–186.
- Krishnadas R, Cavanagh J. Depression: an inflammatory illness? *J Neurol Neurosurg Psychiatry* 2012; **83**: 495–502.
- Muller N, Myint AM, Schwarz MJ. Inflammatory biomarkers and depression. *Neurotox Res* 2011; **19**: 308–318.
- Dantzer R, O'Connor JC, Lawson MA, Kelley KW. Inflammation-associated depression: from serotonin to kynurenine. *Psychoneuroendocrinology* 2011; **36**: 426–436.
- Duijvis HE, de Jonge P, Penninx BW, Na BY, Cohen BE, Whooley MA. Depressive symptoms, health behaviors, and subsequent inflammation in patients with coronary heart disease: prospective findings from the heart and soul study. *Am J Psychiatry* 2011; **168**: 913–920.
- Baune BT, Neuhauser H, Ellert U, Berger K. The role of the inflammatory markers ferritin, transferrin and fibrinogen in the relationship between major depression and cardiovascular disorders—The German Health Interview and Examination Survey. *Acta Psychiatr Scand* 2010; **121**: 135–142.

- 17 Vasse M, Paysant J, Soria J, Collet JP, Vannier JP, Soria C. Regulation of fibrinogen biosynthesis by cytokines, consequences on the vascular risk. *Haemostasis* 1996; **26**(Suppl 4): 331–339.
- 18 Miller AH, Maletic V, Raison CL. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol Psychiatry* 2009; **65**: 732–741.
- 19 Catena-Dell'Osso M, Rotella F, Dell'Osso A, Fagiolini A, Marazziti D. Inflammation, serotonin and major depression. *Curr Drug Targets* 2013; **14**: 571–577.
- 20 Fish RJ, Neerman-Arbez M. Fibrinogen gene regulation. *Thromb Haemost* 2012; **108**: 419–426.
- 21 Yusuf S, Lessem J, Jha P, Lonn E. Primary and secondary prevention of myocardial infarction and strokes: an update of randomly allocated, controlled trials. *J Hypertens Suppl* 1993; **11**: S61–S73.
- 22 Meade TW. Haemostatic function and arterial disease. *Br Med Bull* 1994; **50**: 755–775.
- 23 He S, Blomback M, Yoo G, Sinha R, Henschen-Edman AH. Modified clotting properties of fibrinogen in the presence of acetylsalicylic acid in a purified system. *Ann N Y Acad Sci* 2001; **936**: 531–535.
- 24 Caspary EA. Studies on the acetylation of human fibrinogen. *Biochem J* 1956; **62**: 507–512.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplementary Information accompanies the paper on the Translational Psychiatry website (<http://www.nature.com/tp>)