

# The “*Fungia patella* group” (Scleractinia, Fungiidae) revisited with a description of the mini mushroom coral *Cycloseris boschmai* sp. n.

Bert W. Hoeksema<sup>1,†</sup>

<sup>1</sup> Department of Marine Zoology, Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

<sup>†</sup> <http://zoobank.org/548DBAFD-040B-4773-A043-6FCC466160A1>

Corresponding author: Bert W. Hoeksema (bert.hoeksema@naturalis.nl)

---

Academic editor: L.P. van Ofwegen | Received 25 November 2013 | Accepted 2 January 2014 | Published 17 January 2014

<http://zoobank.org/66723DA9-2152-41F0-96F0-56C0F75C1CBD>

---

**Citation:** Hoeksema BW (2014) The “*Fungia patella* group” (Scleractinia, Fungiidae) revisited with a description of the mini mushroom coral *Cycloseris boschmai* sp. n. ZooKeys 371: 57–84. doi: 10.3897/zookeys.371.6677

---

## Abstract

The recent taxonomic history of extant free-living *Cycloseris* species is briefly reviewed, resulting in the description of *Cycloseris boschmai* sp. n. (Scleractinia, Fungiidae) and a discussion on the validity of two other recently described species. Some *Cycloseris* species were previously considered to belong to the *Fungia patella* group, which also concerned misidentified museum specimens that actually belong to the new species. Other specimens of *C. boschmai* sp. n. were photographed and collected in the course of 30 years of fieldwork. The new mushroom coral is compared with other free-living *Cycloseris* species by means of an identification key. With a maximum diameter of 50 mm, it is the smallest free-living mushroom coral discovered so far. It can also be distinguished by its large primary order costae and variable colouration. Its distribution range is limited to the Coral Triangle, where it can be observed as an uncommon species on lower reef slopes.

## Keywords

Coral reef, free-living, fungiid, solitary, budding, collection, fieldwork, Coral Triangle, Indo-Pacific

## Introduction

Mushroom corals (Scleractinia, Fungiidae) form a common element in the fauna of most Indo-Pacific coral reefs. Depending on the species, full-grown specimens are either attached or free-living, which are character states occurring in various evolutionary lineages and therefore do not necessarily reflect phylogenetic relationships among the Fungiidae (Cairns 1984, Hoeksema 1989, 1991b, 2009, Gittenberger et al. 2011, Benzoni et al. 2012). After settlement, each fungiid individual starts as a small attached coral (anthocaulus). While corals of attached species obtain a foliaceous or encrusting growth form (Hoeksema 1989, 1993a, 2009, Gittenberger et al. 2011, Benzoni et al. 2012), those of free-living species eventually become detached from their stalk, reaching the anthocyathus stage (Wells 1966, Hoeksema 1989, Hoeksema and Yeemin 2011, Hoeksema and Waheed 2012). Owing to their charismatic appearance, abundance and large polyp size, these free-living fungiids are usually easy to find. By growing large and occurring in high densities, they may form mono- or multi-species assemblages covering extensive reef areas (Littler et al. 1997, Hoeksema 2004, 2012a, Elahi 2008, Hoeksema and Koh 2009, Hoeksema and Gittenberger 2010, Hoeksema and Matthews 2011, Hoeksema and Benzoni 2013). They can live on various kinds of reef substrates, ranging from silt to solid rock, from nearshore to offshore and from shallow reef flats to deep reef bases (e.g., Hoeksema and Moka 1989, Hoeksema 1991a, Hoeksema 2012a, 2012b, Waheed and Hoeksema 2013, in press). Mushroom corals themselves may in turn act as habitat to associated faunas consisting of molluscs, crabs, shrimps, acoel flatworms, comb jellies, and fish, some of which are host-specific (Hoeksema and Kleemann 2002, Kleemann and Hoeksema 2002, Gittenberger and Gittenberger 2005, 2011, Hoeksema and Fransen 2011, Owada and Hoeksema 2011, Bos 2012, Hoeksema et al. 2012, 2013a, 2013b, Hoeksema and Farenzena 2012, Gittenberger and Hoeksema 2013, Van der Meij and Hoeksema 2013). The various mushroom coral species vary in size (Hoeksema 1989, 1991b, Gittenberger et al. 2011) and it is obvious that corals with large surface area and thick skeletons offer most habitat space for associated fauna in contrast to small species (Hoeksema et al. 2012).

The smallest species among free-living mushroom corals appear to be among the most difficult to identify because they show relatively few distinguishing characters and much ecophenotypic variation (Hoeksema and Moka 1989, Hoeksema 1993d). Döderlein (1901, 1902) classified them as the “*patella* group” within the genus *Fungia* Lamarck, 1801. He considered them to be the most “primitive” species owing to their small size, imperforate (solid) corallum wall, and rudimentary, hardly discernible costal spines and septal dentations (see Scholtz et al. 2012). The fossil record of this *F. patella* group could be traced back to the Cretaceous, while its distribution ranged from eastern Africa to the west coast of America and its maximum depth was known to be over 100 m (Döderlein 1901).


Previously, species of the *F. patella* group were considered to belong to the genera *Cycloseris* Milne Edwards & Haime, 1849, consisting of complete corals, and *Diaseris* Milne Edwards & Haime, 1849, representing radially fragmenting corals of the

same species (see Hoeksema 1989). Döderlein (1902) already considered complete and fragmenting corals to be different forms that can be found within each species. He recognized the following six species and synonyms: 1. *F. patella* Ellis & Solander, 1786 (including *F. patellaris* Lamarck, 1801, *F. tenuis* Dana, 1846, *F. hexagonalis* Milne Edwards & Haime, 1849, *Diaseris fragilis* Alcock, 1893); 2. *F. erosa* Döderlein, 1901; 3. *F. distorta* Michelin, 1842; 4. *F. cyclolites* Lamarck, 1816; 5. *F. elegans* Verrill, 1868; 6. *F. costulata* Ortmann, 1889. Some authors still consider *Diaseris* a separate genus (e.g. Veron 2000, Claereboudt 2006) while there is morphological evidence for their synonymy (Hoeksema 1989, Hoeksema and Waheed 2011, 2012).

Oddly, both *F. patella* and *F. patellaris* are junior synonyms of *F. fungites* (Linnaeus, 1756), the type species of *Fungia* (see Hoeksema 1989). However, Döderlein's (1902) photographs of his *F. patella* corals show that he referred to complete and fragmenting specimens of the free-living species *Cycloseris fragilis* (Alcock, 1893) and *C. sinensis* Milne Edwards & Haime, 1851 (see Hoeksema 1989). The outlines of Döderlein's (1902: pls. 1–2) smallest specimens are either circular or hexagonal.

Van der Horst (1921) mentioned six specimens of *Fungia patella* (*sensu* Döderlein), which he obtained from the Siboga Expedition collections: “The 3 *Cycloseris*-forms of Stat. 315 have both surfaces flat. Two of these specimens have the ribs obviously only at the edge of the corallum, the centrum of the under surface being irregularly covered by fine granulations. The septa of these two specimens are equal in height and very much grained. They have the appearance of *F. patella* var. *filigrana* Död. In the third specimen (dimensions 52 × 52 m.M.) the ribs reach the centrum, the scar of the attachment is here obvious. The edge of the corallum is slightly undulating.”

Boschma (1923a) had access to only the first two of these specimens and also to mushroom corals collected during the Danish Expedition to the Kei Islands in the Banda Sea, eastern Indonesia, and to other corals from Hawaii, among which were small specimens that according to him belonged to “Döderlein's, *patella*-group”. He recognized eight separate species besides *F. patella* (*sensu* Döderlein 1902), which previously were considered synonyms: 1. *F. patelliformis* Boschma, 1923; 2. *F. fragilis* (Alcock, 1893); 3. *F. marginata* Boschma, 1923; 4. *F. distorta* (Michelin, 1842); 5. *Fungia* sp.; 6. another *Fungia* sp.; 7. *Stephanophyllia neglecta* Boschma, 1923; 8. *F. vaughani* Boschma, 1923. Several species were represented by complete and fragmenting shapes. Among the complete corals, several had a hexagonal outline, such as juvenile specimens of *F. tenuis* (Boschma 1923a: pl. 9), which actually belong to *F. hexagonalis* (see Boschma 1925, Hoeksema 1989). He had many specimens available to him that he identified as *F. marginata*; some of these were illustrated together with the original description (Boschma 1923a) and other ones in a subsequent publication (Boschma 1925). In the diagnosis of *F. marginata*, Boschma (1923a) refers to a circular corallum outline, although he also states that young stages are hexagonal in shape. According to him this species could be distinguished by its thick corallum margin, which might be true if compared with for instance his *Fungia patelliformis* (a synonym of *Cycloseris fragilis* (see Hoeksema 1989) but not with some other species, such as *C. costulata* (see Hoeksema 1989).



**Figure 1.** Lectotype of *Fungia marginata* Boschma, 1923 (ZMA Coel. 604, ethanol), which is a specimen of *Cycloseris costulata* Ortmann, 1889, collected at Siboga Expedition Sta. 315, Anchorage East of Sailus Besar, in the Paternoster islands, Indonesia. **a** Upper side **b** Lower side **c** Collection labels indicating the first identification by Van der Horst (*Fungia patella*) and the later one by Boschma (*Fungia marginata*). Scale bar: 0.5 cm.

The two specimens (ZMA Coel. collection) earlier described by Van der Horst (1921) as *F. patella* from Siboga Expedition Station 315 in the Paternoster Islands, Indonesia were considered syntypes (Van Soest 1979). One of these syntypes is a coral with a circular outline and a diameter of 48.5 mm (ZMA Coel. 604, herein designated lectotype; Figure 1), whereas the other is a smaller specimen with hexagonal outline, a diameter of 13 mm, and relatively thick primary costae (ZMA Coel. 723, herein designated paralectotype; Figure 2). Based on these two types, *F. marginata* is considered a junior synonym of *F. costulata* (Hoeksema 1989). Because of its doubtful identity, the juvenile specimen is not useful as type. Boschma (1923a) assumed that *F. costulata* does not have a solid corallum wall and has more or less equal costae and therefore described *F. marginata* as a species with a solid corallum wall, unequal costae, and thick corallum margin as distinguishing characters. Actually, the type of *F. costulata* does have a solid wall, whereas costae of all mushroom coral species may be dissimilar in size, including *F. costulata* (Hoeksema 1989, Gittenberger and Hoeksema 2006). Furthermore, the





**Figure 2.** Paralectotype of *Fungia marginata* Boschma, 1923 (ZMA Coel. 723, dry), which may be a juvenile specimen of *Cycloseris costulata* Ortmann, 1889 or *C. hexagonalis* Milne Edwards & Haime, 1849, collected at Siboga Expedition Sta. 315, Anchorage East of Sailus Besar, in the Paternoster islands, Indonesia **a** Specimen with collection box and labels; scale bar: 1 cm. Collection labels indicating the first identification by Van der Horst (*Fungia patella*) and the later one by Boschma (*Fungia marginata*) **b** Close-up upper side **c** Close-up lower side.

identity of *F. marginata* has never been really clear, because most subsequent authors confused it with other species (see Hoeksema 1989).

Boschma (1925) had access to many more *F. marginata* specimens according to the accompanying identification labels, but he did not mention them in the original species description. These specimens are still available in the collections of the Zoological Museum - University of Copenhagen (UZMK) and Naturalis Biodiversity Center (RMNH Coel.). Among these corals, several specimens are slightly larger than the syntypes and they actually do show enlarged lower order costae, although this was not clearly illustrated in his plates (Boschma 1925: pls. 5–6). These specimens are re-examined in the present study along with many small mushroom corals collected during various recent field surveys (1983–2013). The last ones were striking because of their outstanding long and thick costae and their colorful appearance in comparison with the usual brown hues found in individuals of related species (Hoeksema 1989, Gittenberger and Hoeksema 2006). Based on this unique set of characters consisting of a relatively small adult size, enlarged primary order costae and variable coloration, these corals are considered to belong to a new species, herein described as *Cycloseris boschmai* sp. n.

All species of the former *Fungia patella*-group belong to *Cycloseris*, previously considered a subgenus of *Fungia* (see Hoeksema 1989). As a consequence of recent molecular phylogenetic studies, so far 13 species have been recognized in the genus *Cycloseris*,

three of which are attached (Gittenberger et al. 2011, Benzoni et al. 2012) and not part of the old *F. patella*-group. The new species, here described as the free-living *Cycloseris boschmai* sp. n., is the fourteenth. Specimens were either available in museum collections or were photographed and sampled during SCUBA diving in the field. In addition to a detailed description, many illustrations are presented to show the phenotypic variation of this small *Cycloseris* species, following in the foot steps of Döderlein (1902) and Boschma (1923a, 1925). For the purpose of specimen identification, a key to the presently known free-living extant *Cycloseris* species is given.

## Abbreviations

BWH = B.W. Hoeksema; Exp. = Expedition; I. = Island; Sta. = Station; MTQ = Museum of Tropical Queensland, Townsville, Australia; RMNH Coel. = Rijksmuseum van Natuurlijke Historie, Coelenterate collection (Naturalis Biodiversity Center), Leiden, the Netherlands; UZMK = Zoological Museum - University of Copenhagen, Denmark; ZMA Coel. = Zoological Museum of Amsterdam Coelenterate collection (Naturalis Biodiversity Center), Leiden, the Netherlands.

## Systematic section

**Order Scleractinia Bourne, 1900**

**Suborder Fungiina Verrill, 1856**

**Superfamily Fungiicae Dana, 1846**

**Family Fungiidae Dana, 1846**

**Genus *Cycloseris* Milne Edwards & Haime, 1849**

<http://species-id.net/wiki/Cycloseris>

**Type species.** *Fungia cyclolites* Lamarck 1816. Designation by monotypy.

**Synonymy.** *Cycloseris* Milne Edwards and Haime 1849: 72; Milne Edwards and Haime 1850: xlix; Milne Edwards and Haime 1851: 111–112; Milne Edwards 1860: 49; Tenison-Woods 1878: 328; Duncan 1883: 149–150; Quelch 1886: 119–120; Gardiner 1899: 171; Gardiner 1905: 944; Vaughan and Wells 1943: 139; Wells 1956: 388; Wells 1966: 235–236; Veron and Pichon 1980: 107–108; Ditlev 1980: 54; Nemenzo 1981: 182; Scheer and Pillai 1983: 74; Nemenzo 1986: 140; Pillai 1986: 153; Veron 1986: 320–321; Chevalier and Beauvais 1987: 710; Veron 1992: 123; Veron 1993: 199; Latypov 1995: 88; Nishihira and Veron 1995: 234; Veron 2000: 236; Suharsono 2004: 191; Claereboudt 2006: 187; Latypov 2006: 178; Suharsono 2008: 215; Wallace et al. 2009: 46.

*Diaseris* Milne Edwards and Haime 1849: 72; Milne Edwards and Haime 1850: xlix; Milne Edwards and Haime 1851: 117; Milne Edwards 1860: 54–55; Duncan 1883: 150; Gardiner 1905: 945; Veron and Pichon 1980: 119–121; Veron 1986: 326–

327; Veron 1992: 127; Veron 1993: 205; Latypov 1995: 95; Nishihira and Veron 1995: 239; Veron 2000: 248; Suharsono 2004: 197; Claereboudt 2006: 190; Latypov 2006: 185; Suharsono 2008: 222. (Type species: *Fungia distorta* Michelin, 1842. Designation by monotypy.

*Fungia* (*Cycloseris*) – Hoeksema 1989: 30–31; Hoeksema and Van Ofwegen 2004.

**Characters.** Adult corals either encrusting and polystomatous or free-living and monostomatous (Gittenberger et al. 2011, Benzoni et al. 2012). Outline of free-living, unfragmented specimens varying from circular to oval. Juveniles may be hexagonal. Free-living corals may fracture repeatedly into regenerating wedge-shaped pieces (Hoeksema 1989, Yamashiro et al. 1989, Yamashiro and Nishihira 1994, 1998, Hoeksema and Waheed 2011, 2012). Fragmenting corals may produce extra stomata along fracture lines. Corallum wall without perforations. Septal margins ornamented by fine, sharp dentations. Costae covered by fine spiny protuberances, which may become granular and blunt in large specimens. Tentacles small and usually translucent in extended state.

***Cycloseris boschmai* sp. n.**

<http://zoobank.org/8FA4CA99-7074-4425-A7ED-D051D6AB3311>

[http://species-id.net/wiki/Cycloseris\\_boschmai](http://species-id.net/wiki/Cycloseris_boschmai)

Figures 3–13

*Fungia marginata* (partim) Boschma 1923: 141–142; 1925: 199–202.

*Fungia* (*Cycloseris*) *costulata* (partim) – Hoeksema 1989: 64–69.

*Fungia* (*Cycloseris*) spec. – Hoeksema et al. 2004: 15; Hoeksema 2008: 11–12; 2010: 24–25.


*Cycloseris* sp. 1 – Gittenberger et al. 2011: 117; Hoeksema 2012a: 188.

*Cycloseris* sp. – Hoeksema et al. 2012: 652.


*Cycloseris* spec. – Waheed and Hoeksema 2013: 41.

**Type material.** Type specimens of *Cycloseris boschmai* from Banda, Moluccas, Indonesia (Danish Exp. to the Kei Islands, 1922), previously identified as *Fungia marginata* by Boschma 1925). **Holotype:** RMNH Coel. 8333 (1 dry specimen: 28 mm; Figure 3), Banda, 1922. **Paratypes:** RMNH Coel. 8334 (1 dry specimen: 29 mm), Banda, Lontor, 12.vi.1922; RMNH Coel. 8335 (5 dry specimens: 20–30 mm), Banda, off Lontor, 10–20 m depth, 4°33'S, 129°52'E, 1922.

**Other material: Indonesia. Bali:** RMNH Coel. 40146 (2 dry specimens: 25, 31 mm), NE Bali, Tulamben, 3–5m, 08°16'36"S, 115°35'37"E, Bali Lombok Strait Exp. Sta. BAL.20, 09.iv.2001, coll. BWH; RMNH Coel. 40147 (1 dry specimen: 18 mm), NE Bali, Tulamben, 5m depth, 08°16'26"S, 115°35'28"E, Bali Lombok Strait Exp. Sta. BAL.21, 11.iv.2001, coll. BWH. **Nusa Tenggara Timur (Lesser Sunda Islands):** RMNH Coel. 21471 (1 dry specimen: 38 mm), NE Komodo, S Gili Lawa Laut, 08°27'00"S, 119°34'24"E, Snellius-II Exp. Sta. 4.253, 27.x.1984,



**Figure 3.** Holotype of *Cycloseris boschmai* sp. n. (RMNH Coel. 8333). Indonesia, Banda, Danish Exp. to the Kei Islands, 1922. **a** Upper side **b** Lower side. Scale bar: 0.5 cm.




**Figure 4.** Specimen of *Cycloseris boschmai* sp. n. (RMNH Coel. 21471). Indonesia, NE Komodo, S Gili Lawa Laut, Snellius-II Exp. Sta. 4.253, 27 October 1984. **a** Upper side **b** Lower side. Scale bar: 0.5 cm.


coll. BWH; RMNH Coel. 40145 (4 dry specimens: 24–32 mm), SE Komodo, N side bay S of Tanjung Lohnamu, 08°38'19"S, 119°28'45" E, TNC Komodo Rapid Ecological Assessment Sta. KOM.16, coll. BWH; RMNH Coel. 31190 (1 dry specimens: 32 mm), N Sumbawa, Bay of Sanggar, 08°19'36"S, 118°15'12"E, Snellius-II Exp. Sta. 4.132, 30.x.1984, coll. BWH. **South Sulawesi.** RMNH Coel. 31188 (1 dry specimen: 31 mm), Spermonde Archipelago, N Bone Tambung Island, 05°01'50"S, 119°16'25"E, 13.vi.1986, coll. BWH; RMNH Coel. 31189 (5 dry specimens: 29–50 mm), Spermonde Archipelago, W Kudingareng Keke I., 9–18 m depth, 05°06'30"S, 119°17'04"E, 6.xii.1984, coll. BWH; RMNH Coel. 31192



(5 dry specimens: 17–31 mm), SW Selayar I., NW Bahuluang I., 06°28'00"S, 120°25'30"E, Snellius-II Exp. Sta. 4.202, 10.x.1984, coll. BWH; RMNH Coel. 31191 (3 dry specimens: 35–37 mm), NE Taka Bone Rate, E Tarupa Besar, 06°28'S, 121°08'E, Snellius-II Exp. Sta. 4.140, 25.ix.1984, coll. BWH. **Central Sulawesi, Tomini Bay, Togian Islands:** RMNH Coel. 24278 (2 dry specimens: 34, 39 mm), S Talatakoh I., 00°26'34"S, 122°06'07"E, Tethyana Exp. Sta. 10, 21.ix.1999, coll. BWH; RMNH Coel. 24291 (1 dry specimen: 48 mm), S Togian I., 00°20'10"S, 121°59'00"E, Tethyana Exp. Sta. 14, 23/24.ix.1999, coll. BWH; RMNH Coel. 24706 (5 dry specimens: 30–42 mm), S Batudaka I., 00°35'25"S, 121°41'38"E, Tethyana Exp. Sta. 15, 24.ix.1999, coll. BWH; RMNH Coel. 31193 (7 dry specimens: 29–50 mm), S Waleabahi I., 00°26'16"S, 122°15'16"E, Tethyana Exp. Sta. 8, 19.ix.1999, coll. BWH. **North Sulawesi:** RMNH Coel. 40156 (4 dry specimens: 36–40 mm), Lembah Strait, Tanjung Mawali, 14 m depth, 01°26'36"N, 125°13'46"E, Lembah Strait Exp. Sta. LEM.04, 31.i.2012, coll. BWH. **SE Sulawesi, Tukang Besi Islands (Wakatobi):** RMNH Coel. 40143 (1 dry specimen: 12 mm), NW Tomia, 05°43'59"S, 123°53'35"E, TNC-WWF Wakatobi Rapid Ecological Assessment Sta. WAK.25, 13.v.2003, coll. BWH; RMNH Coel. 40144 (1 dry specimen: 40 mm), SW Karang Kaledupa, lagoon, 05°51'46"S, 123°43'17"E, TNC-WWF Wakatobi Rapid Ecological Assessment Sta. WAK.28, 14.v.2003, coll. BWH. **Moluccas:** RMNH Coel. 33586 (2 dry specimens: 47, 58 mm), Ambon, N coast near Morela, 03°33'S, 128°12'E, Fauna Malesiana Maluku Exp. Sta. MAL.12, 13.xi.2002, coll. BWH; **Northern Moluccas:** RMNH Coel. 8286 (10 dry specimens, some with buds, 32–50 mm, previously identified as *Fungia marginata*), Banda, off Lontor, to 13 m depth, 4°33'S, 129°52'E, Danish Exp. to the Kei Islands, 15.vi.1922; UZMK (5 specimens in ethanol, 13–40 mm, previously identified as *Fungia marginata*), Gunung Api, 20–25 m depth, Danish Exp. to the Kei Islands, 13.vi.1922; UZMK (9 specimens in ethanol, 26–50 mm, previously identified as *Fungia marginata*), Lontor, 13 m depth, Danish Exp. to the Kei Islands, 15.vi.1922; RMNH Coel. 40096 (1 dry specimen: 29 mm), Halmahera, East coast Teluk Dodinga, Karang Galiasa, 00°50'46"N, 127°35'07"E, Ternate Exp. Sta. TER.38, 14.xi.2009, coll. BWH; RMNH Coel. 40102 (1 dry specimen: 34 mm), Hiri I., Tanjung Ngafauda, 00°54'38"N, 127°19'03"E, Ternate Exp. Sta. TER.14, 16 m depth, 31.x.2009, coll. BWH; RMNH Coel. 40103 (1 dry specimen: 30 mm), Ternate, Sulamadaha I., 00°52'04"N, 127°19'33"E, Ternate Exp. Sta. TER.22, 18 m depth, 6.xi.2009, coll. BWH; RMNH Coel. 40104 (1 dry specimen: 19 mm), Ternate, Dufadufa, Benteng Toloko, 00°48'49"N, 127°23'22"E, Ternate Exp. Sta. TER.24, 8 m depth, 7.xi.2009, coll. BWH; RMNH Coel. 40105 (1 dry specimen: 31 mm), Halmahera, W Pasir Lamo, 00°53'21"N, 127°27'34"E, Ternate Exp. Sta. TER.26, 8.xi.2009, coll. BWH; RMNH Coel. 40106 (1 dry specimen: 48 mm), Tidore, north Pilonnga, 00°42'50"N, 127°28'45"E, Ternate Exp. Sta. TER.34, 12 m depth, 12.xi.2009, coll. BWH; RMNH Coel. 40173 (1 dry specimen: 29 mm), Ternate, Tanjung Tabam, 00°50'05"N, 127°23'10"E, Ternate Exp. Sta. TER.12, 11 m depth, 30.x.2009, coll. BWH; RMNH Coel. 40174 (1 dry




**Figure 5.** Specimen of *Cycloseris boschmai* sp. n. (RMNH Coel. 31190). Indonesia, N Sumbawa, Bay of Sanggar, Snellius-II Exp. Sta. 4.132, 30 October 1984 **a** Upper side **b** Lower side. Scale bar: 0.5 cm.




**Figure 6.** Specimen of *Cycloseris boschmai* sp. n. (RMNH Coel. 31188). Indonesia, South Sulawesi, Spermonde Archipelago, north side of Bone Tambung Island, 13 June 1986. **a** Upper side **b** Lower side. Scale bar: 0.5 cm.

specimen: 17 mm), Ternate, outside harbor to the east, 00°46'55"N, 127°30'20"E, Ternate Exp. Sta. TER.25, 8 m depth, 7.xi.2009, coll. BWH. **East Kalimantan, Berau Islands:** RMNH Coel. 31922 (1 dry specimen: 19 mm), W Derawan I., 7 m depth, 02°16'53"N, 118°13'39"E, East Kalimantan Berau Exp. Sta. BER.02, 4.x.2003, coll. BWH; RMNH Coel. 31923 (1 dry specimen: 15 mm), Berau Delta, Lighthouse-2 Reef, 6 m depth, 02°09'34"N, 118°10'11"E, East Kalimantan Berau Exp. Sta. BER.05, 5.x.2003, coll. BWH; RMNH Coel. 31924 (2 dry specimen: 22, 31 mm), Samama I., 8 m depth, 02°07'32"N, 118°20'10"E, East Kalimantan Berau



**Figure 7.** Specimen of *Cycloseris boschmai* sp. n. (RMNH Coel. 24291). Indonesia, Central Sulawesi, Togian Islands, S Togian I., Tethyana Exp. Sta. 14, 23/24 September 1999 **a** Upper side **b** Lower side. Scale bar: 0.5 cm.


Exp. Sta. BER.10, 7.x.2003, coll. BWH; RMNH Coel. 31925 (1 dry specimen: 42 mm), NE Kakaban I., 14 m depth, 02°08'53" 118°32'32"E, East Kalimantan Berau Exp. Sta. BER.49, 28.x.2003, coll. BWH; RMNH Coel. 40149 (1 dry specimen: 14 mm), S of Samama I., NE Buliulin I., 14 m depth, 02°07'07"N, 118°20'32"E, East Kalimantan Berau Exp. Sta. BER.26, 15.x.2003, coll. BWH; RMNH Coel. 40153 (1 dry specimen: 46 mm), N Maratua I., near entrance lagoon, 9 m depth, 02°14'53"N, 118°37'36"E, East Kalimantan Berau Exp. Sta. BER.29, 17.x.2003, coll. BWH; RMNH Coel. 40154 (1 dry specimen: 21 mm), S Derawan I., 13 m, 02°15'04"N, 118°15'04"E, East Kalimantan Berau Exp. Sta. BER.04a, 18.x.2003, coll. BWH; RMNH Coel. 40155 (1 dry specimen: 50 mm), E Sangalaki I., Lighthouse, 12 m depth, 02°04'54"N, 118°24'30"E, East Kalimantan Berau Exp. Sta. BER.22, 14.x.2003, coll. BWH. **West Papua, Raja Ampat Islands:** RMNH Coel. 40140 (1 dry specimen: 30 mm), S. Mansuar (Sawandarik village), 00°35'26"S, 130°36'12"E, Raja Ampat Exp. Sta. RAJ.06, 20.xi.2007, coll. BWH; RMNH Coel. 40141 (1 dry specimen: 33 mm), Yeffam I., E Penemu I., 8 m depth, 00°35'20"S, 130°17'06"E, Raja Ampat Exp. Sta. RAJ.66, 13.xii.2007, coll. BWH. **Malaysia, Sabah, northern Borneo:** RMNH Coel. 33545 (1 dry specimen: 20 mm), W Sabah, Gaya Islands off Kota Kinabalu, W Sapi I., 06°00'26"N, 116°00'13"E, 28.vii.2005, coll. BWH. **Layang-Layang:** RMNH Coel. 40095 (2 dry specimens: 18 mm attached, 37 mm free-living), Easternmost point, Sta. LAC.14, 15-25 m depth, 07°22'34"N, 113°51'15"E, 28.iii.2013, coll. BWH. **Philippines:** RMNH Coel. 24908 (2 dry specimens: 20, 32 mm), Cebu Strait, West of Bohol, NW Cabilao I., 09°53'20"N, 123°45'53"E, 2.x.1999, coll. BWH. **Palau:** RMNH Coel. 40225 (1 dry specimen), SE off Garreru I., S Goraklbud Passage, 07°19'15"N, 134°35'50"E, 29.vii.2002, coll BWH.




**Figure 8.** Specimens of *Cycloseris boschmai* sp. n. (RMNH Coel. 31193). Indonesia, Central Sulawesi, Togian Islands, S Waleabahi I., Tethyana Exp. Sta. 8, 19 September 1999 **a** Upper side **b** Lower side. Scale bar: 0.5 cm.

**Characters.** (Figures 3–13) The diameter of the examined specimens ranges between 12 and 50 mm. Corals mostly flat, moderately thick and robust. Adult animals unattached and monostomatous with septa-costae extending outside the circular to slightly oval corallum outline. Juvenile specimens vary from round to slightly hexagonal.






**Figure 9.** Two specimens of *Cycloseris boschmai* sp. n. (RMNH Coel. 24278). Indonesia, Central Sulawesi, Togian Islands, S Talatakoh I., Tethyana Exp. Sta. 10, 21 September 1999. Scale bar: 0.5 cm.



**Figure 10.** Two specimens of *Cycloseris boschmai* sp. n. (RMNH Coel. 8286) with marginal buds and sand in the mouths. Indonesia, Banda, off Lontor, Danish Exp. to the Kei Islands, 15 June 1922.

Wedge-shaped, regenerating fragments not known. The length of the fossa, measured at its bottom, is 1/9 to 1/6 of the corallum length. The columella is formed by a mingled mass of tightly to loosely packed trabeculae. Septa densely packed and straight, unequal in thickness and height. The relatively thick and high septa of lower orders are solid; they are flanked by perforated septa of higher orders. Tentacular lobes absent. Septal margins are finely ornamented with sharp and granular dentations. Their number varies




**Figure 11.** Juvenile, attached specimens of *Cycloseris boschmai* sp. n. **a** Papua New Guinea, Bismarck Sea, Madang, June 1992 **b–d** Malaysia, South China Sea, Layang Layang, Easternmost point, (RMNH Coel. 40095), 28 March 2013 **b** In situ (bleached) **c** Lower side **d** Upper side. Scale bars: 0.5 cm.


from 20 to 70 per cm. Septal sides are densely covered by fine granulations, which are irregularly dispersed or arranged in rows perpendicular to the septal margin. Compound synapticalae (fulturae) connecting the septa laterally cannot easily be distinguished because of tight septal arrangement. The solid corallum wall is granulated and may show a detachment scar. The lower side varies from flat to slightly convex. Costae unequal in size, straight and prominent near the corallum margin but less distinct at the centre. Corallum margin may be slightly undulating because of enlarged lower order costae. Costae ornamented with fine granular or acute spines. Their number varies from 15 to 80 per cm. Some individuals have small buds over their surface, especially in the proximity of the corallum margin (Figure 10). Attached juveniles (anthocaulus stage) are rare (Figure 11). The color of the living animal is variable with hues of red or green (Figures 11–13). Tentacles small and transparent with white acrospheres at their tips (Figure 13).

**Geographical distribution** (Figure 14). The distribution range is limited to the Coral Triangle (Hoeksema 2007): eastern Malaysia (Sabah and Layang Layang), east-





**Figure 12.** *Cycloseris boschmai* sp. n. **a** Indonesia, Bali, Tulamben, September 1997 **b** Philippines, Cebu, November 1999 **c–e** Indonesia, Central Sulawesi, Togian Islands, September 1999 **f** Indonesia, South Sulawesi, Spermonde Archipelago, Bone Lola reef, August 1997.



**Figure 13.** *Cycloseris boschmai* sp. n. specimen showing transparent extended tentacles with white acrospheres at their tips; Indonesia, North Sulawesi, Lembah Strait, Lobangbatu, February 2012.

ern Indonesia (from Bali to West Papua), central Philippines (Cebu Strait), Papua New Guinea (Madang Lagoon), and Palau.


**Etymology.** The species is named after the late Prof. Hilbrand Boschma, former director of the Rijksmuseum van Natuurlijke Historie (now Naturalis Biodiversity Center), who devoted much of his research time to the study of mushroom corals, including specimens of the new species.

**Diagnosis.** Adult corals small (< 50 mm) with uneven circular corallum margin owing to enlarged costae. Live specimens with variable, patchy colouration.

**Key to recent free-living *Cycloseris* species (full-grown, unfragmented specimens), partly after Hoeksema (1989)**

- 1a Lower order costae distinctly larger than other ones..... 2
- 1b Costae fine, adjacent ones equal to almost equal ..... 5
- 2a Coralla flat and thin, corallum outline circular ..... 3
- 2b Coralla thick and slightly arched, corallum outline slightly or much oval.... 4





**Figure 14.** Map of the Central Indo-Pacific indicating localities where *Cycloseris boschmai* sp. n. has been recorded.

- 3a Lower order costae thicker and longer than higher order costae, ornamentation fine (20–70 / cm), maximum corallum diameter 5 cm, habitat mostly consisting of reef slopes and sandy reef bases..... *Cycloseris boschmai* sp. n.
- 3b Lower order costae sharp, ornamentation very fine (40–80 / cm) on lower order costae and indistinct on higher order costae, maximum corallum diameter 8 cm, habitat mostly consisting of deep, sandy reef bases.....  
..... *Cycloseris vaughani*
- 4a Corallum outline slightly oval, lower order costae irregularly and roughly ornamented (20–70 / cm), maximum corallum diameter 8.5 cm, habitat consisting of upper reef slopes ..... *Cycloseris tenuis*
- 4b Corallum outline clearly oval, lower order costae sharp, costal ornamentation very fine (40–90 / cm) and nearly absent on higher order costae, maximum corallum diameter 12.5 cm, habitat mostly consisting of deep, sandy reef bases ..... *Cycloseris somervillei*
- 5a Septa densely packed and (almost) equal in height..... 6
- 5b Septa loosely packed, septa of lower orders thicker and more exsert than others ..7
- 6a Central fossa short (< 10% of corallum diameter); all septa perforated, nearly equal in size and tightly packed with little space in between them, maximum corallum diameter 8.5 cm, habitat mostly consisting of deep, sandy reef bases ..... *Cycloseris sinensis*

- 6b Length of central fossa > 10% of corallum diameter, septa of lower order solid and thicker than adjacent septa with distinct space in between them, maximum corallum diameter 7.5 cm, habitat consisting of deeper reef slopes or reef bases ..... *Cycloseris distorta*
- 7a Corallum outline oval ..... 8
- 7b Corallum outline circular or irregularly round with folds or undulations .... 9
- 8a Coralla thick; underside flat or arched, costae equal, maximum corallum diameter 9 cm, habitat consisting of lower reef slopes or sandy reef bases .....  
..... *Cycloseris cyclolites*
- 8b Coralla convex around fossa (humped), costae equal in juveniles, maximum corallum diameter 12.5 cm, habitat mostly consisting of deep, sandy reef bases ..... *Cycloseris somervillei*
- 9a Coralla with folded, undulating margin ..... 10
- 9b Coralla with regular, smooth periphery ..... 11
- 10a Coralla thin, central fossa short (< 10% of corallum diameter), margin undulating (hexagonal in juveniles) maximum corallum diameter 8.5 cm, habitat consisting of sandy reef slopes or sandy reef bases ..... *Cycloseris hexagonalis*
- 10b Coralla thick and usually strongly arched, margin with folds, maximum corallum diameter 8.5 cm, habitat consisting of lower reef slopes or sandy reef bases ..... *Cycloseris curvata*
- 11a Coralla and septa thin, adjacent costae slightly alternating in size maximum corallum diameter 15 cm, habitat mostly consisting of sandy reef bases .....  
..... *Cycloseris fragilis*
- 11b Coralla moderately thick, lower order septa thicker than others, costae nearly similar in size, maximum corallum diameter 12 cm, habitat consisting of lower reef slopes or sandy reef bases ..... *Cycloseris costulata*

## Discussion

Although some material of *Cycloseris boschmai* sp. n. was already available in museum collections (RMNH, UZMK), the species could only be discovered because of much fieldwork (1983-2013) with proportionate opportunities for observations and sampling to enable separation of the new species from resembling ones. Boschma (1923a, 1925) might have had the same species in mind when he described and studied *Fungia marginata*, but his selection of type specimens from the Paternoster Islands and the unclear comparison with other species of Döderlein's, *Fungia patella* group are not convincing. He described this species because of its supposedly thick corallum margin as compared to the other species in the *F. patella* group, but this character is not useful when applied several other *Cycloseris* species.

The *F. marginata* material from Banda is suitable as type material of the new species. Because the Banda specimens were wrongly identified by him, this does not concern a new name for an existing species but an entirely new species (Hoeksema 1993b).

These old museum specimens are not just useful as type material but also because they supply information about habitat (field data) and asexual reproduction by budding. They also constitute the oldest material of *C. boschmai*, which gives them potential historical value as baseline material in studies on changing coral faunas (Hoeksema and Koh 2009, Van der Meij et al. 2010, Hoeksema et al. 2011, Van der Meij and Visser 2011, Hoeksema and Wirtz 2013).

*Cycloseris boschmai* sp. n. is the smallest mushroom coral known so far (see key). Superficially, it resembles *C. costulata*, which has less prominent costae, a more even corallum margin (not undulating), a larger maximum size and less colourful appearance (see Hoeksema 1989, Hoeksema and Van Ofwegen 2004, Gittenberger and Hoeksema 2006). Both species can be found on reef slopes and sandy reef bases. *C. costulata* is common and wide-spread (Hoeksema 1989, 2012a) and has a variable growth form (Hoeksema and Moka 1989, Gittenberger and Hoeksema 2006), which is why much material had to be examined to find sufficient consistency in the diagnostic characters of the new species.

*C. boschmai* sp. n. also resembles *C. tenuis*, which is more oval, less colourful (see Gittenberger and Hoeksema 2006), and with lower order costae that are rougher and more irregularly ornamented. *C. tenuis* is most common on upper reef slopes (Hoeksema 2012a), whereas the rarer *C. boschmai* shows a deeper depth range. The new species also resembles *C. vaughani*, which has sharper lower order costae, a brown colouration (Hoeksema 1989, Hoeksema and Van Ofwegen 2004) and a much deeper depth range (Hoeksema 2012a).

With the inclusion of *C. boschmai* sp. n., 11 free-living *Cycloseris* species are distinguished. Since the taxonomic revision of the Fungiidae by Hoeksema (1989) various other mushroom coral species were reported as new to science (Veron 1990, 2000, 2002, Hoeksema and Dai 1991, Hoeksema 1993a, 1993c, 2009, 2012c, Latypov 1995, 2006, Ditlev 2003, Mondal and Raghunathan 2013). Two of these were originally classified as *Cycloseris* but they appear to be synonyms of previously described species and one of these is not a *Cycloseris*.

*Cycloseris colini* Veron, 2000 is a synonym of *Lithophyllon spinifer* (Claereboudt and Hoeksema 1987). The central dome and upward margins, combined with the large corallum size as described by Veron (2002) are characters commonly found in *L. spinifer* (see Hoeksema 1993a: fig. 14, Veron 2000, Hoeksema and Van Ofwegen 2004, Hoeksema 2008: fig. 9). A specimen from Palau (MTQ G55817) was designated holotype by Veron in 2002, but since the species was described in 2000 this designation was invalid (ICZN 2011). Hence, this specimen is hereby designated lectotype.

*Cycloseris densicolummelus* Latypov, 2006 has not been described in an official publication but in an electronic document that was distributed via a CD-ROM. This work should have contained a clear publication date and a statement naming at least five major publicly accessible libraries in which copies of the optical disc were to have been deposited (ICZN 2012). Since only the year of translation has been mentioned in the introduction and no names of libraries were given, this name is not valid. Although the publication by Latypov (2006) is said to be an English translation of an original

book in Russian, the latter does not mention *C. densicolummelus* but “*Cycloseris* sp. 1” (Latypov 1995: 95, pl. 28 fig. 2). The specimen indicated as “holotype”, spec. 1/95158 deposited in the Museum of Institute Marine Biology, Vladivostok 69041, Russia, is from Mai Rut Island, Gulf of Thailand. The illustration with the species description (Latypov 2006: figs 46–8) shows that it is a specimen of *C. costulata*. The well developed tentacular lobes, the intensive granulation of the lateral septal surfaces and the densely packed trabeculae of columella, which are indicated as diagnostic characters, do not really distinguish *C. densicolummelus* from *C. costulata* (see Hoeksema 1989, Gittenberger and Hoeksema 2006).

Because *C. boschmai* is a rare species (considering that most material was gathered during fieldwork in a time span of 30 years) and its geographic distribution range is restricted to the Coral Triangle, not much can be said about its ecology. Specimens are difficult to find, owing to their small body size compared to other mushroom coral species (Hoeksema 1991b, Gittenberger et al. 2011), which may be restrictive to the settlement of associated fauna and therefore none of its symbionts was reported previously (Hoeksema et al. 2012: “*Cycloseris* sp.”). In the present study, one of the photographed specimens shows a coral barnacle (Figure 11e), which is now the only known associated animal.

Small-sized free-living mushroom corals have been reported to show much mobility (Hoeksema 1988, Yamashiro and Nishihira 1995), which may help them to escape from competition for space with other organisms (Chadwick 1988, Hoeksema and De Voogd 2012). The distinctive large lower order costae may be useful as ridges in stabilizing the corals in order to prevent them to slide too rapidly downslope to deeper reef zones with sandy substrate (Hoeksema 1988). The enlarged ridge-like costae can also be seen in some other *Cycloseris* species (see key) and in the small-sized free-living deep-sea coral *Deltocyathus rotulus* (Alcock, 1898) (see Cairns and Kitahara 2012: fig 18Q).

Although free-living *Cycloseris* species were previously considered primitive, this is not the case according to their phylogeny reconstruction (Gittenberger et al. 2011). Their predominant habitat of deeper sandy substrates can also be considered an advanced trait (Hoeksema 2012b). The sand in the mouth of nearly dead specimens from Banda (Figure 10) suggests that they were collected from a sandy substrate, which may not be their preferred habitat. During the author's fieldwork (1983–2013), no specimens were observed on sandy reef bases. *C. boschmai* corals are small but not thin, which may not facilitate mobility and sediment-shedding as seen in large-polyped corals (Bongaerts et al. 2012, Erfteimeijer et al. 2012). The presence of buds in several of the specimens from Banda, in addition to the sand in their stomata, (Boschma 1925; Figure 10) may also indicate that specimens have been buried (Gilmour 2002). Specimens of *C. hexagonalis* have been observed to show a similar abundance of buds on a sandy slope in eastern Sabah (BWH personal observation 2009). Budding may be a mushroom coral's last resort of survival when its mouth is clogged and not capable of food intake anymore (Boschma 1922, 1923b, 1923c).



## Acknowledgements

I am grateful to the following people, organizations and institutes who assisted during the fieldwork or other parts of the research: Dr. Alfian Noor and Drs Willem Moka (Universitas Hasanuddin, Makassar), Dr. Carden C. Wallace (MTQ, Tethyana Expedition to the Togian Islands), Dr. Filipina B. Sotto and Dr Thomas Heeger (University of San Carlos, Cebu City), Dr Jacob van der Land (Naturalis, Snellius-II Expedition), Dr. Pat Colin and Ms Lori Colin (Coral Reef Research Foundation, Palau), Dr. Phillipa Mansell (Operation Wallacea, Wakatobi, Indonesia), Dr. Jos Pet (The Nature Conservancy, Bali), Dr Suharsono (Research Centre for Oceanography, Indonesian Institute of Sciences, Jakarta), Ms. Zarinah Waheed (Borneo Marine Research Institute, Kota Kinabalu). The research was partly financed by the Netherlands Foundation for the Advancement of Tropical Research (WOTRO, grant W77-96) and by the Schure-Beijerinck-Popping Fonds. I thank Ms. Majken T. Tøttrup and Dr. M.V. Sørensen for the loan of specimens from the UZMK. I thank Dr Carden C. Wallace and Mrs. Barbara Done for the loan of specimens from the MTQ. I am grateful to Mrs. Elly Beglinger and Mr. Koos van Egmond for their help in accessing ZMA and RMNH specimens. Comments given by two anonymous reviewers helped considerably to improve the text.

## References

- Alcock A (1893) On some newly-recorded corals from the Indian Seas. *Journal of the Asiatic Society of Bengal (Natural History)* 62: 138–149, pl. 5.
- Alcock A (1898) An account of the deep-sea Madreporaria collected by the Royal Indian marine survey ship Investigator. Trustees Indian Museum, Calcutta, 29 pp., 3 pls.
- Benzoni F, Arrigoni R, Stefani F, Reijnen BT, Montano S, Hoeksema BW (2012) Phylogenetic position and taxonomy of *Cycloseris explanulata* and *C. wellsi* (Scleractinia: Fungiidae): lost mushroom corals find their way home. *Contributions to Zoology* 81: 125–146.
- Bongaerts P, Hoeksema BW, Hay KB, Hoegh-Guldberg O (2012) Mushroom corals overcome live burial through pulsed inflation. *Coral Reefs* 31: 399. doi: 10.1007/s00338-011-0862-z
- Bos AR (2012) Symbiotic fishes (Gobiidae and Labridae) of the mushroom coral *Heliofungia actiniformis* (Scleractinia; Fungiidae). *Coral Reefs* 31: 133. doi: 10.1007/s00338-011-0834-3
- Boschma H (1922) On budding and coalescence of buds in *Fungia fungites* and *Fungia actiniformis*. *Proceedings Koninklijke Nederlandse Akademie van Wetenschappen*, Amsterdam 24: 257–268.
- Boschma H (1923a) The Madreporaria of the Siboga Expedition IV. *Fungia patella*. Siboga-Expedition XVIId. Brill, Leiden, 129–148, pls. 9–10.
- Boschma H (1923b) Experimental budding in *Fungia fungites*. *Proceedings Koninklijke Nederlandse Akademie van Wetenschappen*, Amsterdam 26: 88–96, pls. 1–2.
- Boschma H (1923c) Knospung und verwandte Erscheinungen bei *Fungia fungites* und *Fungia actiniformis*. *Treubia* 3: 149–179.

- Boschma H (1925) Madreporaria I. Fungiidae. Papers from Dr. Mortensen's, Pacific Expedition 1914–16. Videnskabelige meddelelser fra Dansk Naturhistorisk Forening 79: 185–259, pls. 5–11.
- Cairns SD (1984) An application of phylogenetic analysis to the Scleractinia: Family Fungiidae. *Paleontographica Americana* 54: 49–57.
- Cairns SD, Kitahara MV (2012) An illustrated key to the genera and subgenera of the Recent azooxanthellate Scleractinia (Cnidaria: Anthozoa), with an attached glossary. *ZooKeys* 227: 1–47. doi: 10.3897/zookeys.227.3612
- Chadwick NE (1988) Competition and locomotion in a free-living fungiid coral. *Journal of Experimental Marine Biology and Ecology* 123: 189–200. doi: 10.1016/0022-0981(88)90041-X
- Chevalier JP, Beauvais L (1987) Ordre des Scléractiniaires. In: Grassé PP (Ed) *Traité de Zoologie* tome III, fasc. 3. Masson, Paris, 679–753.
- Claereboudt MR (2006) Reef corals and coral reefs of the Gulf of Oman. Historical Association of Oman, Oman, 344 pp.
- Claereboudt M, Hoeksema BW (1987) *Fungia (Verrillofungia) spinifer* spec. nov., a new scleractinian coral (Fungiidae) from the Indo-Malayan region. *Zoologische Mededelingen, Leiden* 61: 303–309.
- Dana JD (1846-1849) United States Exploring Expedition during the years 1838–1842. *Zoophytes* 7. Lea and Blanchard, Philadelphia, 740 pp.
- Ditlev H (1980) A Field-guide to the Reef-building Corals of the Indo-Pacific. Backhuys, Rotterdam, and Scandinavian Science press, Klampenborg, 291 pp.
- Ditlev H (2003) New scleractinian corals (Cnidaria: Anthozoa) from Sabah, north Borneo. Description of one new genus and eight new species, with notes on their taxonomy and ecology. *Zoologische Mededelingen, Leiden* 77: 193–219.
- Döderlein L (1901) Die Korallengattung *Fungia*. *Zoologischer Anzeiger* 24: 351–360.
- Döderlein L (1902) Die Korallengattung *Fungia*. *Abhandlungen der Senckenbergischen naturforschenden Gesellschaft* 27(1): 1–162, pls. 1–25.
- Duncan PM (1883) Observations on the madreporarian family the Fungidae with special reference to the hard structures. *Journal of the Linnean Society of London, Zoology* 17: 137–163, pls. 5–6.
- Elahi R (2008) Effects of aggregation and species identity on the growth and behavior of mushroom corals. *Coral Reefs* 27: 881–885. doi: 10.1007/s00338-008-0403-6
- Ellis J, Solander D (1786) The natural history of many curious and uncommon zoophytes. White and Son, London, 1–208, pls. 1–63.
- Erftemeijer PLA, Riegl B, Hoeksema BW, Todd PA (2012) Environmental impacts of dredging and other sediment disturbances on corals: a review. *Marine Pollution Bulletin* 64: 1737–1765. doi: 10.1016/j.marpolbul.2012.05.008
- Gardiner JS (1899) On the postembryonic development of *Cycloseris*. In: A. Willey's, *Zoological Results, Zoological Results based on Material from New Britain, New Guinea, Loyalty Islands and Elsewhere*, Cambridge 2: 171–180, pls. 19–20.
- Gardiner JS (1905) Madreporaria III. Fungida IV. Turbinolidae. In: *Fauna and geography of the Maldives and Laccadives Archipelagoes*, Cambridge 2: 933–957, pls. 89–93.

- Gilmour JP (2002) Acute sedimentation causes size-specific mortality and asexual budding in the mushroom coral, *Fungia fungites*. *Marine and Freshwater Research* 53: 805–812. doi: 10.1071/MF01219
- Gittenberger A, Gittenberger E (2005) A hitherto unnoticed adaptive radiation: epitoniid species (Gastropoda: Epitoniidae) associated with corals (Scleractinia). *Contributions to Zoology* 74: 125–204.
- Gittenberger A, Gittenberger E (2011) Cryptic, adaptive radiation of parasitic snails: sibling species of *Leptoconchus* (Gastropoda: Coralliophilidae) in corals. *Organisms, Diversity and Evolution* 11: 21–41. doi: 10.1007/s13127-011-0039-1
- Gittenberger A, Hoeksema BW (2006) Phenotypic plasticity revealed by molecular studies on reef corals of *Fungia (Cycloseris)* spp. (Scleractinia: Fungiidae) near river outlets. *Contributions to Zoology* 75: 195–201.
- Gittenberger A, Hoeksema BW (2013) Habitat preferences of coral-associated wentletrap snails (Gastropoda: Epitoniidae). *Contributions to Zoology* 82: 1–25.
- Gittenberger A, Reijnen BT, Hoeksema BW (2011) A molecularly based phylogeny reconstruction of mushroom corals (Scleractinia: Fungiidae) with taxonomic consequences and evolutionary implications for life history traits. *Contributions to Zoology* 80: 107–132.
- Hoeksema BW (1988) Mobility of free-living fungiid corals (Scleractinia), a dispersion mechanism and survival strategy in dynamic reef habitats. *Proceedings of the 6th International Coral Reef Symposium, Townsville, Australia* 2: 715–720.
- Hoeksema BW (1989) Taxonomy, phylogeny and biogeography of mushroom corals (Scleractinia: Fungiidae). *Zoologische Verhandelingen, Leiden* 254: 1–295.
- Hoeksema BW (1991a) Control of bleaching in mushroom coral populations (Scleractinia: Fungiidae) in the Java Sea: stress tolerance and interference by life history strategy. *Marine Ecology Progress Series* 74: 225–237. doi: 10.3354/meps074225
- Hoeksema BW (1991b) Evolution of body size in mushroom corals (Scleractinia: Fungiidae) and its ecomorphological consequences. *Netherlands Journal of Zoology* 41: 122–139.
- Hoeksema BW (1993a) Mushroom corals (Scleractinia: Fungiidae) of Madang Lagoon, northern Papua New Guinea: an annotated checklist with the description of *Cantharellus jebbi* spec. nov. *Zoologische Mededelingen, Leiden* 67: 1–19.
- Hoeksema BW (1993b) Some misapplied nomina nova in reef coral taxonomy (Scleractinia). *Zoologische Mededelingen, Leiden* 67: 41–47.
- Hoeksema BW (1993c) Historical biogeography of *Fungia (Pleuractis)* spp. (Scleractinia: Fungiidae), including a new species from the Seychelles). *Zoologische Mededelingen, Leiden* 67: 639–654.
- Hoeksema BW (1993d) Phenotypic corallum variability in Recent mobile reef corals. *Courier Forschungs-Institut Senckenberg* 164: 263–272.
- Hoeksema BW (2004) Impact of budding on free-living corals at East Kalimantan, Indonesia. *Coral Reefs* 23: 492. doi: 10.1007/s00338-004-0402-1
- Hoeksema BW (2007) Delineation of the Indo-Malayan Centre of maximum marine biodiversity: the Coral Triangle. In: Renema W (Ed) *Biogeography, Time and Place: Distributions, Barriers and Islands*, Springer, Dordrecht, 117–178. doi: 10.1007/978-1-4020-6374-9\_5

- Hoeksema BW (2008) Stony corals (Fungiidae). In: Hoeksema BW, Van der Meij SET (Eds) Cryptic marine biota of the Raja Ampat Islands group. Naturalis, Leiden, 8–12.
- Hoeksema BW (2009) Attached mushroom corals (Scleractinia: Fungiidae) in sediment-stressed reef conditions at Singapore, including a new species and a new record. Raffles Bulletin of Zoology, Supplement 22: 81–90.
- Hoeksema BW (2010) Stony corals (Fungiidae). In: Hoeksema BW, Van der Meij SET (Eds) Crossing marine lines at Ternate: Capacity building of junior scientists in Indonesia for marine biodiversity assessments. Naturalis, Leiden, 19–24.
- Hoeksema BW (2012a) Distribution patterns of mushroom corals (Scleractinia: Fungiidae) across the Spermonde Shelf, South Sulawesi. Raffles Bulletin of Zoology 60: 183–212.
- Hoeksema BW (2012b) Evolutionary trends in onshore-offshore distribution patterns of mushroom coral species (Scleractinia: Fungiidae). Contributions to Zoology 81: 199–221.
- Hoeksema BW (2012c) Mushroom corals (Scleractinia: Fungiidae) of Espiritu Santo (Vanuatu, West Pacific) with the description of a new species. Zoosystema 34: 429–443. doi: 10.5252/z2012n2a14
- Hoeksema BW, Benzoni F (2013) Multispecies aggregations of mushroom corals in the Gambier Islands, French Polynesia. Coral Reefs 32: 1041. doi: 10.1007/s00338-013-1054-9
- Hoeksema BW, Dai CF (1991) Scleractinia of Taiwan. II Family Fungiidae (with the description of a new species). Bulletin Zoological Institute, Academia Sinica, Taipei 30: 201–226.
- Hoeksema BW, De Voogd NJ (2012) On the run: free-living mushroom corals avoiding interaction with sponges. Coral Reefs 31: 455–459. doi: 10.1007/s00338-011-0856-x
- Hoeksema BW, Farenzena ZT (2012) Tissue loss in corals infested by acoelomorph flatworms (*Waminoa* sp.). Coral Reefs 31: 869. doi: 10.1007/s00338-012-0919-7
- Hoeksema BW, Franssen CHJM (2011) Space partitioning by symbiotic shrimp species cohabitating in the mushroom coral *Heliopora actiniformis* at Semporna, eastern Sabah. Coral Reefs 30: 519. doi: 10.1007/s00338-011-0736-4
- Hoeksema BW, Gittenberger A (2010) High densities of mushroom coral fragments at West Halmahera, Indonesia. Coral Reefs 29: 691. doi: 10.1007/s00338-010-0616-3
- Hoeksema BW, Kleemann K (2002) New records of *Fungiacava eilatensis* Goreau et al., 1968 (Bivalvia: Mytilidae) boring in Indonesian mushroom corals (Scleractinia: Fungiidae). Bacteria 66: 25–30.
- Hoeksema BW, Koh EGL (2009) Depauperation of the mushroom coral fauna (Fungiidae) of Singapore (1860s–2006) in changing reef conditions. Raffles Bulletin of Zoology, Supplement 22: 91–101.
- Hoeksema BW, Matthews JL (2011) Contrasting bleaching patterns in mushroom coral assemblages at Koh Tao, Gulf of Thailand. Coral Reefs 30: 95. doi: 10.1007/s00338-010-0675-5
- Hoeksema BW, Moka W (1989) Species assemblages and ecomorph variation of mushroom corals (Scleractinia: Fungiidae) related to reef habitats in the Flores Sea. Netherlands Journal of Sea Research 23: 149–160. doi: 10.1016/0077-7579(89)90009-4
- Hoeksema BW, Van Ofwegen LP (2004) Indo-Malayan reef corals: a generic overview. World Biodiversity Database, CD-ROM Series, ETI, Amsterdam.
- Hoeksema BW, Waheed Z (2011) Initial phase of autotomy in fragmenting *Cycloseris* corals at Semporna, eastern Sabah, Malaysia. Coral Reefs 30: 1087. doi: 10.1007/s00338-011-0807-6

- Hoeksema BW, Waheed Z (2012) Onset of autotomy in an attached *Cycloseris* coral. *Galaxea Journal of Coral Reef Studies* 14: 25–26. doi: 10.3755/galaxea.14.25
- Hoeksema BW, Wirtz P (2013) Over 130 years of survival by a small, isolated population of *Favia gravida* corals at Ascension Island (South Atlantic). *Coral Reefs* 32: 551. doi: 10.1007/s00338-012-1002-0
- Hoeksema BW, Yeemin T (2011) Late detachment conceals serial budding by the free-living coral *Fungia fungites* in the Inner Gulf of Thailand. *Coral Reefs* 30: 975. doi: 10.1007/s00338-011-0784-9
- Hoeksema BW, Suharsono Cleary DFR (2004) Stony corals. In: Hoeksema BW (Ed) Marine biodiversity of the coastal area of the Berau region, East Kalimantan, Indonesia. Progress report East Kalimantan Program - Pilot phase (October 2003). Naturalis, Leiden, 7–16.
- Hoeksema BW, Van der Land J, Van der Meij SET, Van Ofwegen LP, Reijnen BT, Van Soest RWM, De Voogd NJ (2011) Unforeseen importance of historical collections as baselines to determine biotic change of coral reefs: the Saba Bank case. *Marine Ecology* 32: 135–141. doi: 10.1111/j.1439-0485.2011.00434.x
- Hoeksema BW, Van der Meij SET, Franssen CHJM (2012) The mushroom coral as a habitat. *Journal of the Marine Biological Association of the United Kingdom* 92: 647–663. doi: 10.1017/S0025315411001445
- Hoeksema BW, Scott C, True JD (2013a) Dietary shift in corallivorous *Drupella* snails following a major bleaching event at Koh Tao, Gulf of Thailand. *Coral Reefs* 32: 423–428. doi: 10.1007/s00338-012-1005-x
- Hoeksema BW, Waheed Z, Alamaru A (2013b) Out of sight: aggregations of epizoic comb jellies underneath mushroom corals. *Coral Reefs* 32: 1065. doi: 10.1007/s00338-013-1078-1
- ICZN (2011) Coral taxon names published in ‘Corals of the world’ by J.E.N. Veron (2000): potential availability confirmed under Article 86.1.2. *Bulletin of Zoological Nomenclature* 68: 162–166.
- ICZN (2012) Amendment of Articles 8, 9, 10, 21 and 78 of the International Code of Zoological Nomenclature to expand and refine methods of publication. *ZooKeys* 219: 1–10. doi: 10.3897/zookeys.219.3944
- Kleemann K, Hoeksema BW (2002) *Lithophaga* (Bivalvia: Mytilidae), including a new species, boring in mushroom corals (Scleractinia: Fungiidae) at South Sulawesi, Indonesia. *Basteria* 66: 11–24.
- Lamarck JBPA de M de (1801) *Système des Animaux sans vertèbres, ou Tableau général des classes, des ordres et des genres de ces animaux; présentant leurs caractères essentiels et leur distribution d’après les considérations de leurs rapports naturels et de leur organisation, et suivant l’arrangement établi dans les galeries du Muséum d’Histoire naturelle, parmi leurs dépouilles conservées; précédé du Discours d’ouverture de l’an VIII de la République*. Déterville, Paris, 432 pp.
- Lamarck JBPA de M de (1816) *Histoire naturelle des animaux sans vertèbres*. Vol. 2. Verdier, Paris, 568 pp.
- Latypov YY (1995) Scleractinian corals of Vietnam. Part III. Faviidae, Fungiidae. Nauka, Moscow, 140 pp. [in Russian]
- Latypov YY (2006) The scleractinian corals of Vietnam. Pdf file on CD-ROM, Far East Branch Russian Academy of Science, Institute of Marine Biology, Vladivostok, 405 pp.



- Linnaeus C (1758) *Systema naturae* (ed. 10), 1. Laurentii Salvii, Holmiae, 824 pp.
- Littler MM, Littler DD, Brooks BL, Koven JF (1997) A unique coral reef formation discovered on the Great Astrolabe Reef, Fiji. *Coral Reefs* 16: 51–54. doi: 10.1007/s003380050059
- Michelin H (1842) Description d'une nouvelle espèce de Zoophyte du genre Fongie. *Revue Zoologique par La Société Cuvierienne* 5: 316.
- Milne Edwards H (1860) *Histoire naturelle des coralliaires ou polypes proprement dits* 3. Librairie Encyclopédique de Roret, Paris, 560 pp.
- Milne Edwards H, Haime J (1849) Mémoire sur les polypiers appartenant a la famille des Oculinides, au groupe intermédiaire des Pseudoastréides et a la famille des Fongides. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 29: 67–73.
- Milne Edwards H, Haime J (1850) Classification of polypi. A monograph of the British fossil corals 1. Palaeontographical Society of London, London, viii-lxxxv.
- Milne Edwards H, Haime J (1851) Recherches sur les polypiers. Mémoire 6. Monographie des Fongides. *Annales des Sciences Naturelles, Zoologie, Series 3*, 15: 73–144.
- Mondal T, Raghunathan C (2013) Description of a new coral species (Scleractinia: Fungiidae) from Andaman Islands, India. *Journal of Threatened Taxa* 5: 4653–4659. doi: 10.11609/JoTT.o3194.4653-9
- Nemenzo F (1981) Philippine corals. *Guide to the Philippine Flora and Fauna* 2: 1–320.
- Nemenzo F (1986) Corals. *Guide to the Philippine Flora and Fauna* 5: 1–273.
- Nishihira M, Veron JEN (1995) *Hermatypic corals of Japan*. Kaiyusha Publishers, Tokyo, 439 pp.
- Ortmann A (1889) Beobachtungen an Steinkorallen von der Südküste Ceylons. *Zoologische Jahrbüchern, Abteilung für Systematik, Biologie und Biogeographie der Tiere* 4: 493–590, pls. 11–17.
- Owada M, Hoeksema BW (2011) Molecular phylogeny and shell microstructure of *Fungiacava eilatensis* Goreau et al. 1968, boring into mushroom corals (Scleractinia: Fungiidae), in relation to other mussels (Bivalvia: Mytilidae). *Contributions to Zoology* 80: 169–178.
- Pillai CSG (1986) Recent corals from the south-east coast of India. In: James PSBR (Ed) *Recent Advances in Marine Biology. Today and Tomorrow'S*, Printers and Publishers, New Delhi, 107–198.
- Quelch JJ (1886) Report on the reef-corals collected by H.M.S. Challenger during the years 1873-76. Report on the Scientific Results of the Voyage of H.M.S. Challenger during the years 1873-79. *Zoology* 16(3): 1–203, pls. 1–12.
- Scholtz J, Hoeksema BW, Pawson DL, Ruthensteiner B (2012) Ludwig Döderlein (1855–1936): Some aspects of his life, research, and legacy. *Spixiana* 35: 177–191.
- Scheer G, Pillai CSG (1983) Report on the stony corals from the Red Sea. *Zoologica, Stuttgart* 45(133): 1-198, pls. 1–41.
- Suharsono (2004) *Jenis-jenis karang di Indonesia*. Pusat Penelitian Oseanografi - LIPI, Jakarta, 343 pp.
- Suharsono (2008) *Jenis-jenis karang di Indonesia*. LIPI Press, Jakarta, 372 pp.
- Tenison-Woods JE (1878) On the extra tropical corals of Australia. *Proceedings of the Linnean Society of New South Wales* 2: 292–341, pls. 4–6.
- Van der Horst CJ (1921) The Madreporaria of the Siboga Expedition II. *Madreporaria Fungi-da. Siboga-Expedition XVIIb*. Brill, Leiden, 53-98, pls. 1–6.

- Van der Meij SET, Hoeksema BW (2013) Distribution of gall crabs inhabiting mushroom corals on Semporna reefs, Malaysia. *Marine Biodiversity* 43: 53–59. doi: 10.1007/s12526-012-0135-2
- Van der Meij SET, Visser RR (2011) The *Acropora humilis* group (Scleractinia) of the Snellius expedition (1929–30). *Raffles Bulletin of Zoology* 59: 9–17.
- Van der Meij SET, Suharsono, Hoeksema BW (2010) Long-term changes in coral assemblages under natural and anthropogenic stress in Jakarta Bay (1920–2005). *Marine Pollution Bulletin* 60: 1442–1454. doi: 10.1016/j.marpolbul.2010.05.011
- Van Soest RWM (1979) A catalogue of the Coelenterate type specimens of the Zoological Museum of Amsterdam IV. Gorgonacea, Actinaria, Scleractinia. *Beaufortia* 29: 81–126.
- Vaughan TW, Wells JW (1943) Revision of the suborders, families and genera of the Scleractinia. *Special Papers of the Geological Society of America* 44: 1–363.
- Veron JEN (1986) *Corals of Australia and the Indo-Pacific*. Angus & Robertson, North Ryde (N.S.W.), 644 pp.
- Veron JEN (1990) New Scleractinia from Japan and other Indo-Pacific countries. *Galaxea* 9: 95–173.
- Veron JEN (1992) Hermatypic corals of Japan. *Australian Institute of Marine Science Monograph Series* 9: 1–234.
- Veron JEN (1993) A biogeographic database of hermatypic corals. *Australian Institute of Marine Science Monograph Series* 10: 1–433.
- Veron JEN (2000) *Corals of the World*. Volume 2. Australian Institute of Marine Science, Townsville, 429 pp.
- Veron JEN (2002) New species described in *Corals of the World*. *Australian Institute of Marine Science Monograph Series* 11: 1–209.
- Veron JEN, Pichon M (1980) Scleractinia of Eastern Australia – Part III. Family Agariciidae, Siderastreidae, Fungiidae, Oculinidae, Merulinidae, Mussidae, Pectinidae, Caryophyllidae, Dendrophylliidae. *Australian Institute of Marine Science Monograph Series* 4: 1–459.
- Verrill AE (1868) Notes on the Radiata in the Museum of Yale College, with descriptions of new genera and species. No. 6. Review of the polyps and corals of the West Coast of America. *Transactions of the Connecticut Academy of Arts and Sciences* 1: 377–558, pls. 5–10.
- Waheed Z, Hoeksema BW (2013) A tale of two winds: species richness patterns of reef corals around the Semporna peninsula, Malaysia. *Marine Biodiversity* 43: 37–51. doi: 10.1007/s12526-012-0130-7
- Waheed Z, Hoeksema BW (in press) Diversity patterns of scleractinian corals at Kota Kinabalu, Malaysia, in relation to depth and exposure. *Raffles Bulletin of Zoology*.
- Wallace CC, Fellegara I, Muir PR, Harrison PL (2009) The Scleractinian Corals of Moreton Bay, Eastern Australia: high latitude, marginal assemblages with increasing species richness. *Memoirs of the Queensland Museum – Nature* 54: 1–118.
- Wells JW (1956) Scleractinia. In: Moore RC (Ed) *Treatise on Invertebrate Paleontology*. F. Coelenterata. Geological Society of America & University of Kansas Press, 328–440.
- Wells JW (1966) Evolutionary development in the scleractinian family Fungiidae. In: Rees WJ (Ed) *The Cnidaria and their Evolution*. Symposia of the Zoological Society London 16. Academic Press, London, 223–246, pl. 1.

- Yamashiro H, Nishihira M (1994) Radial skeletal dissolution to promote vegetative reproduction in a solitary coral *Diaseris distorta*. *Experientia* 50: 497–498. doi: 10.1007/BF01920756
- Yamashiro H, Nishihira M (1995) Phototaxis in Fungiidae corals (Scleractinia). *Marine Biology* 124: 461–465. doi: 10.1007/BF00363920
- Yamashiro H, Nishihira M (1998) Experimental study of growth and asexual reproduction in *Diaseris distorta* (Michelin, 1843), a free-living fungiid coral. *Journal of Experimental Marine Biology and Ecology* 225: 253–267. doi: 10.1016/S0022-0981(97)00229-3
- Yamashiro H, Hidaka M, Nishihira M, Pong-In S (1989) Morphological studies on skeletons of *Diaseris fragilis*, a free-living coral which reproduces asexually by natural autotomy. *Galaxea* 8: 283–294.