

# Potential Efficacy of Citicoline as Adjunct Therapy in Treatment of Cerebral Malaria

Fatima El-Assaad,<sup>a</sup> Valery Combes,<sup>a</sup> Georges Emile Raymond Grau,<sup>a</sup> Ronan Jambou<sup>a,b</sup>

Vascular Immunology Unit, Medical Foundation Building, Department of Pathology, Sydney Medical School, The University of Sydney, Sydney, Australia<sup>a</sup>; Immunology Unit, Institut Pasteur de Madagascar, Antananarivo, Madagascar<sup>b</sup>

**Cerebral malaria (CM) is characterized by a dysregulated immune response that results in endothelial membrane destabilization and increased microparticle (MP) production. Citicoline (CTC) is a membrane stabilizer used for the treatment of neurological disorders. We evaluated the efficacy of CTC as adjunct therapy to aid recovery from experimental CM. We show that CTC reduces MP production *in vitro*; in combination with artesunate *in vivo*, confers partial protection against CM; and prolongs survival.**

The incidence of cerebral malaria (CM) is highly dependent on the immune status of the population and the frequency of *Plasmodium falciparum* transmission. This debilitating syndrome is characterized by coma, seizures, and respiratory distress. In children, the coma usually develops rapidly after a seizure, whereas in adults its development is slower and takes 2 to 3 days (1).

During infection, the sequestration of mature-stage-parasitized erythrocytes, platelets, and leukocytes within cerebral microvessels, coupled with cytokine overproduction, leads to hypoxia (2), endothelial activation, and blood-brain barrier disruption (3). During the acute phase of pediatric CM, several pathophysiological mechanisms are involved, including CD8<sup>+</sup> T cell activation, (4) and the complement cascade (5), although disruption of the endothelium seems to be a supported mechanism of cerebral dysfunction.

Evidence of a destabilized endothelium is the increased release of endothelial-cell-derived microparticles (MP). In CM patients, high titers of MP are detected in the blood coinciding with a rise in the tumor necrosis factor (TNF) level, which returns to the baseline postinfection (p.i.) (6). These elevated levels do not occur in uncomplicated malaria or severe anemia (7). TNF plays a major role in this mechanism, as it strongly induces endothelial vesiculation (8).

Current therapies for CM are designed to target the pathogen rather than the underlying pathogenic mechanisms responsible for the manifestation of the pathology (9–11). In the last 5 years, several novel compounds have been tested in animal models or in humans as adjunct therapy to prevent tissue and brain alterations during infection, including erythropoietin (EPO) (2, 12), defibrotide (13), atorvastatin (14, 15), the exogenous nitric oxide (NO) donor dipropylene-triamine-NONOate (16), and others (17, 18). EPO, atorvastatin, and NO have undergone U.S. FDA review and been approved for other indications. However, most of these compounds are new and have a long way to go from the benchtop before being implemented as a line of treatment at the bedside. Targeting membrane alteration in the same approach as for ischemia-reperfusion disorders paves the way for the use of a treatment already used for patients. Citicoline (CTC) is used in the United States and Europe by adults and children to improve brain recovery after ischemic stroke and neurological and vascular disorders (19–22). It works efficiently on damage after focal ischemia (neuronal damage, blood-brain barrier dysfunction, behavioral dysfunction) and brain edema (for a review, see reference 23) in

mice, but its benefit for humans is still controversial (24). CTC is a natural compound found in eukaryotic cells that regulates membrane fluidity and is synthesized mainly by the CDP-choline pathway (23–26).

We evaluated the efficacy of CTC, a membrane-stabilizing agent, as adjunct therapy to enhance recovery from experimental CM. In this study, we used both *in vitro* and *in vivo* models of CM. Mouse models using *Plasmodium berghei* ANKA infection can mimic CM pathology. *P. berghei* ANKA-infected mice develop a lethal syndrome 7 days p.i. with a significant increase in MP in their plasma at the time of onset of the neurological syndrome (27, 28).

**CTC confers protection against neurological syndrome in murine CM.** All of the mice used in this study were handled under University of Sydney Animal Ethics Committee approval (approval number K20/7-2006/3/4434). Mice were infected with *P. berghei* ANKA by following protocols in previously published studies (29). Eight-week-old female CBA mice received intraperitoneal injections of  $1 \times 10^6$  infected red blood cells. The cerebral syndrome usually occurred during the neurological phase, day 7 to 14 p.i. Mice that surpassed the neurological syndrome died during the hyperparasitemia phase (day 14 onward) because of severe anemia. Parasitemia was determined by light microscopy with thin Diff-Quick-stained smears of blood collected from the tail vein. Clinical presentation was monitored daily.

Starting on day 4 p.i., mice received intraperitoneal injections of CTC ( $n = 28$ ), artesunate (ART) (Sigma) ( $n = 21$ ), or a combination of CTC and ART ( $n = 14$ ). CTC was solubilized in phosphate-buffered saline (PBS) and administered daily for 3 days. Two doses (500 and 1,000 mg/kg) were used in separate experiments. The regimens were chosen in accordance with literature on mouse stroke models. A subeffective dose of ART (40 mg/kg) was

Received 22 December 2012 Returned for modification 21 March 2013


Accepted 18 October 2013

Published ahead of print 28 October 2013

Address correspondence to Ronan Jambou, rjambou@pasteur.mg.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.02591-12




**FIG 1** CTC as adjunct therapy for mice with CM. Shown are the parasitemia levels (A) and percentages of survival (B) of infected mice following treatment with CTC alone ( $n = 28$ ; open squares), ART ( $n = 21$ ; closed triangles), a combination of CTC and ART ( $n = 14$ ; open triangles), or PBS ( $n = 28$ ; closed squares). Parasitemia was calculated from Diff-Quick-stained blood smears, and data are presented as means  $\pm$  standard deviations. pRBC, parasitized red blood cells.

administered once at day 4 p.i. Control mice received PBS treatment ( $n = 28$ ).

Brain, lung, liver, and spleen tissue samples were collected from mice that exhibited clinical signs of CM or were moribund. The tissue was prepared for thin paraffin sectioning. The hematoxylin-eosin-stained sections analyzed had no histopathological features associated with CM to widespread hemorrhage, a high degree of edema and leukocyte and parasitized red blood cell adherence to the endothelium throughout the brain, plugging of microvessels, and some necrosis of microvessels (30). In this study, histopathology was mainly used to confirm CM.

At day 7, the parasitemia of the two groups treated with ART differed significantly from that of the control group (5.96, 2.6, and 2.35% in controls, ART, and ART plus CTC, respectively, [ $P = 0.0051$  and  $P < 0.0001$ ]) (Fig. 1). However, no significant difference in parasitemia between controls and the CTC-alone group was found ( $P = 0.4081$ ). After day 7, no significant difference in parasitemia between the ART and ART-plus-CTC groups was found either, suggesting that CTC by itself has no impact on the parasite.

Control mice developed clinical CM between days 6 and 10 p.i. When CTC was administered at 1,000 mg/kg, the onset of clinical CM was delayed by at least 24 h. Thus, when mice were checked at day 7, treatment with CTC alone had enhanced survival from 20.6 to 82.1% but all of the mice died before day 14. After day 14,



**FIG 2** Effect of CTC on MP production by D3 HBEC. HBEC were stimulated with 100 ng/ml of TNF. The MP in the supernatant were counted in a flow cytometer after annexin V-FITC labeling. (A) Incubation with CTC 2 h prior to TNF stimulation. (B) Incubation with CTC concomitantly with TNF stimulation. (C) Incubation with CTC 6 h after TNF stimulation.

survival increased from 0% (controls and CTC) or 23.8% (ART alone) to 64.2% in the combined ART-plus-CTC group. Combination of CTC and ART prolonged survival by more than 6 days. When used at 500 mg/kg, the trend of the CTC effect was similar to that observed at the 1,000-mg/ml treatment but statistical tests fell under the threshold of significance. Statistics were performed with GraphPad Prism software. Survival curves were compared by using the Kaplan-Meier (log rank) test;  $P < 0.05$  was considered significant. Parasitemia in the control and other groups was compared by using the Mann-Whitney  $t$  test.

**CTC decreases the production of MP by TNF-stimulated HBEC.** MP production by human brain endothelial cells (HBEC) was studied *in vitro* (Fig. 2). Immortalized HBEC (D3 line) (31)

were grown to confluence in EBM-2 medium (Lonza CC-3156) in 24-well plates at 37°C. MP production was induced by incubation with TNF (100 ng/ml) overnight. CTC was added 2 h prior to, concomitantly with, or 6 h after stimulation with TNF (doses, 0.1, 1, 10, 30, and 100 µM). Culture supernatants were collected, and MP were labeled with annexin V-fluorescein isothiocyanate (FITC), a known marker of MP, prior to flow cytometric analysis.

Negative controls for the experiment were obtained from resting cells. Positive controls were obtained from cells stimulated with TNF (100 ng/ml) for 18 h, which induced a 2-fold increase in MP vesiculation compared to that of resting cells. CTC at 0.1 µM showed no apparent inhibition of MP production. However, when added 2 h prior to TNF stimulation, CTC (1 to 10 µM) reduced the formation of MP to the baseline level. When added 6 h after TNF stimulation, CTC (1 to 10 µM) induced a 70% reduction in MP production compared to that of the positive control ( $P < 0.05$  and  $P < 0.01$ , respectively). Again, at the lowest dose, CTC had no significant effect.

In conclusion, data obtained *in vivo* clearly demonstrate that CTC can confer partial protection against experimental CM. This improvement of survival also fits in with a decrease in MP production when HBEC are treated with CTC *in vitro*. These data support a role for CTC in reducing MP production by stabilizing the microvascular endothelium, similar to what is seen in other vascular pathologies (32–35). CTC therapy can be proposed for clinical studies of human patients to improve recovery from CM, as it is already used as a food supplement for malnourished children and has low toxicity.

## ACKNOWLEDGMENTS

This work was supported by funds from the National Health and Medical Research Council of Australia (NHMRC project grant 464893), the Australian Research Council (DP0774425), and the Rebecca L. Cooper Medical Research Foundation, Sydney, Australia.

We are grateful to P. O. Couraud for providing hCMEC/D3 cells and to M. J. Jambou for her technical assistance on this project.

We have no conflict of interest to declare.

## REFERENCES

1. von Seidlein L, Olaosebikan R, Hendriksen IC, Lee SJ, Adedoyin OT, Agbenyega T, Nguah SB, Bojang K, Deen JL, Evans J, Fanello CI, Gomes E, Pedro AJ, Kahabuka C, Karema C, Kivaya E, Maitland K, Mokuolu OA, Mtowe G, Mwanga-Amumpaire J, Nadjm B, Nansumba M, Ngum WP, Onyamboko MA, Reyburn H, Sakulthaw T, Silamut K, Tshetu AK, Umulisa N, Gesase S, Day NP, White NJ, Dondorp AM. 2012. Predicting the clinical outcome of severe falciparum malaria in African children: findings from a large randomized trial. *Clin. Infect. Dis.* 54:1080–1090. <http://dx.doi.org/10.1093/cid/cis034>.
2. Hempel C, Combes V, Hunt NH, Kurtzhals JA, Grau GE. 2011. CNS hypoxia is more pronounced in murine cerebral than noncerebral malaria and is reversed by erythropoietin. *Am. J. Pathol.* 179:1939–1950. <http://dx.doi.org/10.1016/j.ajpath.2011.06.027>.
3. Kim H, Higgins S, Liles WC, Kain KC, Sandra A. 2011. Endothelial activation and dysregulation in malaria: a potential target for novel therapeutics. *Curr. Opin. Hematol.* 18:177–185. <http://dx.doi.org/10.1097/MOH.0b013e328345a4cf>.
4. Villegas-Mendez A, Greig R, Shaw TN, de Souza JB, Gwyer Findlay E, Stumhofer JS, Hafalla JC, Blount DG, Hunter CA, Riley EM, Couper KN. 2012. IFN- $\gamma$ -producing CD4<sup>+</sup> T cells promote experimental cerebral malaria by modulating CD8<sup>+</sup> T cell accumulation within the brain. *J. Immunol.* 189:968–979. <http://dx.doi.org/10.4049/jimmunol.1200688>.
5. Ramos TN, Darley MM, Hu X, Billker O, Rayner JC, Ahras M, Wohler JE, Barnum SR. 2011. Cutting edge: the membrane attack complex of complement is required for the development of murine experimental cerebral malaria. *J. Immunol.* 186:6657–6660. <http://dx.doi.org/10.4049/jimmunol.1100603>.
6. Pankoui Mfonkeu JB, Gouado I, Fotso Kuate H, Zambou O, Amvamo Zollo PH, Grau GE, Combes V. 2010. Elevated cell-specific microparticles are a biological marker for cerebral dysfunctions in human severe malaria. *PLoS One* 5:e13415. <http://dx.doi.org/10.1371/journal.pone.0013415>.
7. Combes V, Taylor TE, Juhan-Vague I, Mege JL, Mwenechanya J, Tembo M, Grau GE, Molyneux ME. 2004. Circulating endothelial microparticles in Malawian children with severe falciparum malaria complicated with coma. *JAMA* 291:2542–2544. <http://dx.doi.org/10.1001/jama.291.21.2542-b>.
8. Combes V, El-Assaad F, Faillie D, Jambou R, Hunt NH, Grau GE. 2010. Microvesiculation and cell interactions at the brain-endothelial interface in cerebral malaria pathogenesis. *Prog. Neurobiol.* 91:140–151. <http://dx.doi.org/10.1016/j.pneurobio.2010.01.007>.
9. Krishna S. 2012. Adjunctive management of malaria. *Curr. Opin. Infect. Dis.* 25:484–488. <http://dx.doi.org/10.1097/QCO.0b013e3283567b20>.
10. Shikani HJ, Freeman B, Lisanti MP, Weiss LM, Tanowitz HB, Desruisseaux MS. 2012. Cerebral malaria: we have come a long way. *Am. J. Pathol.* 25:1484–1492. <http://dx.doi.org/10.1016/j.ajpath.2012.08.010>.
11. Higgins SJ, Kain KC, Liles WC. 2011. Immunopathogenesis of falciparum malaria: implications for adjunctive therapy in the management of severe and cerebral malaria. *Expert Rev. Anti Infect. Ther.* 9:803–819. <http://dx.doi.org/10.1586/eri.11.96>.
12. Hempel C, Hyttel P, Staals A, T, Nyengaard JR, Kurtzhals JA. 2012. Erythropoietin treatment alleviates ultrastructural myelin changes induced by murine cerebral malaria. *Malar. J.* 11:216. <http://dx.doi.org/10.1186/1475-2875-11-216>.
13. Francischetti IM, Oliveira CJ, Ostera GR, Yager SB, Debierre-Grockiego F, Carregaro V, Jaramillo-Gutierrez G, Hume JC, Jiang L, Moretz SE, Lin CK, Ribeiro JM, Long CA, Vickers BK, Schwarz RT, Seydel KB, Iacobelli M, Ackerman HC, Srinivasan P, Gomes RB, Wang X, Monteiro RQ, Kotsyfakis M, Sá-Nunes A, Waisberg M. 2012. Defibrinolytic interferes with several steps of the coagulation-inflammation cycle and exhibits therapeutic potential to treat severe malaria. *Arterioscler. Thromb. Vasc. Biol.* 32:786–798. <http://dx.doi.org/10.1161/ATVBAHA.111.240291>.
14. Parquet V, Briolant S, Torrentino-Madamet M, Henry M, Almeras L, Amalvict R, Baret E, Fusai T, Rogier C, Pradines B. 2009. Atorvastatin is a promising partner for antimalarial drugs in treatment of *Plasmodium falciparum* malaria. *Antimicrob. Agents Chemother.* 53:2248–2252. <http://dx.doi.org/10.1128/AAC.01462-08>.
15. Dormoi J, Briolant S, Desgrouas C, Pradines B. 2013. Impact of methylene blue and atorvastatin combination therapy on the apparition of cerebral malaria in a murine model. *Malar. J.* 12:127. <http://dx.doi.org/10.1186/1475-2875-12-127>.
16. Zanini GM, Martins YC, Cabrales P, Frangos JA, Carvalho LJ. 2012. S-Nitrosoglutathione prevents experimental cerebral malaria. *J. Neuroimmunol. Pharmacol.* 7:477–487. <http://dx.doi.org/10.1007/s11481-012-9343-6>.
17. Dai M, Freeman B, Bruno FP, Shikani HJ, Tanowitz HB, Weiss LM, Reznik SE, Stephani RA, Desruisseaux MS. 2012. The novel ETA receptor antagonist HJP-272 prevents cerebral microvascular hemorrhage in cerebral malaria and synergistically improves survival in combination with an artemisinin derivative. *Life Sci.* 91:687–692. <http://dx.doi.org/10.1016/j.lfs.2012.07.006>.
18. Achtman AH, Pilat S, Law CW, Lynn DJ, Janot L, Mayer ML, Ma S, Kindrachuk J, Finlay BB, Brinkman FS, Smyth GK, Hancock RE, Schofield L. 2012. Effective adjunctive therapy by an innate defense regulatory peptide in a preclinical model of severe malaria. *Sci. Transl. Med.* 4:135ra64. <http://dx.doi.org/10.1126/scitranslmed.3003515>.
19. Hankey GJ. 2012. How effective is citicoline for acute ischaemic stroke? *Lancet* 380:318–320. [http://dx.doi.org/10.1016/S0140-6736\(12\)60912-X](http://dx.doi.org/10.1016/S0140-6736(12)60912-X).
20. Dávalos A, Alvarez-Sabín J, Castillo J, Díez-Tejedor E, Ferro J, Martínez-Vila E, Serena J, Segura T, Cruz VT, Masjuan J, Cobo E, Secades JJ. 2012. International Citicoline Trial on acute Stroke (ICTUS) trial investigators. Citicoline in the treatment of acute ischaemic stroke: an international, randomised, multicentre, placebo-controlled study (ICTUS trial). *Lancet* 380:349–357. [http://dx.doi.org/10.1016/S0140-6736\(12\)60813-7](http://dx.doi.org/10.1016/S0140-6736(12)60813-7).
21. Diederich K, Frauenknecht K, Minnerup J, Schneider BK, Schmidt A, Altach E, Eggert V, Sommer CJ, Schäbitz WR. 2012. Citicoline enhances neuroregenerative processes after experimental stroke in rats. *Stroke* 43:1931–1940. <http://dx.doi.org/10.1161/STROKEAHA.112.654806>.

22. Secades JJ. 2012. Probably role of citicoline in stroke rehabilitation: review of the literature. *Rev. Neurol.* 54:173–179. (In Spanish.) <http://www.revneurol.com/sec/resumen.php?or=pubmed&id=2011490>.
23. Jambou R, El-Assaad F, Combes V, Grau GE. 2009. Citicoline (CDP-choline): what role in the treatment of complications of infectious diseases. *Int. J. Biochem. Cell Biol.* 41:1467–1470. <http://dx.doi.org/10.1016/j.biocel.2009.02.011>.
24. Clark WM, Clark TD. 2012. Stroke: treatment for acute stroke—the end of the citicoline saga. *Nat. Rev. Neurol.* 8:484–485. <http://dx.doi.org/10.1038/nrneurol.2012.166>.
25. Lizasoain I, Cárdenas A, Hurtado O, Romera C, Mallolas J, Lorenzo P, Castillo J, Moro MA. 2006. Targets of cytoprotection in acute ischemic stroke: present and future. *Cerebrovasc. Dis.* 21(Suppl 2):1–8. <http://dx.doi.org/10.1159/000091698>.
26. Weiss GB. 1995. Metabolism and actions of CDP-choline as an endogenous compound and administered exogenously as citicoline. *Life Sci.* 56:637–660. [http://dx.doi.org/10.1016/0024-3205\(94\)00427-T](http://dx.doi.org/10.1016/0024-3205(94)00427-T).
27. Piguet PF, Kan CD, Vesin C. 2002. Thrombocytopenia in an animal model of malaria is associated with an increased caspase-mediated death of thrombocytes. *Apoptosis* 7:91–98. <http://dx.doi.org/10.1023/A:1014341611412>.
28. van der Heyde HC, Nolan J, Combes V, Gramaglia I, Grau GE. 2006. A unified hypothesis for the genesis of cerebral malaria: sequestration, inflammation and hemostasis leading to microcirculatory dysfunction. *Trends Parasitol.* 22:503–508. <http://dx.doi.org/10.1016/j.pt.2006.09.002>.
29. Grau GE, Fajardo LF, Piguet PF, Allet B, Lambert PH, Vassalli P. 1987. Tumor necrosis factor (cachectin) as an essential mediator in murine cerebral malaria. *Science* 237:1210–1212. <http://dx.doi.org/10.1126/science.3306918>.
30. Combes V, Rosenkranz AR, Redard M, Pizzolato G, Lepidi H, Vestweber D, Mayadas TN, Grau GE. 2004. Pathogenic role of P-selectin in experimental cerebral malaria: importance of the endothelial compartment. *Am. J. Pathol.* 164:781–786. [http://dx.doi.org/10.1016/S0002-9440\(10\)63166-5](http://dx.doi.org/10.1016/S0002-9440(10)63166-5).
31. Weksler BB, Subileau EA, Perrière N, Charneau P, Holloway K, Leveque M, Tricoire-Leignel H, Nicotra A, Bourdoulous S, Turowski P, Male DK, Roux F, Greenwood J, Romero IA, Couraud PO. 2005. Blood-brain barrier-specific properties of a human adult brain endothelial cell line. *FASEB J.* 9:1872–1874. <http://dx.doi.org/10.1096/fj.04-3458fje>.
32. Gutiérrez-Fernández M, Rodríguez-Frutos B, Fuentes B, Vallejo-Cremades MT, Alvarez-Grech J, Expósito-Alcaide M, Díez-Tejedor E. 2012. CDP-choline treatment induces brain plasticity markers expression in experimental animal stroke. *Neurochem. Int.* 60:310–317. <http://dx.doi.org/10.1016/j.neuint.2011.12.015>.
33. Alvarez-Sabín J, Román GC. 2011. Citicoline in vascular cognitive impairment and vascular dementia after stroke. *Stroke* 42:S40–3. <http://dx.doi.org/10.1161/STROKEAHA.110.606509>.
34. Baskys A, Hou AC. 2007. Vascular dementia: pharmacological treatment approaches and perspectives. *Clin. Interv. Aging* 2:327–335. <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2685259/>.
35. García-Cobos R, Frank-García A, Gutiérrez-Fernández M, Díez-Tejedor E. 2010. Citicoline, use in cognitive decline: vascular and degenerative. *J. Neurol. Sci.* 299:188–192. <http://dx.doi.org/10.1016/j.jns.2010.08.027>.