

Host Specificity and Source of *Enterocytozoon bienersi* Genotypes in a Drinking Source Watershed

Yaqiong Guo,^{a,b} Kerri A. Alderisio,^c Wenli Yang,^b Vitaliano Cama,^d Yaoyu Feng,^a Lihua Xiao^b

State Key Laboratory of Bioreactor Engineering, School of Resources and Environmental Engineering, East China University of Science and Technology, Shanghai, China^a; Division of Foodborne, Waterborne, and Environmental Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, USA^b; New York City Department of Environmental Protection, Valhalla, New York, USA^c; Division of Parasitic Diseases and Malaria, Centers for Disease Control and Prevention, Atlanta, Georgia, USA^d

To assess the host specificity of *Enterocytozoon bienersi* and to track the sources of *E. bienersi* contamination, we genotyped *E. bienersi* in wildlife and stormwater from the watershed of New York City's source water, using ribosomal internal transcribed spacer (ITS)-based PCR and sequence analyses. A total of 255 specimens from 23 species of wild mammals and 67 samples from stormwater were analyzed. Seventy-four (29.0%) of the wildlife specimens and 39 (58.2%) of the stormwater samples from streams were PCR positive. Altogether, 20 *E. bienersi* genotypes were found, including 8 known genotypes and 12 new ones. Sixteen and five of the genotypes were seen in animals and stormwater from the watershed, respectively, with WL4 being the most common genotype in both animals (35 samples) and stormwater (23 samples). The 20 *E. bienersi* genotypes belonged to five genogroups (groups 1, 3, 4, and 7 and an outlier), with only 23/113 (20.4%) *E. bienersi*-positive samples belonging to zoonotic genogroup 1 and 3/20 genotypes ever being detected in humans. The two genogroups previously considered host specific, groups 3 and 4, were both detected in multiple groups of mammals. Thus, with the exception of the type IV, Peru11, and D genotypes, which were detected in only 7, 5, and 2 animals, respectively, most *E. bienersi* strains in most wildlife samples and all stormwater samples in the watershed had no known public health significance, as these types have not previously been detected in humans. The role of different species of wild mammals in the contribution of *E. bienersi* contamination in stormwater was supported by determinations of host-adapted *Cryptosporidium* species/genotypes in the same water samples. Data from this study indicate that the host specificity of *E. bienersi* group 3 is broader than originally thought, and wildlife is the main source of *E. bienersi* in stormwater in the watershed.

Microsporidiosis is a significant cause of diarrhea, especially in children and immunocompromised persons. Although over a dozen microsporidian parasites have been found in humans, *Enterocytozoon bienersi* is the species responsible for >90% of human microsporidian infections (1–3). It is also commonly found in animals, especially mammals and some birds (4). Although animals have been generally considered a major source of human *E. bienersi* infection, this is based largely on results of genetic characterizations of parasites from humans and some animals (3, 4).

Water potentially plays an important role in the transmission of microsporidiosis. Epidemiological and environmental studies have frequently identified drinking of untreated water as a potential risk factor for microsporidiosis in humans (5, 6). A putative waterborne outbreak of intestinal microsporidiosis caused by *E. bienersi* and *Encephalitozoon intestinalis* was reported in France (7), although the validity of the claim was disputed (8). Spores of *E. bienersi* have been identified in surface water at high frequencies (9–14).

Sequence analysis of the ribosomal internal transcribed spacer (ITS) has been used extensively in molecular epidemiologic studies of *E. bienersi* infection in humans and assessment of the human-infective potential of *E. bienersi* in animals (3, 4). Over 100 *E. bienersi* genotypes have been identified, belonging to at least six phylogenetically distinct genogroups (groups 1 to 5 and the so-called outlier in dogs) (15–17). Host adaptation is apparent among *E. bienersi* genotypes (18–20), with almost all human isolates and some animal isolates belonging to genotypes in group 1 (3, 4). Other genogroups contain mostly animal-adapted genotypes, including group 2 genotypes in cattle, group 3 genotypes in

muskrats, group 4 genotypes in raccoons, group 5 genotypes in primates, and the outlier in dogs (15–17). Sequence characterizations of other genes support the ITS classification of genogroups, as PCR primers based on nucleotide sequences of group 1 largely cannot amplify DNA from other *E. bienersi* genogroups (21). Some additional genogroups, such as groups 6 and 7, have also been identified recently (22, 23), although the host specificity of them is less clear.

Determination of host adaptation of *E. bienersi* genotypes, however, was based largely on ITS sequence characterizations of specimens from only a few domestic and wild animal species (18, 19, 24, 25). Although the bovine-adapted nature of group 2 genotypes was confirmed in subsequent studies (26–29), the host specificity of other genogroups remains to be determined. Indeed, group 4 genotypes, originally shown to be common in raccoons in the United States (18), have recently been found in wastewater in Tunisia, where raccoons do not exist, indicating that some other animals are probably also natural hosts of this group of genotypes (30). More studies on the distribution of *E. bienersi* genotypes in wildlife are needed before we have a better understanding of host adaptation in *E. bienersi*.

Despite the zoonotic potential of *E. bienersi* in wildlife, few

Received 4 September 2013 Accepted 14 October 2013

Published ahead of print 18 October 2013

Address correspondence to Yaoyu Feng, yfeng@ecust.edu.cn.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.02997-13

studies have examined the distribution of *E. bieneusi* genotypes in wild mammals in drinking source watersheds. In a study of fecal specimens from 239 muskrats, 85 beavers, 67 foxes, 55 raccoons, and 19 otters in watersheds in the Chesapeake Bay area, 17 *E. bieneusi* genotypes (WL1 to WL17) were found among 59 PCR-positive specimens. Three *E. bieneusi* genogroups were found, with groups 1, 3, and 4 seen in 43 animals (from 13 beavers, 11 muskrats, 9 foxes, 8 raccoons, and 2 otters), 9 animals (all from muskrats), and 7 animals (all from raccoons), respectively (18). Likewise, only a few studies have assessed the public health potential of *E. bieneusi* in water by using genotyping tools, which showed the presence of group 1 genotypes in a small number of samples genotyped (13, 14).

In this study, *E. bieneusi* in fecal specimens from wildlife living in the watershed of the New York City (NYC) water supply system was detected and genotyped by PCR and sequence analysis of the ITS. The host range of each genotype was used to assess the extent of host adaptation in *E. bieneusi*. Data on the host range of *E. bieneusi* genotypes were used to infer the source and potential public health significance of *E. bieneusi* in storm runoff within the watershed. The latter interpretation was also aided by data on the distribution of host-adapted *Cryptosporidium* species/genotypes present in the water samples. Data from this study indicated that most *E. bieneusi* isolates in wild animals and all isolates from water in the watershed had not been found in humans, although the host range of some of the so-called host-adapted genogroups was broader than previously believed.

MATERIALS AND METHODS

Animal specimens. Fecal specimens were collected between September 2005 and July 2007 from 255 wild mammals in the New York City Department of Environmental Protection (NYCDEP) watershed. They were a part of a previous study on the distribution of *Cryptosporidium* species and genotypes in wildlife (31). The collection areas included reservoir buffer lands (forest, woodland, transition edge, and mixed meadow/shrub), areas along stream corridors, and maintained surfaces (lawns and parks). The collection of animal fecal specimens was described in detail previously (31). All animals were identified by species (except *Peromyscus*) and age, with most animals being adults. Samples were initially collected by NYCDEP staff and then shipped in coolers to the laboratory at the Centers for Disease Control and Prevention (CDC) for pathogen detection by PCR.

Stormwater samples. Sixty-seven stormwater samples were collected during July 2002 to October 2003 from three sites within the NYCDEP Kensico watershed in Valhalla, NY, including the E9, Malcolm Brook, and N5 basins. They were mostly a part of a previous study on the distribution and public health potential of *Cryptosporidium* species and genotypes in stormwater within the New York City drinking source water supply (32). The N5 stream basin is 298 acres, is located on the west side of the watershed, and consists almost entirely of residential lots (91%). The Malcolm Brook basin is 95 acres, is close to the N5 site, and has large wooded areas (39% of the total area) and corporate parks (22%), in addition to relatively high-density suburban residential lots (34%) with public sewer systems. The E9 stream basin, in contrast, is located on the east side of the watershed, is 321 acres, and drains a large wetland area (18.4% of the total area). In addition, it has large wooded areas (56.5%) and some development areas (17.6%) (Fig. 1).

Stormwater samples were collected with preset autosamplers (Isco 6700; Isco, Inc., Lincoln, NE), which were set to collect 20-liter composite samples over a predicted intensity of the storm event. The samples were filtered through an Envirochek HV filter (Pall Gelman Laboratory, Ann Arbor, MI) in the NYCDEP laboratory according to procedures described in U.S. Environmental Protection Agency Method 1623 (33). Material on

the filter was eluted, and a portion of the concentrated pellet from each sample was transported to the CDC laboratory for molecular analysis. The Malcolm Brook and N5 basin sites and stormwater sample collection were described in detail previously (32). All stormwater samples were collected in stream channels prior to entry into the reservoir; therefore, they were not yet subjected to reservoir residence time, treatment with chlorine, or UV disinfection.

DNA extraction. After washing the specimens twice with distilled water by centrifugation, genomic DNA was extracted from 0.2 ml of fecal specimens or 0.5 ml of stormwater concentrates by using the FastDNA Spin kit for soil (MP Biomedicals, Santa Ana, CA) and eluted in 100 μ l of reagent-grade water, as described previously (34). The extracted DNAs from both fecal specimens and stormwater samples were used for PCR detection of *E. bieneusi*, whereas those from stormwater samples were further analyzed for *Cryptosporidium* by PCR. To neutralize residual PCR inhibitors in the extracted DNA, 400 ng/ μ l of nonacetylated bovine serum albumin (Sigma-Aldrich, St. Louis, MO) was used for primary PCR for both *E. bieneusi* and *Cryptosporidium* detection. In a previous study on the distribution of *Cryptosporidium* genotypes in these and other stormwater samples (32), DNA was extracted by using a QIAamp DNA minikit from immunomagnetic separation-purified oocysts present in a separate portion of the water concentrates.

E. bieneusi detection and genotyping. A nested PCR analysis of a ~392-bp fragment of the rRNA gene containing the entire internal transcribed spacer (ITS) was used to detect *E. bieneusi*, using a dog-specific *E. bieneusi* genotype (PtEbIX) as the positive control (18). The primers used included AL4037 (5'-GATGGTCATAGGGATGAAGAGCTT-3') and AL4039 (5'-ACGGATCCAAGTGATCCTGTATT-3') for primary PCR and AL4038 (5'-AGGGATGAAGAGCTTCGGCTCTG-3') and AL4040 (5'-AGTGATCCTGTATTAGGGATATT-3') for secondary PCR. Each animal specimen was analyzed twice by using 2 μ l of extracted DNA per PCR, whereas each stormwater sample was analyzed once by using 3 μ l of extracted DNA per PCR. Positive PCR products were sequenced to determine *E. bieneusi* genotypes. *E. bieneusi* genotypes were named according to an established nomenclature system (35). The genotypes with ITS sequences identical to those in the GenBank database were considered known genotypes. In contrast, genotypes that produced ITS sequences with any single-nucleotide substitutions, deletions, or insertions and were confirmed by DNA sequencing of at least two PCR products were considered new genotypes.

Cryptosporidium detection and genotyping. To aid in the interpretation of the source of *E. bieneusi* genotypes in stormwater samples, we also genotyped *Cryptosporidium* present in the same extracted DNAs by nested PCR amplification of an approximately 830-bp fragment of the small-subunit (SSU) rRNA gene, as *Cryptosporidium* spp. are known to have more strict host specificity. The primers used include AL1687 (5'-TCTAGAGCTAATACATGCG-3') and AL3417 (5'-CCCATTTCCTTCG AACAGGA-3') for primary PCR and AL3032 (5'-GGAAGGGTTGTATTATTAGATAAAG-3') and AL4871 (5'-CTCATAAGGTGCTGAAGGA GTA-3') for secondary PCR (32). DNA of *Cryptosporidium serpentis* was used as the positive control. The *Cryptosporidium* species and genotypes present were differentiated by restriction fragment length polymorphism (RFLP) analysis of the secondary PCR products using the restriction enzymes SspI and VspI (32). All positive PCR products were sequenced to confirm the genotype identification. To detect the occurrence of mixed *Cryptosporidium* species and genotypes, each water sample was analyzed six times by the PCR-RFLP technique, using 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 μ l of DNA per PCR.

Sequence analysis. After purification using Montage PCR filters (Millipore, Bedford, MA), the secondary PCR products were sequenced in both directions with secondary PCR primers by using the ABI BigDye Terminator v3.1 cycle sequencing kit (Applied Biosystems, Foster City, CA) and an ABI3130 genetic analyzer (Applied Biosystems). Nucleotide sequences obtained were aligned with reference sequences downloaded from the GenBank database by using ClustalX (<http://www.clustal.org/>)




FIG 1 Sampling sites in the Kensico watershed of the New York City Department of Environmental Protection (NYCDEP).

to determine *E. bienersi* genotypes and *Cryptosporidium* species or genotypes. For ITS sequences of *E. bienersi*, to confirm the genogroup designation and assess the genetic relationship among genotypes, a neighbor-joining tree was constructed by using Mega 5.2 (<http://www.megasoftware.net/>), based on genetic distances calculated by the Kimura two-parameter model. A sequence (GenBank accession no. [DQ885585](https://www.ncbi.nlm.nih.gov/nuccore/DQ885585)) from the dog-specific genotype (PtEbIX) was used as the outgroup. The reliability of various clusters was evaluated by the bootstrap method with 1,000 replicates. The topology of the tree was supported by a maximum parsimony analysis of the same sequence alignment, with all alignment sites taken into consideration.

Nucleotide sequence accession numbers. Unique *E. bienersi* ITS sequences obtained from wildlife and stormwater during the study were deposited in the GenBank database under accession numbers [KF591677](https://www.ncbi.nlm.nih.gov/nuccore/KF591677) to [KF591688](https://www.ncbi.nlm.nih.gov/nuccore/KF591688).

RESULTS

Prevalence of *E. bienersi* in wild mammals. *E. bienersi* was detected in 74 (29.0%) of the 255 wild mammals sampled (Table 1). These animals belonged to 23 species in six orders. Most of the

positive animals (64/74) were rodents and carnivores, while only 10 were other mammals. Rodents, however, accounted for over one-half (142/255) of the sampled animals and had an *E. bienersi* infection rate (38/142; 26.8%) similar to the average infection rate. Within rodents, members of the squirrel family had the highest infection rate (21/49; 42.9%), while the remaining 17 positive rodents were all from the Cricetidae (deer mice and voles). Among two other groups of animals commonly examined, carnivores had an infection rate (26/42; 61.9%) much higher than the average, whereas ruminants (white-tailed deer) had an infection rate (6/49; 12.2%) lower than the average. The small numbers of insectivores (5 animals), lagomorphs (8), and marsupials (9) sampled did not allow an accurate determination of *E. bienersi* infection rates; only 3/8 eastern cottontails and 1/9 Virginia opossums sampled were positive (Table 1).

***E. bienersi* genotypes in wildlife.** Altogether, 16 *E. bienersi* genotypes were found among the 74 PCR-positive specimens, including 7 known genotypes (D, Peru11, PtEbV, type IV, WL2,

TABLE 1 Prevalence of *Enterocytozoon bieneusi* genotypes in wild mammals in the NYCDEP watershed

Animal	No. of samples	No. of positive samples	Infection rate (%)	Genotype(s) (no. of samples positive)	Genogroup(s) (no. of samples positive)
Rodents	142	38	26.8		
Sciuridae	49	21	42.9		
<i>Sciurus carolinensis</i> (eastern gray squirrel)	34	11	32.4	Type IV (3), WL4 (5), WW6 (2), PtEbV + WL21 (1)	1 (4), 3 (5), 4 (2)
<i>Sciurus vulgaris</i> (red squirrel)	2	0	0		
<i>Glaucomys volans</i> (southern flying squirrel)	1	0	0		
<i>Tamias striatus</i> (eastern chipmunk)	7	5	71.4	Type IV (1), WL4 (3), WL23 (1)	1 (1), 3 (4)
<i>Marmota monax</i> (woodchuck)	5	5	100.0	Type IV + WL20 (1), WL4 (2), WL22 (1), WW6 (1)	1 (1), 3 (3), 4 (1)
Castoridae	16	0	0		
<i>Castor canadensis</i> (beaver)	16	0	0		
Muridae	1	0	0		
<i>Rattus norvegicus</i> (Norway rat)	1	0	0		
Zapodidae	1	0	0		
<i>Napaeozapus insignis</i> (woodland jumping mouse)	1	0	0		
Cricetidae	71	17	23.9		
<i>Peromyscus</i> sp. (deer mouse)	55	13	23.6	WL4 (10), WL23 (2), WL25 (1)	3 (13)
<i>Myodes gapperi</i> (boreal red-backed vole)	5	1	20.0	WL20 + WL21(1)	1 (1)
<i>Microtus pennsylvanicus</i> (meadow vole)	10	3	30.0	Peru11 (1), Peru11 + type IV (1), WL21 + unknown (1)	1 (3)
<i>Ondatra zibethicus</i> (muskrat)	1	0	0		
Erethizontidae	4	0	0		
<i>Erethizon dorsatum</i> (porcupine)	4	0	0		
Carnivores	42	26	61.9		
<i>Mustela vison</i> (mink)	4	0	0		
<i>Mustela erminea</i> (ermine)	1	1	100.0	WL4 (1)	3 (1)
<i>Procyon lotor</i> (raccoon)	22	18	81.8	WL4 (8), WW6 (7), Peru11 (1), WL26 (1), WL24 (1)	3 (8), 4 (8), 1 (1), 7 (1)
<i>Ursus americanus</i> (black bear)	5	2	40.0	Type IV (1), WL4 (1)	1 (1), 3 (1)
<i>Lontra canadensis</i> (river otter)	8	5	62.5	D (2), WL4 (2), WL2 (1)	1 (2), 3 (2), 4 (1)
<i>Mephitis mephitis</i> (striped skunk)	2	0	0		
Insectivores	5	0	0		
<i>Blarina brevicauda</i> (northern short-tailed shrew)	5	0	0		
Lagomorpha	8	3	37.5		
<i>Sylvilagus floridanus</i> (eastern cottontail)	8	3	37.5	Peru11 (1), Peru11 + WL4 (1), mixed (1)	1 (1), 1 + 3 (1), unknown (1)
Ruminants	49	6	12.2		
<i>Odocoileus virginianus</i> (white-tailed deer)	49	6	12.2	WL18 (2), WL19 (2), WL4 (2)	1 (4), 3 (2)
Marsupials	9	1	11.1		
<i>Didelphis virginiana</i> (Virginia opossum)	9	1	11.1	Mixed (1)	Unknown (1)
Total ^a	255	74	29.0		

^a There were 16 total genotypes detected and 4 total genogroups detected.

WL4, and WW6) and 9 new ones (named WL18 to WL26 in this study). WL4 was the most common genotype (35), followed by WW6 (10), type IV (7), and Peru11 (5). Other genotypes were detected in only 1 to 3 animals (Table 1). Seven specimens had mixed genotypes, as reflected by discordant genotype results in replicate PCRs (five samples) or underlying signals in sequence trace files, which prevented determination of genotypes (two samples).

E. *bieneusi* genotypes in stormwater samples. Thirty-nine (58.2%) of the 67 stormwater samples analyzed were PCR positive for *E. bieneusi*, including 3 of 4 (75.0%) samples from the E9 basin, 3 of 15 (20.0%) samples from Malcolm Brook, and 33 of 48 (68.8%) samples from the N5 basin. These isolates belonged to five genotypes, including two known genotypes (WL4 and WL6) and three new genotypes (called SW1, SW2, and SW3). WL4 was the most common genotype, being seen in 23 samples. This was followed by WL6, being seen in five samples. In contrast, the three new genotypes were each seen in only 1 to 2 samples. Seven samples had mixed genotypes, as reflected by the underlying signals in sequence trace files. There were apparent differences in the occurrences and genotype distributions of *E. bieneusi* among the three sampling sites. The N5 and E9 basins had a higher occurrence of *E. bieneusi* than Malcolm Brook, with PCR positivities of 68.8%,

75.0%, and 20.0%, respectively. WL4 was also the dominant *E. bieneusi* genotype in the N5 and E9 basins but was absent in Malcolm Brook (Table 2).

Phylogenetic relationship of *E. bieneusi* isolates. Altogether, 20 *E. bieneusi* genotypes were found in this study. Both the neighbor-joining and maximum parsimony analyses suggested that these genotypes belonged to five genotype groups or genogroups: groups 1, 3, 4, and 7 and an outlier (Fig. 2). Four of the five genogroups, including groups 1, 3, 4, and 7, were seen in 20, 40, 12, and 1 animal, respectively (Table 1). Groups 1 and 3 had a broad host range; each group was detected in 10 animal species belonging to the rodent, carnivore, lagomorph, and ruminant groups. In contrast, group 4 was detected in only four animal species belonging to the carnivore and rodent groups, with raccoons being the major host and members of the squirrel family being minor hosts. The only detection of group 7 was in a raccoon (Table 1). Three of the five genogroups were found in stormwater samples, including groups 1 and 3 and an outlier, with the majority (28/32) of the genotyped samples having group 3 genotypes (Table 2). An almost identical topology was produced by both phylogenetic approaches. The only exception was the placement of the outlier found in one stormwater sample. In the neighbor-joining analysis, the outlier was placed near the base of the tree but

TABLE 2 *Enterocytozoon bieneusi* and *Cryptosporidium* genotypes in stormwater samples from the NYCDEP watershed

Site	<i>E. bieneusi</i>			<i>Cryptosporidium</i>			Possible major animal source(s) for pathogens in water
	No. of samples	No. (%) of positive samples	Genotypes (no. of samples positive)	Major known host(s)	No. (%) of positive samples	Genotypes ^a (no. of replicate PCR-positive samples)	
E9 basin	4	3 (75.0)	WL4 (2), mixed (1)	Deer mice, muskrats, raccoons, squirrel family	3 (75.0)	W7 (4), W10 (2), W11 (2), W16 (1)	Muskrats, birds, snakes
Malcolm Brook	15	3 (20.0)	SW1 (1), SW3 (1), mixed (1)	Unknown	6 (40.0)	W4 (10), W10 (2), W7 (1)	Squirrel family, birds
N5 basin	48	33 (68.8)	SW1 (1), SW2 (1), WL4 (2), WL6 (5), mixed (5)	Deer mice, muskrats, raccoons, squirrel family	35 (72.9)	W4 (23), W1 (15), W13 (14), W10 (8), W7 (7), W19 (5), W16 (3), W17 (2), W8 (2), W12 (2), W15 (2), W5 (1), W11 (1)	Squirrel family, raccoons, muskrats
Total ^b	67	39 (58.2)			44 (65.7)		

^a The abbreviations of *Cryptosporidium* genotypes found in water are as follows: W1, deer mouse genotype III; W4, *C. ubiquitum*; W5, shrew genotype; W7, muskrat genotype I; W8, opossum genotype II; W10, *C. baileyi*; W11, snake genotype; W12, unknown genotype; W13, skunk genotype; W15, vole genotype; W16, muskrat genotype II; W17, chipmunk genotype I; W19, rat genotype IV.
^b There were 5 total genotypes and 2 total genogroups for *E. bieneusi* and 13 total *Cryptosporidium* genotypes detected.

was separated from genotype PtEbIX in dogs (Fig. 2). In the maximum parsimony analysis, the outlier grouped together with genotype PtEbIX, with 94% bootstrap support.

***Cryptosporidium* genotypes in stormwater.** Forty-four (65.7%) of the 67 stormwater samples analyzed were PCR positive for *Cryptosporidium* spp. in at least one of the six replicate PCR analyses, including 3 of 4 (75.0%) samples from the E9 basin, 6 of 15 (40.0%) samples from Malcolm Brook, and 35 of 48 (72.9%) samples from the N5 basin (Table 2). Of the 44 *Cryptosporidium*-positive samples, 9, 15, 15, 24, 20, and 25 samples generated the expected SSU rRNA PCR products when 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 μl of the 100 μl of extracted DNA were used for PCR, respectively. In general, there were good agreements in the occurrences of *Cryptosporidium* and *E. bieneusi*, although the number of samples from one of the study sites, the E9 basin, was small. Thus, the overall prevalence of *Cryptosporidium* in these water samples was similar to that of *E. bieneusi* (72.9% and 68.8%, respectively); sites with higher *E. bieneusi* occurrences also had higher *Cryptosporidium* occurrences, and *E. bieneusi*-positive water samples were almost twice as likely to be positive for *Cryptosporidium* than were *E. bieneusi*-negative samples (32/39 [82.1%] versus 12/28 [42.8%]).

Altogether, 13 *Cryptosporidium* species/genotypes were found in stormwater, including deer mouse genotype III (W1), *C. ubiquitum* (W4), the shrew genotype (W5), muskrat genotype I (W7), opossum genotype II (W8), *C. baileyi* (W10), the snake genotype (W11), an unknown genotype (W12), the skunk genotype (W13), the vole genotype (W15), muskrat genotype II (W16), chipmunk genotype I (W17), and rat genotype IV (W19). The most common genotypes included *C. ubiquitum*, deer mouse genotype III, the skunk genotype, muskrat genotype I, and *C. baileyi*, being found in 33, 15, 14, 12, and 12 PCR replicates, respectively. The remaining genotypes were each found in only 1 to 5 PCR replicates. There were differences in the distributions of common *Cryptosporidium* species/genotypes among study sites. Thus, although *C. baileyi* and muskrat genotype I were found at all three sampling sites, deer mouse genotype III and the skunk genotype were found only in the N5 basin. Likewise, the most commonly detected *Cryptosporidium* species in the study, *C. ubiquitum*, was absent in the E9 basin (Table 2).

DISCUSSION

The results of this study suggest that the host range of group 3 *E. bieneusi* genotypes is much broader than previously believed. This group of genotypes was initially found in muskrats in the Chesapeake area. Subsequent ITS sequence analyses of numerous *E. bieneusi* isolates from domestic animals and humans have failed to detect them thus far. In this study, however, one known genotype and three new genotypes of this phylogenetically unique group were found in 10 species of mammals of four orders. This is about the same host range of eight group 1 genotypes in this study, which are well known to have a broad host range. In fact, one member of group 3, WL4, was by far the most commonly detected genotype in this study, being identified in nearly one-half (35/74) of the *E. bieneusi*-positive animals. Likewise, group 4 genotypes, which were previously considered raccoon specific, were also found in other carnivores in addition to raccoons as well as in several species of rodents in the squirrel family. Nevertheless, these genetically unique genotypes are rarely seen in domestic animals, implying that they do have some host preference. Indeed, in one of the




FIG 2 Phylogenetic relationship among various *E. bieneusi* genotypes in wildlife in the NYCDEP watershed and known *E. bieneusi* genotypes, as inferred by a neighbor-joining analysis of SSU rRNA sequences. Genotypes with open circles are known genotypes found in the NYCDEP watershed, and those with black circles are new genotypes. A similar tree topology was also produced by maximum parsimony analysis, with the exception that the SW3 genotype grouped together with genotype PtEbIX, with 94% bootstrap support. Numbers on branches are percent bootstrap values (>50) using 1,000 replicates, with the first number being generated by neighbor-joining analysis and the second number being generated by maximum parsimony analysis. NS, nonsignificant (bootstrap value of <50%).

most sampled animals in this study, deer mice, only group 3 genotypes were found in the 13 *E. bieneusi*-positive animals.

The dominant *E. bieneusi* genotype in animals, WL4, was also the dominant genotype in stormwater samples, suggesting that wildlife played a significant role in contamination of stormwater by *E. bieneusi* in the watershed. This was especially the case in the N5 basin, which is much larger and less commercially developed

than the neighboring Malcolm Brook. The high usage of lands for commercial parks and high-density residential lots in the Malcolm Brook basin might have reduced the activity of wildlife, thus resulting in the disappearance of the WL4 genotype at this site. Interestingly, although the N5 basin consists mostly of large residential lots and Malcolm Brook has many high-density residential lots, no group 1 genotypes previously detected in humans were

found in any stormwater samples, indicating that humans and companion animals were not major contributors to *E. bienersi* contamination in stormwater in the watershed. This is in agreement with the genotyping results for *Cryptosporidium* oocysts isolated from different aliquots of these samples in a previous study, which showed that wildlife contributed exclusively to *Cryptosporidium* spp. in stormwater from the Malcolm Brook and N5 basins (32).

The lack of strict host specificity makes it difficult to track the source of contamination of *E. bienersi* in stormwater to specific wildlife species or groups, especially when the dominant genotype, WL4, was found in 10 species of mammals of four orders in the watershed. Because WL4 was most often detected in members of the squirrel family, deer mice, and raccoons in this study and was shown to be a common genotype in muskrats, it is likely that these animals were the main source of *E. bienersi* in stormwater. To facilitate the interpretation of contamination in stormwater, we analyzed the same DNA from stormwater samples in this study for *Cryptosporidium* spp. There was good agreement in the occurrences of *Cryptosporidium* and *E. bienersi* in both the overall prevalence and sample-specific positivity. There was also good agreement between the nature of dominant *E. bienersi* and *Cryptosporidium* genotypes at the Malcolm Brook and N5 basins (only four samples were collected from the E9 basin). Over one-half of the *Cryptosporidium*-positive stormwater samples (21/35) and 33 of 108 PCR-positive replicates had *C. ubiquitum* (W4), which was found mostly in members of the squirrel family in the watershed. This, together with the occurrence of another *Cryptosporidium* genotype of the squirrel family, chipmunk genotype I (W17), supports the role of these animals in *E. bienersi* contamination in the watershed, especially in the N5 basin and Malcolm Brook. The role of raccoons in *E. bienersi* contamination in the N5 basin was also supported by the common finding (in 11 samples and 14 PCR replicates) of the *Cryptosporidium* skunk genotype (commonly found in raccoons) in stormwater samples. Likewise, the relatively common finding of *Cryptosporidium* muskrat genotypes I (W7) and II (W16) and the vole genotype (W15) indicates that muskrats contributed to *E. bienersi* contamination in water in the N5 and E9 basins. In contrast, *Cryptosporidium* genotyping results do not support a significant role of deer mice in *E. bienersi* contamination at all these sampling sites. Although *Cryptosporidium* deer mouse genotype III (W1) was commonly seen in stormwater samples (6 samples and 15 PCR replicates), another common *Cryptosporidium* genotype in deer mice, deer mouse genotype IV (W3), was absent. As deer mouse genotype III also occurs in squirrels at a high frequency, members of the squirrel family could be responsible for the occurrence of deer mouse genotype III in stormwater. Taken together, data from *E. bienersi* and *Cryptosporidium* genotyping indicate that muskrats were a major source of pathogen contamination in stormwater in the E9 basin, and squirrel family members were a major source in Malcolm Brook, while squirrel family members, raccoons, and muskrats were all major sources of pathogens in stormwater in the N5 basin (Table 2). These results are in agreement with the environmental ecology and land use at these sampling sites.

Most of the *E. bienersi* isolates found in wildlife and stormwater samples are not from human-pathogenic genotypes. Among the 20 *E. bienersi* genotypes found in this study, only 3 have ever been detected in humans: type IV, Peru11, and D, which were detected in 7, 5, and 2 animals, respectively. None of them have

been found in stormwater. Seven other genotypes in the study also belong to the potentially human-pathogenic group 1, but altogether, group 1 genotypes were found in only 20 of 255 (7.8%) wildlife samples and 20 of 74 (27.0%) *E. bienersi*-positive animals. Likewise, only 3 of 67 (4.5%) stormwater samples and 3 of 39 (7.7%) *E. bienersi*-positive stormwater samples had group 1 genotypes. Some of the group 1 genotypes in wildlife and stormwater samples are newly discovered genotypes. Because host adaptation also exists within group 1 genotypes (36), it is likely that not all group 1 genotypes are human pathogenic. Thus, wildlife may contribute to *E. bienersi* contamination in stormwater but most likely do not have public health significance because they are generally infected with non-human-pathogenic genotypes. Moreover, stormwater samples from this study were collected prior to entry into the reservoir and any treatment with chlorine or UV disinfection, which may further reduce the public health impact of *E. bienersi* in source water.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (grants 31229005 and 31110103901); Fundamental Research Funds for the Central Universities, China; the National Special Fund for the State Key Laboratory of Bioreactor Engineering (no. 2060204); and the Centers for Disease Control and Prevention.

We thank the NYCDEP Water Quality staff for the collection and shipping of wildlife and stormwater samples for this study.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

REFERENCES

- Ghosh K, Weiss LM. 2009. Molecular diagnostic tests for microsporidia. *Interdiscip. Perspect. Infect. Dis.* 2009:926521. <http://dx.doi.org/10.1155/2009/926521>.
- Didier ES, Weiss LM. 2006. Microsporidiosis: current status. *Curr. Opin. Infect. Dis.* 19:485–492. <http://dx.doi.org/10.1097/01.qco.0000244055.46382.23>.
- Matos O, Lobo ML, Xiao L. 2012. Epidemiology of *Enterocytozoon bienersi* infection in humans. *J. Parasitol. Res.* 2012:981424. <http://dx.doi.org/10.1155/2012/981424>.
- Santin M, Fayer R. 2011. Microsporidiosis: *Enterocytozoon bienersi* in domesticated and wild animals. *Res. Vet. Sci.* 90:363–371. <http://dx.doi.org/10.1016/j.rvsc.2010.07.014>.
- Didier ES, Stovall ME, Green LC, Brindley PJ, Sestak K, Didier PJ. 2004. Epidemiology of microsporidiosis: sources and modes of transmission. *Vet. Parasitol.* 126:145–166. <http://dx.doi.org/10.1016/j.vetpar.2004.09.006>.
- Wumba R, Longo-Mbenza B, Menotti J, Mandina M, Kintoki F, Sitakibanza NH, Kakicha MK, Zanga J, Mbanzulu-Makola K, Nseka T, Mukendi JP, Kendjo E, Sala J, Thellier M. 2012. Epidemiology, clinical, immune, and molecular profiles of microsporidiosis and cryptosporidiosis among HIV/AIDS patients. *Int. J. Gen. Med.* 5:603–611. <http://dx.doi.org/10.2147/IJGM.S32344>.
- Cotte L, Rabodonirina M, Chapuis F, Bailly F, Bissuel F, Raynal C, Gelas P, Persat F, Piens MA, Trepo C. 1999. Waterborne outbreak of intestinal microsporidiosis in persons with and without human immunodeficiency virus infection. *J. Infect. Dis.* 180:2003–2008. <http://dx.doi.org/10.1086/315112>.
- Hunter PR. 2000. Waterborne outbreak of microsporidiosis. *J. Infect. Dis.* 182:380–381. <http://dx.doi.org/10.1086/315654>.
- Graczyk TK, Sunderland D, Awantang GN, Mashinski Y, Lucy FE, Graczyk Z, Chomicz L, Breyse PN. 2010. Relationships among bather density, levels of human waterborne pathogens, and fecal coliform counts in marine recreational beach water. *Parasitol. Res.* 106:1103–1108. <http://dx.doi.org/10.1007/s00436-010-1769-2>.
- Graczyk TK, Girouard AS, Tamang L, Nappier SP, Schwab KJ. 2006. Recovery, bioaccumulation, and inactivation of human waterborne

- pathogens by the Chesapeake Bay nonnative oyster, *Crassostrea ariakensis*. *Appl. Environ. Microbiol.* 72:3390–3395. <http://dx.doi.org/10.1128/AEM.72.5.3390-3395.2006>.
11. Coupe S, Delabre K, Pouillot R, Houdart S, Santillana-Hayat M, Derouin F. 2006. Detection of *Cryptosporidium*, *Giardia* and *Enterocytozoon bieneusi* in surface water, including recreational areas: a one-year prospective study. *FEMS Immunol. Med. Microbiol.* 47:351–359. <http://dx.doi.org/10.1111/j.1574-695X.2006.00098.x>.
 12. Graczyk TK, Conn DB, Lucy F, Minchin D, Tamang L, Moura LN, DaSilva AJ. 2004. Human waterborne parasites in zebra mussels (*Dreissena polymorpha*) from the Shannon River drainage area, Ireland. *Parasitol. Res.* 93:385–391. <http://dx.doi.org/10.1007/s00436-004-1142-4>.
 13. Ye J, Xiao L, Ma J, Guo M, Liu L, Feng Y. 2012. Anthroponotic enteric parasites in monkeys in public park, China. *Emerg. Infect. Dis.* 18:1640–1643. <http://dx.doi.org/10.3201/eid1810.120653>.
 14. Galvan AL, Magnet A, Izquierdo F, Fenoy S, Rueda C, Fernandez Vadillo C, Henriques-Gil N, del Aguila C. 2013. Molecular characterization of human-pathogenic microsporidia and *Cyclospora cayetanensis* isolated from various water sources in Spain: a year-long longitudinal study. *Appl. Environ. Microbiol.* 79:449–459. <http://dx.doi.org/10.1128/AEM.02737-12>.
 15. Breton J, Bart-Delabesse E, Biligui S, Carbone A, Seiller X, Okome-Nkoumou M, Nzamba C, Kombila M, Accoceberry I, Thellier M. 2007. New highly divergent rRNA sequence among biodiverse genotypes of *Enterocytozoon bieneusi* strains isolated from humans in Gabon and Cameroon. *J. Clin. Microbiol.* 45:2580–2589. <http://dx.doi.org/10.1128/JCM.02554-06>.
 16. Thellier M, Breton J. 2008. *Enterocytozoon bieneusi* in human and animals, focus on laboratory identification and molecular epidemiology. *Parasite* 15:349–358. <http://dx.doi.org/10.1051/parasite/2008153349>.
 17. Henriques-Gil N, Haro M, Izquierdo F, Fenoy S, del Aguila C. 2010. Phylogenetic approach to the variability of the microsporidian *Enterocytozoon bieneusi* and its implications for inter- and intrahost transmission. *Appl. Environ. Microbiol.* 76:3333–3342. <http://dx.doi.org/10.1128/AEM.03026-09>.
 18. Sulaiman IM, Fayer R, Lal AA, Trout JM, Schaefer FW, III, Xiao L. 2003. Molecular characterization of microsporidia indicates that wild mammals harbor host-adapted *Enterocytozoon* spp. as well as human-pathogenic *Enterocytozoon bieneusi*. *Appl. Environ. Microbiol.* 69:4495–4501. <http://dx.doi.org/10.1128/AEM.69.8.4495-4501.2003>.
 19. Sulaiman IM, Fayer R, Yang C, Santin M, Matos O, Xiao L. 2004. Molecular characterization of *Enterocytozoon bieneusi* in cattle indicates that only some isolates have zoonotic potential. *Parasitol. Res.* 92:328–334. <http://dx.doi.org/10.1007/s00436-003-1049-5>.
 20. Widmer G, Akiyoshi DE. 2010. Host-specific segregation of ribosomal nucleotide sequence diversity in the microsporidian *Enterocytozoon bieneusi*. *Infect. Genet. Evol.* 10:122–128. <http://dx.doi.org/10.1016/j.meegid.2009.11.009>.
 21. Feng Y, Li N, Dearen T, Lobo ML, Matos O, Cama V, Xiao L. 2011. Development of a multilocus sequence typing tool for high-resolution genotyping of *Enterocytozoon bieneusi*. *Appl. Environ. Microbiol.* 77:4822–4828. <http://dx.doi.org/10.1128/AEM.02803-10>.
 22. Li N, Xiao L, Wang L, Zhao S, Zhao X, Duan L, Guo M, Liu L, Feng Y. 2012. Molecular surveillance of *Cryptosporidium* spp., *Giardia duodenalis*, and *Enterocytozoon bieneusi* by genotyping and subtyping parasites in wastewater. *PLoS Negl. Trop. Dis.* 6:e1809. <http://dx.doi.org/10.1371/journal.pntd.0001809>.
 23. Akinbo FO, Okaka CE, Omoregie R, Adamu H, Xiao L. 2013. Unusual *Enterocytozoon bieneusi* genotypes and *Cryptosporidium hominis* subtypes in HIV-infected patients on highly active antiretroviral therapy. *Am. J. Trop. Med. Hyg.* 89:157–161. <http://dx.doi.org/10.4269/ajtmh.12-0635>.
 24. Santin M, Cortes Vecino JA, Fayer R. 2008. *Enterocytozoon bieneusi* genotypes in dogs in Bogota, Colombia. *Am. J. Trop. Med. Hyg.* 79:215–217. <http://www.ajtmh.org/content/79/2/215.long>.
 25. Li W, Kiulia NM, Mwenda JM, Nyachio A, Taylor MB, Zhang X, Xiao L. 2011. *Cyclospora papionis*, *Cryptosporidium hominis*, and human-pathogenic *Enterocytozoon bieneusi* in captive baboons in Kenya. *J. Clin. Microbiol.* 49:4326–4329. <http://dx.doi.org/10.1128/JCM.05051-11>.
 26. Santin M, Fayer R. 2009. A longitudinal study of *Enterocytozoon bieneusi* in dairy cattle. *Parasitol. Res.* 105:141–144. <http://dx.doi.org/10.1007/s00436-009-1374-4>.
 27. Fayer R, Santin M, Trout JM. 2007. *Enterocytozoon bieneusi* in mature dairy cattle on farms in the eastern United States. *Parasitol. Res.* 102:15–20. <http://dx.doi.org/10.1007/s00436-007-0746-x>.
 28. Santin M, Trout JM, Fayer R. 2005. *Enterocytozoon bieneusi* genotypes in dairy cattle in the eastern United States. *Parasitol. Res.* 97:535–538. <http://dx.doi.org/10.1007/s00436-005-1482-8>.
 29. Lee JH. 2007. Prevalence and molecular characteristics of *Enterocytozoon bieneusi* in cattle in Korea. *Parasitol. Res.* 101:391–396. <http://dx.doi.org/10.1007/s00436-007-0468-0>.
 30. Ben Ayed L, Yang W, Widmer G, Cama V, Ortega Y, Xiao L. 2012. Survey and genetic characterization of wastewater in Tunisia for *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bieneusi*, *Cyclospora cayetanensis* and *Eimeria* spp. *J. Water Health* 10:431–444. <http://dx.doi.org/10.2166/wh.2012.204>.
 31. Feng Y, Alderisio KA, Yang W, Blancero LA, Kuhne WG, Nadareski CA, Reid M, Xiao L. 2007. *Cryptosporidium* genotypes in wildlife from a New York watershed. *Appl. Environ. Microbiol.* 73:6475–6483. <http://dx.doi.org/10.1128/AEM.01034-07>.
 32. Jiang J, Alderisio KA, Xiao L. 2005. Distribution of *Cryptosporidium* genotypes in storm event water samples from three watersheds in New York. *Appl. Environ. Microbiol.* 71:4446–4454. <http://dx.doi.org/10.1128/AEM.71.8.4446-4454.2005>.
 33. US EPA. 2001. Method 1623: *Cryptosporidium* and *Giardia* in water by filtration/IMS/FAEPA 821-R-01–025. Office of Water, US Environmental Protection Agency, Washington, DC.
 34. Jiang J, Alderisio KA, Singh A, Xiao L. 2005. Development of procedures for direct extraction of *Cryptosporidium* DNA from water concentrates and for relief of PCR inhibitors. *Appl. Environ. Microbiol.* 71:1135–1141. <http://dx.doi.org/10.1128/AEM.71.3.1135-1141.2005>.
 35. Santin M, Fayer R. 2009. *Enterocytozoon bieneusi* genotype nomenclature based on the internal transcribed spacer sequence: a consensus. *J. Eukaryot. Microbiol.* 56:34–38. <http://dx.doi.org/10.1111/j.1550-7408.2008.00380.x>.
 36. Li W, Cama V, Akinbo FO, Ganguly S, Kiulia NM, Zhang X, Xiao L. 2013. Multilocus sequence typing of *Enterocytozoon bieneusi*: lack of geographic segregation and existence of genetically isolated sub-populations. *Infect. Genet. Evol.* 14:111–119. <http://dx.doi.org/10.1016/j.meegid.2012.11.021>.