

Environmental Determinants of Transformation Efficiency in *Helicobacter pylori*

Mary E. Moore,^a Anna Lam,^a Srijak Bhatnagar,^b Jay V. Solnick^{a,b,c}

Center for Comparative Medicine,^a Department of Microbiology and Immunology,^b and Department of Medicine,^c University of California, Davis, Davis, California, USA

***Helicobacter pylori* uses natural competence and homologous recombination to adapt to the dynamic environment of the stomach mucosa and maintain chronic colonization. Although *H. pylori* competence is constitutive, its rate of transformation is variable, and little is known about factors that influence it. To examine this, we first determined the transformation efficiency of *H. pylori* strains under low O₂ (5% O₂, 7.6% CO₂, 7.6% H₂) and high O₂ (15% O₂, 2.9% CO₂, 2.9% H₂) conditions using DNA containing an antibiotic resistance marker. *H. pylori* transformation efficiency was 6- to 32-fold greater under high O₂ tension, which was robust across different *H. pylori* strains, genetic loci, and bacterial growth phases. Since changing the O₂ concentration for these initial experiments also changed the concentrations of CO₂ and H₂, transformations were repeated under conditions where O₂, CO₂, and H₂ were each varied individually. The results showed that the increase in transformation efficiency under high O₂ was largely due to a decrease in CO₂. An increase in pH similar to that caused by low CO₂ was also sufficient to increase transformation efficiency. These results have implications for the physiology of *H. pylori* in the gastric environment, and they provide optimized conditions for the laboratory construction of *H. pylori* mutants using natural transformation.**

H. pylori is a naturally competent (1), Gram-negative bacterium that colonizes the human gastric mucosa and causes peptic ulcer or gastric cancer in about 10% of those infected (2). Strains isolated from different individuals demonstrate extraordinary genomic diversity (3). Like other organisms with small genomes (4), *H. pylori* has relatively few regulatory networks (5) and instead exploits natural competence and genetic changes to adapt to changing conditions (6). Since infection is chronic over the lifetime of the host, sometimes with more than one resident strain (7–9), frequent genetic exchange enables *H. pylori* to better adapt to the changing environment of the host stomach (10, 11) and to acquire antibiotic resistance (12). Thus, natural competence is essential to the lifestyle of *H. pylori*.

H. pylori natural competence is mediated by a two-step process: uptake of double-stranded DNA across the outer membrane followed by transport of single-stranded DNA through the inner membrane (13). DNA is transported across the *H. pylori* outer membrane and into the periplasm via a ComB type IV secretion system (T4SS) similar to the Vir system of *Agrobacterium tumefaciens*. ComB components are expressed from two operons and have been named according to homology with Vir proteins. Several have been shown to be essential for *H. pylori* natural competence. For example, ComB9, homologous to VirB9, is thought to be an important structural component, tightly anchored to the outer membrane within the periplasmic space (1), and ComB4 is an ATPase putatively responsible for function of the ComB structure (13). Both proteins are thought to be involved in the biogenesis of the ComB apparatus (1). Once inside the periplasm, double-stranded DNA is processed to single-stranded DNA and transported across the bacterial inner membrane by a ComEC channel (14). Single-stranded DNA can then become incorporated into the bacterial chromosome via RecA-mediated homologous recombination (13).

Transformation efficiency in *H. pylori* is variable and can be affected by the length and similarity of DNA sequences and restriction-modification systems (15–17). Variation also occurs during *H. pylori* growth phases, with peaks at mid-log- and late-

stationary-phase growth (18). Since natural transformation is an important mechanism for bacterial adaptation to the gastric environment, certain conditions of stress may increase *H. pylori* competence. For example, induction of double-strand breaks by treatment with ciprofloxacin can increase expression of *comB* genes and enhance competence (19). Taken together, these studies suggest that *H. pylori* has evolved to increase natural competence when this is beneficial for its survival.

Another stress condition that might alter *H. pylori* competence is a change in the partial pressure of O₂ or CO₂. CO₂ is highly prevalent in the gastric environment, while O₂ tension is low. In the live mouse, for example, the partial O₂ pressure (pO₂) in the stomach is estimated at 58 ± 15 torr (~7.6%) (20). O₂ and CO₂ concentrations are variable within the gastric environment and can be affected by disease state, food consumption, and location within the gastric mucosa (21, 22). For example, the CO₂ concentration tends to be low nearest the epithelium (where the pH is high) and higher toward the gastric lumen (where the pH is low). Although, *H. pylori* is generally regarded as a microaerophile (2), it expresses high levels of catalase and superoxide dismutase, which facilitate the elimination of toxic oxygen intermediates (23, 24). More recent studies have shown that many strains can tolerate and even thrive in the presence of high O₂ levels as long as CO₂ levels are high (25). These studies indicate that *H. pylori* is a capnophile whose growth is promoted by atmospheric O₂ as long as sufficient CO₂ is present.

Here we show that low levels of CO₂ markedly increase *H.*

Received 29 May 2013 Accepted 28 October 2013

Published ahead of print 1 November 2013

Address correspondence to Jay V. Solnick, jvsolnick@ucdavis.edu.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/JB.00633-13>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.00633-13

TABLE 1 Transformation conditions and defined gas mixtures

Transformation conditions (%, vol/vol)			Defined gas mixture (%, vol/vol)		
Oxygen	Carbon dioxide	Hydrogen	Carbon dioxide	Hydrogen	Nitrogen
5.0	7.6	7.6	10.0	10.0	80.0
15.0	2.9	2.9	10.0	10.0	80.0
15.0	7.7	7.7	27.0	27.0	46.0
5.0	2.9	7.6	3.8	10.0	86.2
5.0	7.6	2.9	10.0	3.8	86.2

pylori transformation efficiency and induce expression of the ComB T4SS. This change in competence seems to be a result of CO₂ influence on the pH of the bacterial culture medium. These results are important for understanding the physiology of *H. pylori* in the gastric environment and also have practical implications for using natural transformation to generate *H. pylori* mutants in the laboratory.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *Helicobacter pylori* strains J166 and SS1 were cultured on brucella agar (Becton, Dickinson and Company, Sparks, MD) supplemented with 5% heat-inactivated newborn calf serum (Gibco, Grand Island, NY), 5 µg/ml trimethoprim, 10 µg/ml vancomycin, 2.5 units/ml polymyxin B, and 2.5 µg/ml amphotericin B (Sigma-Aldrich, Inc., St. Louis, MO). Cultures were grown at 37°C using the Anoxomat system (Advanced Instruments, Inc., Norwood, MA), which is a sealed jar connected to a pressurized tank that contained one of several defined gas mixtures (Table 1). Specific gas conditions in the sealed jar were achieved by removing a defined volume of ambient air and replacing it with the appropriate defined gas. Transformation efficiency was initially studied under 5% O₂ (7.6% CO₂, 7.6% H₂) or 15% O₂ (2.9% CO₂, 2.9% H₂), where O₂ was manipulated by changing the volume of a single defined gas mixture, allowing the other gases to vary. Separate defined gas mixtures were then used to manipulate O₂, CO₂, and H₂ individually, using N₂ as a replacement gas (Table 1).

Plasmid construction. Two plasmids were used to measure transformation efficiency, one that deleted an outer membrane protein (pOMP) and a second that deleted an intergenic region (pIG) between oppositely oriented genes and did not disrupt a known open reading frame. A third plasmid (pRecA) was developed to disrupt the *recA* gene to determine the influence of RecA expression on natural competence genes under relevant experimental conditions. Construction of the plasmids is shown schematically in Fig. 1; primer sequences are given in Table 2. To construct pOMP for strain J166 (pOMPJ166), a chloramphenicol resistance cassette (26) was amplified and ligated to DNA sequences from upstream and downstream of the promoter region in the *babA* gene (designated HP1243 in *H. pylori* 26695), which encodes the blood group antigen binding protein (27). A similar plasmid was constructed for strain SS1 (pOMPSS1), except that upstream and downstream sequences were amplified so as to delete the entire HP1243 locus, which in SS1 carries a *babA* paralog, *babC*. To construct pIG, a nonpolar kanamycin cassette (*aphA3*) was amplified from pUC18K (28) and ligated between HP0512 and HP0519 amplified from strain J166 (pIGJ166) or SS1 (pIGSS1), which in both strains are transcribed in opposite directions and contain no intervening open reading frames. These features, along with the insertion of a nonpolar cassette, minimized potential effects on the strain phenotype. Construction of pRecA was the same as for pIG except that *aphA3* was ligated into the *recA* open reading frame of *H. pylori* SS1. PCR-amplified fragments were digested with the appropriate enzymes (New England BioLabs, Ipswich, MA), ligated into NotI/XhoI-digested pBluescript SK- vector (Stratagene, La Jolla, CA) using T4 DNA ligase (New England BioLabs, Ipswich,

MA) according to the manufacturer's instructions, and cloned into *Escherichia coli* Top10 (Invitrogen, Carlsbad CA) with selection on chloramphenicol (5 µg/liter) or kanamycin (25 µg/liter) as appropriate.

Transformations. Transformations were performed as previously described (29). Briefly, overnight (18- to 24-h) *H. pylori* cultures were harvested from agar plates using a sterile swab and suspended in brucella broth to an optical density at 600 nm (OD₆₀₀) of 5, and 10 µl was spread evenly in a 1.5-cm-diameter circle on a fresh agar plate. Five replicate circles of these bacterial cells were then incubated at 37°C in an Anoxomat jar adjusted to contain the appropriate atmospheric conditions. After 3 to 6 h, 0.3 to 3 µg of plasmid or genomic DNA suspended in deionized water was mixed with each of the circles using a sterile loop. The cells with DNA were then returned to 37°C in the same gas mixture with which they had been conditioned. After overnight incubation, the bacteria were removed from the 1.5-cm circle using a sterile loop and suspended in 300 µl brucella broth. Dilutions were plated on nonselective and selective plates with 5 µg/ml chloramphenicol or 25 µg/ml kanamycin. Transformation efficiency was calculated as the number of CFU present on selective medium divided by the number of CFU on nonselective medium divided by the amount (µg) of DNA used for transformation.

Effects of growth phase on competence. *H. pylori* SS1 harvested from an overnight agar culture was plated on brucella agar in 1.5-cm-diameter circles as described for transformations. Plates were incubated under 5% or 15% O₂ conditions, and at 6-h intervals from 0 to 30 h, 3.0 µg of pIGSS1 was added to three replicate circles and mixed with a sterile loop. After 6 h of additional incubation with DNA, bacteria were removed from the circles with a sterile loop, suspended in brucella broth, and then plated and analyzed for both CFU and transformation efficiency.

RNA extraction and quantitative reverse transcription-PCR (qRT-PCR). A 100-µl suspension in brucella broth (OD₆₀₀ = 5) was harvested from an overnight brucella agar culture of *H. pylori* SS1, spread evenly on a brucella agar plate, and incubated at 37°C with 5% or 15% O₂ for 18 to 24 h. Bacterial cells were suspended in 1 ml TRIzol reagent (Ambion,




FIG 1 Schematic of plasmid construction. pOMP was constructed by inserting a chloramphenicol resistance cassette (CAT) into the promoter region (*H. pylori* J166) or the open reading frame of *babA* (HP1234) (*H. pylori* SS1). pIG was constructed by insertion of a kanamycin resistance cassette (*aphA3*) into an intergenic region between HP0519 and HP0512 in *H. pylori* J166 and SS1. pRecA was constructed by insertion of *aphA3* into the *recA* gene in *H. pylori* SS1. The primers used for amplification of plasmid components for each strain are represented by small numbered arrows and correspond to the primers in Table 2. Gene designations are for *H. pylori* 26695.

TABLE 2 Primers used for plasmid construction and RT-PCR

Number	Primer name	Restriction site	Sequence (5' → 3') ^a
1	aphA3_F	SacI	AAC <u>GAGCTC</u> GGT ACC CGG GTG AC
2	aphA3_R	HincII	AAC <u>GTCGAC</u> TCT AGA GGA TCC CC
3	J166_IG_F1	HincII	AAT <u>GTCGAC</u> AAA TCT TAT TAT CGA CAC TCT ATT AGA A
4	J166_IG_R1	XhoI	AAT <u>CTCGAG</u> CCC TTT GTT GGC CGT CAA
5	J166_IG_F2	NotI	AAT <u>GCGGCCGC</u> TAG CAC CGC TCA AGA CAC AA
6	J166_IG_R2	SacI	AAT <u>GAGCTC</u> CCA AGT GAT AGT AAA GAT TAG GTA A
7	SS1_IG_F1	XhoI	AAT <u>CTCGAG</u> ATC ATA CCC CTA TTT CTA GGA
8	SS1_IG_R1	HincII	AAT <u>GTCGAC</u> AAA TCT TAT CAT CGA CAC TCT ATT AAA G
9	SS1_IG_R2	SacI	AAT <u>GAGCTC</u> TCA AGT GAT AGT AAA GAT TAG GTA G
10	cat_F	SacI	AAC <u>GAGCTC</u> GAT GCT TTA TAA CTA TGG ATT AAA CAC
11	cat_R	BamHI	AAC <u>GGATCC</u> TTA TCA GTG CGA CAA ACT GGG AT
12	J166_OMP_F1	BamHI	AAC <u>GGATCC</u> TTG CTC CAC GCT GAA GAC
13	J166_OMP_R1	XhoI	AAC <u>CTCGAG</u> GAC GCT CGT TTG ATT GAC CA
14	J166_OMP_F2	NotI	AAC <u>GCGGCCGC</u> AGC CAC AAA ACC TCT AAA GA
15	J166_OMP_R2	SacI	AAC <u>GAGCTC</u> GGG GTA TTT TGA AAT AAC TCT C
16	SS1_OMP_F1	XhoI	AAC <u>CTCGAG</u> TTT TGA GCC GTT GGA TAT ATT AG
17	SS1_OMP_R1	SacI	AAC <u>GAGCTC</u> GCA AGC TCT TTT ATT ATT TAT CTT A
18	SS1_OMP_F2	BamHI	AAC <u>GGATCC</u> CGT TAA AAC CTT TTG TGA AAC T
19	SS1_OMP_R2	NotI	AAA <u>GCGGCCGC</u> TGT GGA TCT AGC GGT TCT T
20	aphA3_F2	PstI	AAT <u>CTGCAG</u> GGT ACC CGG GTG AC
21	SS1_RecA_F1	NotI	AAT <u>GCGGCCGC</u> TCG TTA CTG CCC TTA ATG AGC TC
22	SS1_RecA_R1	PstI	AAT <u>CTGCAG</u> CAA TTT GTT TGA TCG CTA AAG AAA TCG C
23	SS1_RecA_F2	HincII	AAT <u>GTCGAC</u> AAT GAA GAG ATC ATG CCC TTA CCC
24	SS1_RecA_R1	XhoI	AAT <u>CTCGAG</u> AAAAGACAATCAGGGAGCTATGGC
25	comB4_F1		TGT TAG AAA AGC TTT TAA GCG C
26	comB4_R1		TTG CTC TGT GCT TAA ATG GGT G
27	comB9_F		AAG CCG ATG ATT TTT TAG AAG AAG C
28	comB9_R		AAA GCT CCC TTG AAT GGC GTT TAA
29	recA_F		GTG CGC CTT GGG GAT AAG CA
30	recA_R		CTT AGA GTG GTC TTC CCG CTT GAC
31	16S_F		GGA GTA CGG TCG CAA GAT TAA A
32	16S_F		CTA GCG GAT TCT CTC AAT GTC AA

^a Underlining indicates a restriction site.

Carlsbad, CA), and total RNA was isolated and then treated with DNase I (Roche Applied Science, Mannheim, Germany) according to the manufacturer's directions. RNA was purified using an RNeasy Minikit (Qiagen Sciences, Germantown, MD). Gene-specific primers (Table 2) were used to synthesize cDNA with the SuperScript III first-strand synthesis system for RT-PCR (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. A 1- μ l aliquot of a 1:5 dilution of cDNA was used to perform quantitative PCR with the same primers using SYBR green PCR master mix with standard cycling parameters (Applied Biosystems, Warrington, WA) in 20- μ l reaction mixtures on an ABI 7900HT 96 well block.

Determination of CO₂ effects on brucella medium pH. The pH of 25 ml sterile brucella broth (Becton, Dickinson and Company, Sparks, MD) was measured using a pH meter (Beckman Coulter) after overnight incubation under 7.6% CO₂ (5% O₂) or 2.9% CO₂ (15% O₂) at 37°C. Brucella agar plates, supplemented with 11 mg/liter phenol red (Matheson, Coleman & Bell, Norwood, OH), were incubated under 7.6% CO₂ (5% O₂) or 2.9% CO₂ (15% O₂) at 37°C overnight. The pH of brucella agar plates from each condition was visually estimated by comparison to a standard curve, prepared by supplementing 25-ml aliquots of brucella broth with 11 mg/liter phenol red and adjusting the pH in increments of 0.2 from 5.8 to 6.8 using 6 N HCl or NaOH. The agar pHs of two to four brucella agar plates from each condition were independently estimated by four observers, and the average values are reported.

Statistical analysis. Pairwise comparisons were performed using the Student *t* test with Welch's correction on GraphPad Prism software. Data involving bacterial counts were log transformed prior to statistical analysis. Error bars in all figures represent standard error.

RESULTS

***H. pylori* transformation efficiency under 5% and 15% O₂.** To determine the transformation efficiency of *H. pylori* under different atmospheric conditions, we first incubated *H. pylori* SS1 with plasmid DNA directed to a *babA* paralog (pOMPSS1) and compared the results under 5% O₂ to 15% O₂ using a single defined gas mixture (Table 1). The transformation efficiency was increased approximately 6-fold in 15% O₂ (Fig. 2A). To determine if these results were strain specific, we repeated the experiment with *H. pylori* J166 using the corresponding plasmid, pOMPJ166. This resulted in approximately 8-fold-greater transformation efficiency in 15% O₂ than in 5% O₂ (Fig. 2B). The transformation efficiency could be increased under high O₂ conditions because deletion of an OMP increases *H. pylori* fitness under high O₂, though it seems unlikely since two different OMPs were targeted (*babA* in J166 and *babC* in SS1). In other words, the effect of O₂ on transformation efficiency could be specific to one genetic locus. To test this, we next transformed *H. pylori* SS1 and J166 with plasmid DNA containing a kanamycin resistance cassette directed to an intergenic region (pIGSS1 and pIGJ166) that would not be expected to disrupt an open reading frame. Under 15% O₂, the transformation efficiencies in strains SS1 and J166 increased 32-fold and 10-fold, respectively (Fig. 2C and D). These results show that a change in atmospheric conditions can cause a robust change




FIG 2 Transformation efficiency in *H. pylori* is increased under conditions of high O_2 and low CO_2 . *H. pylori* SS1 and J166, as indicated, were transformed with pOMPSS1 (A), pOMPJ166 (B), pIGSS1 (C), pIGJ166 (D), genomic DNA from pIGSS1-transformed *H. pylori* SS1 (E) and genomic DNA from pIGJ166-transformed *H. pylori* J166 (F) at 5% O_2 (7.6% CO_2 , 7.6% H_2) or 15% O_2 (2.9% CO_2 , 2.9% H_2). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$.

in *H. pylori* natural competence that is independent of strain or genetic locus.

To ensure that this phenomenon was not specific to plasmid DNA, genomic DNA was isolated from *H. pylori* SS1 and J166 which had been transformed with pIGSS1 and pIGJ166, respectively. This DNA was then used to perform transformation efficiency experiments as described above. Under 15% O_2 , the transformation efficiency increased 24-fold and 34-fold in strains SS1 and J166, respectively (Fig. 2E and F), illustrating that both plasmid and genomic DNAs are taken up by the bacteria with increased efficiency under 15% O_2 conditions.

Atmospheric effects on transformation efficiency are independent of growth phase. High O_2 conditions that increased transformation efficiency (Fig. 2) also resulted in a trend of decreased bacterial growth (Fig. 3). Since it has previously been shown that transformation efficiency is variable throughout *H. pylori* growth phases (18), we considered the possibility that the effects of changing atmospheric conditions on competence were due to altered growth. To address this, we measured the transformation efficiency of *H. pylori* SS1 at different growth phases under both 5% and 15% O_2 . pIGSS1 DNA was added at 6-h intervals under 5% and 15% O_2 , and the transformation efficiency was determined 6 h later. As expected, 15% O_2 caused a trend of decreased bacterial growth (Fig. 4A). No transformants were detected at 0 h, but when DNA was added after 6 to 30 h of growth (mid-log to stationary phase), the transformation efficiency was markedly higher at 15% O_2 at each point in the growth curve (Fig. 4B). Thus, although 15% O_2 moderately impaired *H. pylori* growth, it also increased competence at every point in the growth




FIG 3 A trend of reduced *H. pylori* growth is observed under conditions of high O_2 and low CO_2 . CFU for *H. pylori* SS1 (A, C, and E) and J166 (B, D, and F) under high and low O_2 conditions are shown. The data correspond to the transformation efficiency results shown in Fig. 2. *, $P < 0.05$; ***, $P < 0.001$; n.s., not significant.




FIG 4 *H. pylori* growth and transformation efficiency are affected by atmospheric conditions throughout the growth curve. (A) Total CFU present under 5% O_2 (solid line) or 15% O_2 (dotted line) when transformation efficiency was measured for each time point shown. (B) Transformation efficiency of *H. pylori* SS1 under 5% O_2 (filled bars) or 15% O_2 (open bars), as measured 6 h after the addition of pIGSS1, which was added at the time points shown. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; n.s. = not significant.




FIG 5 ComB expression is induced by conditions of high O₂ and low CO₂ independent of RecA. *H. pylori* SS1 (A) or SS1 Δ*recA* (B) mRNA expression levels of the *comB4*, *comB9*, *recA*, and 16S rRNA genes after overnight incubation with 5% O₂ (filled bars) or 15% O₂ (open bars) are shown. In both SS1 and SS1 Δ*recA*, *comB4* and *comB9* were significantly induced under 15% O₂. Induction of *recA* did not occur. 16S rRNA was used as a loading control. *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001; ****, *P* < 0.0001; n.s. = not significant.

curve, indicating that growth phase is not responsible for the increase in natural competence at 15% O₂ (Fig. 4).

The ComB T4SS is induced under 15% O₂ conditions independent of RecA. *H. pylori* natural competence occurs primarily through expression of the ComB T4SS apparatus. RecA is also an important protein involved in homologous recombination following uptake of exogenous DNA. Therefore, we examined expression of the *comB* and *recA* genes under both high and low O₂ conditions. Since the ComB apparatus is composed of elements present on two operons, *comB2-4* and *comB6-10* (30), we examined expression of one *comB* gene from each operon (*comB4* and *comB9*). qRT-PCR analysis of RNA isolated from *H. pylori* SS1 incubated under either high or low O₂ conditions showed significantly increased expression of both *comB4* and *comB9* under 15% O₂ conditions (Fig. 5A). No changes in expression of *recA* were seen. This result is in agreement with previous studies (19) indicating that *recA* expression was not increased during induction of other natural competence genes. 16S rRNA, which served as a loading control, was also unchanged. These results suggest that increased *H. pylori* competence under high O₂ is mediated by the ComB T4SS.

It has previously been shown that RecA is required for *H. pylori* induction of *comB* genes in response to DNA double-strand breaks (19). To determine whether the induction of *comB* genes in response to increased O₂ was also dependent on RecA, we developed a *recA* deletion mutant of *H. pylori* SS1 (SS1 Δ*recA*) using p*RecA*. qRT-PCR was performed on RNAs from three individual SS1 Δ*recA* clones incubated under either 5% or 15% O₂ conditions. Significant induction of both *comB4* and *comB9* was detected under 15% O₂ (Fig. 5B). These data indicate that induction




FIG 6 *H. pylori* transformation efficiency is induced by low CO₂ rather than high O₂ levels. Transformation efficiency as measured after the addition of pIGSS1 (A) and total CFU (B) of *H. pylori* SS1 under defined atmospheric conditions are shown. *, *P* < 0.05; **, *P* < 0.01; ****, *P* < 0.0001; n.s. = not significant.

of the ComB apparatus under 15% O₂ conditions occurs by a novel mechanism, independent of the response to double-strand break induction.

***H. pylori* natural competence is induced in response to low CO₂.** In all of the above-described experiments in which a single defined gas mixture was used, CO₂ and H₂ concentrations were decreased to accommodate an increased O₂ concentration. Because recent work shows that *H. pylori* is a capnophile (25, 31), we asked whether the change in CO₂, rather than O₂, was responsible for the increase in transformation efficiency. Transformation efficiency in *H. pylori* SS1 was examined using pIGSS1 under a series of conditions (Table 1) in which we individually manipulated O₂, H₂, or CO₂, leaving the other gases at the same levels found under low O₂ conditions. Nitrogen was used as a replacement gas. When O₂ was increased to 15% with no change in CO₂ or H₂, a small (1.8-fold) increase in competence was observed. In contrast, reducing CO₂ without changing O₂ or H₂ caused a marked (40-fold) increase in transformation efficiency (Fig. 6A). Reduction of H₂ alone had no effect. Additionally, the least amount of growth was measured under conditions of low CO₂, which resulted in the highest transformation efficiency (Fig. 6B). These results indicate that the increase in *H. pylori* transformation efficiency under different atmospheric conditions is predominantly a result of a decreased CO₂ concentration.

Increased pH mimics the increase in transformation efficiency caused by decreased CO₂. Since CO₂ reacts with water to form carbonic acid, differences in CO₂ concentration could affect the pH of the unbuffered culture medium. The pH of autoclaved brucella broth was 6.87 (±0.01), which decreased to 6.31 (±0.02) after overnight incubation under 7.6% CO₂ (5% O₂). Conditions




FIG 7 A change in pH is sufficient to reproduce the effects of CO₂ on *H. pylori* transformation efficiency. (A) The transformation efficiency of *H. pylori* SS1 was increased both by a decrease in CO₂ (2.9%) and by an increase in pH under buffered conditions (7.6% CO₂). (B) Increased transformation efficiency was accompanied by small decreases in CFU. **, $P < 0.01$; ***, $P < 0.001$; n.s. = not significant.

of low (2.9%) CO₂ resulted in a lesser decrease in pH, to 6.47 (± 0.03), after overnight incubation. Since all transformations were done on agar-grown bacteria, phenol red was added to brucella agar medium prior to autoclaving in order to verify that a similar effect occurred in brucella agar plates. Brucella agar plates containing phenol red were incubated under 7.6% CO₂ or 2.9% CO₂ overnight. The color of the plates before and after incubation at each of the conditions was estimated by comparison to a phenol red standard curve. Under aerobic conditions, the pH of brucella agar was 6.76 (± 0.04), and this decreased to 6.55 (± 0.08) after incubation at 7.6% CO₂. Low (2.9%) CO₂ conditions again resulted in less of a decrease in pH, to 6.63 (± 0.03). Thus, incubation under conditions of increased CO₂ caused a decrease in the pH of unbuffered brucella medium that was relative to the amount of CO₂ present.

To determine whether increased pH could be responsible for the increase in competence under low CO₂ conditions, transformation efficiency was determined on buffered brucella agar plates adjusted to pH 6.5 or 7.0 with 20 mM phosphate buffer. After incubation at 7.6% CO₂, the pH of the plates buffered at 6.5 or 7.0 decreased slightly to 6.46 (± 0.06) and 6.69 (± 0.02), respectively. *H. pylori* SS1 was transformed with pIGSS1 on these buffered plates under 7.6% CO₂. The transformation efficiency of *H. pylori* SS1 was about 35-fold higher with pIGSS1 at pH 6.69 than at pH 6.46 under 7.6% CO₂, which mimicked the increase in competence caused by conditions of low CO₂ and high O₂ (Fig. 7A). This was also accompanied by decreased growth (Fig. 7B). Incubation of *H. pylori* in liquid medium prepared in the same way as the plates described above showed that although the presence of *H.*

pylori could moderately affect the absolute pH of the buffered medium, it did not significantly affect the difference in pH between the two buffers (see Fig. S1 in the supplemental material). Together these results show that increased pH is sufficient to increase transformation efficiency, and they suggest that this may explain the increased competence under conditions of low CO₂.

DISCUSSION

Genomic diversity is a signature feature of *H. pylori* that arises from mutation, recombination, and importation of exogenous DNA via natural transformation, which allows *H. pylori* to adapt to the hostile environment of the human stomach. Variability in competence among different *H. pylori* strains is well documented (16, 32, 33), but little is known about conditions that affect transformation efficiency. This is important, not only for understanding the physiology of *H. pylori* in the gastric environment but also for optimizing methods that are commonly exploited in the laboratory to generate isogenic mutants. Here we systematically examined the effects of environmental growth conditions on *H. pylori* transformation efficiency.

Competence in naturally transformable bacteria is commonly induced by physiological stress (34). Since *H. pylori* is considered a microaerophile that is subject to toxic oxygen species under ambient O₂ conditions (35), we first examined the impact of O₂ concentration on natural competence. Culturing *H. pylori* under 15% versus 5% O₂ decreased bacterial growth, increased transformation efficiency, and induced the *comB* operon, which is required for natural competence. These effects were robust across different *H. pylori* strains, genetic loci, and bacterial growth phases (Fig. 2 to 5). However, in these experiments manipulation of O₂ was accomplished using a single defined gas mixture, so the increase in O₂ was also accompanied by decreases in CO₂ and H₂. Since some studies have suggested that *H. pylori* is a capnophile (25, 31), we considered the possibility that low CO₂ rather than high O₂ was the relevant environmental variable responsible for increasing competence. Independent manipulation of O₂ and CO₂ revealed that in fact low CO₂ was primarily responsible for the increased transformation efficiency, though there was also a small effect of increased O₂, which can induce the *comB* operon (31). An increase in pH similar to that caused by incubating unbuffered agar plates under low CO₂ also increased transformation efficiency. Thus, increased pH is itself sufficient to enhance *H. pylori* competence. Increased pH has also been observed to enhance competence in *Campylobacter jejuni* (36, 37) and *Streptococcus pneumoniae* (38), which suggests that this response may be a convergent evolutionary adaptation of naturally competent, opportunistic mucosal pathogens.

It may at first seem paradoxical that *H. pylori* transformation efficiency increased under high pH (low CO₂), since the stomach is commonly thought of as a harsh, acidic environment in which the acidity might be expected to induce a stress response and increase competence. However, several lines of evidence are consistent with the observation that increased pH may induce stress in *H. pylori*. First, studies of gene expression *in vitro* and *in vivo* suggest that the natural habitat of *H. pylori* within the gastric mucosa is actually quite acidic, with a pH of ≤ 4.0 (39). In fact, *H. pylori* cannot survive neutral pH in the presence of physiological concentrations of urea because the pH rapidly rises (40). The stringent response is also required for *H. pylori* survival under high O₂ conditions (41), which can be abrogated by addition of

CO₂ (25). *Ex vivo* measurements of guinea pig gastric tissue indicate that a pH gradient exists across the mucosa, with a pH of approximately 6.1 immediately adjacent to the epithelial cell surface that gradually decreases toward the gastric lumen (42). Therefore, *in vivo* it is likely that increased pH and reduced CO₂ would be experienced immediately adjacent to the host epithelial cells (43), while *H. pylori* is thought to more commonly localize just offshore in the mucous layer (44). It may be that *H. pylori* undergoes a stress response when located close to the gastric epithelium, with consequent upregulation of ComB and increased competence.

We conclude that the increased transformation efficiency observed *in vitro* under elevated pH and decreased CO₂ provides insight into *H. pylori* natural competence regulation within the gastric mucosa. Our results indicate that *H. pylori* competence is highly sensitive to subtle changes that could be experienced within microenvironments of the mucous layer. These conditions of increased pH and reduced CO₂ can also be exploited to optimize transformation efficiency for generating isogenic mutants in the laboratory.

ACKNOWLEDGMENTS

The *H. pylori* SS1 sequence, used for the construction of pOMPSS1 and pIGSS1, was generously provided by Jenny Draper and Karen Ottemann. We also thank Lori Hansen for designing several primers.

This work was supported by grants R01 AI081037 and R01 AI070803 from the National Institutes of Health to Jay V. Solnick.

REFERENCES

- Hofreuter D, Odenbreit S, Haas R. 2001. Natural transformation competence in *Helicobacter pylori* is mediated by the basic components of a type IV secretion system. *Mol. Microbiol.* 41:379–391. <http://dx.doi.org/10.1046/j.1365-2958.2001.02502.x>.
- Kusters JG, van Vliet AH, Kuipers EJ. 2006. Pathogenesis of *Helicobacter pylori* infection. *Clin. Microbiol. Rev.* 19:449–490. <http://dx.doi.org/10.1128/CMR.00054-05>.
- Falush D, Wirth T, Linz B, Pritchard JK, Stephens M, Kidd M, Blaser MJ, Graham DY, Vacher S, Perez-Perez GI, Yamaoka Y, Megraud F, Otto K, Reichard U, Katzowitz E, Wang X, Achtman M, Suerbaum S. 2003. Traces of human migrations in *Helicobacter pylori* populations. *Science* 299:1582–1585. <http://dx.doi.org/10.1126/science.1080857>.
- Galperin MY. 2006. Structural classification of bacterial response regulators: diversity of output domains and domain combinations. *J. Bacteriol.* 188:4169–4182. <http://dx.doi.org/10.1128/JB.01887-05>.
- Tomb JF, White O, Kerlavage AR, Clayton RA, Sutton GG, Fleischmann RD, Ketchum KA, Klenk HP, Gill S, Dougherty BA, Nelson K, Quackenbush J, Zhou L, Kirkness EF, Peterson S, Loftus B, Richardson D, Dodson R, Khalak HG, Glodek A, McKenney K, Fitzgerald LM, Lee N, Adams MD, Hickey EK, Berg DE, Gocayne JD, Utterback TR, Peterson JD, Kelley JM, Cotton MD, Weidman JM, Fujii C, Bowman C, Watthey L, Wallin E, Hayes WS, Borodovsky M, Karp PD, Smith HO, Fraser CM, Venter JC. 1997. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 388:539–547. <http://dx.doi.org/10.1038/41483>.
- Baltrus DA, Guillemin K, Phillips PC. 2008. Natural transformation increases the rate of adaptation in the human pathogen *Helicobacter pylori*. *Evolution* 62:39–49. <http://dx.doi.org/10.1111/j.1558-5646.2007.00271.x>.
- Taylor NS, Fox JG, Akopyants NS, Berg DE, Thompson N, Shames B, Yan L, Fonham E, Janney F, Hunter FM, et al. 1995. Long-term colonization with single and multiple strains of *Helicobacter pylori* assessed by DNA fingerprinting. *J. Clin. Microbiol.* 33:918–923.
- Morales-Espinosa R, Castillo-Rojas G, Gonzalez-Valencia G, Ponce de Leon S, Cravioto A, Atherton JC, Lopez-Vidal Y. 1999. Colonization of Mexican patients by multiple *Helicobacter pylori* strains with different *vacA* and *cagA* genotypes. *J. Clin. Microbiol.* 37:3001–3004.
- Patra R, Chattopadhyay S, De R, Ghosh P, Ganguly M, Chowdhury A, Ramamurthy T, Nair GB, Mukhopadhyay AK. 2012. Multiple infection and microdiversity among *Helicobacter pylori* isolates in a single host in India. *PLoS One* 7:e43370. <http://dx.doi.org/10.1371/journal.pone.0043370>.
- Dorer MS, Cohen IE, Sessler TH, Fero J, Salama NR. 2013. Natural competence promotes *Helicobacter pylori* chronic infection. *Infect. Immun.* 81:209–215. <http://dx.doi.org/10.1128/IAI.01042-12>.
- Amundsen SK, Fero J, Hansen LM, Cromie GA, Solnick JV, Smith GR, Salama NR. 2008. *Helicobacter pylori* AddAB helicase-nuclease and RecA promote recombination-related DNA repair and survival during stomach colonization. *Mol. Microbiol.* 69:994–1007. <http://dx.doi.org/10.1111/j.1365-2958.2008.06336.x>.
- Smeets LC, Arents NL, van Zwet AA, Vandenbroucke-Grauls CM, Verboom T, Bitter W, Kusters JG. 2003. Molecular patchwork: chromosomal recombination between two *Helicobacter pylori* strains during natural colonization. *Infect. Immun.* 71:2907–2910. <http://dx.doi.org/10.1128/IAI.71.5.2907-2910.2003>.
- Stingl K, Muller S, Scheidgen-Kleyboldt G, Clausen M, Maier B. 2010. Composite system mediates two-step DNA uptake into *Helicobacter pylori*. *Proc. Natl. Acad. Sci. U. S. A.* 107:1184–1189. <http://dx.doi.org/10.1073/pnas.0909955107>.
- Chen I, Dubnau D. 2004. DNA uptake during bacterial transformation. *Nat. Rev. Microbiol.* 2:241–249. <http://dx.doi.org/10.1038/nrmicro844>.
- Humbert O, Dorer MS, Salama NR. 2011. Characterization of *Helicobacter pylori* factors that control transformation frequency and integration length during inter-strain DNA recombination. *Mol. Microbiol.* 79:387–401. <http://dx.doi.org/10.1111/j.1365-2958.2010.07456.x>.
- Levine SM, Lin EA, Emara W, Kang J, DiBenedetto M, Ando T, Falush D, Blaser MJ. 2007. Plastic cells and populations: DNA substrate characteristics in *Helicobacter pylori* transformation define a flexible but conservative system for genomic variation. *FASEB J.* 21:3458–3467. <http://dx.doi.org/10.1096/fj.07-8501.com>.
- Kumar R, Mukhopadhyay AK, Ghosh P, Rao DN. 2012. Comparative transcriptomics of *H. pylori* strains AM5, SS1 and their hpyAVIBM deletion mutants: possible roles of cytosine methylation. *PLoS One* 7:e42303. <http://dx.doi.org/10.1371/journal.pone.0042303>.
- Baltrus DA, Guillemin K. 2006. Multiple phases of competence occur during the *Helicobacter pylori* growth cycle. *FEMS Microbiol. Lett.* 255:148–155. <http://dx.doi.org/10.1111/j.1574-6968.2005.00066.x>.
- Dorer MS, Fero J, Salama NR. 2010. DNA damage triggers genetic exchange in *Helicobacter pylori*. *PLoS Pathog.* 6:e1001026. <http://dx.doi.org/10.1371/journal.ppat.1001026>.
- He G, Shankar RA, Chzhan M, Samouilov A, Kuppusamy P, Zweier JL. 1999. Noninvasive measurement of anatomic structure and intraluminal oxygenation in the gastrointestinal tract of living mice with spatial and spectral EPR imaging. *Proc. Natl. Acad. Sci. U. S. A.* 96:4586–4591. <http://dx.doi.org/10.1073/pnas.96.8.4586>.
- Steer H. 2009. The source of carbon dioxide for gastric acid production. *Anat. Rec. (Hoboken)* 292:79–86. <http://dx.doi.org/10.1002/ar.20762>.
- Taylor DE, Gutierrez G, Clark C, Hainley S. 1997. Measurement of gastric mucosal carbon dioxide tension by saline and air tonometry. *J. Crit. Care* 12:208–213. [http://dx.doi.org/10.1016/S0883-9441\(97\)90034-4](http://dx.doi.org/10.1016/S0883-9441(97)90034-4).
- Krieg NR, Hoffman PS. 1986. Microaerophily and oxygen toxicity. *Annu. Rev. Microbiol.* 40:107–130. <http://dx.doi.org/10.1146/annurev.mi.40.100186.000543>.
- Kelly DJ, Hughes NJ, Poole RK. 2001. Microaerobic physiology: aerobic respiration, anaerobic respiration, and carbon dioxide metabolism, p 113–124. *In* Mobley HLT, Mendz GL, Hazell SL (ed), *Helicobacter pylori: physiology and genetics*. ASM Press, Washington, DC.
- Park SA, Ko A, Lee NG. 2011. Stimulation of growth of the human gastric pathogen *Helicobacter pylori* by atmospheric level of oxygen under high carbon dioxide tension. *BMC Microbiol.* 11:96. <http://dx.doi.org/10.1186/1471-2180-11-96>.
- Styer CM, Hansen LM, Cooke CL, Gundersen AM, Choi SS, Berg DE, Benghezal M, Marshall BJ, Peek RM, Jr, Boren T, Solnick JV. 2010. Expression of the BabA adhesin during experimental infection with *Helicobacter pylori*. *Infect. Immun.* 78:1593–1600. <http://dx.doi.org/10.1128/IAI.01297-09>.
- Ilver D, Arqvist A, Ogren J, Frick IM, Kersulyte D, Incecik ET, Berg DE, Covacci A, Engstrand L, Boren T. 1998. *Helicobacter pylori* adhesin binding fucosylated histo-blood group antigens revealed by retagging. *Science* 279:373–377. <http://dx.doi.org/10.1126/science.279.5349.373>.
- Menard R, Sansonetti PJ, Parsot C. 1993. Nonpolar mutagenesis of the

- ipa genes defines IpaB, IpaC, and IpaD as effectors of *Shigella flexneri* entry into epithelial cells. *J. Bacteriol.* 175:5899–5906.
29. Wang Y, Roos KP, Taylor DE. 1993. Transformation of *Helicobacter pylori* by chromosomal metronidazole resistance and by a plasmid with a selectable chloramphenicol resistance marker. *J. Gen. Microbiol.* 139:2485–2493. <http://dx.doi.org/10.1099/00221287-139-10-2485>.
 30. Karnholz A, Hoefler C, Odenbreit S, Fischer W, Hofreuter D, Haas R. 2006. Functional and topological characterization of novel components of the comB DNA transformation competence system in *Helicobacter pylori*. *J. Bacteriol.* 188:882–893. <http://dx.doi.org/10.1128/JB.188.3.882-893.2006>.
 31. Park SA, Lee NG. 2013. Global regulation of gene expression in the human gastric pathogen *Helicobacter pylori* in response to aerobic oxygen tension under a high carbon dioxide level. *J. Microbiol. Biotechnol.* 23:451–458. <http://dx.doi.org/10.4014/jmb.1209.09064>.
 32. Donahue JP, Israel DA, Peek RM, Blaser MJ, Miller GG. 2000. Overcoming the restriction barrier to plasmid transformation of *Helicobacter pylori*. *Mol. Microbiol.* 37:1066–1074. <http://dx.doi.org/10.1046/j.1365-2958.2000.02036.x>.
 33. Lin EA, Zhang XS, Levine SM, Gill SR, Falush D, Blaser MJ. 2009. Natural transformation of *Helicobacter pylori* involves the integration of short DNA fragments interrupted by gaps of variable size. *PLoS Pathog.* 5:e1000337. <http://dx.doi.org/10.1371/journal.ppat.1000337>.
 34. Lorenz MG, Wackernagel W. 1994. Bacterial gene transfer by natural genetic transformation in the environment. *Microbiol. Rev.* 58:563–602.
 35. Park AM, Li Q, Nagata K, Tamura T, Shimono K, Sato EF, Inoue M. 2004. Oxygen tension regulates reactive oxygen generation and mutation of *Helicobacter pylori*. *Free Radic. Biol. Med.* 36:1126–1133. <http://dx.doi.org/10.1016/j.freeradbiomed.2004.02.001>.
 36. Wilson DL, Bell JA, Young VB, Wilder SR, Mansfield LS, Linz JE. 2003. Variation of the natural transformation frequency of *Campylobacter jejuni* in liquid shake culture. *Microbiology* 149:3603–3615. <http://dx.doi.org/10.1099/mic.0.26531-0>.
 37. Vegge CS, Brondsted L, Ligowska-Marzeta M, Ingmer H. 2012. Natural transformation of *Campylobacter jejuni* occurs beyond limits of growth. *PLoS One* 7:e45467. <http://dx.doi.org/10.1371/journal.pone.0045467>.
 38. Chen JD, Morrison DA. 1987. Modulation of competence for genetic transformation in *Streptococcus pneumoniae*. *J. Gen. Microbiol.* 133:1959–1967.
 39. Scott DR, Marcus EA, Wen Y, Oh J, Sachs G. 2007. Gene expression in vivo shows that *Helicobacter pylori* colonizes an acidic niche on the gastric surface. *Proc. Natl. Acad. Sci. U. S. A.* 104:7235–7240. <http://dx.doi.org/10.1073/pnas.0702300104>.
 40. Clyne M, Labigne A, Drumm B. 1995. *Helicobacter pylori* requires an acidic environment to survive in the presence of urea. *Infect. Immun.* 63:1669–1673.
 41. Mouery K, Rader BA, Gaynor EC, Guillemin K. 2006. The stringent response is required for *Helicobacter pylori* survival of stationary phase, exposure to acid, and aerobic shock. *J. Bacteriol.* 188:5494–5500. <http://dx.doi.org/10.1128/JB.00366-06>.
 42. Schreiber S, Nguyen TH, Stuben M, Scheid P. 2000. Demonstration of a pH gradient in the gastric gland of the acid-secreting guinea pig mucosa. *Am. J. Physiol. Gastrointest. Liver Physiol.* 279:G597–G604.
 43. Bury-Mone S, Mendz GL, Ball GE, Thibonnier M, Stingl K, Eco-bichon C, Ave P, Huerre M, Labigne A, Thiberge JM, De Reuse H. 2008. Roles of alpha and beta carbonic anhydrases of *Helicobacter pylori* in the urease-dependent response to acidity and in colonization of the murine gastric mucosa. *Infect. Immun.* 76:497–509. <http://dx.doi.org/10.1128/IAI.00993-07>.
 44. Schreiber S, Konradt M, Groll C, Scheid P, Hanauer G, Werling HO, Josenhans C, Suerbaum S. 2004. The spatial orientation of *Helicobacter pylori* in the gastric mucus. *Proc. Natl. Acad. Sci. U. S. A.* 101:5024–5029. <http://dx.doi.org/10.1073/pnas.0308386101>.