

Relationship of Serum Magnesium and Vitamin D Levels in a Nationally-Representative Sample of Iranian Adolescents: The CASPIAN-III Study

Roya Kelishadi, Ehsan Ataei, Gelayol Ardalan¹, Maryam Nazemian, Mohammadhasan Tajadini², Ramin Heshmat³, Mojtaba Keikha, Mohammad Esmaeil Motlagh^{1,4}

Department of Pediatrics, Child Growth and Development Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, ¹Department of School Health, Bureau of Population, Family and School Health, Ministry of Health and Medical Education, Tehran, Iran, ²Department of Biotechnology, School of Pharmacy and Isfahan Pharmaceutical Sciences Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, ³Department of Epidemiology, Chronic Diseases Research Center, Endocrinology and Metabolism Population Sciences Institute, Endocrinology and Metabolism Research Institute, Tehran University of Medical Sciences, Tehran, Iran, ⁴Department of Pediatrics, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Correspondence to:

Prof. Mohammad Esmaeil Motlagh,
Ahvaz Jundishapur University of Medical
Sciences, Ahvaz, Iran.
E-mail: mohammad-motlagh1389@
yahoo.com

Date of Submission: Jul 25, 2013

Date of Acceptance: Oct 21, 2013

How to cite this article: Kelishadi R, Ataei E, Ardalan G, Nazemian M, Tajadini M, Heshmat R, *et al.* Relationship of serum magnesium and vitamin D levels in a nationally-representative sample of Iranian adolescents: The CASPIAN-III study. *Int J Prev Med* 2014;5:99-103.

ABSTRACT

Background: This study aims to assess the relationship of serum Mg and vitamin D levels in a nationally-representative sample of Iranian adolescents.

Methods: The study participants consisted of 330 students, aged range from 10 to 18 years, consisting of an equal number of individuals with and without hypovitaminosis D. The correlation between serum 25 hydroxy vitamin D (25(OH) D) and magnesium (Mg) concentrations was determined.

Results: The mean age of participants was 14.74 ± 2.587 years, without significant difference between those with hypovitaminosis D and those without it. The mean 25(OH) D level was 6.34 ± 1.47 ng/ml in the group with hypovitaminosis D and 39.27 ± 6.42 ng/ml in the group without it. The mean Mg level was 0.80 ± 0.23 mg/dl with lower level in the group with hypovitaminosis D than in others (0.73 ± 0.22 mg/dl vs. 0.87 ± 0.22 mg/dl, respectively) and according to *t*-test analysis, significant lower levels in the deficient group was observed ($P = 0.0001$). The linear regression analysis showed the meaningful relationship between Mg and 25(OH) D serum levels ($P = 0.0001$).

Conclusions: Our study revealed significant associations between serum Mg and 25(OH) D levels. This finding may be of use for further studies on the prevention and management of hypovitaminosis D in children and adolescents. Further longitudinal studies shall evaluate the underlying mechanisms and the clinical significance of the current findings.

Keywords: Adolescents, iran, magnesium, vitamin D

INTRODUCTION

Vitamin D is a lipid soluble vitamin with steroidal structure. It has an extensive range of cellular and molecular functions. It plays key roles in different organs as bone mineralization by maintaining serum calcium and phosphorus concentrations at required levels, cellular differentiation and replication in many body organs, glucose homeostasis, muscle and skeletal function

etc., Adequate intake of this vitamin can prevent the risk of some skeletal as well as non-skeletal disorders as autoimmunity, atopic conditions, some types of cancers, chronic hepatitis C and cardiometabolic diseases.^[1-4] It seems that blood levels of vitamin D during childhood encompass a great value to prevent the abovementioned disorders in adulthood.^[5]

The main source of vitamin D is its synthesis in the skin by sun exposure; however it is well-documented that vitamin D deficiency is prevalent in all regions of the world^[6-9] even in sunny regions, as in the Middle East and North Africa (MENA).^[7]

The population of MENA region has one of the greatest prevalence rates of hypovitaminosis D with lower bone density than in Western countries. Different factors as low dietary intake, lack of enough sun exposure, air pollution, latitude and season can be attributed as the reasons of hypovitaminosis D.^[10-14] However, the effects of some essential elements like magnesium shall be taken into account in this regard.^[1,3,5,15,16]

Magnesium (Mg) as the second most plentiful intracellular cation and an important part of bone mineralization has a crucial function in synthesis and metabolism of vitamin D.^[15,17]

As hypovitaminosis D is prevalent in the pediatric age group of the MENA region, this study aims to assess the relationship of serum Mg and vitamin D levels in a nationally-representative sample of Iranian adolescents.

METHODS

This national population-based study was performed as a sub-study of the “national survey of school student high risk behaviors”, which was conducted as the third survey of the school-based surveillance system entitled Childhood and Adolescence Surveillance and Prevention of Adult Non-communicable disease study. This school-based nationwide health survey was conducted among 5528 students, aged 10-18 years, living in 27 provinces in Iran. Details of data collection and sampling are reported previously^[18] and here it is reported in brief.

Participants

The study protocols were reviewed and approved by ethical committees and other relevant

national regulatory organizations. After complete explanation of the study objectives and protocols for students and their parents, written informed consent and oral assent were obtained from the parents and the students, respectively.^[18] The current survey was done as a sub-study among a randomly selected sample of frozen sera (-70°C) of participants in whom 25-hydroxy vitamin D (25(OH) D) was determined.^[19] Sera of two groups of adolescents with or without vitamin D deficiency were randomly selected. The Research and Ethics Committee of Isfahan University of Medical Sciences, Isfahan, Iran, approved the current study.

Serum concentration of 25(OH) D was analyzed quantitatively by direct competitive immunoassay chemiluminescence method applying LIASON[®] 25 OH vitamin D assay TOTAL (DiaSorin, Inc.), with a coefficient of variation of 9.8%.

25(OH) D level of less than 10 ng/ml was considered as vitamin D deficiency and levels between 10 and 30 ng/ml as vitamin D insufficiency.^[20]

Magnesium level was determined by atomic absorption spectrophotometer by using hollow cathode lamps of Mg.

The relationship between serum vitamin D and Mg levels was assessed through linear regression analysis purposing Mg as independent variable and 25(OH) D as dependent. Mean Mg serum level in deficient and normal group was compared by *t*-test analysis. Data were analyzed by SPSS statistical software (version 18.0; SPSS Inc., Chicago, IL, USA); the significance level was set at $P < 0.05$.

RESULTS

The study participants consisted of 330 students consisting of an equal number of participants with and without hypovitaminosis D. The mean age of participants was 14.74 ± 2.587 years, without significant difference between those with hypovitaminosis D and those without it (14.75 ± 2.52 vs. 14.74 ± 2.66 years, respectively, $P > 0.05$). Baseline characteristics of participants are shown in Table 1. The mean 25(OH) D level was 6.34 ± 1.47 ng/ml in the group with hypovitaminosis D and 39.27 ± 6.42 ng/ml in the group without it. The mean Mg level was 0.80 ± 0.23 mg/dl with lower level in the group with hypovitaminosis D than in others

Table 1: Baseline characteristics of participants

Parameters	Normal group	Deficient group
Age	14.74±2.66	14.74±2.52
BMI	19.58±4.56	19.64±5.41
SBP	102.33±12.55	102.55±14.47
DBP	63.87±9.52	65.37±9.90
HDL-C	48.55±14.95	48.69±14.10
LDL-C	82.34±26.49	80.81±29.47
Total cholesterol	148.19±32.22	147.15±30.685
TG	97.46±44.81	95.07±40.20
FBG	87.85±12.26	87.93±12.09
AST	24.51±9.48	23.49±10.39
ALT	18.21±14.10	17.82±9.20

All variables are reported in mean±SD. BMI=Body mass index, SBP=Systolic blood pressure, DBP=Diastolic blood pressure, HDL-C=High density lipoprotein cholesterol, LDL-C=Low density lipoprotein cholesterol, TG=Triglyceride, FBG=Fasting blood glucose, AST=Aspartate aminotransaminase, ALT=Alanine aminotransaminase, SD=Standard deviation

(0.73 ± 0.22 mg/dl vs. 0.87 ± 0.22 mg/dl, respectively), with significant lower levels in the deficient group according to *t*-test analysis ($P = 0.0001$). The linear regression analysis showed that as one unit increase in Mg level, 25(OH) D level raises 0.276, which shows a significant relationship ($P = 0.0001$).

DISCUSSION

This study, which to the best of our knowledge is the first of its kind, investigated the association of serum 25(OH) D levels with serum Mg in a large nationally-representative sample of the pediatric population. We found that serum 25(OH) D and Mg concentrations had a positive significant correlation.

It is well-documented that hypovitaminosis D is highly prevalent in children and adolescents and shall be considered a public health priority in many parts of the world.^[9,21-30] Vitamin D deficiency is important from different aspects due to key roles of this vitamin in different organs and also mechanisms in body tissues.

Different studies revealed the skeletal^[21,29,30] and extra skeletal health effects of this vitamin.^[5,8,31] In a triple masked clinical controlled trial among 50 obese children aged 10-16 years, vitamin D supplementation was effective on reducing insulin resistance and cardiometabolic risk factors.^[32] Mg,

as the second intracellular positive cation may effect on this process. Deng *et al.* in their study surveyed the relationship between Mg and vitamin D status and explored Mg intake alone or its interaction with vitamin D intake may involve in vitamin D status.^[15] Furthermore, Vojtková *et al.* found that deficient vitamin D type 1 diabetic children have lower Mg level in their blood levels.^[20] This relationship may be the result of the central role of that in mineral homeostasis, regulating parathormone secretion and vitamin D action and activation.^[17,33] Sahota *et al.* found that Mg is a crucial contributing factor in patients with established osteoporosis with vitamin D deficiency and blunted parathormone level.^[34]

Study limitations and strengths

The main limitation of this study was its cross-sectional nature, so the associations of diverse variables should be considered with caution. The study strengths are its novelty in the pediatric age group and using data of a nationally representative group of adolescents, which would increase the generalizability of the study findings.

CONCLUSIONS

Our study revealed significant associations between serum Mg and 25(OH) D levels. This finding may be of use for interventional actions to prevent and manage the highly prevalent problem of hypovitaminosis D in children and adolescents. Further longitudinal studies are required to find the underlying mechanisms and the clinical significance of this association.

REFERENCES

1. Nucci AM, Russell CS, Luo R, Ganji V, Olabopo F, Barbara H, *et al.* The effectiveness of a short food frequency questionnaire in determining vitamin D intake in children. *Dermatoendocrinol* 2013;5:205-10.
2. Ceglia L. Vitamin D and its role in skeletal muscle. *Curr Opin Clin Nutr Metab Care* 2009;12:628-33.
3. Cosenza L, Pezzella V, Nocerino R, Di Costanzo M, Coruzzo A, Passariello A, *et al.* Calcium and vitamin D intakes in children: A randomized controlled trial. *BMC Pediatr* 2013;13:86.
4. Torun E, Gönüllü E, Ozgen IT, Cindemir E, Oktem F. Vitamin D deficiency and insufficiency in obese children and adolescents and its relationship with insulin resistance. *Int J Endocrinol* 2013;2013:631845.

5. Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M, Drug and Therapeutics Committee of the Lawson Wilkins Pediatric Endocrine Society. Vitamin D deficiency in children and its management: Review of current knowledge and recommendations. *Pediatrics* 2008;122:398-417.
6. Andıran N, Çelik N, Akça H, Doğan G. Vitamin D deficiency in children and adolescents. *J Clin Res Pediatr Endocrinol* 2012;4:25-9.
7. Bassil D, Rahme M, Hoteit M, Fuleihan Gel-H. Hypovitaminosis D in the Middle East and North Africa: Prevalence, risk factors and impact on outcomes. *Dermatoendocrinol* 2013;5:274-98.
8. Olson ML, Maalouf NM, Oden JD, White PC, Hutchison MR. Vitamin D deficiency in obese children and its relationship to glucose homeostasis. *J Clin Endocrinol Metab* 2012;97:279-85.
9. Zhu Z, Zhan J, Shao J, Chen W, Chen L, Li W, *et al.* High prevalence of vitamin D deficiency among children aged 1 month to 16 years in Hangzhou, China. *BMC Public Health* 2012;12:126.
10. Kelishadi R, Sharifi-Ghazvini F, Poursafa P, Mehrabian F, Farajian S, Yousefy H, *et al.* Determinants of hypovitaminosis d in pregnant women and their newborns in a sunny region. *Int J Endocrinol* 2013;2013:460970.
11. Kelishadi R, Moeini R, Poursafa P, Farajian S, Yousefy H, Okhovat-Souraki AA. Independent association between air pollutants and vitamin D deficiency in young children in Isfahan, Iran. *Paediatr Int Child Health* 2013; [Epub ahead of print].
12. Ardestani PM, Salek M, Keshteli AH, Nejadnik H, Amini M, Hosseini SM, *et al.* Vitamin D status of 6-to 7-year-old children living in Isfahan, Iran. *Endokrynol Pol* 2010;61:377-82.
13. Salek M, Hashemipour M, Aminorroaya A, Gheiratmand A, Kelishadi R, Ardestani PM, *et al.* Vitamin D deficiency among pregnant women and their newborns in Isfahan, Iran. *Exp Clin Endocrinol Diabetes* 2008;116:352-6.
14. Maalouf G, Gannagé-Yared MH, Ezzedine J, Larijani B, Badawi S, Rached A, *et al.* Middle East and North Africa consensus on osteoporosis. *J Musculoskelet Neuronal Interact* 2007;7:131-43.
15. Deng X, Song Y, Manson JE, Signorello LB, Zhang SM, Shrubsole MJ, *et al.* Magnesium, vitamin D status and mortality: Results from US National Health and Nutrition Examination Survey (NHANES) 2001 to 2006 and NHANES III. *BMC Med* 2013;11:187.
16. Wagner CL, Greer FR, American Academy of Pediatrics Section on Breastfeeding, American Academy of Pediatrics Committee on Nutrition. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics* 2008;122:1142-52.
17. Carpenter TO, DeLucia MC, Zhang JH, Bejnerowicz G, Tartamella L, Dziura J, *et al.* A randomized controlled study of effects of dietary magnesium oxide supplementation on bone mineral content in healthy girls. *J Clin Endocrinol Metab* 2006;91:4866-72.
18. Kelishadi R, Heshmat R, Motlagh ME, Majdzadeh R, Keramatian K, Qorbani M, *et al.* Methodology and early findings of the third survey of CASPIAN study: A national school-based surveillance of students' High Risk Behaviors. *Int J Prev Med* 2012;3:394-401.
19. Kelishadi R, Ardalan G, Motlagh ME, Shariatinejad K, Heshmat R, Poursafa P, *et al.* National report on the association of serum vitamin D with cardiometabolic risk factors in the pediatric population of the Middle East and North Africa (MENA): The CASPIAN-III Study 2014;30:33-8.
20. Vojtková J, Ciljaková M, Vojarová L, Janíková K, Michnová Z, Sagiová V. Hypovitaminosis D in children with type 1 diabetes mellitus and its influence on biochemical and densitometric parameters. *Acta Medica (Hradec Kralove)* 2012;55:18-22.
21. Cheng S, Tylavsky F, Kröger H, Kärkkäinen M, Lyytikäinen A, Koistinen A, *et al.* Association of low 25-hydroxyvitamin D concentrations with elevated parathyroid hormone concentrations and low cortical bone density in early pubertal and prepubertal Finnish girls. *Am J Clin Nutr* 2003;78:485-92.
22. El-Hajj Fuleihan G, Nabulsi M, Choucair M, Salamoun M, Hajj Shahine C, Kizirian A, *et al.* Hypovitaminosis D in healthy schoolchildren. *Pediatrics* 2001;107:E53.
23. Looker AC, Dawson-Hughes B, Calvo MS, Gunter EW, Sahyoun NR. Serum 25-hydroxyvitamin D status of adolescents and adults in two seasonal subpopulations from NHANES III. *Bone* 2002;30:771-7.
24. Munns C, Zacharin MR, Rodda CP, Batch JA, Morley R, Cranswick NE, *et al.* Prevention and treatment of infant and childhood vitamin D deficiency in Australia and New Zealand: A consensus statement. *Med J Aust* 2006;185:268-72.
25. Rabbani A, Alavian SM, Motlagh ME, Ashtiani MT, Ardalan G, Salavati A, *et al.* Vitamin D insufficiency among children and adolescents living in Tehran, Iran. *J Trop Pediatr* 2009;55:189-91.
26. Rockell JE, Green TJ, Skeaff CM, Whiting SJ, Taylor RW, Williams SM, *et al.* Season and ethnicity are determinants of serum 25-hydroxyvitamin D concentrations in New Zealand children aged 5-14 y. *J Nutr* 2005;135:2602-8.
27. Tylavsky FA, Cheng S, Lyytikäinen A, Viljakainen H, Lamberg-Allardt C. Strategies to improve vitamin D status in northern European children: Exploring the merits of vitamin D fortification and supplementation. *J Nutr* 2006;136:1130-4.

28. Zamora SA, Rizzoli R, Belli DC, Slosman DO, Bonjour JP. Vitamin D supplementation during infancy is associated with higher bone mineral mass in prepubertal girls. *J Clin Endocrinol Metab* 1999;84:4541-4.
29. Ardawi MS, Sibiany AM, Bakhsh TM, Qari MH, Maimani AA. High prevalence of vitamin D deficiency among healthy Saudi Arabian men: Relationship to bone mineral density, parathyroid hormone, bone turnover markers, and lifestyle factors. *Osteoporos Int* 2012;23:675-86.
30. Schnabel C, Jett K, Friedman JM, Frieling I, Kruse HP, Mautner V. Effect of vitamin D3 treatment on bone density in neurofibromatosis 1 patients: A retrospective clinical study. *Joint Bone Spine* 2013;80:315-9.
31. Polidoro L, Properzi G, Marampon F, Gravina GL, Festuccia C, Di Cesare E, *et al.* Vitamin D protects human endothelial cells from H₂O₂ oxidant injury through the Mek/Erk-Sirt1 axis activation. *J Cardiovasc Transl Res* 2013;6:221-31.
32. Kelishadi R, Saleka S, Saleka M, Hashemipoura M, Movahedian M. Effects of vitamin D supplementation on insulin resistance and cardiometabolic risk factors in children with metabolic syndrome: A triple-masked controlled trial. *J Pediatr (Rio J)* 2013; [Epub ahead of print].
33. Zofková I, Kancheva RL. The relationship between magnesium and calciotropic hormones. *Magnes Res* 1995;8:77-84.
34. Sahota O, Munday MK, San P, Godber IM, Hosking DJ. Vitamin D insufficiency and the blunted PTH response in established osteoporosis: The role of magnesium deficiency. *Osteoporos Int* 2006;17:1013-21.

Source of Support: This study was conducted as a thesis funded by Isfahan University of Medical Sciences, Isfahan, Iran,
Conflict of Interest: None declared.