

Published in final edited form as:

Science. 2010 September 3; 329(5996): 1201–1205. doi:10.1126/science.1191241.

Signaling Kinase AMPK Activates Stress-Promoted Transcription via Histone H2B Phosphorylation

David Bungard¹, Benjamin J. Fuerth^{2,3}, Ping-Yao Zeng^{1,4}, Brandon Faubert^{2,3}, Nancy L. Maas¹, Benoit Viollet^{5,6}, David Carling⁷, Craig B. Thompson⁸, Russell G. Jones^{2,3,8,*}, and Shelley L. Berger^{1,9,10,*}

¹Department of Cellular and Developmental Biology, University of Pennsylvania Medical School, Philadelphia, PA 19104, USA

²Rosalind and Morris Goodman Cancer Research Centre, McGill University, Montreal, Quebec H3G 1Y6, Canada

³Department of Physiology, McGill University, Montreal, Quebec H3G 1Y6, Canada

⁴Institutes of Biomedical Sciences Epigenetics Program, Mingdao Building, Room 511, Fudan University, Mail Box 281, 138 Yixue Yuan Road, Shanghai 200032, P.R. China

⁵Institut Cochin, Université Paris Descartes, CNRS (UMR 8104), 75014 Paris, France

⁶INSERM U1016, 75014 Paris, France

⁷Cellular Stress Group, MRC Clinical Sciences Centre, Imperial College, Hammersmith Hospital, London W12 0NN, UK

⁸Abramson Cancer Center and Abramson Family Cancer Research Institute, University of Pennsylvania, Philadelphia, PA 19104, USA

⁹Department of Genetics, University of Pennsylvania Medical School, Philadelphia, PA 19104, USA

¹⁰Department of Biology, School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA 19104, USA

Abstract

The mammalian adenosine monophosphate-activated protein kinase (AMPK) is a serine-threonine kinase protein complex that is a central regulator of cellular energy homeostasis. However, the mechanisms by which AMPK mediates cellular responses to metabolic stress remain unclear. We found that AMPK activates transcription through direct association with chromatin and phosphorylation of histone H2B at serine 36. AMPK recruitment and H2B Ser³⁶ phosphorylation colocalized within genes activated by AMPK-dependent pathways, both in promoters and in transcribed regions. Ectopic expression of H2B in which Ser³⁶ was substituted by alanine reduced transcription and RNA polymerase II association to AMPK-dependent genes, and lowered cell survival in response to stress. Our results place AMPK-dependent H2B Ser³⁶ phosphorylation in a direct transcriptional and chromatin regulatory pathway leading to cellular adaptation to stress.

Signaling pathways often involve cascades of protein phosphorylation that terminate in regulation of nuclear transcription. However, the central kinases in these pathways are

Copyright 2010 by the American Association for the Advancement of Science; all rights reserved

*To whom correspondence should be addressed. russell.jones@mcgill.ca (R.G.J.); bergers@mail.med.upenn.edu (S.L.B.).

Supporting Online Material www.sciencemag.org/cgi/content/full/science.1191241/DC1

generally not thought to directly modulate transcription. One such signaling kinase, adenosine monophosphate-activated protein kinase (AMPK) (1), is activated by high substrate adenosine monophosphate levels under conditions of energetic stress, such as nutrient deprivation or hypoxia. AMPK activation initiates a program of metabolic adaptation to preserve cellular energy (adenosine triphosphate conservation) and maintain cellular viability (2).

We investigated possible direct transcriptional mechanisms in the AMPK pathway. First, we examined whether AMPK broadly responds to cellular stress. Mouse embryonic fibroblasts (MEFs) were stressed by ultraviolet (UV) irradiation, γ irradiation, the topoisomerase inhibitor camptothecin, the DNA intercalating agent adriamycin, the alkylating agent *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine, or hydrogen peroxide. AMPK was activated [Thr¹⁷² phosphorylation in the AMPK activation loop and phosphorylation of substrate ACC α (acetyl coenzyme A-carboxylase- α)] by all agents (Fig. 1A). Liver kinase B1 (LKB1) is one of three mammalian upstream AMPK kinases and activates AMPK upon bioenergetic stress (3). LKB1 also activated AMPK in response to either genotoxic (UV irradiation) or metabolic stress [the glycolytic inhibitor 2-deoxyglucose (2-DG)] (fig. S1).

We next used wild-type or AMPK α 1- and AMPK α 2-deficient MEFs (*ampk α ^{-/-}*) to examine whether AMPK influences cellular survival during stress (4). The *ampk α ^{-/-}* MEFs displayed decreased survival in the presence of metabolic stressors (Fig. 1B and fig. S2), UV irradiation (Fig. 1B), or hydrogen peroxide (fig. S2), indicating that AMPK promotes cell survival in response to various metabolic and genotoxic stresses.

Our previous work implicated AMPK and LKB1 in tumor suppressor p53-dependent metabolic and genotoxic stress responses (2, 5). We examined the involvement of AMPK in transcriptional regulation of p53-responsive genes. *ampk α ^{-/-}* MEFs showed decreased induction of *p21*, a well-characterized p53 target gene, in response to both glucose withdrawal and UV irradiation (Fig. 1C and fig. S3). The reduction in glucose-regulated gene expression was similar in magnitude to knockdown of p53 (Fig. 1C). Other p53-dependent genes, including *reprimo*, *cyclinG*, and *cpt1c*, showed reduced expression in *ampk α ^{-/-}* cells deprived of glucose, whereas expression of a control gene, *gapdh*, was unaffected (Fig. 1C and fig. S4). A similar defect in stress-dependent transcription was observed in human cancer cells (HCT116 colon carcinoma) (fig. S5). The transcriptional defect observed in UV-treated cells occurred downstream of the DNA-damage checkpoint, as phosphorylation of Chk1, Rad17, and p53 in both *lkb1^{-/-}* and *ampk α ^{-/-}* cell lines was similar to that of wild-type controls (fig. S6). Thus, in cell culture, the LKB1-AMPK pathway responds to a wide variety of metabolic and genotoxic stresses and is required for maximal stress-induced transcription of multiple p53-dependent genes that promote survival.

These and previous results (6) implicate AMPK in gene activation, although the underlying mechanisms have remained elusive. We used chromatin immunoprecipitation (ChIP) and quantitative polymerase chain reaction (qPCR) to investigate whether AMPK plays a direct role at p53-regulated genes in MEFs. We found that LKB1 coimmunoprecipitated with AMPK in response to UV irradiation, and this interaction was dependent on AMPK kinase activity (fig. S7). ChIP showed ectopic myc-AMPK associated with the *cpt1c* and *p21* gene promoters in response to glucose withdrawal and UV treatment (Fig. 1D and fig. S8). Ectopic kinase-deficient myc-AMPK α or FLAG-tagged LKB1 abrogated the binding of AMPK to chromatin (Fig. 1D and fig. S8). FLAG-LKB1 similarly associated with the *p21* promoter (fig. S9). Binding of AMPK or LKB1 to promoters in unstressed cells was low (Fig. 1D and figs. S8 and S9). AMPK was also detected at the *p21* promoter in human HCT116 cells after UV treatment (fig. S10). This AMPK activation appears to be specific

for survival in response to stress, because AMPK was not detected at the PUMA promoter, a p53-dependent, pro-apoptotic gene (fig. S11). The immunoprecipitation and ChIP results suggest a functional, kinase-dependent interaction and promoter association of LKB1 and AMPK.

ChIP of endogenous AMPK, using an AMPK α 2 antibody that immunoprecipitated the protein (fig. S12), showed increased AMPK α 2 binding at p53 sites upstream of *cpt1c* in cells treated with 2-DG and UV light (Fig. 1E; at the *p21* promoter, fig. S13). The binding was reduced in both *ampk α ^{-/-}* cells (Fig. 1F) and *p53^{-/-}* cells (Fig. 1G). Thus, in response to cellular stress, LKB1 and AMPK physically localize to chromatin in an interdependent and p53-dependent fashion to regulate gene transcription.

Posttranslational modifications of core histones, including phosphorylation, function to regulate transcription (7, 8). To examine histones as substrates of AMPK, we epitope-tagged the predominantly nuclear isoform AMPK α 2 (9) and immunoprecipitated wild-type AMPK α 2 or a catalytic mutant (Asp¹⁵⁷ → Ala) from transfected *ampk α ^{-/-}* MEFs after low glucose treatment to activate AMPK. The wild-type AMPK α 2 complex specifically phosphorylated histone H2B above the nonspecific background activity of all four histones observed in the catalytic mutant and mock immunoprecipitations (Fig. 2A). AMPK α 2 from untreated cells exhibited low activity on H2B relative to AMPK α 2 from stressed cells (Fig. 2B and fig. S15). H2A was also phosphorylated in these assays; however, phospho-H2A was detected in the mock immunoprecipitation and did not respond to stress, which suggested that the activity is nonspecific. These data indicate that activated AMPK α 2 specifically phosphorylates H2B.

Analysis of histone H2B revealed a potential AMPK target motif (10, 11) at Ser³⁶ in the N terminus, similar to the AMPK substrates endothelial nitric oxide synthase (eNOS) and phosphofructokinase 2 (PFK2) (H2B, Arg-Ser-Arg-Lys-Glu-Ser; eNOS, Arg-Ile-Arg-Thr-Gln-Ser; PFK2, Arg-Met-Arg-Arg-Asn-Ser), with conserved Arg residues at P-5 and P-3 relative to the phosphoacceptor Ser³⁶ (fig. S14) (12). We therefore compared peptides corresponding to residues 1 to 20 or 21 to 42 of H2B; only the H2B 21–42 peptide was phosphorylated by immunoprecipitated AMPK α 2. Moreover, an H2B 21–42 peptide harboring a Ser³⁶ → Ala (S36A) substitution was not phosphorylated, whereas S32A and S38A peptides were phosphorylated (Fig. 2C). In addition, myc-AMPK α 2 phosphorylated the H2B 21–42 peptide more efficiently than did SAMS (His-Met-Arg-Ser-Ala-Met-Ser-Gly-Leu-His-Leu-Val-Lys-Arg-Arg) peptide, a canonical AMPK target sequence (Fig. 2D) (13). Thus, H2B Ser³⁶ is a robust *in vitro* AMPK substrate.

We next examined whether AMPK is involved in the regulation of H2B Ser³⁶ phosphorylation *in vivo*. We generated an antibody against an H2B pS36 peptide and extensively characterized the affinity-purified antibody *in vitro* and *in vivo* (fig. S16). H2B pS36 occurs in cells treated with 2-DG, which induced metabolic stress shown by increased phosphorylation of AMPK α and ACC α (Fig. 2E), whereas other histone marks, including H3, K9me3 H3, and pS10 H3, did not significantly change (fig. S17). In *ampk α ^{-/-}* MEFs, H2B pS36 did not increase after 2-DG treatment (Fig. 2E). We detected similar induction of H2B pS36 in cells with two well-characterized AMPK activators, aminoimidazole carboxamide ribonucleotide (AICAR) and phenformin (Fig. 2F). In addition, H2B pS36 was not detected in *lkb1^{-/-}* MEFs but increased after reintroduction of LKB1, and 2-DG stimulated H2B pS36 only when LKB1 was present (fig. S18). These results suggest that H2B Ser³⁶ phosphorylation is downstream of the LKB1-AMPK signaling pathway *in vivo* in response to a broad range of metabolic stresses, and that AMPK is likely the direct kinase in this pathway. This global increase of H2B pS36 may not be limited to p53 target genes (Fig. 1G), because H2B pS36 levels were not reduced in *p53^{-/-}* cells (fig. S19).

We found that transfected myc-AMPK and FLAG-H2B associate in 293T cells in response to the AMPK agonist AICAR (Fig. 3A). This interaction was ablated by an H2B S36A mutation (Fig. 3A), suggesting that Ser³⁶ is critical for the interaction of AMPK and H2B. We next performed ChIP experiments in wild type and *ampkα*^{-/-} MEFs to determine whether H2B pS36 is associated with AMPK-dependent genes, specifically at the p53 binding site in the *cpt1c* promoter. The H2B pS36 ChIP signal increased in response to both 2-DG and UV treatment, and the ChIP signal was reduced to background levels in AMPKα-deficient cells (Fig. 3B).

To compare the locations of endogenous AMPK and H2B pS36, we performed ChIP at *cpt1c* under normal and glucose-free conditions, using primers along the promoter, transcribed region, and upstream or downstream nontranscribed regions (Fig. 3C, primer sets 1 to 7). Upon stress, an increase in H2B pS36 occurred at the p53 binding site (p53bs, primer set 2), a larger increase at the transcription start site (TSS, primer set 4), and only background levels upstream (primer set 1) and between (primer set 3) these sites, and downstream of the gene (primer set 7) (Fig. 3C). Notably, H2B pS36 associated throughout the transcribed region of *cpt1c* (primer sets 5 and 6) (Fig. 3C). We then used the AMPKα2 antibody and discovered a positive correlation between locations of endogenous AMPK and H2B pS36, again with localization throughout the transcribed region (Fig. 3C). The *p21* gene showed a similar positive correlation between H2B pS36 and endogenous AMPK localization at the p53 binding sites and along the transcribed region, but not around the gene and only in response to low glucose (Fig. 3D). As a control, we tested the *gapdh* gene, which showed no difference in ChIP signal between the stressed and unstressed cells (fig. S20). Similar results were observed at the *cpt1c* gene in response to UV treatment (fig. S21). Ectopic myc-AMPK also localized along the *p21* transcribed gene in response to low glucose and required AMPK catalytic activity (fig. S22). This correspondence in the location of AMPK and H2B pS36 suggests that the role of AMPK in transcriptional regulation is linked to the phosphorylation of H2B along transcribed regions of target genes.

We generated clonal MEF cell lines with stable ectopic expression of FLAG-H2B or mutant FLAG-H2B S36A, and, as shown above, only FLAG-H2B was detected by the H2B pS36 antibody (fig. S16). The level of FLAG-tagged wild-type or mutant histone H2B was lower than that of endogenous H2B as measured by Western blot (fig. S23); however, both ectopic wild-type and H2B S36A similarly incorporated into chromatin, as determined by ChIP and qPCR at the promoter and transcribed region of the *cpt1c* gene (fig. S24) and by coimmunoprecipitation of endogenous H3 and H2B with FLAG-H2B as shown by Western blot (fig. S25). We isolated several equivalently expressing FLAG-H2B- and FLAG-H2B S36A-expressing clonal lines (fig. S26). The cell lines treated with 2-DG showed comparable levels of AMPK and induction of AMPK Thr¹⁷² and pACCα Ser⁷⁹ phosphorylation (fig. S27), suggesting that signaling to AMPK is unaffected by the S36A mutation. In response to glucose withdrawal, elevated transcription of the AMPK target genes *p21*, *cpt1c*, *reprimin*, and *cyclinG* was detected in wild-type FLAG-H2B-expressing cells; however, FLAG-H2B S36A-expressing cells showed significantly reduced induction of these genes (Fig. 4A). Expression of the control genes *gapdh* and *hadh* was unaffected by expression of the S36A mutant (fig. S28). Thus, although expression of FLAG-H2B is low relative to endogenous H2B (figs. S23 and S25), we observed a significant and specific transcriptional defect in the cells ectopically expressing S36A mutant H2B.

We next tested the viability of these cells in response to various metabolic stresses including glucose withdrawal, 2-DG, and phenformin. The level of cell death induced by these stresses was significantly greater in the presence of ectopic H2B S36A, relative to cells expressing ectopic wild-type H2B (Fig. 4B).

Finally, we examined whether RNA polymerase II (Pol II) occupancy at genes is altered by ectopic expression of H2B S36A. ChIP analysis showed that Pol II was present through the *cpt1c* gene, and Pol II increased in glucose withdrawal (Fig. 4C). In FLAG–H2B S36A cells, the Pol II ChIP signal was reduced along the transcribed region of the *cpt1c* gene specifically under conditions of glucose withdrawal (Fig. 4C). These data point to AMPK-mediated phosphorylation of H2B at Ser³⁶ as a critical transcriptional regulatory step in the cellular adaptation response to stress, to optimize cell survival.

Our results show that AMPK directly associates with chromatin to regulate transcriptional programs required to survive a wide variety of metabolic and environmental stresses. Mammalian AMPK is a histone kinase that preferentially phosphorylates histone H2B at Ser³⁶. Phosphorylation of this site in vivo is acutely dependent on the LKB1-AMPK energy-sensing pathway, and through their direct action at genes.

AMPK and H2B pS36 appear to regulate transcription by direct interaction between the kinase and the H2B substrate. AMPK appears to be recruited to specific promoters by transcription factors such as p53. AMPK may similarly influence other stress-induced transcriptional activators, including Foxo3a, peroxisome proliferator-activated receptor γ coactivator 1 α (PGC-1 α), and transducer of regulated CREB activity 2 (TORC2) (2, 14–16). AMPK colocalizes with and phosphorylates H2B at p53 binding sites and along the transcribed *cpt1c* and *p21* genes. This pattern may be due to activator recruitment of the enzyme, followed by enzyme transit across transcribed regions, as seen for the SAGA-Gcn5-Ubp8 complex and in the acetylation of H3 with deubiquitination of H2B (17, 18). Moreover, histone methylases associate with RNA Pol II to modify H3 across transcribed regions (19). Our findings represent evidence for histone phosphorylation across transcribed regions, which suggests that AMPK, via H2B Ser³⁶ phosphorylation, may play a role in transcriptional elongation.

Ectopic expression of the H2B S36A mutant, even in the presence of higher levels of endogenous wild-type H2B, significantly reduces transcription of AMPK target genes and stress survival. Hence, the modification may function processively to facilitate transcription elongation. Indeed, H2B Ser³⁶ phosphorylation promotes RNA Pol II association to transcribed regions. Ectopic expression of mutant histone genes may be of broad utility to unravel functions of other histone modifications in mammalian cells.

As part of the central energy-sensing pathway in eukaryotes, AMPK functions to maintain cellular energy homeostasis in the face of metabolic perturbations. Our data suggest that modification of H2B by AMPK may be a general stress-response pathway to tune specific transcriptional responses, regulate cellular metabolism, and promote cell survival.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by NIH grants CA078831 (S.L.B.) and CA105463 (C.B.T.), NIH Training Program in Basic Cancer Research grant CA09171 (D.B.), Canadian Institutes of Health Research grant MOP-93799 (R.G.J.), the McGill Integrated Cancer Research Training Program (B.F.), and the Leukemia and Lymphoma Society, NIH, National Cancer Institute, and American Association for Cancer Research (AACR)/Stand Up to Cancer (C.B.T.). C.B.T. reports the filing of patent applications exploiting the use of AMPK activators in cancer therapy. He is also a member of the board of directors of Merck, the scientific advisory board of Agios Pharmaceuticals, and the medical advisory board of the Howard Hughes Medical Institute.

References and Notes

1. Hardie DG. *Nat. Rev. Mol. Cell Biol.* 2007; 8:774. [PubMed: 17712357]
2. Jones RG, et al. *Mol. Cell.* 2005; 18:283. [PubMed: 15866171]
3. Hardie DG. *Curr. Opin. Cell Biol.* 2005; 17:167. [PubMed: 15780593]
4. Laderoute KR, et al. *Mol. Cell. Biol.* 2006; 26:5336. [PubMed: 16809770]
5. Zeng PY, Berger SL. *Cancer Res.* 2006; 66:10701. [PubMed: 17108107]
6. McGee SL, Hargreaves M. *Front. Biosci.* 2008; 13:3022. [PubMed: 17981775]
7. Berger SL. *Nature.* 2007; 447:407. [PubMed: 17522673]
8. Ivaldi MS, Karam CS, Corces VG. *Genes Dev.* 2007; 21:2818. [PubMed: 17942706]
9. Salt CJ, et al. *Biochem. J.* 1998; 334:177. [PubMed: 9693118]
10. Gwinn DM, et al. *Mol. Cell.* 2008; 30:214. [PubMed: 18439900]
11. Weekes J, Ball KL, Caudwell FB, Hardie DG. *FEBS Lett.* 1993; 334:335. [PubMed: 7902296]
12. Towler MC, Hardie DG. *Circ. Res.* 2007; 100:328. [PubMed: 17307971]
13. Hardie DG, Salt IP, Davies SP. *Methods Mol. Biol.* 2000; 99:63. [PubMed: 10909077]
14. Shaw RJ, et al. *Science.* 2005; 310:1642. [PubMed: 16308421]
15. Greer EL, et al. *J. Biol. Chem.* 2007; 282:30107. [PubMed: 17711846]
16. Jäger S, Handschin C, -Pierre J, St. Spiegelman BM. *Proc. Natl. Acad. Sci. U.S.A.* 2007; 104:12017. [PubMed: 17609368]
17. Govind CK, Zhang F, Qiu H, Hofmeyer K, Hinnebusch AG. *Mol. Cell.* 2007; 25:31. [PubMed: 17218269]
18. Wyce A, et al. *Mol. Cell.* 2007; 27:275. [PubMed: 17643376]
19. Shilatifard A. *Annu. Rev. Biochem.* 2006; 75:243. [PubMed: 16756492]




Fig. 1. AMPK is required for stress-dependent transcription and localizes to stress-responsive genes. **(A)** Western blot of AMPK activation (AMPK α pThr¹⁷²) and ACC α phosphorylation (ACC α pSer⁷⁹) in MEFs in response to UV light, γ irradiation (IR), camptothecin (CPT), adriamycin (Adr), *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG), or hydrogen peroxide (H₂O₂). NT, no treatment; Glc, glucose. **(B)** Viability of wild-type (WT) and *ampk α ^{-/-}* MEFs in response to indicated stresses. Left panel: glucose (Glc) withdrawal; right panel, UV irradiation (UVC). **(C)** Expression (qPCR) of *p21*, *cpt1c*, and *gapdh* mRNA in wild-type, AMPK α -deficient, or p53 RNAi-expressing MEFs after glucose withdrawal, relative to untreated cells. **(D)** ChIP of WT or dominant negative (MUT) myc-AMPK after glucose withdrawal in *lkb1^{-/-}* MEFs cotransfected with either WT or catalytically inactive (MUT) FLAG-LKB1, both able to be immunoprecipitated [see (A), fig. S7, and (5)]. Data represent means \pm SEM for *n* = 3. **(E to G)** Endogenous AMPK α 2 ChIP in untreated MEFs (-), 2-DG (15 min), or UV (6 hours) (E); WT or *ampk α ^{-/-}* MEFs (F); and WT or *p53^{-/-}* MEFs (G). Data represent means \pm SEM for *n* = 3. #*P* < 0.06, **P* < 0.05, ***P* < 0.01, ****P* < 0.03.




Fig. 2. H2B is an AMPK target. (A to D) In vitro phosphorylation by myc-AMPK immunoprecipitated from *ampka*^{-/-} MEFs of recombinant human histones by beads alone, WT myc-AMPK, or catalytic mutant (D157A) myc-AMPK (right: myc Western blot) (A); recombinant human histones using myc-AMPK from glucose-treated cells (+glucose) or untreated cells (-glucose) (B); H2B peptides (C); and H2B 21–42 and SAMS peptide (D). (E) Western blots of wild-type or *ampka*^{-/-} MEFs treated with 2-DG (25 mM for 10 min) or untreated. (F) Western blots of MEFs treated for 1 hour with AICAR (AIC, 2 mM) or phenformin (Ph, 3 mM) or untreated.




fig. 3. AMPK and H2B pS36 are localized to chromatin in response to stress. **(A)** Western blots showing myc-AMPK immunoprecipitation from WT H2B- or H2B S36A-expressing cells after AICAR treatment (1 mM, 24 hours). WCL, whole-cell lysate. **(B)** ChIP in WT or *ampkα^{-/-}* MEFs. Data represent means ± SEM for *n* = 3. **(C and D)** ChIP along *cpt1c* gene (C) and *p21* gene (D) with glucose (white bars) or without glucose (black bars). Data represent means ± SEM for *n* = 3. **P* < 0.01, ***P* < 0.05.




Fig. 4. H2B Ser³⁶ is essential for transcription and survival in response to metabolic stress. Data for MEF cell lines expressing wild-type H2B or H2B S36A are shown. **(A)** Relative expression (qPCR) of indicated genes after glucose withdrawal in individual MEF clones (H2B, clones 5 and 6; H2B S36A, clones 4 and 6). **(B)** Viability of H2B (open squares) or H2B S36A (solid squares) MEFs in no glucose, 2-DG (10 mM), or phenformin (3 mM). **(C)** ChIP of RNA Pol II across the *cpt1c* gene. Data represent means \pm SEM for $n = 3$. * $P < 0.05$, ** $P < 0.01$, # $P < 0.03$, ## $P < 0.08$.