

Published in final edited form as:

Nat Cell Biol. 2005 November ; 7(11): 1066–1073. doi:10.1038/ncb1315.

Distinct enzyme combinations in AKAP signalling complexes permit functional diversity

Naoto Hoshi¹, Lorene K. Langeberg¹, and John D. Scott^{1,2}

¹Howard Hughes Medical Institute, Vollum Institute, Oregon Health and Science University, 3181 S.W. Sam Jackson Park Road, Portland, OR 97239, USA

Abstract

Specificity in cell signalling can be influenced by the targeting of different enzyme combinations to substrates. The A-kinase anchoring protein AKAP79/150 is a multivalent scaffolding protein that coordinates the subcellular localization of second-messenger-regulated enzymes, such as protein kinase A, protein kinase C and protein phosphatase 2B. We developed a new strategy that combines RNA interference of the endogenous protein with a protocol that selects cells that have been rescued with AKAP79/150 forms that are unable to anchor selected enzymes. Using this approach, we show that AKAP79/150 coordinates different enzyme combinations to modulate the activity of two distinct neuronal ion channels: AMPA-type glutamate receptors and M-type potassium channels. Utilization of distinct enzyme combinations in this manner provides a means to expand the repertoire of cellular events that the same AKAP modulates.

Cellular regulation must be accomplished through the synchronized actions of a limited number of gene products, as the number of mammalian genes that are required to sustain life is significantly less than was originally anticipated^{1,2}. Signal-transduction pathways are created when enzymes, often with broad substrate specificities, act sequentially to evoke cellular responses³. Restricting the subcellular localization of these enzymes with scaffolding proteins contributes to the fidelity of each response⁴. Prototypic examples of these are the A-kinase anchoring proteins (AKAPs) that target the cyclic-AMP-dependent protein kinase (protein kinase A, PKA) and other enzymes to defined subcellular locations⁵.

AKAP signalling complexes often include signal-transduction and signal-termination enzymes to regulate the forward and backward steps of a given process. The notion of multivalent anchoring proteins was first proposed for the AKAP79 family, which consists of a group of three structurally similar orthologues: human AKAP79, murine AKAP150 and bovine AKAP75 (ref. 6). These AKAPs contain binding sites for PKA, the calcium/phospholipid-dependent kinase (protein kinase C, PKC) and the calcium/calmodulin-dependent phosphatase (protein phosphatase 2B, PP2B)⁷. They are tethered to the inner face of the plasma membrane, where they can respond to the generation of second messengers, such as cAMP, calcium and phospholipid⁷. Functional studies have shown that this AKAP family controls the phosphorylation status and action of several ion channels, including AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid)-type glutamate receptors, L-type calcium channels, aquaporin water channel, and M-type potassium channels^{8–10}. One theory that accounts for this diversity of action is that unique combinations of anchored

© 2005 Nature Publishing Group

²Correspondence should be addressed to J.D.S. (scott@ohsu.edu).

Note: Supplementary Information is available on the Nature Cell Biology website.

Competing Financial Interests: The authors declare that they have no competing financial interests.

enzymes are recruited to individual ion channels. We tested this hypothesis in cells in which the endogenous anchoring protein was silenced and replaced with AKAP forms that were unable to anchor selected binding partners.

Results

Functional characterization of AKAP79-depleted cells

Plasmid-based RNA interference (RNAi) of *AKAP79* was initially performed in HEK293 cells. The effect was maximal 3–5 d after transfection of the plasmid when AKAP79 levels were reduced to ~25% of the control (Fig. 1a). The pSAKAP79i-positive cultures were enriched by co-expression of the cell-surface marker CD4, followed by magnetic sorting with anti-CD4-coupled magnetic beads (Fig. 1b). Cell extracts from pSAKAP79i/CD4-positive cultures exhibited almost complete loss of AKAP79 compared with controls when assessed by immunoblot (Fig. 1c, top panel). Control immunoblots confirmed that both samples contained equivalent amounts of a standard protein (Fig. 1c, bottom panel). Next, the CD4-positive cells were transfected with plasmids encoding ion channels, modified AKAP forms and a green fluorescent protein (GFP) marker (Fig. 1b). Functional confirmation of *AKAP79* knockdown was provided by whole-cell patch-clamp recording experiments from GFP cells expressing the GluR1 subunit of the AMPA-type glutamate receptor channel (Fig. 1b). *AKAP79*-silenced cells did not exhibit agonist-dependent downregulation of current (Fig. 1d, red circles) compared with recordings from control cells (Fig. 1d, black triangles). Importantly, downregulation of the GluR1 current was rescued following expression of the murine orthologue AKAP150 that is refractory to pSAKAP79i RNAi (Fig. 1c, middle panel; Fig. 1d, green squares). Comparison of the GluR1 response 5 min post-agonist to the response at time zero showed statistically significant recovery of current rundown with expression of AKAP150 (representative traces and amalgamated data, Fig. 1e, f). In control experiments, RNAi knockdown with pSAKAP150i did not prevent AKAP79 expression (Fig. 1c, lane 3) or agonist-dependent rundown of GluR1 current (Fig. 1e, f)

Signalling to AMPA channels requires anchored PP2B and SAP97

AKAP-bound PKA enhances the phosphorylation of Ser 845 in the cytoplasmic tail of GluR1 and is believed to stabilize AMPA currents^{11,12}. Conversely, the AKAP-bound phosphatase PP2B may mediate the rundown of GluR1 currents¹³ (Fig. 2a). We tested this hypothesis by measuring GluR1 currents from *AKAP79*-silenced HEK293 cells that had been reconstituted with modified AKAP150 forms (Fig. 2b). Agonist-dependent rundown of GluR1 currents was sustained in cells expressing AKAP150 Δ PKA (Δ 675–697) and AKAP150 Δ PKC (Δ 31–51) deletion mutants that are unable to anchor PKA or PKC, respectively^{14–16} (Fig. 2c, d). By contrast, expression of AKAP150 Δ PP2B (Δ 605–647), which is unable to anchor PP2B, prevented current rundown, indicating that dephosphorylation is a predominant regulatory event (Fig. 2c, d). This is believed to occur because the PP2B-AKAP complex is optimally positioned in relation to the channel by the MAGUK adapter protein SAP97 (ref 17; Fig. 2a). Accordingly, gene silencing of *SAP97*, which is endogenously expressed in HEK293 cells (Fig. 3a), uncoupled the AKAP-mediated regulation of GluR1 currents (Fig. 3b–d). Normal regulation was restored following rescue with mouse *SAP97* that is refractory to RNAi (Fig. 2c–e).

Function of AMPA channels with modified AKAP complexes

A reciprocal RNAi-rescue procedure was developed in cultured rat hippocampal neurons, where the endogenous *AKAP150* gene was silenced following electroporation of plasmids and was rescued with AKAP79 forms. *In situ* changes in AKAP150 protein expression were detected by immunofluorescence, whereas co-expression of GFP was used as a transfection

marker and to identify positive neurons for whole-cell electrophysiological recording (Fig. 4a–f). Gene silencing of *AKAP150* abolished glutamate-dependent attenuation of AMPA currents (Fig. 4g, red triangles) compared with controls (Fig. 4g, black diamonds), whereas ectopic expression of AKAP79 rescued this phenomenon (Fig. 4g, green circles). Representative current traces were taken 5 min after the application of glutamate (Fig. 4h), and amalgamated data are presented (Fig. 4i). Functional rescue of AMPA-channel regulation was obtained with AKAP79 complexes that lacked PKA or PKC, whereas reconstruction with AKAP79 forms that were unable to bind PP2B were ineffective (Fig. 4j–l). Collectively, the data so far (Fig. 1–4) indicates that downregulation of AMPA channels involves two AKAP-mediated binding events: the cross-linking of AKAP to GluR1 and the anchoring of PP2B.

We were intrigued by evidence that the AKAP79(Δ PKA)-rescued neurons showed minimal effects on the downregulation of the AMPA current (Fig. 4k–l). Therefore, we wondered if the PKA inhibitor peptide (PKI), which was included in the pipette solution, masked the anchored-PKA-deficient phenotype. To address this question, we examined the AMPA current with or without PKI. The downregulation of AMPA currents is induced by inclusion of PKI in the patch pipette solution in control neurons (Fig. 5a). However PKI is not required to downregulate the channel in cells in which AKAP79(Δ PKA) replaced the endogenous anchoring protein (Fig. 5b). Collectively, these findings strongly indicate that anchored PKA is crucial for maintaining AMPA currents during glutamate stimulation.

AKAP150 coordinates muscarinic suppression of M currents

Our RNAi/rescue approach was used to examine a second AKAP79/150-mediated event: the suppression of ‘M current’, a K^+ conductance that negatively regulates neuronal excitability in a variety of brain regions^{18,19}. Muscarinic agonists inhibit this channel via a G_q -coupled pathway that evokes phosphoinositide turnover^{20,21} (Fig. 6a). Although a role for PKC in this process has been elusive²², recent pharmacological studies imply that an anchored pool of the kinase facilitates this suppression of M current in superior cervical ganglion (SCG) neurons¹⁰. PKC phosphorylation of Ser 534 and Ser 541 on the KCNQ2 channel subunit has been implicated in the suppression of recombinant M currents¹⁰. The net result is a hyperexcitability of neurons because firing of action potentials is favoured when M currents are suppressed^{18,19}.

AKAP79-silenced HEK293 cells expressing a KCNQ2 subunit, a core component of the M current, showed reduced muscarinic inhibition by acetylcholine (see Supplementary Information, Fig. S1). RNAi knockdown of *AKAP150* in cultured rat SCG neurons (see Supplementary Information, Fig. S2) permitted examination of the role of AKAP150 anchoring in endogenous M-current suppression. Accordingly, gene silencing of *AKAP150* in cultured rat SCG neurons reduced suppression of M currents in response to the muscarinic agonist oxotremorine-M (Oxo-M) (Fig. 6b, red circles) compared with controls (Fig. 6b, black circles). Ectopic expression of AKAP79 rescued this phenomenon (Fig. 6b, green squares). Representative traces and amalgamated data using 1 μ M Oxo-M as agonist are presented (Fig. 6c–d). Moreover, the anchoring protein is required for the muscarinic suppression of M current as the *AKAP150*-silenced SCG neurons were refractory to higher doses of Oxo-M (see Supplementary Information, Fig. S3). Surprisingly, loss of AKAP150 did not alter the response to bradykinin, another physiological agonist that suppresses M currents via a G_q -coupled pathway that also involves phosphoinositides and calcium²³ (Fig. 6e; and see Supplementary Information, Fig. S4). Further support for this unexpected selectivity in AKAP action was provided by evidence that the recombinant m_1 muscarinic receptor co-precipitated AKAP150, whereas the B_2 bradykinin receptor did not (Fig. 6f).

AKAP150 binds to the KCNQ2 subunit, a core component of the M current, presumably to direct PKC towards phosphorylation sites in the cytoplasmic tail of the channel¹⁰ (Fig. 6a). Therefore, RNAi/rescue experiments tested the hypothesis that PKC is recruited into a muscarine-sensitive signalling network that includes the m₁ receptor, the anchoring protein and the M channel. Initially, the complement of AKAP binding partners that contribute to M-current regulation was examined by expression of various AKAP79 deletion mutants in *AKAP150*-silenced rat SCG neurons. Only the PKC-anchoring-deficient form of AKAP79 failed to rescue muscarinic suppression of the M current (Fig. 7a–c). In complementary experiments, muscarinic inhibition of M current was restored following expression of AKAP79 amino acids (aa) 1–153 that encompasses the phosphoinositide binding sites and PKC anchoring region²⁴ (Fig. 7a–c). Importantly, this effect was lost when AKAP79 aa 1–153 was co-expressed with a catalytically inactive PKC β II isoform²⁵ (Fig. 7a–c). AKAP79 aa 1–153 shares 77% homology with the corresponding region of AKAP150. These studies imply that AKAP150 facilitates muscarinic suppression of M current by bringing PKC and the m₁ muscarinic receptor in close proximity to the channel.

Discussion

The important components of M-channel configuration (Fig. 7d) are distinct from the anchored signalling complex that functions to downregulate AMPA channels (Fig. 7e). Apparently, the only common functional element is the anchoring protein itself. This raises the intriguing possibility that these AKAP complexes, built on the same scaffold, use different components to modulate individual signalling events. There may be contextual cues that drive the preferential assembly of different AKAP complexes, such as the initial binding event of the anchoring protein with its target substrate. For example, direct association of AKAP150 with the M channel may provide a configuration that retains the membrane tethering and anchoring of PKC. Similarly, a GluR1–MAGUK–AKAP ternary complex may be established before PKA and PP2B can be oriented towards the channel. Other factors that could influence the composition of these signalling networks include co-translational assembly of protein complexes through localized protein synthesis²⁶ and either species-specific or cell-type-specific expression of particular binding partners²⁷. However, whether or not every enzyme is engaged on the scaffold simultaneously, our results show that AKAPs not only regulate the spatial restriction of broad-specificity enzymes, but also may regulate different signalling pathways by using only a subset of the AKAP signalling complex. Use of different signalling molecules on the same AKAP is a mechanism that permits maximum flexibility from a limited gene pool. This may be particularly important in a highly polarized cell, such as the neuron, in which the synaptic environment could take advantage of an array of AKAP79/150 signalling complexes in the regulation of distinct molecules and cellular events. For example, we show that downregulation of AMPA currents requires anchored PP2B, but we are unable to demonstrate a role for the same enzyme in the repression of muscarine-dependent M-current inhibition. This observation is supported by a detailed biochemical analysis showing that PP2B is not present in KCNQ2–AKAP79 complexes that are isolated from mammalian neurons²⁸. It is, therefore, reasonable to propose that parallel AKAP signalling complexes may recruit different combinations of signalling enzymes. This hitherto unappreciated function of AKAPs could resolve an apparent paradox in second-messenger signalling — namely, that parallel pathways that are composed of broad-specificity enzymes frequently trigger precise, localized cellular events.

Methods

Antibodies

The following primary antibodies were used for immunoblotting and immunoprecipitation: rabbit polyclonal antibody to AKAP150 (VO88), rabbit polyclonal antibody to AKAP79 (ICOS), mouse monoclonal MAGUK-family antibody (Upstate Biotechnology, Lake Placid, NY), mouse monoclonal HA (Sigma, St Louis, MO), rabbit polyclonal HA (Sigma), mouse monoclonal tubulin and mouse monoclonal FLAG M2 (Sigma).

Cell cultures

HEK293 cells were grown in Dulbecco's modified Eagle medium with 10% fetal bovine serum. SCG neurons were isolated from 2–3-week-old rats and cultured as described previously¹⁰. Hippocampal neurons were isolated from 1–2-d-old rats and cultured as described previously²⁹. Acquisition of primary cultures was under the regulation of the Institutional Animal Care and Use Committee at Oregon Health and Science University.

RNAi constructs and induction

Sequences from AKAP79 or AKAP150 were ligated into pSilencer U6 vectors (Ambion, Austin, TX) according to the manufacturers' instructions. The sequences were as follows: for AKAP79 (pSAKAP79i), 5'-GACTCAGACTGCAGCATCAAAGTTCAAGAGACTTTGATGCTGCAGTCTGAGTCTTTTTT-3'; for AKAP150 (pSAKAP150i), 5'-GAAGAAGACAAAATCCAAACTTTCAAGAGAAGTTTGGATTTTGTCTTCTTCTTTTTTGG-3'; for hSAP97 (pSSAP97i-1), 5'-GAGCATATTTCCCCACAAATTCAAGAGATTTGTGGGGAAATATGCTCTTTTTTGG-3', and (pSSAP97i-2), 5'-GAACTTATCAGAGATTGAGAATTCAAGAGATTCTCAATCTCTGATAAGTCTTTTTTGG-3'. To induce RNAi in HEK293 cells, the appropriate construct in the *pSilencer* vector (pSAKAP79i or a mixture of pSSAP97i-1 and pSSAP97i-2) and the CD4 expression vector (pJPACD4) were co-transfected using Fugene6 (Roche Diagnostics, Indianapolis, IN). At 48 h after transfection, these cultures were enriched for transfected cells by selecting the CD4-positive cells with anti-CD4-magnetic beads (Dyna, Oslo, Norway) (Fig. 1b). Pilot experiments using pDsRED2 as a fluorescent marker and pJPACD4 showed that the magnetic selection process increased DsRED-positive cells from 45% ± 4.1% ($n = 3$ fields, 841 cells) to 82% ± 1.7% ($n = 3$ fields, 393 cells). Enrichment of transfected cells facilitated re-transfection and further experiments on a gene-silenced background (Fig. 1b). Re-transfection was done with AKAP150 full-length wild type and mutants in pcDNA3, GluR1_{flip} in pRK5 vector and pEGFP-N1. Cells positive for both CD4 beads and GFP were visually selected for electrophysiological recording 2 d after re-transfection. For experiments in *AKAP150*-silenced hippocampal neurons, suspensions of hippocampal cells from newborn rats were transfected with pSAKAP150i, various AKAP79 forms in pcDNA3, and either pEGFP-N1 or pDsRED2 using a Nucleofector kit (Amaxa, Koeln, Germany). Treated cells were overlaid onto a glial feeder layer and cultured for 4–7 d. For *AKAP150*-silenced SCG neurons, the same constructs were introduced by nuclear injection as described previously¹⁰ and cultured for 3 d.

Immunohistochemistry

Rat hippocampal neurons cultured on coverslips were fixed by 3.7% formaldehyde and permeabilized with 0.2% Triton X-100 in phosphate-buffered saline (PBS). Following incubation in blocking buffer (0.2% bovine serum albumin in PBS) for 1 h, they were incubated with AKAP150 antibody (1:1000 dilution). After three washes with blocking buffer, the coverslips were incubated in goat anti-rabbit Cy5 (1:100 dilution; Jackson

Immunoresearch, West Grove, PA) for 1 h, washed as before and mounted with ProLong mounting medium (Molecular Probes, Eugene, OR). Cells were imaged with a Bio-Rad MRC1024 UV/VIS confocal microscope.

Immunoprecipitation

HA-tagged receptor/AKAP150 complexes were immunoprecipitated from one 10 cm dish of HEK293 cells that had been transfected with 1 μ g pcDNA3–AKAP150 and 9 μ g of either HA-tagged bradykinin B₂ receptor or muscarinic m₁ receptor cDNA (University of Missouri-Rolla, cDNA Resource Center). Cells were harvested 48 h after transfection and lysed in 500 μ l HSE buffer (150 mM NaCl, 5 mM EDTA, 5 mM EGTA, 20 mM HEPES, pH 7.4, 1% Triton X-100 and Roche protease inhibitor cocktail). Supernatants were incubated with 1 μ g mouse anti-HA and 40 μ l protein G agarose beads. Following overnight incubation at 4°C, immunoprecipitates were washed twice in HSE, twice in HSE with 650 mM NaCl, twice in HSE and once in TE buffer. Bound proteins were analysed by SDS–PAGE and immunoblotted with rabbit antibody to AKAP150 (1:5000 dilution) and HA-tag (1:1000 dilution).

Electrophysiological measurements

Whole-cell recordings were performed on isolated cells using an Axopatch 200B patch-clamp amplifier (Axon Instruments, Sunnyvale, CA). Signals were sampled at 2 kHz, filtered at 1 kHz, and acquired using pClamp software (version 7, Axon Instruments). Series resistance (90–95%) and whole-cell capacitance compensation were used. To measure the GluR1 response in HEK293 cells, patch pipettes (2–4 M Ω) were filled with an intracellular solution containing 140 mM Cs methanesulphonate, 5 mM ATP, 1 mM BAPTA, 5 mM MgCl₂, 300 μ M CaCl₂, 10 mM HEPES (pH 7.3) and 10 μ M PKI. The external solution consisted of 150 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 10 mM glucose, 100 μ M cyclothiazide and 10 mM HEPES (pH 7.4). Rapid solution exchanges were accomplished through a two-barrel pipe controlled by a solution stimulus delivery device, SF-77B (Warner Instruments, Hamden, CT). GluR1 receptor currents were evoked by a 500-ms application of 1 mM glutamate at 30-s intervals at a holding potential of –70 mV. For recording AMPA currents in hippocampal neurons, the pipette solution was the same as above except that it contained no calcium. Hippocampal cells were perfused with a solution containing 150 mM NaCl, 3 mM KCl, 1 mM CaCl₂, 11 mM glucose, 10 mM HEPES (pH 7.4), 100 μ M cyclothiazide, 10 μ M bicuculline, 10 μ M ZD7288 and 1 μ M TTX. Neuronal glutamate responses were evoked as above, but with a lower glutamate concentration (100 μ M) and a slower washout because the cells were not lifted. Adding cyclothiazide increased the responses to glutamate from 0.43 ± 0.11 nA to 3.9 ± 0.52 nA ($n = 4$), which indicates that the majority of the recorded currents were through the AMPA receptor. Control measurements without PKI revealed a relatively stable AMPA response ($84.2\% \pm 3.3\%$ at 10 min, $n = 4$). Data are presented as mean \pm SEM. For recording M currents in SCG neurons, the perforated patch method was used as described previously¹⁰. Briefly, amphotericin B (0.1–0.2 mg ml^{–1}) was dissolved in an intracellular solution containing 130 mM potassium acetate, 15 mM KCl, 3 mM MgCl₂, 6 mM NaCl and 10 mM HEPES (pH 7.3). When filled with this internal solution, pipette resistance was 1.5–2 M Ω . Access resistances after permeabilization ranged from 7–22 M Ω . The external solution consisted of 120 mM NaCl, 6 mM KCl, 1.5 mM MgCl₂, 2.5 mM CaCl₂, 11 mM glucose and 10 mM HEPES (pH 7.4). Amplitudes of the M currents were measured as deactivating currents during 500-ms test pulses to –50 mV from a holding potential of –20 mV. Data are presented as mean \pm SEM.

Statistical analysis

All statistical tests were carried out using one-way ANOVA, followed by two-tailed unpaired Student's *t*-tests using Excel (Microsoft) and InStat (Graphpad software). All alpha levels were set to 0.05. For each data set, Gaussian distribution was confirmed by the method used by Kolmogorov and Smirnov (InStat, Graphpad software).

BIND identifiers

One BIND identifier (www.bind.ca) is associated with this manuscript: 335775.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank J. J. Carlisle Michel and R. Goodman for insightful comments on the manuscript and R. Mouton for technical assistance. This work was supported by grant no. GM48231 from the National Institutes of Health to J.D.S.

References

1. Venter JC, et al. The sequence of the human genome. *Science*. 2001; 291:1304–1351. [PubMed: 11181995]
2. Lander ES, et al. Initial sequencing and analysis of the human genome. *Nature*. 2001; 409:860–921. [PubMed: 11237011]
3. Hunter T. Signaling — 2000 and beyond. *Cell*. 2000; 100:113–127. [PubMed: 10647936]
4. Pawson T, Nash P. Assembly of cell regulatory systems through protein interaction domains. *Science*. 2003; 300:445–452. [PubMed: 12702867]
5. Wong W, Scott JD. AKAP signalling complexes: Focal points in space and time. *Nature Rev Mol Cell Biol*. 2004; 5:959–971. [PubMed: 15573134]
6. Carr DW, Stofko-Hahn RE, Fraser IDC, Cone RD, Scott JD. Localization of the cAMP-dependent protein kinase to the postsynaptic densities by A-kinase anchoring proteins: characterization of AKAP79. *J Biol Chem*. 1992; 267:16816–16823. [PubMed: 1512224]
7. Klauck TM, et al. Coordination of three signaling enzymes by AKAP79, a mammalian scaffold protein. *Science*. 1996; 271:1589–1592. [PubMed: 8599116]
8. Gao T, et al. cAMP-dependent regulation of cardiac L-type Ca²⁺ channels requires membrane targeting of PKA and phosphorylation of channel subunits. *Neuron*. 1997; 19:185–196. [PubMed: 9247274]
9. Fraser ID, Scott JD. Modulation of ion channels: a ‘current’ view of AKAPs. *Neuron*. 1999; 23:423–426. [PubMed: 10433254]
10. Hoshi N, et al. AKAP150 signaling complex promotes suppression of the M-current by muscarinic agonists. *Nature Neurosci*. 2003; 6:564–571. [PubMed: 12754513]
11. Banke TG, et al. Control of GluR1 AMPA receptor function by cAMP-dependent protein kinase. *J Neurosci*. 2000; 20:89–102. [PubMed: 10627585]
12. Esteban JA, et al. PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity. *Nature Neurosci*. 2003; 6:136–143. [PubMed: 12536214]
13. Tavalin SJ, et al. Regulation of GluR1 by the A-kinase anchoring protein 79 (AKAP79) signaling complex shares properties with long-term depression. *J Neurosci*. 2002; 22:3044–3051. [PubMed: 11943807]
14. Carr DW, Hausken ZE, Fraser ID, Stofko-Hahn RE, Scott JD. Association of the type II cAMP-dependent protein kinase with a human thyroid RII-anchoring protein Cloning and characterization of the RII-binding domain. *J Biol Chem*. 1992; 267:13376–13382. [PubMed: 1618839]

15. Faux MC, Scott JD. Regulation of the AKAP79-protein kinase C interaction by Ca²⁺/calmodulin. *J Biol Chem.* 1997; 272:17038–17044. [PubMed: 9202019]
16. Dell'Acqua ML, Dodge KL, Tavalin SJ, Scott JD. Mapping the protein phosphatase-2B anchoring site on AKAP79 Binding and inhibition of phosphatase activity are mediated by residues 315–360. *J Biol Chem.* 2002; 277:48796–48802. [PubMed: 12354762]
17. Garner CC, Nash J, Haganir RL. PDZ domains in synapse assembly and signalling. *Trends Cell Biol.* 2000; 10:274–280. [PubMed: 10856930]
18. Brown DA, Adams PR. Muscarinic suppression of a novel voltage-sensitive K⁺ current in a vertebrate neurone. *Nature.* 1980; 283:673–676. [PubMed: 6965523]
19. Marrion NV. Control of M-current. *Annu Rev Physiol.* 1997; 59:483–504. [PubMed: 9074774]
20. Suh BC, Hille B. Recovery from muscarinic modulation of M current channels requires phosphatidylinositol 4,5-bisphosphate synthesis. *Neuron.* 2002; 35:507–520. [PubMed: 12165472]
21. Zhang H, et al. PIP(2) activates KCNQ channels, and its hydrolysis underlies receptor-mediated inhibition of M currents. *Neuron.* 2003; 37:963–975. [PubMed: 12670425]
22. Bosma MM, Hille B. Protein kinase C is not necessary for peptide-induced suppression of M current or for desensitization of the peptide receptors. *Proc Natl Acad Sci USA.* 1989; 86:2943–2947. [PubMed: 2468164]
23. Cruzblanca H, Koh DS, Hille B. Bradykinin inhibits M current via phospholipase C and Ca²⁺ release from IP₃-sensitive Ca²⁺ stores in rat sympathetic neurons. *Proc Natl Acad Sci USA.* 1998; 95:7151–7156. [PubMed: 9618554]
24. Dell'Acqua ML, Faux MC, Thorburn J, Thorburn A, Scott JD. Membrane-targeting sequences on AKAP79 bind phosphatidylinositol-4, 5- bisphosphate. *EMBO J.* 1998; 17:2246–2260. [PubMed: 9545238]
25. Edwards AS, Newton AC. Phosphorylation at conserved carboxyl-terminal hydrophobic motif regulates the catalytic and regulatory domains of protein kinase C. *J Biol Chem.* 1997; 272:18382–18390. [PubMed: 9218480]
26. Smith WB, Starck SR, Roberts RW, Schuman EM. Dopaminergic stimulation of local protein synthesis enhances surface expression of GluR1 and synaptic transmission in hippocampal neurons. *Neuron.* 2005; 45:765–779. [PubMed: 15748851]
27. Marrion NV. Calcineurin regulates M channel modal gating in sympathetic neurons. *Neuron.* 1996; 16:163–173. [PubMed: 8562080]
28. Cooper EC, et al. Colocalization and coassembly of two human brain M-type potassium channel subunits that are mutated in epilepsy. *Proc Natl Acad Sci USA.* 2000; 97:4914–4919. [PubMed: 10781098]
29. Westphal RS, et al. Regulation of NMDA receptors by an associated phosphatase-kinase signaling complex. *Science.* 1999; 285:93–96. [PubMed: 10390370]




Figure 1.

RNA interference of *AKAP79* in HEK293 cells. **(a)** Immunoblot showing *AKAP79* (top panel) and tubulin (loading control, middle panel) expression levels in cell lysates from cells transfected with control or pSAKAP79i plasmids (indicated above lanes). The time (days) post-transfection is indicated above each lane. (Bottom panel) *AKAP79* expression levels from control (black circles) and gene-silenced (red circles) cells were quantified by densitometry from immunoblots using an NIH image. Amalgamated data from five experiments is presented. **(b)** Flowchart depicting the selection protocol used to isolate CD4/pSAKAP79i double-positive cells and green fluorescent protein (GFP) cells expressing modified *AKAP* forms and ion-channels. **(c)** Characterization of *AKAP79* knockdown and rescue with recombinant *AKAP150* in CD4/pSAKAP79i double-positive cells (lane 2). Immunoblot detection of *AKAP79* (top panel), recombinant *AKAP150* (middle panel) and tubulin (bottom panel) in cell lysates is shown. The reciprocal experiment with *AKAP150* knockdown in CD4/pSAKAP150i double-positive cells and rescue with recombinant *AKAP79* is shown in lane 3. **(d)** Electrophysiological recording of recombinant GluR1 channels time course from control (black triangles, $n = 8$), *AKAP79*-silenced (red circles, $n = 9$) and CD4/pSAKAP79i double-positive cells expressing recombinant *AKAP150* (*AKAP150* rescue, green squares, $n = 12$). Normalized GluR1 currents (relative to time 0, 1 mM glutamate) over a time course of 10 min are presented. The PKA inhibitor PKI was included in the pipette solution. **(e)** Representative traces at time 0 and 5 min after the first application of 1 mM glutamate are presented. A representative trace from an additional control group of cells transfected with the pSAKAP150i construct specific for the murine orthologue is shown ($n = 6$). **(f)** Amalgamated data depicting the level of normalized current for all experimental groups taken 5 min after the application of agonist. The statistical significance of electrophysiology data shown in this figure, part **f**, was calculated using one-

way ANOVA followed by two-tailed Student's *t*-test. Statistical significance of control versus pSAKAP79i is $P = 0.0050$ and of pSAKAP79i + AKAP150 rescue versus pSAKAP79i is $P = 0.0050$. Error bars indicate SEM.




Figure 2.

PP2B is a fundamental component of the GluR1–AKAP9/150 signalling network in HEK293 cells. **(a)** Diagram depicting a proposed configuration of the GluR1–AKAP signalling network. **(b)** Schematic representation of AKAP150 deletion mutants lacking enzyme-binding sites for protein kinase A (PKA; green), protein kinase C (PKC; blue) or protein phosphatase 2B (PP2B; red) used to reconstitute modified GluR1–AKAP signalling networks. **(c)** Representative traces at time 0 and 5 min after the initial application of 1 mM glutamate and **(d)** amalgamated data depicting the level of normalized current for each experimental group taken 5 min after the application of agonist (n values are indicated). The PKA inhibitor PKI was included in the pipette solution. The statistical significance of electrophysiology data shown in this figure, part **d**, was calculated using one-way ANOVA followed by two-tailed Student's t -test. Statistical significance of AKAP150ΔPP2B rescue versus AKAP150 rescue is $P = 0.0090$. Error bars indicate SEM.




Figure 3. RNA interference of *SAP97* in HEK293 cells. **(a)** Plasmid-based RNA interference knockdown of the adapter protein *SAP97*. (Top panel) Immunoblot detection of *SAP97* and (Bottom panel) immunoblot detection of tubulin in lysates from control and pSSAP97i-transfected cells 2 and 3 d after transfection. **(b)** Schematic representation depicting the disruption of the GluR1–AKAP79 signalling network following removal of *SAP97*. **(c)** Representative traces at time 0 and 5 min after the application of 1 mM glutamate and **(d)** amalgamated data depicting the level of normalized current for each experimental group taken 5 min after the application of agonist. The protein kinase A (PKA) inhibitor PKI was included in the pipette solution. **(e)** Schematic depicting the reformation of the GluR1–AKAP79 signalling network following the expression of rat *SAP97*. The statistical significance of electrophysiology data shown in this figure, part **d**, was calculated using one-way ANOVA followed by two-tailed Student's *t*-test. Statistical significance of *SAP97* rescue versus pSSAP97i knockdown is $P = 0.00011$. Error bars indicate SEM.




Figure 4. Modulation of AMPA channels with modified AKAP complexes. Gene silencing of *AKAP150* was achieved by electroporation of cultured hippocampal neurons with the pSAKAP150i plasmid. (**a, d**) Immunofluorescent detection of AKAP150; (**b, e**) fluorescent detection of green fluorescent protein (GFP); (**c, f**) composite images of AKAP150 and the outline of GFP-positive cells in cultured hippocampal neurons transfected with pSAKAP150i or a control plasmid. Fluorescent emission from GFP and anti-AKAP150 labelled with Cy5-conjugated secondary antibodies was detected using a confocal microscope (scale bar, 20 μm). (**g**) Whole-cell electrophysiological recording of AMPA

currents in control (black diamonds, $n = 11$), *AKAP150*-silenced (red triangles, $n = 7$) or *AKAP79*-rescued (green circles, $n = 8$) neurons. The protein kinase A (PKA) inhibitor PKI was included in the pipette solution. Time courses of normalized AMPA currents are presented for 0–10 min after the application of 100 μM glutamate. **(h)** Representative traces at time 0 and 5 min after the application of 100 μM glutamate. **(i)** Amalgamated data depicting the level of normalized current for each experimental group taken 5 min after the application of agonist. **(j)** Diagram of *AKAP79* deletion mutants lacking enzyme-binding sites for PKA (green), protein kinase C (PKC; blue) or protein phosphatase 2B (PP2B; red) used to reconstitute modified signalling complexes in *AKAP150*-silenced neurons. **(k)** Representative traces at time 0 and 5 min after the application of 100 μM glutamate and **(l)** amalgamated data depicting the level of normalized current for each experimental group taken 5 min after the application of agonist (n values are indicated). The statistical significance of electrophysiology data shown in this figure, parts **i** and **l**, was calculated using one-way ANOVA followed by two-tailed Student's t -test. Statistical significance of pSAKAP150i versus control is $P = 0.000093$, and of *AKAP79* Δ PP2B rescue versus *AKAP79* rescue is $P = 0.018$. Error bars indicate SEM.





Figure 5. PKA activation is required for maintaining AMPA currents. **(a)** Whole-cell electrophysiological recording of AMPA currents in control hippocampal neurons without (–PKI, $n = 5$, black diamonds) and with (+PKI, $n = 7$, blue circles) PKI ($10 \mu\text{M}$) in the pipette solution. Time courses of normalized AMPA currents are presented for 0–10 min after the application of $100 \mu\text{M}$ glutamate. Representative traces from both –PKI (black) and +PKI (blue) at time 0 and 5 min after the application of $100 \mu\text{M}$ glutamate. **(b)** Similar recordings of AMPA currents in *AKAP150*-silenced and *AKAP79*(ΔPKA)-rescued neurons (–PKI, $n = 9$, black diamonds; +PKI, $n = 5$, green circles).

**Figure 6.**

AKAP150 facilitates selective M-current suppression. **(a)** Schematic diagram of receptor-mediated G-protein-coupled pathways that modulate M channels. **(b)** Muscarinic suppression of M channels in dissociated superior cervical ganglion (SCG) neurons. The muscarinic agonist oxotremorine-M (Oxo-M; 1 μM) was applied through the flow pipette for 1 min. Whole-cell electrophysiological recording was continued for another 4 min in control (black circles, $n = 8$), *AKAP150*-silenced (red circles, $n = 12$) and AKAP79-rescued (green squares, $n = 6$) SCG neurons. **(c)** Representative traces at time 0 and 1 min after the application of 1 μM Oxo-M and **(d)** amalgamated data depicting the level of normalized current 1 min after the application of agonist are presented for each experimental group. An additional control group of cells transfected with the pSAKAP79i construct specific for the human orthologue is also shown ($n = 10$). **(e)** M currents recorded from control (black circles, $n = 7$) and *AKAP150*-silenced (red squares, $n = 9$) neurons following Oxo-M (1 μM) treatment. After a recovery period of 10 min, a second agonist, bradykinin (100 nM), was applied to the same cells. Representative traces from control (top) and *AKAP150*-silenced (below) neurons at time zero and 1 min after the application of Oxo-M (left inset) and bradykinin (right inset). **(f)** AKAP150 and the m₁ muscarinic receptor interact. HEK293 cells were co-transfected with plasmids encoding AKAP150 and HA-tagged m₁ muscarinic receptor or HA-tagged B₂ bradykinin receptor, followed by immunoprecipitation with monoclonal antibodies against the HA epitope tag. (Top panel) Immunoblot detection of AKAP150 and (middle panel) immunoblot detection of HA-tagged receptors present in immune complexes isolated with anti-HA. (Bottom panel) Shows immunoblot detection of AKAP150 presence in cell lysates. The statistical significance of electrophysiology data shown in this figure, part **d**, was calculated using one-way ANOVA followed by two-tailed Student's *t*-test. Statistical significance of pSAKAP150i versus control is $P = 0.00021$. Error bars indicate SEM.




Figure 7.

Functional dissection of an M-channel–AKAP complex. **(a)** Schematic diagram of AKAP79 deletion mutants and fragments used in functional rescue experiments. Enzyme binding sites for protein kinase C (PKC; blue), protein phosphatase 2B (PP2B; red) and protein kinase A (PKA; green) are indicated. **(b)** Representative traces at time 0 and 1 min after the application of 10 μM Oxo-M and **(c)** amalgamated data depicting the level of normalized current for each experimental group taken 1 min after the application of agonist. Summary diagrams depicting the functionally relevant components required for **(d)** muscarinic suppression of M channels in superior cervical ganglion neurons and **(e)** glutamate-mediated downregulation of AMPA channels in hippocampal neurons. The statistical significance of electrophysiology data shown in part **c** was calculated using one-way ANOVA followed by two-tailed Student's *t*-test. Statistical significance of pSAKAP150i versus control is $P = 0.0000047$, AKAP79ΔPKC versus AKAP79 rescue is $P = 0.0022$, AKAP79 aa 1–153 + kinase-dead PKC versus AKAP79 rescue is $P = 0.00069$, AKAP79 aa 100–427 versus AKAP79 rescue is $P = 0.0058$. Error bars indicate SEM.