

Burnout and Career Satisfaction Among US Oncologists

Tait D. Shanafelt, William J. Gradishar, Michael Kosty, Daniel Satele, Helen Chew, Leora Horn, Ben Clark, Amy E. Hanley, Quyen Chu, John Phippen, Jeff Sloan, and Marilyn Raymond

A B S T R A C T

Purpose

To evaluate the personal and professional characteristics associated with career satisfaction and burnout among US oncologists.

Methods

Between October 2012 and March 2013, the American Society of Clinical Oncology conducted a survey of US oncologists evaluating burnout and career satisfaction. The survey sample included equal numbers of men and women and represented all career stages.

Results

Of 2,998 oncologists contacted, 1,490 (49.7%) returned surveys (median age of respondents, 52 years; 49.6% women). Among the 1,117 oncologists (37.3% of overall sample) who completed full-length surveys, 377 (33.8%) were in academic practice (AP) and 482 (43.2%) in private practice (PP), with the remainder in other settings. Oncologists worked an average of 57.6 hours per week (AP, 58.6 hours per week; PP, 62.9 hours per week) and saw a mean of 52 outpatients per week. Overall, 484 oncologists (44.7%) were burned out on the emotional exhaustion and/or depersonalization domain of Maslach Burnout Inventory (AP, 45.9%; PP, 50.5%; $P = .18$). Hours per week devoted to direct patient care was the dominant professional predictor of burnout for both PP and AP oncologists on univariable and multivariable analyses. Although a majority of oncologists were satisfied with their career (82.5%) and specialty (80.4%) choices, both measures of career satisfaction were lower for those in PP relative to AP (all $P < .006$).

Conclusion

Overall career satisfaction is high among US oncologists, albeit lower for those in PP relative to AP. Burnout rates among oncologists seem similar to those described in recent studies of US physicians in general. Those oncologists who devote the greatest amount of their professional time to patient care seem to be at greatest risk for burnout.

J Clin Oncol 32:678-686. © 2014 by American Society of Clinical Oncology

INTRODUCTION

Although rewarding, caring for patients with cancer is demanding and stressful.¹ Oncologists work long hours, supervise the administration of highly toxic therapy, and are continually exposed to death and suffering.¹⁻³ These characteristics place oncologists at risk for burnout, a syndrome characterized by emotional exhaustion, treating people as if they are objects (ie, depersonalization), and loss of meaning or purpose in work.^{1,4} In addition to potentially profound personal consequences (eg, anxiety, depression, alcohol/substance use, suicide),⁵⁻⁸ burnout among physicians seems to have important professional consequences, including adverse effects on quality of care and professionalism.⁹⁻¹³ Studies also suggest that physicians experiencing burnout are more likely to reduce their work hours and/or pursue early retirement,¹⁴ with potential manpower

implications for the physician workforce. Although isolated studies have explored burnout in national samples of US oncologists (most recently in 2003),^{15,16} little is known about personal and professional characteristics associated with burnout and professional satisfaction.^{1,17,18}

METHODS

Participants

A sample of 3,000 oncologists was assembled from the 8,998 US oncologists in the American Society of Clinical Oncology (ASCO; Alexandria, VA) membership file. To ensure adequate representation of oncologists at different career stages and of both sexes, oncologists in the membership file were classified by sex and categorized into three groups according to years in practice (< 10, 10 to 19, and > 20 years). Oncologists were then selected at random to construct a sample evenly distributed by career stage ($n = 1,000$ from each of career stage category) and sex (1,500 men; 1,500 women).

Tait D. Shanafelt, Daniel Satele, Jeff Sloan, Mayo Clinic, Rochester, MN; William J. Gradishar, Northwestern University, Evanston, IL; Michael Kosty, Scripps Clinic, La Jolla; Helen Chew, University of California Davis, Davis, CA; Leora Horn, Vanderbilt University Medical Center, Nashville, TN; Ben Clark, Amy E. Hanley, Marilyn Raymond, American Society of Clinical Oncology, Alexandria, VA; Quyen Chu, Louisiana State University Health Sciences Center, Shreveport, LA; and John Phippen, Texas Oncology-Baylor Charles A. Sammons Cancer Center, Dallas, TX.

Published online ahead of print at www.jco.org on January 27, 2014.

Supported by the American Society of Clinical Oncology, National Institutes of Health (American College of Surgeons Oncology Group Community Clinical Oncology Program Grant No. U10 CA149950), and Mayo Clinic Department of Medicine Program on Physician Well-Being.

Authors' disclosures of potential conflicts of interest and author contributions are found at the end of this article.

Corresponding author: Tait D. Shanafelt, MD, Mayo Clinic, Department of Internal Medicine, 200 First St SW, Rochester, MN 55905; e-mail: shanafelt.tait@mayo.edu.

© 2014 by American Society of Clinical Oncology

0732-183X/14/3207w-678w/\$20.00

DOI: 10.1200/JCO.2013.51.8480

Career Satisfaction of US Oncologists

Table 1. Personal Characteristics for Oncologists in AP Versus PP

Characteristic	All (N = 1,117)		AP (n = 377)		PP* (n = 482)		P†
	No.	%	No.	%	No.	%	
Age, years							
Median		52		50		52	.0037
Missing		32		10		12	.0380
< 40	63	5.8	29	7.9	24	5.1	
40-49	369	34.0	150	40.9	161	34.3	
50-59	343	31.6	113	30.8	165	35.1	
≥ 60	310	28.6	75	20.4	120	25.5	
Sex							< .001
Missing		18		5		3	
Male	554	50.4	158	42.5	260	54.3	
Female	545	49.6	214	57.5	219	45.7	
Children							< .001
Missing		17		4		3	
Yes	946	86.0	299	80.2	431	90.0	
No	154	14.0	74	19.8	48	10.0	
Youngest child age, years							.0532
Missing		173		79		51	
< 5	119	12.6	55	18.5	47	10.9	
5-12	249	26.4	85	28.5	121	28.1	
13-18	159	16.8	51	17.1	83	19.3	
19-22	106	11.2	34	11.4	53	12.3	
> 22	311	32.9	73	24.5	127	29.5	
Relationship status							.1721
Missing		16		3		3	
Single	98	8.9	40	10.7	32	6.7	
Married	949	86.2	317	84.8	427	89.1	
Partnered	34	3.1	12	3.2	12	2.5	
Widowed/widower	20	1.8	5	1.3	8	1.7	
Ever gone through divorce							.1962
Missing		20		6		5	
Yes	193	17.6	60	16.2	79	16.6	
No	896	81.7	310	83.6	391	82.0	
Currently going through one	8	0.7	1	0.3	7	1.5	
Current student loan debt							.0742
Missing		19		2		6	
No debt	985	89.7	324	86.4	432	90.8	
Debt < \$25,000	26	2.4	13	3.5	9	1.9	
\$25,000-\$49,999	13	1.2	9	2.4	2	0.4	
\$50,000-\$74,999	28	2.6	14	3.7	11	2.3	
\$75,000-\$99,999	16	1.5	6	1.6	8	1.7	
\$100,000-\$125,000	10	0.9	2	0.5	6	1.3	
> \$125,000	20	1.8	7	1.9	8	1.7	

Abbreviations: AP, academic practice; PP, private practice.

*Including single-specialty group, multispecialty group, and health maintenance organization.

†AP to PP.

The 3,000 individuals in the sample were sent an e-mail stating the purpose of the study (eg, to better understand factors contributing to career satisfaction among US oncologists) and providing a link to an electronic survey in October 2012. Three reminder requests were sent over the ensuing 3 weeks. Two individuals sent surveys were deceased, yielding a final sample of 2,998. Individuals not responding to the electronic survey were mailed an identical paper version of the survey in November 2012. Those not responding by January 2013 were sent a brief postcard survey. As an incentive to participate, oncologists who completed the full-length survey received a free ASCO educational product. Participation was voluntary, and all data were deidentified before analysis. ASCO commissioned the study with human subject oversight provided by the Institutional Review Board of the Mayo Clinic (Scottsdale, AZ).

Study Measures

Full-length survey. The full-length survey included 60 questions exploring a variety of personal and professional characteristics and using standardized instruments to measure burnout and career satisfaction. The full survey is available by request.

Burnout was measured using the Maslach Burnout Inventory (MBI), a 22-item questionnaire considered the gold-standard tool for measuring burnout.^{4,19-21} The MBI has three subscales to evaluate each domain of burnout: emotional exhaustion, depersonalization, and low personal accomplishment. In the standard scoring for health care workers, physicians with scores ≥ 27 on the emotional exhaustion subscale, ≥ 10 on the depersonalization subscale, or < 33 on the personal accomplishment subscale are considered to have a high degree of burnout in that dimension.⁴ In keeping with previous

Table 2. Practice Characteristics for AA Versus PP

Characteristics	All (N = 1,117)		AP (n = 377)		PP* (n = 482)		P†
	No.	%	No.	%	No.	%	
Years in practice‡							
Mean		21.7		19.4		21.0	.0262
Median		20.0		18.0		20.0	
< 10	148	14.5	64	18.7	62	13.9	.0599
10-19	337	33.1	126	36.8	152	34.0	
≥ 20	533	52.4	152	44.4	233	52.1	
Practice setting							
Academic medical center	377	34.0	377		—		
PP single-specialty group	335	30.2	—		335	69.5	
PP multispecialty group	124	11.2	—		124	25.7	
PP health maintenance organization	23	2.1	—		23	4.8	
Veterans hospital	20	1.8	—		—		
Active military practice	2	0.2	—		—		
Industry	59	5.3	—		—		
Not in practice or retired	31	2.8	—		—		
Other	138	12.4	—		—		
Time devoted to patient care, %							
Missing		10		1		2	< .001
None	82	7.5	3	.8	1	0.2	
1-25	94	8.6	43	11.4	8	1.7	
26-50	121	11.0	99	26.3	7	1.5	
51-75	199	18.0	140	37.2	39	8.1	
76-100	611	55.2	91	24.2	425	88.5	
Focus on specific type cancer							
Missing		45		11		16	< .001
Yes	418	39.0	295	80.6	81	17.4	
No	654	61.0	71	19.4	385	82.6	
Time supervising physicians in training, %							
Missing		78		25		25	< .001
0	385	37.1	9	2.6	241	52.7	
< 5	232	22.3	48	13.6	141	30.9	
5-10	183	17.6	103	29.3	56	12.3	
11-20	130	12.5	106	30.1	12	2.6	
> 20	109	10.5	86	24.4	7	1.5	
Hours and call schedule							
Median nights on call/week		1		1		2	< .001
Hours seeing patients at work/week							< .001
Mean	34.0		29.2		43.4		
SD	17.2		14.1		11.9		
Hours on administrative tasks at work/week							< .001
Mean	11.5		14.6		8.9		
SD	10.5		11.0		6.9		
Hours spent at home on work tasks/week							< .001
Mean	8.5		10.8		7.2		
SD	8.7		8.5		7.2		
Hours at home to keep abreast of developments/week							.4064
Mean	4.6		4.6		4.3		
SD	4.0		3.8		3.3		
Mean total hours/week§							< .001
Median	57.6		58.6		62.9		
SD	20.8		17.7		16.2		
Outpatient practice							
Outpatients in clinic/week							< .001
Mean	51.7		37.4		74.2		
SD	34.6		21.0		31.0		
Minutes allocated/new outpatient							.0011
Mean	49.1		53.9		51.5		
SD	20.3		17.0		14.8		
Minutes allocated/return outpatient							< .001
Mean	18.2		20.7		17.8		
SD	8.2		6.8		6.1		

(continued on following page)

Table 2. Practice Characteristics for AA Versus PP (continued)

Characteristics	All (N = 1,117)		AP (n = 377)		PP* (n = 482)		Pt
	No.	%	No.	%	No.	%	
Hospital practice							
Hospital rounding							< .001
Missing	45		5		18		
Round own patients when hospitalized	162	15.1	20	5.4	118	25.4	
Share rounding with partners in blocks	171	16.0	58	15.6	92	19.8	
Share rounding with partners on weekends	307	28.6	35	9.4	218	47.0	
Attend oncology teaching service	249	23.2	224	6.2	7	1.5	
Do not round in hospital	183	17.1	35	9.4	29	6.3	
No. of inpatients on average hospital day							< .001
Mean	7.0		11.9		5.1		
SD	7.3		8.1		5.3		
No. of weekends rounding in hospital/year							< .001
Mean	9.8		7.2		13.0		
SD	9.8		5.5		10.9		
Compensation method							< .001
Missing	102		25		26		
Salary no incentive	336	33.1	134	38.1	95	20.8	
Salary with bonus	466	45.9	207	58.8	182	39.9	
Pure incentive	213	21.0	11	3.1	179	39.3	

Abbreviations: AP, academic practice; PP, private practice; SD, standard deviation.
 *Including single-specialty group, multispecialty group, and health maintenance organization.
 †Comparison of AP to PP.
 ‡Since completion of fellowship training.
 §Sum of above four categories.

studies²²⁻²⁴ and convention,²⁵ we considered physicians with high scores on the depersonalization and/or emotional exhaustion subscales as having at least one manifestation of professional burnout.⁴ Career satisfaction was assessed using two questions from previous physician surveys regarding career and specialty choice.^{17,23,26-29}

Postcard survey. To gain insight into participation bias, oncologists not completing the full-length survey were sent a six-question postcard survey that collected information on age, sex, years in practice, and career satisfaction, along with a validated two-item measure of burnout shown to be an accurate proxy measure of burnout.³⁰⁻³²

Statistical Analysis

All full-length and postcard surveys received by March 15, 2013, were included in the analysis. Standard descriptive statistics were used to characterize responding oncologists. Associations between variables were evaluated using the Kruskal-Wallis (continuous variables) or χ^2 test (categorical variables) as appropriate. All tests were two sided with type I error rates of 0.05. With the 1,117 responses to the full-length survey, the percentage estimates are accurate to 2.9% with 95% confidence. Comparisons between men and women oncologists were tested using Wilcoxon-Mann-Whitney and Fisher's exact tests. Comparisons with 554 men and 545 women have 80% power to detect an average difference of 17% times the standard deviation, a relatively small effect size.^{33,34} Multivariable analysis to identify demographic and professional characteristics associated with the dependent outcomes was performed using logistic regression (Appendix, online only). All analyses used SAS software (version 9; SAS Institute, Cary, NC).

RESULTS

Personal and Professional Characteristics

Of 2,998 oncologists who received an invitation to participate, 1,490 (49.7%) responded. Of these, 1,117 oncologists (75.0%) completed the full-length survey (613 electronic; 504 paper version), and 373 (25.0%) completed postcard surveys. Participants were represen-

tative of the overall sample with respect to sex; however, early-career oncologists (in practice < 10 years) were somewhat less likely to respond than later-career oncologists (in practice \geq 20 years). Comparison of full-length survey responders with those completing only the postcard survey (a standard approach for evaluating response bias) did not identify any statistically significant differences with respect to age, sex, years in practice, or satisfaction with specialty choice (Appendix Table A1, online only). Validated single-item measures^{34,35} of the emotional exhaustion or depersonalization domains of burnout^{30,31} also failed to identify significant differences. Subsequent analysis focused on participants completing full-length surveys.

The median age of participants was 52 years, with approximately 40% of participants younger than age 50 years (Table 1). Participants were evenly divided by sex. A majority of oncologists (86.2%) were currently married. Independent of relationship status, 86.0% of oncologists reported having children, and nearly half of these oncologists (527; 47.2%) had a child age \leq 18 years (ie, school age).

With respect to practice setting (Table 2), most oncologists were in private practice (PP; 43.2%) or academic practice (AP; 33.8%), with smaller proportions working at a veterans' hospital, in active military practice, or in other settings. Of the 482 oncologists in PP, 335 (69.5%) were in a single-specialty practice, 124 (25.7%) a multispecialty practice, and 23 (4.8%) a health maintenance organization (HMO).

Oncologists spent 57.6 hours per week devoted to professional activities, including an average of 34.0 hours per week on direct patient care, 11.5 hours per week on administrative tasks at work, and 8.5 hours per week performing work tasks at home (completing paperwork, preparing talks, writing grants/manuscripts, and so on), plus 4.6 hours per week keeping abreast of developments in the field and

maintenance of certification. On average, oncologists cared for 52 patients in the outpatient setting each week.

Comparison of PP and AP

Extensive differences in both demographic and practice characteristics were observed between PP and AP oncologists (Tables 1 and 2). Oncologists working in AP settings were slightly younger (median age, 50 v 52 years; $P = .0037$), more likely to be women (57.5% v 45.7%; $P < .001$), and less likely to have children (80.2% v 90.0%; $P < .001$). On average, oncologists in AP worked 4.3 fewer hours each week (58.6 v 62.9 hours; $P < .001$) and devoted less professional effort to direct patient care, with 38.5% in AP spending $\leq 50\%$ of their effort on patient care compared with 3.4% in PP. Oncologists in AP were more likely to focus on treating patients with one specific type of cancer (80.6% v 17.4%; $P < .001$) and spent a greater proportion of their time supervising physicians in training.

Oncologists in PP saw nearly twice as many patients each week, on average, as those in AP (74.2 v 37.4; $P < .001$). Despite the difference in the total number of patients seen per week, the amount of time allocated for each new patient (PP, 52 minutes; AP, 54 minutes; $P = .0011$) and return patient (PP, 18 minutes; AP, 21 minutes; $P < .001$) differed only slightly. The relationship between the number of patients seen per week and percentage of professional effort devoted to clinical care and the number of hours devoted to patient care each week is shown in Figures 1A and 1B.

The method of compensation differed for AP compared with PP, with a larger proportion of PP oncologists in a purely incentive-based model (PP, 39.3% v AP, 3.1%; $P < .001$) and fewer in a salary-only (PP, 20.8% v AP, 38.1%; $P < .001$) or salary-plus-productivity bonus model (PP, 39.9% v AP, 58.8%; $P < .001$). PP oncologists were more likely to report a $> 10\%$ decline in compensation in 2012 relative to 2011 (PP, 35.2% v AP, 8.0%; $P < .001$). Other differences between PP and AP are summarized in Table 2. A subanalysis of PP oncologists according to practice setting (ie, single specialty, multispecialty, HMO) can be found in Appendix Tables A2 and A3 (online only).

Oncologist Well-Being

Table 3 summarizes burnout, fatigue, and career satisfaction among participating oncologists. When assessed using the full MBI, 38.3% of oncologists had high emotional exhaustion, 24.9% had high depersonalization, and 13.2% had a low sense of personal accomplishment. In aggregate, 44.7% of oncologists had at least one symptom of burnout (high emotional exhaustion score and/or high depersonalization). Demographic characteristics associated with burnout on univariable analysis included younger age, being a woman, relationship status, not having children, and greater student loan debt (Appendix Table A4, online only). Professional characteristics associated with burnout on univariable analysis (Appendix Table A5, online only) included hours worked per week, number of hours spent seeing patients per week (Figs 2A and 2B), devoting more time to patient care, seeing a larger number of patients per week, and method of compensation (burnout rates: salary only, 40.7%; salary with bonus, 47.1%; pure incentive, 53.8%; $P = .011$). Although oncologists in PP had higher median emotional exhaustion and depersonalization scores than did those in AP, no difference in the overall burnout rate was observed by practice setting (PP, 50.5% v AP, 45.9%; $P = .177$). A subanalysis of well-being among PP oncologists based on practice setting can be found in Appendix Table A6 (online only).




Fig 1. Hours and effort devoted to clinical care and patients seen per week. The relationship between the number of outpatients seen per week (y-axis) and (A) percentage of effort devoted to clinical care or (B) hours spent seeing patients each week on x-axis.

A majority of oncologists indicated they would choose to become a physician (82.5%) and oncologist (80.5%) again if they could revisit their career and specialty choices. Career satisfaction, as measured by these items, was higher for oncologists in AP than PP (Table 3).

Multivariable Analysis

We performed multivariable analysis to identify personal and professional characteristics associated with burnout and career satisfaction. In addition to an overall model, separate models were developed by practice setting because of the profound differences in personal and professional characteristics of oncologists in PP and AP (Table 4). Younger age and greater number of hours spent seeing patients each week were independently associated with burnout in all models. Each year older reduced the risk of burnout by approximately 4% to 5% (eg, 10 years older, 40% to 50% lower risk), whereas each additional hour spent seeing patients each week increased the risk of burnout by approximately 2% to 4% (eg, 20% to 40% higher risk for

Table 3. Career Satisfaction and Burnout

Characteristic	All (N = 1,117)		AP (n = 377)		PP (n = 482)		P
	No.	%	No.	%	No.	%	
Burnout indices*							
Emotional exhaustion†							
Median		22		22		24	.0895
Low score	433	40.1	146	39.0	157	33.0	.1798
Intermediate score	233	21.6	78	20.9	113	23.7	
High score	413	38.3	150	40.1	206	43.3	
Depersonalization†							
Median		5		5		6	.0124
Low score	558	52.3	191	51.3	220	46.1	.0165
Intermediate score	243	22.8	99	26.6	110	23.1	
High score	265	24.9	82	22.0	147	30.8	
Personal accomplishment							
Median		42		41		42	.0415
High score	660	63.0	225	61.0	304	64.0	.3109
Intermediate score	249	23.8	89	24.1	117	24.6	
Low score‡	138	13.2	55	14.9	54	11.4	
Burned out§	484	44.7	172	45.9	241	50.5	.1769
Career satisfaction							
Would become physician again (career choice)	908	82.5	328	87.5	378	79.2	.0016
Would become oncologist again (specialty choice)	877	80.5	314	85.1	368	77.5	.0053

Abbreviations: AP, academic practice; MBI, Maslach Burnout Inventory; PP, private practice.

*As assessed using the full MBI.

†Per the standard scoring of the MBI for health care workers, physicians with scores ≥ 27 on the emotional exhaustion subscale, ≥ 10 on the depersonalization subscale, or < 33 on the personal accomplishment subscale are considered to have a high degree of burnout in that dimension.

‡Low scores on the personal accomplishment subscale are less favorable.

§High score on emotional exhaustion and/or depersonalization subscales of the MBI (see Methods).

each additional 10 hours). In the overall model, each additional hour per week spent on work-related tasks while at home also increased the risk of burnout by approximately 2% (eg, 10% higher risk for each additional 5 hours per week), and focusing on a specific type of cancer increased the risk of burnout by approximately 40%.

Risks specific to practice setting were also observed. Having children was associated with an approximately 55% decreased risk of burnout among PP oncologists (odds ratio [OR], 0.45) but was not a significant factor for oncologists in AP. In contrast, being a woman was associated with an approximately 65% increased risk of burnout among oncologists in AP (OR, 1.68) but was not a significant factor for oncologists in PP. Each additional hour per week spent on administrative tasks at work increased risk of burnout by approximately 5% among PP oncologists (eg, 5 more hours per week, approximately 25% higher risk), whereas each additional hour per week spent on work tasks at home increased risk by approximately 3.5% among AP oncologists (eg, 5 more hours per week, approximately 17.5% higher risk). Among AP oncologists, focusing on one type of cancer was associated with an increased risk of burnout of 320% (OR, 3.24). For those in AP, having less time allocated for each return patient visit (return slots of 20 minutes in length had a 36% increased risk of burnout compared with return slots of 30 minutes in length) and each additional weekend on call per year also increased risk of burnout.

DISCUSSION

This is the first national study of US oncologists evaluating burnout and career satisfaction to our knowledge since 2003 and is the only

national study to our knowledge to evaluate burnout in US oncologists using standardized instruments. Approximately 45% of oncologists had at least one symptom of burnout at the time of the survey. Although burnout was strongly related to a variety of personal characteristics on univariate analysis, younger age was the only demographic factor independently associated with risk on multivariable analysis adjusting for professional characteristics. In contrast, a variety of professional characteristics were independently associated with burnout. Hours per week devoted to direct patient care was the dominant professional factor associated with burnout. The number of hours per week spent performing work tasks at home and focusing clinical practice on a specific type of cancer were also independently associated with burnout risk.

The strong, incremental relationship between time devoted to patient care and burnout is concerning, especially given the projected shortage in the supply of oncologists during the coming decades. Medical oncologists already work more hours than physicians in most other disciplines.³⁵ Reducing clinical work hours or the volume of patients seen may be a strategy to decrease burnout for individual oncologists but at the societal level could exacerbate the projected oncologist workforce shortage.^{1,14} The findings also suggest that productivity-based compensation models designed to increase the volume of care oncologists provide are associated with higher burnout and may be self-defeating in the long run.

Although the qualitative differences in AP and PP are recognized, the data collected here provide granular information about these differences and explore associations with burnout and career satisfaction. Oncologists in AP were younger, more likely to be women, and less




Fig 2. Patient care activity and burnout. The relationship between patient care hours devoted to patient care and burnout among the 985 oncologists who completed the full-length survey and provided information on both hours and burnout is shown. The number of hours spent seeing patients each week is shown on the x-axis. Mean burnout score in the (A) emotional exhaustion (EE) and (B) depersonalization (DP) domains is shown in the y-axis. Horizontal lines indicate the standardized thresholds to categorize scores for physicians as low, intermediate, or high degree of burnout according to the Maslach Burnout Inventory.

likely to have children. Oncologists in PP saw nearly twice as many patients each week, were more likely to be compensated in a purely incentive-based model, and were less likely to focus their practice on a specific area of oncology. AP oncologists spent far more time on work tasks when at home and dedicated more of their effort to supervising physicians in training. Although no difference in the overall prevalence of burnout was observed by practice setting on multivariable analysis, many of the risk factors for burnout differed between AP and PP oncologists, suggesting that efforts to reduce burnout will need to be tailored to practice setting.

How does the prevalence of burnout among US oncologists compare with that among US adults and physicians in other specialties? A recent national study exploring the prevalence of physician

Table 4. Factors Associated With Burnout on Multivariable Analysis

Predictors	OR	95% CI	P
All oncologists*†‡			
Age (for each additional year older)	0.961	0.947 to 0.975	< .001
Hours/week spent seeing patients (OR each additional hour)	1.032	1.022 to 1.042	< .001
Hours/week at home spent on work tasks (OR each additional hour)	1.019	1.001 to 1.037	.0392
Focus on one certain type of cancer (v multiple focus)	1.422	1.050 to 1.925	.0227
Private practice*†§			
Age (OR each additional year older)	0.953	0.932 to 0.974	< .001
Has children (v not)	0.447	0.210 to 0.950	.0363
Hours/week spent seeing patients (OR each additional hour)	1.041	1.020 to 1.063	< .001
Hours/week spent administrative tasks at work/week (OR each additional hour)	1.054	1.018 to 1.092	.0032
Nights on call/week (OR each additional night)	0.877	0.788 to 0.975	.0152
Academic practice*† 			
Age (OR each additional year older)	0.961	0.935 to 0.987	.0036
Female (v male)	1.678	1.020 to 2.762	.0416
Hours/week spent seeing patients (OR each additional hour)	1.023	1.004 to 1.042	.0190
Hours/week spent at home on work tasks (OR each additional hour)	1.035	1.002 to 1.069	.0363
Minutes allotted for a return outpatient appointment (OR each additional minute)	0.964	0.929 to 1.000	.0494
No. of weekends on call/year (for each additional weekend)	1.071	1.015 to 1.130	.0122
Focus on one certain type of cancer (v multiple focus)	3.244	1.556 to 6.673	.0017

NOTE. Three multivariable analyses were conducted to identify personal and professional factors associated with burnout. The first model included all oncologists. Given substantial differences in professional characteristics, separate models were also created for PP oncologists and AP oncologists. Abbreviations: AP, academic practice; OR, odds ratio; PP, private practice. *Personal characteristics in all models: age, sex, children, youngest child, relationship status, and student loan debt. †Professional characteristics in all models: hours spent seeing patients/week, hours spent on administrative tasks/week, hours spent working at home performing work tasks/week, No. of nights on call per week, No. of outpatients seen/week, focus on certain type of cancer (yes/no), minutes allocated per new outpatient visit, minutes allocated per return outpatient visit, No. of weekends rounding in hospital/year, and method compensation (salary, salary plus bonus, pure incentive). ‡Additional professional characteristics in all oncologist models: practice setting. §Additional professional characteristics in PP model: practice setting (single specialty, multispecialty, health maintenance organization). ||Additional professional characteristics in AP model: percentage of time spent supervising physicians in training.

burnout found that approximately 46% of US physicians were experiencing symptoms of burnout at the time of the study and that the rate of burnout was markedly higher in physicians than in a probability-based sample of US workers.³² Although a subanalysis from that study suggested oncologists may actually have a lower rate of burnout (prevalence of approximately 38%) than other internal medicine physicians, only 87 medical oncologists were included in that analysis.¹ The prevalence of burnout (approximately 45%) in our sample of more than 1,000 oncologists was similar to that of US physicians overall. It was also consistent with rates observed in other internal medicine subspecialists (approximately 44%) and lower than

rates in general internists (approximately 54%).³² Notably, satisfaction with career and specialty choice among oncologists in our study (both > 80%) were the highest of any group of physicians we have studied.^{26,32,36,37}

How do these findings compare with those of previous studies of oncologists? In 1990, Whippen et al¹⁵ sent a 12-item survey with a single question about burnout to 1,000 oncologists who subscribed to *Journal of Clinical Oncology*. Among the 598 respondents, 56% subjectively reported that they felt burned out. In 2003, Allegra et al¹⁶ administered a similar survey to approximately 7,700 US oncologists. Among the 1,740 (23%) who responded, 61.7% endorsed a yes/no question asking, “Do you feel that you are experiencing any signs of burnout?” These historical studies are difficult to interpret because they did not use standardized metrics to assess burnout. The prevalence of high emotional exhaustion (22% to 53%) and high depersonalization (11% to 30%) as measured by the MBI in studies of oncologists from other countries are consistent with the rates of emotional exhaustion (38.3%) and depersonalization (24.9%) observed in our study.^{3,38-41}

Our study is subject to a number of limitations. Although our participation rate of approximately 50% is consistent with⁴² or even higher than^{26,32,36} physician surveys in general, response bias remains a possibility. We found no statistically significant differences with respect to age, sex, years in practice, or career satisfaction among oncologists who completed the postcard survey, further supporting that responders were representative of US oncologists. It should be noted that several previous cross-sectional studies have failed to identify significant differences between responding and nonresponding physicians.⁴³ Because our survey was cross-sectional, we were unable to determine causality or the potential direction of effect for the associations observed. A survival bias may account for some associations such as age (ie, unsatisfied people leave the field). Although we were able to compare differences between oncologists in PP and AP, there were too few participants working in other practice settings to make meaningful comparisons.

Our study also has several important strengths. The oncologists in the sample were drawn from the ASCO oncologist registry, a comprehensive list of US oncologists. The survey included oncologists from all career stages and practice types, as well as a large sample of female oncologists. Our mixed-methods survey design (ie, electronic survey, full-length paper survey, postcard survey) led to a high participation rate relative to other national studies of physicians.^{16,26,36} The

survey collected extensive information on personal and practice characteristics, providing granular insights into relationships among these variables and burnout/career satisfaction.

Given the prevalence of burnout and evidence that it erodes physicians’ personal health^{6,7,44} and the quality of care they provide,^{9-11,45-50} future studies need to focus on how to address this problem. There is currently limited evidence on what interventions reduce the risk of burnout; most available information focuses on individual^{17,18,51-53} rather than system approaches.^{5,54} The high prevalence of burnout suggests that studies evaluating practice models (team-based care) and structural characteristics in the practice environment that may reduce burnout are needed.

In conclusion, the prevalence of burnout among US oncologists seems similar to or lower than that of physicians in other disciplines. Although approximately 45% of oncologists are experiencing burnout, their career and specialty satisfaction are high. The volume of patient care provided seems to be a dominant contributor to burnout for both AP and PP oncologists; however, a number of other contributing factors seem to differ by practice setting. A better understanding of the factors that sustain career satisfaction and studies testing interventions to reduce oncologist burnout are needed.

AUTHORS’ DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The author(s) indicated no potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Conception and design: Tait D. Shanafelt, William J. Gradishar, Michael Kosty, Helen Chew, Leora Horn, Quyen Chu, John Phippen, Jeff Sloan, Marilyn Raymond

Financial support: Tait D. Shanafelt, Jeff Sloan, Marilyn Raymond

Administrative support: Tait D. Shanafelt, Ben Clark, Amy E. Hanley, Jeff Sloan, Marilyn Raymond

Collection and assembly of data: Tait D. Shanafelt, William J. Gradishar, Michael Kosty, Daniel Satele, Helen Chew, Leora Horn, Ben Clark, Amy E. Hanley, Quyen Chu, Jeff Sloan, Marilyn Raymond

Data analysis and interpretation: Tait D. Shanafelt, William J. Gradishar, Michael Kosty, Daniel Satele, Helen Chew, Leora Horn, Quyen Chu, John Phippen, Jeff Sloan, Marilyn Raymond

Manuscript writing: All authors

Final approval of manuscript: All authors

REFERENCES

- Shanafelt T, Dyrbye L: Oncologist burnout: Causes, consequences, and responses. *J Clin Oncol* 30:1235-1241, 2012
- Shanafelt T, Adjei A, Meyskens FL: When your favorite patient relapses: Physician grief and well-being in the practice of oncology. *J Clin Oncol* 21:2616-2619, 2003
- Ramirez AJ, Graham J, Richards MA, et al: Burnout and psychiatric disorder among cancer clinicians. *Br J Cancer* 71:1263-1269, 1995
- Maslach C, Jackson S, Leiter M: *Maslach Burnout Inventory Manual* (ed 3). Palo Alto, CA, Consulting Psychologists Press, 1996
- Shanafelt T, Sloan J, Habermann T: The well-being of physicians. *Am J Med* 114:513-517, 2003

- Shanafelt TD, Balch CM, Dyrbye L, et al: Special report: Suicidal ideation among American surgeons. *Arch Surg* 146:54-62, 2011
- Oreskovich MR, Kaups KL, Balch CM, et al: Prevalence of alcohol use disorders among American surgeons. *Arch Surg* 147:168-174, 2012
- Center C, Davis M, Detre T, et al: Confronting depression and suicide in physicians: A consensus statement. *JAMA* 289:3161-3166, 2003
- Firth-Cozens J, Greenhalgh J: Doctors’ perceptions of the links between stress and lowered clinical care. *Soc Sci Med* 44:1017-1022, 1997
- Shanafelt TD, Balch CM, Bechamps G, et al: Burnout and medical errors among American surgeons. *Ann Surg* 251:995-1000, 2010
- West CP, Huschka MM, Novotny PJ, et al: Association of perceived medical errors with resi-

- dent distress and empathy: A prospective longitudinal study. *JAMA* 296:1071-1078, 2006
- West CP, Tan AD, Habermann TM, et al: Association of resident fatigue and distress with perceived medical errors. *JAMA* 302:1294-1300, 2009
- Dyrbye LN, Massie FS Jr, Eacker A, et al: Relationship between burnout and professional conduct and attitudes among US medical students. *JAMA* 304:1173-1180, 2010
- Shanafelt T, Sloan J, Satele D, et al: Why do surgeons consider leaving practice? *J Am Coll Surg* 212:421-422, 2011
- Whippen DA, Canellos GP: Burnout syndrome in the practice of oncology: Results of a random survey of 1,000 oncologists. *J Clin Oncol* 9:1916-1920, 1991
- Allegra C, Hall R, Yothers G: Prevalence of burnout in the U.S. oncology community: Results of a 2003 survey. *J Oncol Pract* 1:140-147, 2005

17. Shanafelt T, Novotny P, Johnson ME, et al: The well-being and personal wellness promotion practices of medical oncologists in the North Central Cancer Treatment Group. *Oncology (Karger)* 68:23-32, 2005
18. Shanafelt T, Chung H, White H, et al: Shaping your career to maximize personal satisfaction in the practice of oncology. *J Clin Oncol* 24:4020-4026, 2006
19. Rafferty JP, Lemkau JP, Purdy RR, et al: Validity of the Maslach Burnout Inventory for family practice physicians. *J Clin Psychol* 42:488-492, 1986
20. Lee RT, Ashforth BE: A meta-analytic examination of the correlates of the three dimensions of job burnout. *J Appl Psychol* 81:123-133, 1996
21. Leiter M, Durup J: The discriminant validity of burnout and depression: A confirmatory factor analytic study. *Anxiety Stress Coping* 7:357-373, 1994
22. Thomas NK: Resident burnout. *JAMA* 292:2880-2889, 2004
23. Shanafelt TD, Bradley KA, Wipf JE, et al: Burnout and self-reported patient care in an internal medicine residency program. *Ann Intern Med* 136:358-367, 2002
24. Rosen IM, Gimotty PA, Shea JA, et al: Evolution of sleep quantity, sleep deprivation, mood disturbances, empathy, and burnout among interns. *Acad Med* 81:82-85, 2006
25. Dyrbye LN, West CP, Shanafelt TD: Defining burnout as a dichotomous variable. *J Gen Intern Med* 24:440, 2009
26. Kuerer HM, Eberlein TJ, Pollock RE, et al: Career satisfaction, practice patterns and burnout among surgical oncologists: Report on the quality of life of members of the Society of Surgical Oncology. *Ann Surg Oncol* 14:3043-3053, 2007
27. Frank E, McMurray JE, Linzer M, et al: Career satisfaction of US women physicians: Results from the Women Physicians' Health Study—Society of General Internal Medicine Career Satisfaction Study Group. *Arch Intern Med* 159:1417-1426, 1999
28. Lemkau J, Rafferty J, Gordon R Jr: Burnout and career-choice regret among family practice physicians in early practice. *Fam Pract Res J* 14:213-222, 1994
29. Goitein L, Shanafelt TD, Wipf JE, et al: The effects of work-hour limitations on resident well-being, patient care, and education in an internal medicine residency program. *Arch Intern Med* 165:2601-2606, 2005
30. West CP, Dyrbye LN, Sloan JA, et al: Single item measures of emotional exhaustion and depersonalization are useful for assessing burnout in medical professionals. *J Gen Intern Med* 24:1318-1321, 2009
31. West CP, Dyrbye LN, Satele DV, et al: Concurrent validity of single item measures of emotional exhaustion and depersonalization in burnout assessment. *J Gen Intern Med* 27:1445-1445, 2012
32. Shanafelt TD, Boone S, Tan L, et al: Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med* 172:1377-1385, 2012
33. Sloan JA, Cella D, Hays RD: Clinical significance of patient-reported questionnaire data: Another step toward consensus. *J Clin Epidemiol* 58:1217-1219, 2005
34. Sloan JA: Assessing the minimally clinically significant difference: Scientific considerations, challenges and solutions. *COPD* 2:57-62, 2005
35. Leigh JP, Tancredi D, Jerant A, et al: Annual work hours across physician specialties. *Arch Intern Med* 171:1211-1213, 2011
36. Shanafelt TD, Balch CM, Bechamps GJ, et al: Burnout and career satisfaction among American surgeons. *Ann Surg* 250:463-471, 2009
37. Balch CM, Shanafelt TD, Sloan JA, et al: Distress and career satisfaction among 14 surgical specialties, comparing academic and private practice settings. *Ann Surg* 254:558-568, 2011
38. Grunfeld E, Whelan TJ, Zitzelsberger L, et al: Cancer care workers in Ontario: Prevalence of burnout, job stress, and job satisfaction. *Cmaj* 163:166-169, 2000
39. Grunfeld E, Zitzelsberger L, Coristine M, et al: Job stress and job satisfaction of cancer care workers. *Psychooncology* 14:61-69, 2005
40. Arigoni F, Bovier PA, Mermillod B, et al: Prevalence of burnout among Swiss cancer clinicians, paediatricians and general practitioners: Who are most at risk? *Support Care Cancer* 17:75-81, 2009
41. Asai M, Morita T, Akechi T, et al: Burnout and psychiatric morbidity among physicians engaged in end-of-life care for cancer patients: A cross-sectional nationwide survey in Japan. *Psychooncology* 16:421-428, 2007
42. Asch D, Jedrzejewski M, Christakis N: Response rates to mail surveys published in medical journals. *J Clin Epidemiol* 50:1129-1136, 1997
43. Kellerman S, Herold J: Physician response to surveys: A review of the literature. *Am J Prev Med* 20:61-71, 2001
44. Warde CM, Moonesinghe K, Allen W, et al: Marital and parental satisfaction of married physicians with children. *J Gen Intern Med* 14:157-165, 1999
45. Wallace JE, Lemaire JB, Ghali WA: Physician wellness: A missing quality indicator. *Lancet* 374:1714-1721, 2009
46. Grol R, Mookink H, Smits A, et al: Work satisfaction of general practitioners and the quality of patient care. *Fam Pract* 2:128-135, 1985
47. Linn LS, Brook RH, Clark VA, et al: Physician and patient satisfaction as factors related to the organization of internal medicine group practices. *Med Care* 23:1171-1178, 1985
48. Haas JS, Cook EF, Puopolo AL, et al: Is the professional satisfaction of general internists associated with patient satisfaction? *J Gen Intern Med* 15:122-128, 2000
49. Melville A: Job satisfaction in general practice: Implications for prescribing. *Soc Sci Med [Med Psychol Med Sociol]* 14A:495-499, 1980
50. DiMatteo MR, Sherbourne CD, Hays RD, et al: Physicians' characteristics influence patients' adherence to medical treatment: Results from the Medical Outcomes Study. *Health Psychol* 12:93-102, 1993
51. Shanafelt TD, Oreskovich MR, Dyrbye LN, et al: Avoiding burnout: The personal health habits and wellness practices of US surgeons. *Ann Surg* 255:625-633, 2012
52. Krasner MS, Epstein RM, Quill TE, et al: Association of an educational program in mindful communication with burnout, empathy, and attitudes among primary care physicians. *JAMA* 302:1284-1293, 2009
53. Shanafelt TD, West CP, Sloan JA, et al: Career fit and burnout among academic faculty. *Arch Intern Med* 169:990-995, 2009
54. Dunn PM, Arnetz BB, Christensen JF, Homer L: Meeting the imperative to improve physician well-being: Assessment of an innovative program. *J Gen Intern Med* 22:1544-1552, 2007



Appendix

Multivariable Analysis

Predictors included in the logistic model of all oncologists were age, sex, parental status, age of youngest child, relationship status, student loan debt, hours spent per week seeing patients, hours spent per week on administrative tasks, hours spent working at home per week, number of nights on call per week, number of outpatients seen per week, whether oncologist focused on one specific type of cancer, minutes allocated to a new outpatient visit, minutes allocated to a return outpatient visit, number of weekends spent on rounds in hospital per year, method of compensation, and practice setting. The logistic model for private practice oncologists included type of private setting (eg, health maintenance organization, single specialty, multispecialty). The logistic model for academic practice oncologists included percentage of time dedicated to supervising physicians in training.

Table A1. Comparison of Oncologists Who Completed Full-Length With Those Who Completed Postcard Surveys

Characteristic	Full-Length Survey (n = 1,117)		Postcard Survey (n = 373)		P
	No.	%	No.	%	
Median age, years		52		52	.8693
< 40	63	5.8	18	4.9	.8453
40-49	369	34.0	132	36.1	
50-59	343	31.6	115	31.4	
≥ 60	310	28.6	101	27.6	
Sex					.3017
Male	554	50.4	176	47.3	
Female	545	49.6	196	52.7	
Years in practice					
Median		20		19	.6753
< 10	148	14.5	61	16.4	.5950
10-19	337	33.1	126	33.9	
≥ 20	533	52.4	185	49.7	
Would become oncologist again					.9028
Definitely not	30	2.8	11	2.9	
Probably not	77	7.1	27	7.2	
Neutral	105	9.6	37	9.9	
Probably	316	29.0	98	26.3	
Definitely yes	561	51.5	200	53.6	
EE*†					.2930
Never	157	14.5	46	12.4	
A few times a year	286	26.4	100	27.0	
Once a month or less	149	13.8	60	16.2	
A few times a month	174	16.1	73	19.7	
Once a week	108	10.0	29	7.8	
A few times a week	131	12.1	44	11.9	
Every day	78	7.2	19	5.1	
DP*‡					.2936
Never	439	40.8	154	41.3	
A few times a year	301	28.0	120	32.2	
Once a month or less	109	10.1	28	7.5	
A few times a month	83	7.7	33	8.8	
Once a week	65	6.0	14	3.8	
A few times a week	70	6.5	22	5.9	
Every day	9	0.8	2	0.5	

Abbreviations: DP, depersonalization; EE, emotional exhaustion; MBI, Maslach Burnout Inventory; ROC, receiver operating characteristic.

*As assessed using the single-item measures for EE and DP adapted from the full MBI. Area under the ROC curve for the EE and DP single items relative to that of their respective full MBI domain scores in previous studies were 0.94 and 0.93 and the positive predictive values of the single-item thresholds for high levels of EE and DP were 88.2% and 89.6%, respectively.^{30,31}

†Individuals indicating symptoms of EE weekly or more often have median EE scores of > 30 on the full MBI and have a > 75% probability of having a high EE score as defined by the MBI (≥ 27).

‡Individuals indicating symptoms of DP weekly or more often have median DP scores on the full MBI of > 13 and have a > 85% probability of having a high DP score as defined by the MBI (≥ 10).

Table A2. Personal Characteristics Among Different PP Settings

Characteristic	Single Specialty (n = 335)		Multispecialty (N = 124)		HMO (n = 23)		P
	No.	%	No.	%	No.	%	
Age, years							
Median		52.5		52.0		50.0	.1222
Missing		6		5		1	.0505
< 40	10	3.0	12	10.1	2	9.1	
40-49	114	34.7	39	32.8	8	36.4	
50-59	113	34.3	43	36.1	9	40.9	
≥ 60	92	28.0	25	21.0	3	13.6	
Sex							.0070
Missing		1		1		1	
Male	194	58.1	60	48.8	6	27.3	
Female	140	41.9	63	51.2	16	72.7	
Children							.3329
Missing		1		1		1	
Yes	305	91.3	107	87.0	19	86.4	
No	29	8.7	16	13.0	3	13.6	
Age of youngest child, years							.4001
Missing		30		17		4	
< 5	27	8.9	18	16.8	2	10.5	
5-12	86	28.2	29	27.1	6	31.6	
13-18	58	19.0	21	19.6	4	21.1	
19-22	38	12.5	11	10.3	4	21.1	
> 22	96	31.5	28	26.2	3	15.8	
Relationship status							.4851
Missing		1		1		1	
Single	20	6.0	10	8.1	2	9.1	
Married	302	90.4	105	85.4	20	90.9	
Partnered	6	1.8	6	4.9	0	0.0	
Widowed/widower	6	1.8	2	1.6	0	0.0	
Ever gone through divorce							.8326
Missing		3		1		1	
Yes	55	16.6	22	17.9	2	9.1	
No	272	81.9	99	80.5	20	90.9	
Currently going through one	5	1.5	2	1.6	0	0.0	
Current student loan debt							.3009
Missing		4		1		1	
No debt	304	91.8	108	87.8	20	90.9	
Debt < \$25,000	6	1.8	3	2.4	0	0.0	
\$25,000-\$49,999	1	0.3	1	0.8	0	0.0	
\$50,000-\$74,999	8	2.4	1	0.8	2	9.1	
\$75,000-\$99,999	5	1.5	3	2.4	0	0.0	
\$100,000-\$125,000	4	1.2	2	1.6	0	0.0	
> \$125,000	3	0.9	5	4.1	0	0.0	

Abbreviations: HMO, health maintenance organization; PP, private practice.

Career Satisfaction of US Oncologists

Table A3. Comparison of Practice Characteristics Among Different PP* Settings

Characteristic	Single Specialty (n = 335)		Multispecialty (n = 124)		HMO (n = 23)		P
	No.	%	No.	%	No.	%	
Mean years in practice	21.7		19.5		18.8		.1131
General							
Time devoted to patient care, %							.5415
Missing	2		0		0		
None	0	0.0	1	0.8	0	0.0	
1-25	4	1.2	3	2.4	1	4.3	
26-50	4	1.2	2	1.6	1	4.3	
51-75	26	7.8	12	9.7	1	4.3	
76-100	299	89.8	106	85.5	20	87.0	
Focus on specific type cancer							.3162
Missing	12		4		0		
Yes	54	16.7	25	20.8	2	8.7	
No	269	83.3	95	79.2	21	91.3	
Time supervising physicians in training, %							.5871
Missing	19		5		1		
0	166	52.5	63	52.9	12	54.5	
< 5	101	32.0	36	30.3	4	18.2	
5-10	38	12.0	15	12.6	3	13.6	
11-20	7	2.2	3	2.5	2	9.1	
> 20	4	1.3	2	1.7	1	4.5	
Hours and call schedule							
Median nights on call/week	2		2		1		.0309
Hours spent seeing patients at work/week							.4721
Mean	43.7		43.1		40.7		
SD	11.7		12.7		10.0		
Hours spent on administrative tasks at work/week							.5956
Mean	8.9		8.9		7.8		
SD	7.0		6.6		6.9		
Hours spent at home on work tasks/week							.3192
Mean	7.3		7.0		6.0		
SD	7.6		6.1		7.7		
Hours spent at home to keep abreast of developments/week							.2431
Mean	4.4		3.9		3.9		
SD	3.5		3.0		3.1		
Total hours/week†							.2408
Mean	63.7		61.7		57.8		
SD	15.6		18.1		14.1		
Outpatient practice							
Outpatients in clinic per week							< .001
Mean	77.5		68.0		59.5		
SD	33.3		24.0		20.7		
Minutes allocated/new outpatient visit							.2961
Mean	51.2		52.0		53.5		
SD	15.2		14.2		12.7		
Minutes allocated/return outpatient visit							< .001
Mean	16.9		19.5		22.6		
SD	5.5		6.6		6.9		
Hospital practice							
Hospital rounding							< .001
Missing	13		5		0		
Round own patients when hospitalized	93	28.9	25	21.0	0	0.0	
Share rounding with partners in blocks	53	16.5	26	21.8	13	56.5	
Share rounding with partners on weekends	156	48.4	55	46.2	7	30.4	
Attend oncology teaching service	3	0.9	4	3.4	0	0.0	
Do not round in hospital	17	5.3	9	7.6	3	13.0	
Inpatients on average hospital day	5.2	5.5	4.9	4.8	6.3	5.5	.4964
No. of weekends rounding in hospital/year	13.6	12.0	12.0	7.7	8.6	5.3	.0338

(continued on following page)

Table A3. Comparison of Practice Characteristics Among Different PP* Settings (continued)

Characteristic	Single Specialty (n = 335)		Multispecialty (n = 124)		HMO (n = 23)		P
	No.	%	No.	%	No.	%	
Compensation method							< .001
Missing		21		4		1	
Salary no incentive	56	17.8	24	20.0	15	68.2	
Salary with bonus	128	40.8	48	40.0	6	27.3	
Pure incentive	130	41.4	48	40.0	1	4.5	

Abbreviations: HMO, health maintenance organization; PP, private practice; SD, standard deviation.

*Including single-specialty group, multispecialty group, and HMO.

†Sum of above four categories.

Table A4. Personal Characteristics and Burnout

Characteristic	Burned Out (n = 484)		No Burnout (n = 599)		P
	No.	%	No.	%	
Age, years					
Median		50		54	< .001
Missing		10		16	< .001
< 40	36	57.1	27	42.9	
40-49	194	52.9	173	47.1	
50-59	172	51.0	165	49.0	
≥ 60	72	24.8	218	75.2	
Sex					< .001
Missing		2		10	
Male	212	39.8	321	60.2	
Female	270	50.2	268	49.8	
Children					.0057
Missing		2		8	
Yes	398	43.2	523	56.8	
No	84	55.3	68	44.7	
Age of youngest child, years					< .001
Missing		87		78	
< 5	61	51.7	57	48.3	
5-12	125	51.0	120	49.0	
13-18	80	51.0	77	49.0	
19-22	46	43.8	59	56.2	
> 22	85	29.0	208	71.0	
Relationship status					.0456
Missing		1		9	
Single	52	53.6	45	46.4	
Married	410	44.4	514	55.6	
Partnered	17	51.5	16	48.5	
Widowed/widower	4	21.1	15	78.9	
Ever gone through divorce					.3932
Missing		4		10	
Yes	78	41.7	109	58.3	
No	397	45.4	477	54.6	
Currently going through one	5	62.5	3	37.5	
Current student loan debt					.0023
Missing		4		9	
No debt	410	42.8	547	57.2	
Debt < \$25,000	15	57.7	11	42.3	
\$25,000-\$49,999	6	46.2	7	53.8	
\$50,000-\$74,999	21	75.0	7	25.0	
\$75,000-\$99,999	12	75.0	4	25.0	
\$100,000-\$125,000	5	50.0	5	50.0	
> \$125,000	11	55.0	9	45.0	

Career Satisfaction of US Oncologists

Table A5. Practice Characteristics and Burnout

Characteristic	Burned Out (n = 484)		No Burnout (n = 599)		P
	No.	%	No.	%	
General					
Time devoted to patient care, %					< .001
Missing		1		5	
None	10	15.9	53	84.1	
1-25	24	25.8	69	74.2	
26-50	38	31.4	83	68.6	
51-75	99	50.3	98	49.7	
76-100	312	51.7	291	48.3	
Focus on specific type cancer					.3804
Missing		15		19	
Yes	192	46.4	222	53.6	
No	277	43.6	358	56.4	
Time supervising physicians in training, %					.2539
Missing		22		46	
0	154	41.5	217	58.5	
< 5	114	50.2	113	49.8	
5-10	88	48.6	93	51.4	
11-20	60	46.2	70	53.8	
> 20	46	43.4	60	56.6	
Hours and call schedule					
Mean nights spent on call/week		2.2		1.8	< .001
Hours spent seeing patients at work/week					< .001
Mean		39.3		30.6	
SD		15.3		17.1	
Hours spent on administrative tasks at work/week					.1508
Mean		11.4		11.7	
SD		9.2		11.4	
Hours spent at home on work tasks/week					.0081
Mean		9.0		8.2	
SD		8.4		8.9	
Hours spent at home keeping abreast of development/week					< .001
Mean		4.1		4.9	
SD		3.6		4.1	
Total hours/week*					< .001
Mean		63.4		54.3	
SD		16.7		21.5	
Outpatient practice					
Outpatients in clinic/week					< .001
Mean		60.0		46.2	
SD		32.8		33.7	
Minutes allocated/new outpatient visit					.0021
Mean		52.3		47.7	
SD		16.2		21.7	
Minutes allocated/return outpatient visit					.5737
Mean		18.8		18.0	
SD		6.6		8.9	
Hospital practice					
Hospital rounding					< .001
Missing		17		20	
Round own patients when hospitalized	77	48.1	83	51.9	
Share rounding with partners in blocks	92	54.1	78	45.9	
Share rounding with partners on weekends	155	51.0	149	49.0	
Attend oncology teaching service	106	43.1	140	56.9	
Do not round in hospital	37	22.3	129	77.7	
Inpatients on average hospital day	7.9	7.4	6.5	7.2	< .001
No. of weekends rounding in hospital/year	10.9	9.0	9.2	10.3	< .001
Compensation method					
Missing		21		63	.0105
Salary no incentive	135	40.7	197	59.3	
Salary with bonus	216	47.1	243	52.9	
Pure incentive	112	53.8	96	46.2	

Abbreviation: SD, standard deviation.
*Sum of above four categories.

Table A6. Career Satisfaction, Burnout, and Quality of Life Among Different PP Settings

Characteristic	Single Specialty (n = 334)		Multispecialty (n = 122)		HMO (n = 23)		P
	No.	%	No.	%	No.	%	
Burnout indices*							
Emotional exhaustion†							
Median		24		26		26	.5692
Low score	114	34.5	36	29.3	7	30.4	.7380
Intermediate score	80	24.2	28	22.8	5	21.7	
High score	136	41.2	59	48.0	11	47.8	
Depersonalization†							
Median		6		7		5	.2251
Low score	158	47.7	49	39.8	13	56.5	.4034
Intermediate score	71	21.5	34	27.6	5	21.7	
High score	102	30.8	40	32.5	5	21.7	
Personal accomplishment‡							
Median		42		42		41	.5931
Low score	215	65.3	76	61.8	13	56.5	.6827
Intermediate score	75	22.8	35	28.5	7	30.4	
High score	39	11.9	12	9.8	3	13.0	
Burned out§	162	48.9	68	55.3	11	47.8	.4692
Career satisfaction							
Would become physician again (career choice)	260	78.3	101	82.1	17	77.3	.6561
Would become oncologist again (specialty choice)	262	78.9	92	76.0	14	63.6	.2284

Abbreviations: AP, academic practice; HMO, health maintenance organization; MBI, Maslach Burnout Inventory; PP, private practice.

*As assessed using the full MBI.

†Per the standard scoring of the MBI for health care workers, physicians with scores ≥ 27 on the emotional exhaustion subscale, ≥ 10 on the depersonalization subscale, or < 33 on the personal accomplishment subscale are considered to have a high degree of burnout in that dimension.

‡Low scores on the personal accomplishment subscale are less favorable.

§High score on emotional exhaustion and/or depersonalization subscales of the MBI (see Methods).