



Published in final edited form as:

Biochim Biophys Acta. 2014 January ; 1845(1): 84–89. doi:10.1016/j.bbcan.2013.12.002.

Doxorubicin, DNA torsion, and chromatin dynamics

Fan Yang^a, Sheila S. Teves^a, Christopher J. Kemp^{b,*}, and Steven Henikoff^{a,c,*}

^aBasic Science Division, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA

^bHuman Biology Division, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA

^cHoward Hughes Medical Institute, Seattle, Washington 98109, US

Abstract

Doxorubicin is one of the most important anti-cancer chemotherapeutic drugs, being widely used for the treatment of solid tumors and acute leukemias. The action of doxorubicin and other anthracycline drugs has been intensively investigated during the last several decades, but the mechanisms that have been proposed for cell killing remain disparate and controversial. In this review, we examine the proposed models for doxorubicin action from the perspective of the chromatin landscape, which is altered in many types of cancer due to recurrent mutations in chromatin modifiers. We highlight recent evidence for effects of anthracyclines on DNA torsion and chromatin dynamics that may underlie basic mechanisms of doxorubicin-mediated cell death and suggest new therapeutic strategies for cancer treatment.

Keywords

doxorubicin; anthracycline; cancer; DNA torsion; chromatin dynamics; chemotherapy

1. Introduction

Doxorubicin (also called adriamycin) belongs to a class of compounds with similar structures, called anthracyclines. Like daunorubicin, the first anthracycline compound to be described, doxorubicin was isolated from *Streptomyces peucetius*, a soil bacterium [1, 2]. Doxorubicin has shown great efficacy in cancer cell killing for both solid and liquid tumors, but the emergence of drug resistance and potential side effects such as heart muscle damage after doxorubicin treatment are major limitations for successful cancer treatment [3]. Despite the extensive usage in the clinics, the molecular mechanism(s) by which doxorubicin causes cell death or cardiotoxicity remains unclear. Thus, understanding the action of doxorubicin and related anthracycline drugs may provide clues for enhancing cancer cell killing and reducing side effects. A number of models have been proposed for doxorubicin-mediated cell death, including topoisomerase II poisoning, DNA adduct formation, oxidative stress, and ceramide overproduction [4–6]. However, the models remain disparate and controversial. A deeper understanding of the basic molecular interactions of doxorubicin

*Corresponding authors: C.J. Kemp: Tel.: +1 (206) 6674252; fax: +1 (206) 6675815. S. Henikoff: Tel.: +1 (206) 6674515; fax: +1 (206) 6675889, cjtemp@fhrc.org (C.J. Kemp); steveh@fhrc.org (S. Henikoff).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

within the cell is required to understand how doxorubicin kills cancer cells and causes side effects.

Anthracycline drugs such as doxorubicin are mostly planar molecules that preferentially intercalate between neighboring DNA base pairs, anchored on one side by one or more sugar moieties that sit in the DNA minor groove (Fig. 1). When DNA is topologically constrained, as in the case of plasmid circles, the strand separation that occurs during intercalation unwinds the double helix and produces DNA supercoils, resulting in increased torsional stress. Linear genomes of eukaryotes are partitioned into independent topological domains by protein factors such as insulator binding protein CTCF [7], so each domain is topological constrained. *In vitro* studies suggest that torsional stress can affect the structure and dynamics of nucleosomes, the repeating unit of chromatin composed of DNA wrapped around octameric histone cores [8, 9]. Interestingly, recent *in vivo* studies implicate doxorubicin in nucleosome eviction and replacement [10, 11]. Taken together, torsion-induced nucleosome destabilization is emerging as a significant molecular mechanism for the action of doxorubicin and related anthracycline drugs.

2. Models for doxorubicin-mediated cell death

A number of mechanisms have been proposed for doxorubicin-mediated cell death. However, some of these such as inhibition of DNA and RNA synthesis are only seen at doses higher than the clinical dose (~ 40 to 60 mg/m²) [4] (Table 1). Here, we examine the proposed mechanisms for doxorubicin action in clinically relevant drug doses.

2.1. Topoisomerase II poisoning

Topoisomerases are highly conserved enzymes that are present in virtually all life forms, from bacteria to humans, and they regulate DNA topology to facilitate DNA replication, transcription, and other nuclear processes. Many anticancer and antibacterial drugs target topoisomerases for cell killing, such as camptothecins, etoposide, and quinolones [12]. The most parsimonious model for doxorubicin action involves topoisomerase II poisoning, resulting in double-strand DNA breaks and cell death at clinically relevant drug concentrations [3, 4]. Topoisomerase II is an ATP-dependent enzyme that exists in two isoforms in humans, topoisomerase II α and topoisomerase II β . The enzyme binds DNA supercoils and entangled DNA, breaks both strands of one DNA duplex, passes the other duplex through the resulting gap and reseals the break. This process results in the release of torsional stress formed during biological processes such as DNA replication and transcription (discussed below) [12]. In addition, topoisomerase II is essential for decatenation of DNA during mitosis, and deficiency in topoisomerase II prevents normal cytokinesis resulting in cell death [13]. Etoposide, a topoisomerase II poison, traps topoisomerase II at breakage sites, stabilizes the cleavage complex and impedes DNA resealing [14]. Doxorubicin has been hypothesized to function in a similar way [15] and it has been shown that topoisomerase II levels determine the effectiveness of doxorubicin treatment in a mouse model of lymphoma [16]. However, there are many examples in which doxorubicin-mediated cell killing is independent of topoisomerase II. For example, doxorubicin was shown to cause cell death independent of topoisomerase II in a promyelocytic leukemic cell line [17]. In addition, doxorubicin as well as another anthracycline drug, aclarubicin, which does not trap topoisomerase II, evicts histones independent of topoisomerase II leading to cell death [10, 18]. These findings suggest that anthracycline-induced topoisomerase II poisoning by trapping topoisomerase II at cleavage sites is unlikely to be the only mechanism of cancer cell killing by anthracycline drugs.

The anti-cancer activity of doxorubicin is attributable to killing of dividing cells, where topoisomerase II α is the major form of the enzyme. However, heart muscle failure is a side

effect that results from damage to non-dividing cells, where topoisomerase II β is the major form. Indeed, cardiomyocyte-specific deletion of topoisomerase II β has been shown to protect mice from developing doxorubicin-induced heart failure [19]. Inhibitors of topoisomerase II have also been shown to protect cardiomyocytes from doxorubicin-induced toxicity [20]. These findings suggest that trapping topoisomerase II β by doxorubicin in non-dividing heart cells underlies doxorubicin-induced cardiotoxicity.

2.2. DNA adduct formation

As a DNA intercalator, doxorubicin prefers the intercalation site containing adjacent GC base pairs, probably due to specific hydrogen-bond formation between doxorubicin and guanine (Fig. 1a) [21–23]. Formation of doxorubicin-DNA adducts has been shown to activate DNA damage responses and induce cell death independent of topoisomerase II [17, 24]. Importantly, doxorubicin-DNA adducts are detectable at clinically relevant drug concentrations, suggesting that doxorubicin-DNA adducts form during chemotherapy [25]. The interaction between doxorubicin and DNA can be stabilized by a covalent bond mediated by cellular formaldehyde that is generated by free radical reactions from carbon sources such as lipids and spermine [26, 27]. This interaction involves formation of a covalent bond between doxorubicin and guanine on one strand of DNA mediated by formaldehyde and of a hydrogen bond between doxorubicin and guanine on the opposing strand (Fig. 1a). Interestingly, higher levels of formaldehyde have been detected in doxorubicin-sensitive tumor cells compared to resistant tumor cells and normal cells [28, 29]. The formation of more doxorubicin-DNA adducts might contribute to increased effectiveness of the drug in doxorubicin-sensitive tumor cells.

The discovery of the covalent doxorubicin-DNA adduct has led to a new approach to improve the anticancer activity of doxorubicin. Compounds that release formaldehyde upon hydrolysis, such as pivaloyloxymethyl butyrate (AN-9), butyroyloxymethyl-diethyl phosphate (AN-7), and hexamethylenetetramine (HMTA), have been developed and utilized in combination with doxorubicin. AN-9 showed a synergy with doxorubicin in cancer cell killing as well as in overcoming doxorubicin resistance to varying degrees by increasing DNA adduct levels [30, 31]. In addition, the combination of AN-7 and doxorubicin has been shown to enhance its anticancer activity, to protect against doxorubicin-induced toxicity in neonatal rat cardiomyocytes, and to prevent weight loss in mice [32, 33].

Despite the evidence that DNA adducts form during doxorubicin treatment, DNA adduct formation is unlikely to be the major mechanism of doxorubicin action, because clinical doses result in only 4.4 ± 1.0 adducts/ 10^7 base pair DNA, which accounts for just a small fraction of total doxorubicin [25].

2.3. Oxidative stress

The quinone structure of doxorubicin can be oxidized to a semiquinone radical through addition of one electron, mediated by a number of NAD(P)H-oxidoreductases [5, 34]. Semiquinone radicals quickly react with oxygen to generate superoxide and hydrogen peroxide causing DNA damage. Additionally, doxorubicin is an iron chelator and the doxorubicin-iron complex catalyzes the conversion of hydrogen peroxide to highly reactive hydroxyl radicals [35]. Thus, doxorubicin-induced release of free radicals may cause oxidative stress, resulting in DNA damage and cell death [3].

A study using gas chromatography/mass spectrometry has shown that after 72 to 96 hours of slow intravenous infusion of doxorubicin with a steady level of $0.1 \mu\text{M}$ in the plasma, DNA base oxidation increased up to 4-fold in peripheral blood mononuclear cells from breast cancer patients [36]. Oxidized DNA bases were also detected in a cardiac cell line and

normal breast epithelial cells [37, 38]. These findings suggest that free radical-induced DNA damage indeed happens early after doxorubicin treatment. If free radical formation accounts for doxorubicin-mediated cell killing, free radical scavengers would rescue cell death after doxorubicin treatment. Free radical scavengers have been used in the clinic as protectants against doxorubicin-induced cardiotoxicity but many of them failed, suggesting that free radical formation is not the only mechanism of cardiotoxicity [39].

In addition to direct DNA damage by free radical formation, administration of low doses of doxorubicin can result in increased levels of oxidative metabolism [40], which might have multiple effects on components of the chromatin landscape. For example, 2-oxoglutarate is involved in redox cycling reactions in cellular metabolism, and is also the cofactor for JmJc domain-containing histone lysine demethylases [41]. Another enzyme co-factor that is regulated by oxidative metabolism is NAD(P), which is the substrate for ADP-ribosylation reactions, including ADP-ribosylation of histones [42]. Although it is an intriguing possibility that the effects on chromatin by alterations in these enzymes resulting from doxorubicin-induced oxidative changes can lead to DNA damage, other possibilities need to be considered. For example, when a histone acetyl group is removed by a histone deacetylase (HDAC), the acetate anion that is released is co-exported with a proton out of the cell, thus incrementally raising the intracellular pH (pH_i) [43]. The hyperacetylation of histones that results from administration of histone deacetylase inhibitors prevents release of acetate and protons, thus lowering pH_i and potentially decreasing cancer cell survivability in an acidic environment. So although HDAC inhibitors have dramatic effects on global histone lysine acetylation levels and can alter the chromatin landscape, these chromatin effects might not be relevant to the anti-cancer effect of the drug. By the same reasoning, potential effects of doxorubicin on histone modification might have as-yet undiscovered metabolic effects that are independent of their effects on the chromatin landscape.

2.4. Ceramide overproduction

In addition to the generation of free radicals and the increase in oxidative metabolism, doxorubicin treatment increases ceramide levels [44–47]. Ceramide is a lipid molecule consisting of a sphingosine and a fatty acid that is involved in a variety of cellular processes including growth arrest, apoptosis, and senescence [6]. Interestingly, exogenous cell-permeable ceramide sensitizes cancer cells to doxorubicin-induced cell death [48]. In addition, doxorubicin treatment increases ceramide levels in doxorubicin-sensitive MCF-7 cells but not in doxorubicin-resistant MCF-7-AdrR cells, suggesting that ceramide levels might mediate doxorubicin resistance. Indeed, up-regulation of glucosylceramide synthase (GCS), an enzyme that converts ceramide to glucosylceramide, is associated with cellular resistance to doxorubicin, whereas suppression of GCS restores sensitivity to doxorubicin resulting in cell death [49–52]. Recently, doxorubicin has been reported to have a new role in blocking proliferation of cancer cells through stimulation of ceramide synthesis and enhancement of proteolysis of a membrane-bound protein CREB3L1 [53]. Interestingly, CREB3L1 expression level is associated with cellular sensitivity to doxorubicin, although its expression is not required for other anti-cancer drugs such as etoposide, bleomycin, or paclitaxel to inhibit cell growth. Surprisingly, no cell death was observed after doxorubicin treatment in this study [53]. It is possible that inhibition of cell proliferation makes cells less susceptible to doxorubicin, as doxorubicin preferentially kills dividing cells. These findings suggest that ceramide overproduction might be specifically involved in sensitizing cancer cells to doxorubicin treatment, but this upstream effect is unlikely to involve changes in the chromatin landscape.

3. Doxorubicin, DNA torsion, and chromatin structure

The disparate models described above underscore the controversies surrounding the multiple modes of doxorubicin action within the cell. Next we consider the possibility that the intercalation of doxorubicin between DNA bases has a direct effect on chromatin that ultimately results in cancer cell killing. Specifically, we focus on the consequences of doxorubicin intercalation into DNA and ask what effect if any this event might have on chromatin structure and dynamics.

3.1. DNA topology and chromatin

When an anthracycline molecule enters the cell, it diffuses into the nucleus and intercalates into DNA. Anthracyclines are bound to DNA so stably that daunorubicin autofluorescence has been used by cytogenetics for Q-banding chromosomes, whereby AT-rich regions fluoresce brightly and GC-rich regions quench [54]. Intercalation of anthracyclines pushes apart the flanking base pairs (Fig. 1b). Because each base pair is stacked in 36° counterclockwise rotation relative to the pair below, intercalation has topological consequences. In covalently closed circular DNA, the two strands are wound around each other a certain number of times, called the linking number (Lk). When the DNA circle is relaxed, Lk is equal to the number of turns in the double helix, or the twist (Tw), which is roughly 1 turn per 10.5 base pairs. Changes in the relaxed value of Lk results in torsional stress that manifests itself as changes in Tw and/or writhe (Wr), which is the number of times the double-stranded DNA crosses itself, commonly known as supercoiling. Writhe can be described as positive or negative, depending on whether supercoiling occurs in right- or left-handed direction, respectively.

These topological aspects are most evident in bacteria, whose circular genomes exist in a tightly regulated topological state. However, linear genomes of eukaryotes can also be described using the same topological terms, as the genome is partitioned into independent topological domains whose borders are restricted [7]. Thus, changes in Lk, Tw, and Wr in one domain do not transfer to another. The wrapping of the DNA around the octameric histones in a left-handed direction generates one negative Wr per nucleosome. Thus, nucleosomes constrain negative supercoiling.

3.2. Torsional stress and nucleosome destabilization

Cellular processes that require access to DNA inevitably alter its topological state, producing torsional stress. These processes include DNA replication, transcription, recombination, and repair. During replication, helicases that separate the two strands alter the Lk and produce torsional stress, resulting in waves of positive supercoiling downstream of the helicase. Similarly, transcription by RNA Polymerase II (RNAPII) results in denaturation of DNA at the transcription bubble. The subsequent rotation of the DNA relative to RNAPII during transcription creates domains of positive and negative supercoils downstream and upstream, respectively, as predicted by the twin-supercoiled domain model [55, 56]. Furthermore, enzymes that remove or replace nucleosomes alter the supercoiling levels. For example, the nucleosome remodeler SWI/SNF is known to generate negative supercoils [57, 58].

Torsional stress can be detrimental, affecting the overall structure and integrity of DNA. Furthermore, changes in Tw and/or Wr can deregulate processes such as replication, transcription, and nucleosome stability (discussed below). For example, accumulation of positive torsion leads to transcriptional inhibition in over 80% of all genes in yeast [59, 60]. To counteract torsional stresses generated during various DNA-based processes, cells utilize topoisomerases, thereby altering Lk. As discussed above, doxorubicin can inhibit

topoisomerase II in the cleaved form and thus directly cause double-strand breaks. However, as a DNA intercalator, doxorubicin primarily alters DNA topology. Single molecule measurements show that intercalation of one doxorubicin molecule relaxes the natural twist of the double helix by -27° [61], suggesting significant underwinding in the presence of the drug. Such a change in Tw in the negative direction introduces compensatory positive torsional strain on the DNA. Increases in torsional stress can affect many different processes, but because over 80% of eukaryotic DNA is complexed in nucleosomes, perhaps the most immediate effect is directly on nucleosome structure and dynamics.

Nucleosomes are quite sensitive to torsional changes. Single molecule *in vitro* studies show that nucleosome assembly stalls when DNA is under positive torsional stress [62] whereas negative supercoiling promotes assembly [63]. Another *in vitro* study showed a preferential exchange of nucleosomes from positively to negatively supercoiled DNA [64]. Furthermore, the presence of positive torsional stress seems to induce a structural change in the nucleosome, presumably resulting in a more open complex where the H2A/H2B dimers somewhat dissociate from the tetramer core [8, 9]. This kind of restructuring may render the nucleosome less stable.

3.3. Doxorubicin and nucleosome dynamics

The unwinding of DNA upon doxorubicin intercalation may produce sufficient positive torsional stress to destabilize nucleosomes, which would serve as a direct mechanism for action of the drug. Indeed, a study from our laboratory tested this hypothesis using a recently developed strategy to measure nucleosome turnover, the disassembly and subsequent reassembly of nucleosomes. Using metabolic labeling of newly synthesized proteins followed by affinity purification, newly incorporated H3/H4 core particles can be analyzed by tiling array or next generation sequencing [65, 66]. The method is called covalent attachment of tags to capture histones and identify turnover, CATCH-IT. By comparing nucleosome turnover profiles in mouse squamous cell carcinoma cells before and after doxorubicin treatment at a concentration of $0.34 \mu\text{M}$, we found that doxorubicin enhances nucleosome turnover around active gene promoters, despite a minor effect on gene expression level. This enhancement is independent of the DNA damage response proteins, p53 and ATM. This latter point is important as p53 is mutated in many human cancers and both ATM and p53 have been implicated in the response of tumors to doxorubicin therapy [67, 68]. Interestingly, a similar effect on nucleosome turnover was observed by treating the cells with another anthracycline drug, aclarubicin, which inhibits topoisomerase II without causing DNA double strand breaks [18] indicating that anthracycline drug intercalation into DNA may play a direct role in enhancing nucleosome turnover [11]. Another study showed that anthracycline drugs including doxorubicin and aclarubicin, but not etoposide, evict histones from regions of accessible chromatin in both human melanoma cell lines and acute myeloid leukemia blasts from patients, leading to impairment of DNA repair and apoptosis [10]. Additionally, this study found that $9 \mu\text{M}$ doxorubicin evicts histones in topoisomerase II α -depleted cells, suggesting that topoisomerase II α is not required for doxorubicin-induced histone eviction [10]. Taken together, these studies suggest that inhibition of topoisomerase II could further exacerbate torsional strain by anthracycline drug intercalation, leading to enhancement of nucleosome turnover downstream of promoters.

Topoisomerase inhibitors can have dramatic effects on chromatin-associated processes, including replication and transcription. The positive torsion generated ahead of RNA polymerase [55] can potentially unwrap nucleosomes, which are negatively supercoiled, and destabilize them [69]. We have found that inhibitors of both topoisomerase I and II cause both increased positive torsion and increased nucleosome turnover in gene bodies and vice-versa in intergenic regions of *Drosophila* cells [70]. The net effect of topoisomerase-

mediated enhancement of torsion in gene bodies is to further destabilize nucleosomes during RNA polymerase transit, thus increasing exposure of DNA to processes that can cause DNA breaks, such as free radicals, and ultimately cell death. Nucleosome destabilization might also be promoted by SWI/SNF-class remodelers which act to evict nucleosomes at promoters, countered by ISWI/CHD-class remodelers that act to stabilize nucleosomes in gene bodies [71]. In this way, doxorubicin intercalation into promoters and genes might act by interfering with the balance between maintaining promoters free of nucleosomes while preventing loss of nucleosomes during transcription. This model may account for the observation that anthracycline drugs, including aclarubicin, increase somatic recombination in a *Drosophila in vivo* assay for DNA damage [72].

4. Future directions

Over the last several decades, many different modes of doxorubicin action have been reported, which is consistent with the broad spectrum of activity of the drug in cancer treatment. As discussed above, intercalation of doxorubicin into DNA leading to torsional stress and nucleosome destabilization may account for the basic mechanism of anthracycline-mediated cell killing. Surveying the effects of other intercalating drugs on chromatin structure and dissecting the mechanistic link between this and cancer cell killing may lead to the development of better anticancer drugs. Interestingly, another intercalating drug, voreloxin, a quinolone derivative, has been recently found to act similarly to anthracyclines in killing cancer cells, and intercalation is required for voreloxin-mediated cell death [73]. Voreloxin holds promise as an alternative to anthracyclines, as it does not generate high levels of reactive oxygen species, which can contribute to anthracycline-mediated cardiotoxicity [39, 73].

The fact that relatively high doses of anthracyclines (9 μM) used in one study caused histone eviction and apoptosis [10], whereas low sublethal doses (0.1–0.4 μM) used in another study nevertheless caused global nucleosome turnover and a DNA damage response [11], raises the possibility that downward dosage adjustments might be considered for current clinical protocols based on the maximum tolerated dose. The effect of anthracyclines on destabilizing nucleosomes also suggests that combining anthracyclines and other drugs that destabilize nucleosomes may have synergistic effects in cancer cell killing. For example, valproic acid, an HDAC inhibitor approved for clinical use as an anticonvulsant and mood-stabilizing drug, increases the activity of doxorubicin and leads to tumor regression and chromatin decondensation when used in combination with another anthracycline drug, epirubicin [74–76]. The development of new drugs that destabilize nucleosomes have the potential of enhancing the therapeutic efficacy of anthracycline drugs while reducing side effects.

Acknowledgments

We thank S. Ramachandran for helping generate the figure of structure of doxorubicin-DNA adduct. This work was supported by NIH grant R01 ES020116 (S.H. and C.J.K.), NIH grant U54 CA143862 (S.H.), the Howard Hughes Medical Institute (S.H.), and NSF Graduate Research Fellowship DGE-0718124 (S.S.T.).

References

1. Arcamone F, Cassinelli G, Fantini G, Grein A, Orezzi P, Pol C, Spalla C. Adriamycin, 14-hydroxydaunomycin, a new antitumor antibiotic from *S. peucetius* var. *caesius*. *Biotechnol Bioeng.* 1969; 11:1101–1110. [PubMed: 5365804]
2. Di Marco A, Gaetani M, Scarpinato B. Adriamycin (NSC-123,127): a new antibiotic with antitumor activity. *Cancer Chemother Rep.* 1969; 53:33–37. [PubMed: 5772652]

3. Thorn CF, Oshiro C, Marsh S, Hernandez-Boussard T, McLeod H, Klein TE, Altman RB. Doxorubicin pathways: pharmacodynamics and adverse effects. *Pharmacogenet Genomics*. 2011; 21:440–446. [PubMed: 21048526]
4. Gewirtz DA. A critical evaluation of the mechanisms of action proposed for the antitumor effects of the anthracycline antibiotics adriamycin and daunorubicin. *Biochem Pharmacol*. 1999; 57:727–741. [PubMed: 10075079]
5. Minotti G, Menna P, Salvatorelli E, Cairo G, Gianni L. Anthracyclines: molecular advances and pharmacologic developments in antitumor activity and cardiotoxicity. *Pharmacol Rev*. 2004; 56:185–229. [PubMed: 15169927]
6. Senchenkov A, Litvak DA, Cabot MC. Targeting ceramide metabolism—a strategy for overcoming drug resistance. *J Natl Cancer Inst*. 2001; 93:347–357. [PubMed: 11238696]
7. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B. Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature*. 2012; 485:376–380. [PubMed: 22495300]
8. Bancaud A, Conde e Silva N, Barbi M, Wagner G, Allemand JF, Mozziconacci J, Lavelle C, Croquette V, Victor JM, Prunell A, Viovy JL. Structural plasticity of single chromatin fibers revealed by torsional manipulation. *Nat Struct Mol Biol*. 2006; 13:444–450. [PubMed: 16622406]
9. Bancaud A, Wagner G, Conde ESN, Lavelle C, Wong H, Mozziconacci J, Barbi M, Sivolob A, Le Cam E, Mouawad L, Viovy JL, Victor JM, Prunell A. Nucleosome chiral transition under positive torsional stress in single chromatin fibers. *Mol Cell*. 2007; 27:135–147. [PubMed: 17612496]
10. Pang B, Qiao X, Janssen L, Velds A, Groothuis T, Kerkhoven R, Nieuwland M, Ovaa H, Rottenberg S, van Tellingen O, Janssen J, Huijgens P, Zwart W, Neefjes J. Drug-induced histone eviction from open chromatin contributes to the chemotherapeutic effects of doxorubicin. *Nat Commun*. 2013; 4:1908. [PubMed: 23715267]
11. Yang F, Kemp CJ, Henikoff S. Doxorubicin Enhances Nucleosome Turnover around Promoters. *Curr Biol*. 2013; 23:782–787. [PubMed: 23602475]
12. Pommier Y, Leo E, Zhang H, Marchand C. DNA topoisomerases and their poisoning by anticancer and antibacterial drugs. *Chem Biol*. 2010; 17:421–433. [PubMed: 20534341]
13. Carpenter AJ, Porter AC. Construction, characterization, and complementation of a conditional-lethal DNA topoisomerase II α mutant human cell line. *Mol Biol Cell*. 2004; 15:5700–5711. [PubMed: 15456904]
14. Wu CC, Li TK, Farh L, Lin LY, Lin TS, Yu YJ, Yen TJ, Chiang CW, Chan NL. Structural basis of type II topoisomerase inhibition by the anticancer drug etoposide. *Science*. 2011; 333:459–462. [PubMed: 21778401]
15. Nitiss JL. Targeting DNA topoisomerase II in cancer chemotherapy. *Nat Rev Cancer*. 2009; 9:338–350. [PubMed: 19377506]
16. Burgess DJ, Doles J, Zender L, Xue W, Ma B, McCombie WR, Hannon GJ, Lowe SW, Hemann MT. Topoisomerase levels determine chemotherapy response in vitro and in vivo. *Proc Natl Acad Sci U S A*. 2008; 105:9053–9058. [PubMed: 18574145]
17. Swift LP, Rephaeli A, Nudelman A, Phillips DR, Cutts SM. Doxorubicin-DNA adducts induce a non-topoisomerase II-mediated form of cell death. *Cancer Res*. 2006; 66:4863–4871. [PubMed: 16651442]
18. Sorensen BS, Sinding J, Andersen AH, Alsner J, Jensen PB, Westergaard O. Mode of action of topoisomerase II-targeting agents at a specific DNA sequence. Uncoupling the DNA binding, cleavage and religation events. *J Mol Biol*. 1992; 228:778–786. [PubMed: 1335085]
19. Zhang S, Liu X, Bawa-Khalfe T, Lu LS, Lyu YL, Liu LF, Yeh ET. Identification of the molecular basis of doxorubicin-induced cardiotoxicity. *Nat Med*. 2012; 18:1639–1642. [PubMed: 23104132]
20. Vavrova A, Jansova H, Mackova E, Machacek M, Haskova P, Tichotova L, Sterba M, Simunek T. Catalytic Inhibitors of Topoisomerase II Differently Modulate the Toxicity of Anthracyclines in Cardiac and Cancer Cells. *PLoS One*. 2013; 8:e76676. [PubMed: 24116135]
21. Chaires JB, Herrera JE, Waring MJ. Preferential binding of daunomycin to 5'ATCG and 5'ATGC sequences revealed by footprinting titration experiments. *Biochemistry*. 1990; 29:6145–6153. [PubMed: 2207063]

22. Chaires JB, Fox KR, Herrera JE, Britt M, Waring MJ. Site and sequence specificity of the daunomycin-DNA interaction. *Biochemistry*. 1987; 26:8227–8236. [PubMed: 2831939]
23. Chen KS, Gresh N, Pullman B. A theoretical investigation on the sequence selective binding of adriamycin to double-stranded polynucleotides. *Nucleic Acids Res*. 1986; 14:2251–2267. [PubMed: 3960721]
24. Forrest RA, Swift LP, Rephaeli A, Nudelman A, Kimura K, Phillips DR, Cutts SM. Activation of DNA damage response pathways as a consequence of anthracycline-DNA adduct formation. *Biochem Pharmacol*. 2012; 83:1602–1612. [PubMed: 22414726]
25. Coldwell KE, Cutts SM, Ognibene TJ, Henderson PT, Phillips DR. Detection of Adriamycin-DNA adducts by accelerator mass spectrometry at clinically relevant Adriamycin concentrations. *Nucleic Acids Res*. 2008; 36:e100. [PubMed: 18632763]
26. Taatjes DJ, Gaudiano G, Resing K, Koch TH. Alkylation of DNA by the anthracycline, antitumor drugs adriamycin and daunomycin. *J Med Chem*. 1996; 39:4135–4138. [PubMed: 8863788]
27. Taatjes DJ, Gaudiano G, Resing K, Koch TH. Redox pathway leading to the alkylation of DNA by the anthracycline, antitumor drugs adriamycin and daunomycin. *J Med Chem*. 1997; 40:1276–1286. [PubMed: 9111302]
28. Kato S, Burke PJ, Koch TH, Bierbaum VM. Formaldehyde in human cancer cells: detection by preconcentration-chemical ionization mass spectrometry. *Anal Chem*. 2001; 73:2992–2997. [PubMed: 11467545]
29. Kato S, Burke PJ, Fenick DJ, Taatjes DJ, Bierbaum VM, Koch TH. Mass spectrometric measurement of formaldehyde generated in breast cancer cells upon treatment with anthracycline antitumor drugs. *Chem Res Toxicol*. 2000; 13:509–516. [PubMed: 10858324]
30. Cutts SM, Rephaeli A, Nudelman A, Hmelnitsky I, Phillips DR. Molecular basis for the synergistic interaction of adriamycin with the formaldehyde-releasing prodrug pivaloyloxymethyl butyrate (AN-9). *Cancer Res*. 2001; 61:8194–8202. [PubMed: 11719450]
31. Cutts SM, Nudelman A, Pillay V, Spencer DM, Levovich I, Rephaeli A, Phillips DR. Formaldehyde-releasing prodrugs in combination with adriamycin can overcome cellular drug resistance. *Oncol Res*. 2005; 15:199–213. [PubMed: 17822280]
32. Engel D, Nudelman A, Levovich I, Gruss-Fischer T, Entin-Meer M, Phillips DR, Cutts SM, Rephaeli A. Mode of interaction between butyroyloxymethyl-diethyl phosphate (AN-7) and doxorubicin in MCF-7 and resistant MCF-7/Dx cell lines. *J Cancer Res Clin Oncol*. 2006; 132:673–683. [PubMed: 16826403]
33. Rephaeli A, Waks-Yona S, Nudelman A, Tarasenko I, Tarasenko N, Phillips DR, Cutts SM, Kessler-Icekson G. Anticancer prodrugs of butyric acid and formaldehyde protect against doxorubicin-induced cardiotoxicity. *Br J Cancer*. 2007; 96:1667–1674. [PubMed: 17473824]
34. Berlin V, Haseltine WA. Reduction of adriamycin to a semiquinone-free radical by NADPH cytochrome P-450 reductase produces DNA cleavage in a reaction mediated by molecular oxygen. *J Biol Chem*. 1981; 256:4747–4756. [PubMed: 6262301]
35. Myers C. The role of iron in doxorubicin-induced cardiomyopathy. *Semin Oncol*. 1998; 25:10–14. [PubMed: 9768818]
36. Doroshow JH, Synold TW, Somlo G, Akman SA, Gajewski E. Oxidative DNA base modifications in peripheral blood mononuclear cells of patients treated with high-dose infusional doxorubicin. *Blood*. 2001; 97:2839–2845. [PubMed: 11313279]
37. L'Ecuyer T, Sanjeev S, Thomas R, Novak R, Das L, Campbell W, Heide RV. DNA damage is an early event in doxorubicin-induced cardiac myocyte death. *Am J Physiol Heart Circ Physiol*. 2006; 291:H1273–1280. [PubMed: 16565313]
38. Gajewski E, Gaur S, Akman SA, Matsumoto L, van Balgooy JN, Doroshow JH. Oxidative DNA base damage in MCF-10A breast epithelial cells at clinically achievable concentrations of doxorubicin. *Biochem Pharmacol*. 2007; 73:1947–1956. [PubMed: 17445777]
39. De Beer EL, Bottone AE, Voest EE. Doxorubicin and mechanical performance of cardiac trabeculae after acute and chronic treatment: a review. *Eur J Pharmacol*. 2001; 415:1–11. [PubMed: 11245845]

40. Strigun A, Wahrheit J, Niklas J, Heinzle E, Noor F. Doxorubicin increases oxidative metabolism in HL-1 cardiomyocytes as shown by ¹³C metabolic flux analysis. *Toxicol Sci.* 2012; 125:595–606. [PubMed: 22048646]
41. Rotili D, Mai A. Targeting Histone Demethylases: A New Avenue for the Fight against Cancer. *Genes Cancer.* 2011; 2:663–679. [PubMed: 21941621]
42. Kirkland JB. Niacin status impacts chromatin structure. *J Nutr.* 2009; 139:2397–2401. [PubMed: 19812221]
43. McBrien MA, Behbahan IS, Ferrari R, Su T, Huang TW, Li K, Hong CS, Christofk HR, Vogelauer M, Seligson DB, Kurdistani SK. Histone acetylation regulates intracellular pH. *Mol Cell.* 2013; 49:310–321. [PubMed: 23201122]
44. Kawase M, Watanabe M, Kondo T, Yabu T, Taguchi Y, Umehara H, Uchiyama T, Mizuno K, Okazaki T. Increase of ceramide in adriamycin-induced HL-60 cell apoptosis: detection by a novel anti-ceramide antibody. *Biochim Biophys Acta.* 2002; 1584:104–114. [PubMed: 12385893]
45. Lucci A, Han TY, Liu YY, Giuliano AE, Cabot MC. Multidrug resistance modulators and doxorubicin synergize to elevate ceramide levels and elicit apoptosis in drug-resistant cancer cells. *Cancer.* 1999; 86:300–311. [PubMed: 10421266]
46. Delpy E, Hatem SN, Andrieu N, de Vaumas C, Henaff M, Rucker-Martin C, Jaffrezou JP, Laurent G, Levade T, Mercadier JJ. Doxorubicin induces slow ceramide accumulation and late apoptosis in cultured adult rat ventricular myocytes. *Cardiovasc Res.* 1999; 43:398–407. [PubMed: 10536670]
47. Chen Q, Denard B, Huang H, Ye J. Epigenetic silencing of antiviral genes renders clones of Huh-7 cells permissive for hepatitis C virus replication. *J Virol.* 2013; 87:659–665. [PubMed: 23115279]
48. Ji C, Yang B, Yang YL, He SH, Miao DS, He L, Bi ZG. Exogenous cell-permeable C6 ceramide sensitizes multiple cancer cell lines to Doxorubicin-induced apoptosis by promoting AMPK activation and mTORC1 inhibition. *Oncogene.* 2010; 29:6557–6568. [PubMed: 20802518]
49. Uchida Y, Itoh M, Taguchi Y, Yamaoka S, Umehara H, Ichikawa S, Hirabayashi Y, Holleran WM, Okazaki T. Ceramide reduction and transcriptional up-regulation of glucosylceramide synthase through doxorubicin-activated Sp1 in drug-resistant HL-60/ADR cells. *Cancer Res.* 2004; 64:6271–6279. [PubMed: 15342415]
50. Liu YY, Yu JY, Yin D, Patwardhan GA, Gupta V, Hirabayashi Y, Holleran WM, Giuliano AE, Jazwinski SM, Gouaze-Andersson V, Consoli DP, Cabot MC. A role for ceramide in driving cancer cell resistance to doxorubicin. *Faseb J.* 2008; 22:2541–2551. [PubMed: 18245173]
51. Liu YY, Han TY, Giuliano AE, Cabot MC. Ceramide glycosylation potentiates cellular multidrug resistance. *Faseb J.* 2001; 15:719–730. [PubMed: 11259390]
52. Liu YY, Patwardhan GA, Bhinge K, Gupta V, Gu X, Jazwinski SM. Suppression of glucosylceramide synthase restores p53-dependent apoptosis in mutant p53 cancer cells. *Cancer Res.* 2011; 71:2276–2285. [PubMed: 21278235]
53. Denard B, Lee C, Ye J. Doxorubicin blocks proliferation of cancer cells through proteolytic activation of CREB3L1. *Elife.* 2012; 1:e00090. [PubMed: 23256041]
54. Comings DE, Drets ME. Mechanisms of chromosome banding. IX. Are variations in DNA base composition adequate to account for quinacrine, Hoechst 33258 and daunomycin banding? *Chromosoma.* 1976; 56:199–211. [PubMed: 61106]
55. Liu LF, Wang JC. Supercoiling of the DNA template during transcription. *Proc Natl Acad Sci U S A.* 1987; 84:7024–7027. [PubMed: 2823250]
56. Wu HY, Shyy SH, Wang JC, Liu LF. Transcription generates positively and negatively supercoiled domains in the template. *Cell.* 1988; 53:433–440. [PubMed: 2835168]
57. Havas K, Flaus A, Phelan M, Kingston R, Wade PA, Lilley DM, Owen-Hughes T. Generation of superhelical torsion by ATP-dependent chromatin remodeling activities. *Cell.* 2000; 103:1133–1142. [PubMed: 11163188]
58. Thoma NH, Czyzewski BK, Alexeev AA, Mazin AV, Kowalczykowski SC, Pavletich NP. Structure of the SWI2/SNF2 chromatin-remodeling domain of eukaryotic Rad54. *Nat Struct Mol Biol.* 2005; 12:350–356. [PubMed: 15806108]
59. Gartenberg MR, Wang JC. Positive supercoiling of DNA greatly diminishes mRNA synthesis in yeast. *Proc Natl Acad Sci U S A.* 1992; 89:11461–11465. [PubMed: 1333610]

60. Joshi RS, Pina B, Roca J. Topoisomerase II is required for the production of long Pol II gene transcripts in yeast. *Nucleic Acids Res.* 2012; 40:7907–7915. [PubMed: 22718977]
61. Salerno D, Brogioli D, Cassina V, Turchi D, Beretta GL, Seruggia D, Ziano R, Zunino F, Mantegazza F. Magnetic tweezers measurements of the nanomechanical properties of DNA in the presence of drugs. *Nucleic Acids Res.* 2010; 38:7089–7099. [PubMed: 20601682]
62. Gupta P, Zlatanova J, Tomschik M. Nucleosome assembly depends on the torsion in the DNA molecule: a magnetic tweezers study. *Biophys J.* 2009; 97:3150–3157. [PubMed: 20006952]
63. Hizume K, Yoshimura SH, Takeyasu K. Atomic force microscopy demonstrates a critical role of DNA superhelicity in nucleosome dynamics. *Cell Biochem Biophys.* 2004; 40:249–261. [PubMed: 15211026]
64. Clark DJ, Felsenfeld G. Formation of nucleosomes on positively supercoiled DNA. *Embo J.* 1991; 10:387–395. [PubMed: 1991452]
65. Teves SS, Deal RB, Henikoff S. Measuring genome-wide nucleosome turnover using CATCH-IT. *Methods Enzymol.* 2012; 513:169–184. [PubMed: 22929769]
66. Deal RB, Henikoff JG, Henikoff S. Genome-wide kinetics of nucleosome turnover determined by metabolic labeling of histones. *Science.* 2010; 328:1161–1164. [PubMed: 20508129]
67. Jiang H, Reinhardt HC, Bartkova J, Tommiska J, Blomqvist C, Nevanlinna H, Bartek J, Yaffe MB, Hemann MT. The combined status of ATM and p53 link tumor development with therapeutic response. *Genes Dev.* 2009; 23:1895–1909. [PubMed: 19608766]
68. Jackson JG, Pant V, Li Q, Chang LL, Quintas-Cardama A, Garza D, Tavana O, Yang P, Manshoury T, Li Y, El-Naggar AK, Lozano G. p53-mediated senescence impairs the apoptotic response to chemotherapy and clinical outcome in breast cancer. *Cancer Cell.* 2012; 21:793–806. [PubMed: 22698404]
69. Lee MS, Garrard WT. Transcription-induced nucleosome ‘splitting’: an underlying structure for DNase I sensitive chromatin. *EMBO J.* 1991; 10:607–615. [PubMed: 2001676]
70. Teves SS, Henikoff S. Transcription-generated torsional stress destabilizes nucleosomes. *Nat Struct Mol Biol.* 2013 In press.
71. Clapier CR, Cairns BR. The biology of chromatin remodeling complexes. *Annu Rev Biochem.* 2009; 78:273–304. [PubMed: 19355820]
72. Lehmann M, de Vilar SK, Franco A, Reguly ML, Rodrigues de Andrade HH. Activity of topoisomerase inhibitors daunorubicin, idarubicin, and aclarubicin in the *Drosophila* Somatic Mutation and Recombination Test. *Environ Mol Mutagen.* 2004; 43:250–257. [PubMed: 15141364]
73. Hawtin RE, Stockett DE, Byl JA, McDowell RS, Nguyen T, Arkin MR, Conroy A, Yang W, Osheroff N, Fox JA. Voreloxin is an anticancer quinolone derivative that intercalates DNA and poisons topoisomerase II. *PLoS One.* 2010; 5:e10186. [PubMed: 20419121]
74. Marchion DC, Bicaku E, Daud AI, Sullivan DM, Munster PN. In vivo synergy between topoisomerase II and histone deacetylase inhibitors: predictive correlates. *Mol Cancer Ther.* 2005; 4:1993–2000. [PubMed: 16373714]
75. Catalano MG, Fortunati N, Pugliese M, Poli R, Bosco O, Mastrocola R, Aragno M, Boccuzzi G. Valproic acid, a histone deacetylase inhibitor, enhances sensitivity to doxorubicin in anaplastic thyroid cancer cells. *J Endocrinol.* 2006; 191:465–472. [PubMed: 17088416]
76. Marchion DC, Bicaku E, Daud AI, Sullivan DM, Munster PN. Valproic acid alters chromatin structure by regulation of chromatin modulation proteins. *Cancer Res.* 2005; 65:3815–3822. [PubMed: 15867379]
77. Cutts SM, Nudelman A, Rephaeli A, Phillips DR. The power and potential of doxorubicin-DNA adducts. *IUBMB Life.* 2005; 57:73–81. [PubMed: 16036566]
78. Wu S, Ko YS, Teng MS, Ko YL, Hsu LA, Hsueh C, Chou YY, Liew CC, Lee YS. Adriamycin-induced cardiomyocyte and endothelial cell apoptosis: in vitro and in vivo studies. *J Mol Cell Cardiol.* 2002; 34:1595–1607. [PubMed: 12505058]
79. Wang S, Konorev EA, Kotamraju S, Joseph J, Kalivendi S, Kalyanaraman B. Doxorubicin induces apoptosis in normal and tumor cells via distinctly different mechanisms. Intermediacy of H₂O₂- and p53-dependent pathways. *J Biol Chem.* 2004; 279:25535–25543. [PubMed: 15054096]




Fig. 1. Structure of the doxorubicin-DNA complex. (a) Doxorubicin forms a covalent bond (shown in red) with guanine on one strand of DNA mediated by formaldehyde and hydrogen bonds with guanine on the opposing strand [77]. (b) A structure of intercalation of doxorubicin into DNA. Doxorubicin intercalates into DNA and pushes apart the flanking base pairs with the sugar moiety sitting in the minor groove.

Table 1

Actions of doxorubicin and their corresponding drug dose

	Doxorubicin dose ^a	Reference
Topoisomerase II poisoning	0.4 μ M	[4]
DNA adduct formation	0.025 μ M	[25]
Free radical formation	0.1 μ M	[36]
Ceramide overproduction	0.3 μ M	[44]
Histone eviction	0.34 μ M	[11]
Cardiomyocyte apoptosis	0.1 μ M	[78]
Ovarian cancer cell apoptosis	0.5 μ M	[79]

^aThe drug dose varies between different cell types.