

Insulin-producing cells in the brain of adult *Drosophila* are regulated by the serotonin 5-HT_{1A} receptor

Jiangnan Luo · Jaime Becnel · Charles D. Nichols ·
Dick R. Nässel

Received: 5 May 2011/Revised: 1 July 2011/Accepted: 25 July 2011/Published online: 5 August 2011
© Springer Basel AG 2011

Abstract Insulin signaling regulates lifespan, reproduction, metabolic homeostasis, and resistance to stress in the adult organism. In *Drosophila*, there are seven insulin-like peptides (DILP1–7). Three of these (DILP2, 3 and 5) are produced in median neurosecretory cells of the brain, designated IPCs. Previous work has suggested that production or release of DILPs in IPCs can be regulated by a factor secreted from the fat body as well as by neuronal GABA or short neuropeptide F. There is also evidence that serotonergic neurons may regulate IPCs. Here, we investigated mechanisms by which serotonin may regulate the IPCs. We show that the IPCs in adult flies express the 5-HT_{1A}, but not the 5-HT_{1B} or 5-HT₇ receptors, and that processes of serotonergic neurons impinge on the IPC branches. Knockdown of 5-HT_{1A} in IPCs by targeted RNA interference (RNAi) leads to increased sensitivity to heat, prolonged recovery after cold knockdown and decreased resistance to starvation. Lipid metabolism is also affected, but no effect on growth was seen. Furthermore, we show that DILP2-immunolevels in IPCs increase after 5-HT_{1A}

knockdown; this is accentuated by starvation. Heterozygous 5-HT_{1A} mutant flies display the same phenotype in all assays, as seen after targeted 5-HT_{1A} RNAi, and flies fed the 5-HT_{1A} antagonist WAY100635 display reduced lifespan at starvation. Our findings suggest that serotonin acts on brain IPCs via the 5-HT_{1A} receptor, thereby affecting their activity and probably insulin signaling. Thus, we have identified a second inhibitory pathway regulating IPC activity in the *Drosophila* brain.

Keywords 5-hydroxytryptamine · Insulin signaling · G-protein-coupled receptor · Lifespan · Stress resistance

Abbreviations

5-HT	5-hydroxytryptamine
5-HTP	5-hydroxytryptophan
BDSC	Bloomington <i>Drosophila</i> stock center
CNS	Central nervous systems
DILP	<i>Drosophila</i> insulin-like peptide
GABA	Gamma-aminobutyric acid
GFP	Green fluorescent protein
GIRK	G-protein-coupled inwardly rectifying potassium channel
GPCR	G-protein-coupled receptor
IPCs	Insulin-producing cells
MNCs	Median neurosecretory cells
NPF	Neuropeptide F
NS3	Nucleostemin 3
OAMB	Octopamine receptor (mushroom bodies)
PCR	Polymerase chain reaction
PKA	Protein kinase A
RNAi	RNA interference
sNPF	Short neuropeptide F
VDRC	Vienna <i>Drosophila</i> RNAi center

Electronic supplementary material The online version of this article (doi:10.1007/s00018-011-0789-0) contains supplementary material, which is available to authorized users.

J. Luo · D. R. Nässel (✉)
Department of Zoology, Stockholm University,
Svante Arrhenius väg 18B, 10691 Stockholm, Sweden
e-mail: dnassel@zoologi.su.se

J. Becnel · C. D. Nichols
Department of Pharmacology and Experimental Therapeutics,
Louisiana State University Health Sciences Center,
New Orleans, LA, USA

Introduction

Insulin-like peptides regulate growth, reproduction, metabolism and lifespan in both invertebrates and mammals [1–7]. In *Drosophila*, seven insulin-like peptides (DILPs) and a single insulin receptor have been identified [2, 8–10]. DILP signaling in adult fruitflies is important in the regulation of metabolic homeostasis, resistance to stress of different kinds and regulation of lifespan [3, 11–14]. Three of the DILPs (DILP2, 3 and 5) are produced by a set of median neurosecretory cells in the brain of *Drosophila* and are thought to be released into the circulation from axon terminals in neurohemal areas in the corpora cardiaca and anterior aorta [2, 15, 16]. In experiments where the insulin-producing cells (IPCs) in the brain were genetically ablated, it was shown that growth was retarded, lifespan extended, levels of carbohydrate increased in the circulation, storage of lipids increased and resistance to various stresses increased [11, 16]. Also, mutations in genes encoding DILP2, 3 and 5 produce these phenotypes, in addition to defects in growth and fertility [9, 14]. Thus, DILP2, 3 and 5 display pleiotropic functions, and experiments to reveal the roles of individual DILPs by targeted mutations suggested partly redundant functions of the three peptides [9, 17]. It is known, however, that the production of the three DILPs in the IPCs can be individually regulated [13, 17, 18].

We are interested in factors regulating the production and release of DILPs from the IPCs in the adult *Drosophila* brain. In addition to circulating nutritional signals derived from the fat body [19], it has been suggested that a brain-derived short neuropeptide F (sNPF) may stimulate DILP production in IPCs as well as feeding and growth [20, 21]. Another neurotransmitter that seems to stimulate signaling in the IPCs is octopamine [22]. In this study, the IPCs were found to be involved in regulation of sleep-wakefulness, and under stimulatory control of octopamine via the OAMB receptor. Recently, GABA and its metabotropic GABA_B receptor were shown to inhibit IPCs and insulin signaling at metabolic stress, but seemed not to affect growth [23]. A fourth neurotransmitter has been implicated in regulation of insulin signaling, the monoamine serotonin [24]. This study demonstrates that the GTPase nucleostemin 3 (NS3) in serotonergic neurons is required for normal growth of *Drosophila* and that it regulates serotonin levels. Feeding flies the precursor of serotonin, 5-Hydroxytryptophan (5-HTP) mimicks the developmental delay seen in the *ns3* mutant flies. Since *ns3* mutants feed normally, the developmental effect of NS3 and serotonin was sought in a pathway known to regulate growth, the insulin signaling. It was found that the *ns3* mutants have increased levels of DILP2 protein, but not *dilp2* transcript and that this may be

associated with a decrease of DILP2 release since insulin signaling was reduced [24]. Thus, it was concluded that serotonin-producing neurons, which were found to have axon terminations close to the IPCs, regulate release of DILPs and thus growth. However, the specific receptor type mediating the serotonergic signaling to the IPCs was not determined and a direct action of serotonin on IPCs was not established.

There are four different serotonin receptors in *Drosophila*, designated 5-HT_{1A}DRO, 5-HT_{1B}DRO, 5-HT₂DRO, and 5-HT₇DRO, all of which are G-protein-coupled receptors (GPCRs) [25–28]. We henceforth refer to these receptors without the DRO suffix. Of these, 5-HT_{1A} and 5-HT_{1B} are known to inhibit adenylate cyclase and 5-HT₇ stimulates it, whereas 5-HT₂ activation has not yet been investigated in *Drosophila* (reviewed in [25, 28]). Specific roles of the 5-HT_{1A}, 5-HT_{1B} and 5HT₂ receptors in sleep and circadian activity of *Drosophila* has been investigated [29–31]. Furthermore, it has been shown that the 5-HT_{1A} and 5-HT₂ receptors each modulate aggressive behavior [32] and that 5-HT₇ is required for normal courtship behavior [33]. Other studies of serotonergic signaling in *Drosophila* have not specified the receptor type, but indicate pleiotropic roles of this neurotransmitter both in development, physiological processes and specific behaviors (see [34–39]). Here, we undertook a study to determine which of the serotonin receptors mediate the regulation of insulin signaling from the brain IPCs of *Drosophila*.

Using a 5-HT_{1A} promoter Gal4 line combined with antiserum to DILP2, we could show that all the brain IPCs in the adult fly express this receptor, whereas Gal4 drivers for the other serotonin receptors did not display expression in any of the IPCs. Using a *Dilp2*-Gal4 line to drive UAS-mediated RNA interference with 5-HT_{1A} expression in the IPCs, we found that insulin signaling increased with reduced receptor levels. Thus, flies with reduced 5-HT_{1A} in IPCs are less resistant to starvation and heat knockdown, and display delayed recovery from cold coma. Also, the levels of DILP-immunofluorescence in the IPCs and body lipid are affected by receptor knockdown. Similar results were obtained for 5-HT_{1A} mutant flies, and after feeding flies a 5-HT_{1A} antagonist. Knockdown of 5-HT_{1A} had no clear effect on growth of the flies, however. Our data therefore strongly suggest that the serotonin signaling to the neurosecretory cells producing insulin-like peptides is mediated mainly by the 5-HT_{1A} receptor and that this signaling may convey stress signals. However, it is not clear whether all the effects of 5-HT_{1A} knockdown that are shown can be attributed to increased insulin signaling, or if some effects are caused by other signaling from the IPCs.

Materials and methods

Fly strains

Adult white-eyed flies *Drosophila melanogaster* (*w*¹¹¹⁸ strain) were used for some immunocytochemistry and control experiments. For some experiments, early and late third instar larvae were utilized. All flies were kept at 25°C on a 12:12 h light/dark cycle and maintained on a diet of standard *Drosophila* medium.

The following Gal4 lines were used to drive the expression of green fluorescent protein (GFP) and for crosses to induce RNA interference (RNAi): *TRH*-Gal4 [34] (a gift from O.V. Alekseyenko and E. Kravitz, Boston, MA), *5-HT₇*-Gal4 was generated and characterized as described in [33], *5-HT_{1A}*-Gal4 is described below, *5-HT_{1B}*-Gal4 (a gift from A. Sehgal, Philadelphia, PA, USA), *Dilp2*-Gal4 (2nd chromosome) [40] (a gift from P. Shen, Athens, GA, USA), *Dilp2*-Gal4 (3rd chromosome) [18] (a gift from S. Broughton, Univ. Lancaster, UK). OK107-Gal4 [41] and *Act5c*-Gal4 were from The Bloomington *Drosophila* Stock Center (BDSC) at Indiana University, Bloomington, IN, USA. Two different UAS-*5-HT_{1A}*-RNAi lines were used in the experiments: one from the Vienna *Drosophila* RNAi Center (VDRC) and the other from BDSC. UAS-*5-HT₇* [42] (a gift from J.A.T. Dow, Glasgow, UK) was used to ectopically express the receptor. UAS-*mcd8-gfp* or UAS-*s65t-gfp* flies from (BDSC) were used to visualize Gal4 expression. A *5-HT_{1A}* mutant generated by imprecise P element excision was obtained from BDSC (stock number 27640). This mutant was generated and characterized by [30]. The genotype of this mutant is *w*^{*}; *5-HT_{1A}*^{Δ5kb}/CyO, P{ActGFP}JMR1.

Preparation of *5-HT_{1A}*Dro promoter region

Genomic DNA from adult Canton-S flies was prepared as previously described [33]. To isolate putative 5' enhancer regions, which are normally contained within the first few kb of genomic DNA upstream of the RNA transcription start site, 5 kb of genomic DNA immediately upstream of the ATG start codon within the *5-HT_{1A}*Dro locus was amplified from genomic DNA using Platinum Pfx DNA Polymerase (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions (Fig. 1). Primers corresponding to the *5-HT_{1A}*Dro promoter region containing Not I restriction sites at their 5' end were ordered from Integrated DNA Technologies (Coralville, IA, USA). Forward primer = 5'-gcgccgcATGGCCCAAGTATCAGGAATCTGC-3'; reverse primer = 5'-gcgccgcAAGATGCGAATGTACGTCCAGTTG-3'; annealing T = 62.0°C, elongation T = 68°C. The amplification product consisted of a single band of 5 kb, which was gel purified using the Wizard SV gel

cleanup system (Promega, Madison, WI, USA) following the manufacturer's instructions.

Generation of the *5-HT_{1A}* construct and transgenic flies

Both the purified *5-HT_{1A}* PCR product and the pERGP P-element insertion vector [33] were digested with Not I and gel purified. Digested pERGP was dephosphorylated using Apex Heat-Labile Alkaline Phosphatase (Epicentre, Madison, WI, USA) following the manufacturer's directions. The *5-HT_{1A}* promoter fragment was ligated into the Not I site of the pERGP vector using the Fast-Link DNA Ligation Kit (Epicentre) following the manufacturer's directions. The final construct was verified using a panel of restriction enzymes, as well by sequence analysis of the cloning site junctions. Transgenic lines were generated from this final product using the services of BestGene (Chino Hills, CA, USA).

Antisera and immunocytochemistry

For immunocytochemistry, adult *Drosophila* heads or central nervous systems (CNS) of third instar larvae were dissected in 0.01 M phosphate-buffered saline with 0.25% Triton X-100, pH 7.2 (PBS-Tx) and fixed in ice-cold 4% paraformaldehyde in 0.1 M sodium phosphate buffer pH 7.4 (PB) for 2–4 h. Following rinsing with 0.1 M PB, adult brains or larval CNS were either dissected out for whole-mount immunocytochemistry or whole heads were incubated overnight in 20% sucrose in 0.1 M PB at 4°C as cryoprotection. Cryostat sections (50 μm thick) of the heads were cut on a cryostat at –23°C.

Incubation with primary antiserum for whole-mount tissues was performed for 72 h, while sections were incubated overnight, both at 4°C. The following primary antisera were used: a rabbit antiserum to *Drosophila* insulin-like peptide 2 (anti-DILP2) [15], known to recognize *Drosophila* insulin-producing cells at a dilution of 1:4,000. A monoclonal mouse antibody to serotonin (Clone 5HT-H209; Dako, Copenhagen, Denmark) was used at a dilution of 1:80. An antiserum raised to a peptide (from the third intracellular loop) of the honeybee *5-HT_{1A}* receptor [43], kindly donated by M. Thamm and W. Blenau (Potsdam, Germany), was tested at a dilution of 1:1,000. For detection of primary antisera, Cy3-tagged goat anti-rabbit antiserum (Jackson Immuno Research) and Alexa goat anti-mouse 488 were used at a dilution of 1:1,000. Tissues or sections were rinsed thoroughly with PBS-Tx, followed by a final wash in PBS and then mounted in 80% glycerol in PBS. For each experiment, at least 10 adult brains and 5 larval CNS were analyzed.




Fig. 1 The 5-HT_{1A}Dro receptor locus and 5 kb region used for Gal4 construct. The genomic region of the 5-HT_{1A}Dro locus (CG16720) is on the right arm of the second chromosome (*light bars* 5' and 3' untranslated regions, *dark bars* coding regions; *arrows* indicate mRNA transcription start sites). The 5-kb region of genomic DNA

5-HT_{1A} antagonist

To inactivate the 5-HT_{1A} receptors in the adult flies, we used the 5-HT_{1A} antagonist WAY100635 {N-[2-[4-(2-Methoxyphenyl)-1-piperazinyl]ethyl]-N-2-pyr-idinylcyclohexanecarboxamide maleate salt; Sigma, St Louis, MO, USA}. We dissolved 5 mg WAY100635 in a solution of 0.25 g agarose and 50 ml Milli-Q water, which yielded 0.18 mM (0.1 g/L). This agarose/antagonist solution was put in glass tubes in 0.5-ml aliquots for starvation tests (see below).

Stress assays

Male flies, 4–6 days old, were used for starvation assays. All flies were kept in an incubator at 25°C with 12:12 h light:dark (LD) conditions, and controlled humidity. For the starvation experiments, flies were placed individually in 2-ml glass vials with 500 µl of 0.5% aqueous agarose. The vials were checked for dead flies every 12 h. These starvation experiments were run in three replicates with at least 40 flies of each genotype per replicate.

For the heat knockdown and chill coma recovery experiments, newly emerged adults were collected, sexed under light CO₂ anesthesia, and sorted into groups of 20–30 males in fresh vials with *Drosophila* medium. Male flies were used at 4–5 days of age, and were not anaesthetized subsequent to the initial sorting. For the heat knockdown experiments, groups of 20–30 males were placed in glass vials (without food), which were completely immersed in a temperature-regulated water bath to induce heat knockdown. Time to knockdown was measured for flies exposed to 39°C at 5-min intervals.

For the chill coma recovery experiments, groups of 20–30 males were placed in glass vials (without food), which were put into a box full of ice inside a refrigerator (2°C) to induce chill coma. The flies were exposed to the environmental stress for 4 h and the recovery at room temperature was recorded in food vials at 5-min intervals.

Lipid measurements

Lipid content was measured in flies of different genotypes after 0, 12 and 24 h of starvation. The lipid content was

used to make the GAL4 construct is indicated immediately upstream of the mRNA start site. The adjacent CG15109 locus is 7 kb upstream of the first exon of the 5-HT_{1A}Dro gene. The CG15109 transcript has been reported to be exclusively expressed in the male testis

determined according to the method of Service [44]. Groups of five male flies were weighed on a Mettler MT5 Microbalance (Mettler Toledo, Switzerland) to obtain wet weight and subsequently dried at 65°C for 24 h. Flies were then weighed again to obtain dry weight. Lipids were extracted by placing intact dry flies in glass vials containing diethyl ether for 24 h with gentle agitation at room temperature. The diethyl ether was removed and flies were dried for another 24 h and then weighed to obtain lean dry weight. The difference between dry weight and lean dry weight was considered the total lipid content of the flies. Totals of 40 flies of each genotype were tested in three replicates.

Image analysis

Specimens were imaged with Zeiss LSM 510 META confocal microscope (Jena, Germany) using ×20 or ×40 oil immersion objectives and a Leica TCS-SP2 confocal microscope (Buffalo Grove, IL, USA) using a ×20 dry or ×63 water immersion objectives. Confocal images were obtained at an optical section thickness of 0.2–0.5 µm and were processed with Zeiss LSM or Leica software. Images were edited for contrast and brightness in Adobe Photoshop CS3 Extended version 10.0.

Quantification of immunofluorescence

Immunocytochemistry with DILP2 antiserum was performed on adult brains from starved and fed flies of different genotypes for quantification of immunofluorescence in IPCs. The brains were imaged in a Zeiss LSM 510 confocal microscope with fixed exposure time, using LSM software. The immunofluorescence was quantified in each cell, using Image J 1.40 from NHI, Bethesda, MD, USA (<http://rsb.info.nih.gov/ij/>).

Statistical analysis

All statistical analyses were performed by using Prism GraphPad 6.0. Survival data were analyzed by Log rank test with Mantel–Cox post-test, for quantification of immunofluorescence, lipid values and body weights we used one-way ANOVA with Tukeys comparison or

two-way ANOVA depending on analysis (see Figure legends for details). Data are presented as means \pm SEM.

Results

5-HT_{1A} is expressed in brain insulin producing cells

A cluster of 16 insulin-producing cells (IPCs) in pars intercerebralis of the *Drosophila* brain produce DILP2, 3 and 5 [2, 15, 18]. These IPCs have arborizations in three regions of the brain: branches extending laterally in dorsal protocerebrum, shorter branches along the initial IPC neurites (Fig. 2a), and extensive arborizations in the tritocerebrum. The main IPC axons terminate in neurohemal areas of the corpora cardiaca, in the anterior aorta and in the anterior intestine and crop [15, 23, 45]. To investigate which of the 5-HT receptors is expressed on the IPCs, we utilized different 5-HT receptor Gal4 lines to drive GFP expression in combination with DILP2 immunocytochemistry. An earlier account on 5-HT₂ distribution, using Gal4-driven LacZ expression, described no neurons in the median neurosecretory cell group [31].

For this study, a 5-HT_{1A}-Gal4 line was produced as detailed in “Materials and methods”. A promoter fragment consisting of 5 kb of genomic DNA immediately upstream of the ATG start codon within the 5-HT_{1A} locus was used for this construct (Fig. 1). This Gal4 was used for driving UAS-cd8-GFP or UAS-s65t-GFP to display the neuronal expression. Our attempts to raise antisera to a sequence of the *Drosophila* 5-HT_{1A} protein were unsuccessful, and our tests of antiserum to 5-HT_{1A} of the honey bee [43] did not result in any specific immunolabeling in the fruitfly. Therefore, it was necessary to resort to comparing the distribution of 5-HT_{1A}-Gal4 expression to that of serotonin. As seen in Supporting material (S Fig. 1), there is a good correlation between 5-HT_{1A}-Gal4 expression and neuronal processes displaying serotonin immunoreactivity both in the adult and larval CNS.

Significantly, in adult flies, we found the 5-HT_{1A}-Gal4 to be expressed in median neurosecretory cells (MNCs) similar in location and morphology to the IPCs (Fig. 2b). Application of antiserum against DILP2 to the GFP-labeled brains revealed that most, if not all, of the DILP2-immunolabeled cells also display 5-HT_{1A}-Gal4 expression (Fig. 2b–d). Some additional neurons in the same region express 5-HT_{1A}, but do not display DILP immunoreactivity. These neurons are just adjacent to the IPCs and likely to be other MNCs or interneurons (Fig. 2d). It can be noted that antiserum to the 5-HT₁ receptor of the cockroach *Periplaneta americana* labeled MNCs of the cockroach brain with axons extending to the corpora cardiaca, similar to the *Drosophila* neurons expressing 5-HT_{1A} and DILPs

[46]. In the cockroach, it remains to be determined if these MNCs also express insulin-like peptide.

We also examined 5-HT_{1A} expression in relation to IPCs in larval brains. Surprisingly, we found no 5-HT_{1A}-Gal4-driven GFP in IPCs of feeding third instar larvae (Fig. 2e, f). Analysis of non-feeding late third instar larvae revealed 5-HT_{1A} expression in one or two of the 16 IPCs in some, but not all, specimens (Fig. 2g–k). Thus, it appears as if the majority of the IPCs start expressing the 5-HT_{1A} receptor only after completion of the larval stages.

Similar analysis of 5-HT_{1B} and 5-HT₇, where we used promoter Gal4 drivers combined with anti-DILP2 labeling, revealed that these receptors are not expressed on IPCs in larvae or adults (S Fig. 2). Also, the original reports on 5-HT_{1B} and 5-HT₇ distribution show no expression of these receptors in IPCs [29, 33]. Therefore, our findings suggest that only the 5-HT_{1A} is expressed on *Drosophila* IPCs and that onset of expression occurs late in non-feeding larvae or in pupae.

Processes from serotonergic neurons superimpose IPC branches

In recent papers, it was shown that serotonin-immunoreactive neuron processes impinge on IPC branches in the larval brain [24, 47]. However, these studies provided no information on serotonin distribution in relation to IPCs in the adult brain. Here, we applied a monoclonal antibody to serotonin to brains with IPCs marked by *Dilp2*-Gal4-driven GFP. In the adult brain, serotonergic neuron branches can be found close to those of the IPCs in the pars intercerebralis (Fig. 3). The superposition between branches is seen in the dorsolateral area as well as in the median region along initial IPC neurites. The individual serotonin-immunoreactive neurons that supply branches to the IPC dendrites were not identified in the adult brain due to the dense packing of neuronal processes. In third instar larvae, serotonin-immunoreactive neuron processes also superimpose branches of the IPCs (not shown; see [24, 47]).

Knockdown of 5-HT_{1A} causes elevation of DILP2 levels in IPCs

It was shown in an earlier report that manipulation of NS3 in serotonergic neurons affects DILP2 peptide levels in IPCs, but a direct functional connection between serotonergic neurons and IPCs was not demonstrated [24]. We therefore tested whether serotonergic regulation of DILP levels is directly mediated by the 5-HT_{1A} receptor on IPCs. To obtain an estimate of DILP protein levels, we measured the intensity of DILP2 immunofluorescence in the cell bodies of the IPCs after knockdown of 5-HT_{1A} by the transgene *Dilp2*-Gal4/UAS-5-HT_{1A}-RNAi. The *Dilp2*-Gal4




Fig. 2 Expression of 5-HT_{1A} receptor on IPCs in the *Drosophila* brain. All images of adult brains are frontal views (dorsal is up) and of larval brain horizontal views (anterior is up). **a** Anatomy of the IPCs seen with *Dilp2*-Gal4-driven GFP (projection of optic sections). A cluster of 16 cell bodies (*cb*) give rise to axons with branches in two regions of the pars intercerebralis: wide dorsal lateral branches (*asterisks*) and shorter median branches (*arrow*). **b–d** In the adult brain, the 5-HT_{1A}-Gal4 driver is expressed in a set of median neurosecretory cells (*green*), most of which also express DILP2-immunolabeling (*magenta*). In the merged image (**d**), it can be seen that all but one of the 5-HT_{1A}-expressing cells display DILP2

driver used here and in all but one (see S Fig. 3) of the experiments has the insert on the 2nd chromosome [40]. We also tested a loss of function mutation generated previously in the 5-HT_{1A} by imprecise P-element excision [30]; used here as heterozygous mutant flies. The DILP2 levels were measured both in normally fed flies and flies that had been starved for 12 h. It was found that the DILP2-immunolabeling increased in the IPCs of flies with diminished 5-HT_{1A}, both globally and in IPCs only, and that starvation further increased DILP2 (Fig. 4). Since 5-HT_{1A} is likely to act as an inhibitory receptor, the insulin increase in the IPCs after knockdown could be explained by diminished inhibitory signaling to these cells. A very similar increase in DILP2 levels were seen after knockdown of the inhibitory GABA_B receptor on IPCs [23] and

immunolabeling (*whitish*). **e, f** In the feeding third instar larval brain, there is no 5-HT_{1A} expression in the DILP2-immunolabeled IPCs (projection of several optic sections). The weakly labeled cell body in (**e**) (*asterisk*) does not express DILP2. **g** In the wandering (non-feeding) third instar larva, 5-HT_{1A} expression starts. Here, one IPC (*arrow*) coexpresses the receptor and DILP2. **h** In another specimen of the same age, two IPCs on the *left* side coexpress receptor and DILP2, but no cells on the *right* (*asterisk*). This image is a projection of several optic sections. **i–k** A single optic section of the same specimen showing colocalization of markers in one cell body (*arrow*) to the *left* and none on the *right* (*asterisk*)

in the study of the NS3 mutant (reversed by *ns3* rescue in serotonergic neurons) [24]. The increase of DILP2 both in fed and starved flies with diminished 5-HT_{1A}, also suggest some regulation of IPCs under non-stress conditions.

Knockdown of 5-HT_{1A} in IPCs and globally causes decreased resistance to starvation

It is known that flies with decreased insulin signaling display an increased resistance to starvation [11], and a recent study demonstrated that knockdown of the inhibitory GABA_B receptor on IPCs decreased this resistance [23]. Thus, we tested whether the knockdown of 5-HT_{1A} on IPCs affect survival at starvation. In all experiments described in this paper, we used only adult (4–6 days old) male flies,

Fig. 3 Relationships between processes of IPCs and serotonin-immunoreactive neurons in the adult brain. We utilized the *Dilp2-Gal4* to drive GFP in IPCs (green) and a mouse monoclonal antibody to serotonin (magenta) to visualize relationships between the two neuron types in the pars intercerebralis (shown in frontal view; dorsal up). **a–c** IPCs at the level with wide lateral branches dorsally. These IPC branches superimpose those of varicose serotonin immunoreactive ones (even the small set of branches more ventrally, at asterisk). **d** The framed area in (a) is shown at higher magnification. **e–g** The IPCs at the level of the median short branches (other specimen). Again, the branches of the two neuron types superimpose (e.g., at arrows). **h** The framed area in (e) is shown at higher magnification





Fig. 4 Knockdown of 5-HT_{1A} leads to an increase in DILP-immunolabeling in insulin-producing cells (IPCs). **a** Relative DILP immunofluorescence in IPCs in fed and starved flies with and without 5-HT_{1A} receptor knockdown (5-HT_{1A}Ri) in IPCs (*Dilp2-Gal4/UAS-5-HT_{1A}-RNAi*). The DILP2 antiserum used is likely to cross react with DILP2, 3 and 5. Control flies (*w¹¹¹⁸-5-HT_{1A}*) display significantly lower levels of DILP-immunofluorescence than the flies with diminished 5-HT_{1A} (*Dilp2-5-HT_{1A}Ri*), both in fed flies ($***p < 0.001$; One-way ANOVA with Tukey's comparison) and after starvation ($***p < 0.001$). Also, the increases of fluorescence in controls (grey bars) and knockdown flies (red bars) when comparing

fed and starved flies are significant ($p < 0.001$ for both). For each genotype and condition, IPCs of 5–7 brains were investigated. **b** Relative DILP immunofluorescence in IPCs in fed and starved mutant (5-HT_{1A} mut) and wild type (*w¹¹¹⁸*) flies. Wild-type flies display significantly lower levels of DILP-immunofluorescence than the 5-HT_{1A} mutant flies, both in fed flies ($***p < 0.001$) and starved ones ($*p < 0.05$). Again, the increases in immunolabeling in fed and starved controls and fed and starved mutants are significant ($p < 0.001$ for both). IPCs of 7–10 brains of each genotype and condition were investigated

unless otherwise specified. Statistical data are given in figure legends. Flies were kept on aqueous agarose, and it was found that flies with 5-HT_{1A}-RNAi driven by the

Dilp2-Gal4 display a significantly reduced median and maximal lifespan compared to parental controls (Fig. 5). We tested two different 5-HT_{1A}-RNAi lines with the same

result (Fig. 5). As a control, we also utilized a different *Dilp2*-Gal4 driver (insert on 3rd chromosome; [18]) crossed with one of the RNAi lines, and obtained the same phenotype at starvation (S Fig. 3). In all the following experiments, we utilized the *Dilp2*-Gal4 inserted on the 2nd chromosome.

Another Gal4 driver, OK107 [41], also known to be expressed in IPCs [23], was used to drive 5-HT_{1A}-RNAi. This cross resulted in flies with the same abbreviated lifespan at starvation (Fig. 6a). Finally, the heterozygous 5-HT_{1A} mutant flies also displayed reduced resistance to starvation (Fig. 6b). Thus, fly crosses from three different Gal4 drivers and two 5-HT_{1A}-RNAi lines gave the same phenotype at starvation as a 5-HT_{1A} mutant strain. The reduced lifespan at starvation in these experiments indicates an increase in insulin signaling or other activity of IPCs, likely to be caused by loss of inhibition of the IPCs by diminished 5-HT_{1A} expression.

As a further test of serotonin signaling to the IPCs during stress, we ectopically expressed the 5-HT₇ receptor on these cells by crossing *Dilp2*-Gal4 flies to a UAS-5-HT₇ line [42]. Since the 5-HT₇ receptor is known to stimulate adenylate cyclase via G_s [25, 26, 33], we expected the ectopic receptor to increase DILP signaling. Indeed, the flies with 5-HT₇ expressed on IPCs displayed a significantly reduced lifespan at starvation (Fig. 7). Probably, the ectopic 5-HT₇ receptor couples to endogenous G_s and adenylate cyclase. These, and protein kinase A (PKA), are known to be present in the IPCs from a study of octopamine signaling to these cells [22].

We finally tested knockdown of the 5-HT_{1B} and 5-HT₇ receptors in IPCs by crossing different relevant receptor RNAi lines to the *Dilp2*-Gal4 driver. Neither of these fly

crosses resulted in altered lifespan in response to starvation (S Fig 4). Thus, it seems that of the three tested receptors only 5-HT_{1A} plays a role in regulation of IPCs at starvation.

Flies fed a 5-HT_{1A} antagonist display reduced resistance to starvation

To induce conditional interference with 5-HT_{1A} signaling in adults we fed wild-type flies with the 5-HT_{1A} antagonist WAY100635 [32, 48]. This antagonist has also been shown to act as an antagonist at the honeybee 5-HT_{1A} receptor [43] and as an inverse agonist at the 5-HT₁ receptor of the cockroach, *Periplaneta americana* [46]. WAY100635 was dissolved in aqueous agarose that flies were kept on while we monitored lifespan during starvation. Flies fed antagonist displayed significantly reduced lifespan compared to control flies (Fig. 8a). We also fed 5-HT_{1A} mutant flies the antagonist and found that their lifespan was further reduced compared to both wild-type flies and mutants that were not fed the drug (Fig. 8b, c). This additive effect can be explained by the 5-HT_{1A} mutant flies being heterozygous and thus possessing reduced receptor levels, not null levels.

5-HT_{1A} knockdown in IPCs and 5-HT_{1A} mutation affect responses to temperature stress

To further investigate 5-HT_{1A} signaling to IPCs in stress responses, we performed tests of heat and cold tolerance (see [11]). For heat tolerance the time to knockdown at increased temperature (39°C) was monitored. Both the 5-HT_{1A} mutant flies and flies with 5-HT_{1A}-RNAi targeted to IPCs displayed a decreased tolerance to heat (Fig. 9a, b). Likewise these flies required increased time to recover




Fig. 5 Knockdown of 5-HT_{1A} receptor in IPCs increases sensitivity to starvation. We performed GABA_B 5-HT_{1A} receptor knockdown in IPCs with two different RNAi constructs, one from VDRC [5-HT_{1A}Ri(V)] (a) and another from Bloomington Stock Center [5-HT_{1A}Ri (B)] (b). These flies were crossed to a *Dilp2*-Gal4 driver [40] used in all experiments, unless other specified. Male flies were kept on aqueous agarose (to induce starvation) and their survival monitored over time. All experiments were run in three replicates. **a** Using a *Dilp2*-Gal4 driver to knockdown the 5-HT_{1A} receptor in IPCs [Dilp2-5-HT_{1A}Ri (V)], we obtained flies that display

significantly reduced survival at starvation ($p < 0.0001$ to both parental controls, Log rank test, Mantel-Cox; $n = 111$ –215 for each genotype). **b** Flies obtained from crossing *Dilp2*-Gal4 with the other strain 5-HT_{1A}Ri (B) also displayed significantly reduced survival at starvation ($p < 0.0001$ and $p = 0.0002$ to the two controls; $n = 159$ –209 for each genotype). In all subsequent graphs with 5-HT_{1A}RNAi, we used the (V) strain. We also used a different *Dilp2*-Gal4 driver (on the 3rd chromosome) [18] to drive the UAS-5-HT_{1A}Ri(V) and obtained the same significantly reduced life span at starvation (see S Fig. 3)




Fig. 6 Knockdown of 5-HT_{1A} receptor in IPCs or globally increases sensitivity to starvation. Using the same experimental conditions, we tested lifespan at starvation with another Gal4 driver that includes the IPCs and a 5-HT_{1A} mutant. **a** The enhancer trap Gal4 OK107 includes the IPCs in its expression pattern [23] and was used here to drive UAS-5-HT_{1A}Ri. The lifespan of OK107-5-HT_{1A}Ri flies is

significantly reduced at starvation ($p < 0.0001$ to both controls, Log rank test, Mantel-Cox; $n = 84$ –121 for each genotype, experiment run in two replicates). **b** Heterozygous 5-HT_{1A} mutant flies also display significantly reduced survival at starvation compared to wild-type (w^{1118}) and controls ($p < 0.0001$ to wild-type controls; $n = 165$ for each genotype, experiment run in three replicates)




Fig. 7 Targeted expression of 5-HT₇ in IPCs decreases lifespan at starvation. The 5-HT₇ receptor couples via Gs to stimulate adenylate cyclase. Therefore, we ectopically expressed this receptor by means of the Dilp2 Gal4 driver (Dilp2-UAS-5-HT₇) to test the effect on starvation resistance. Indeed, this ectopic expression produced flies with decreased the lifespan at starvation ($p < 0.001$ to both controls, Log rank test, Mantel-Cox; $n = 93$ –122 for each genotype; experiment in two replicates). The decreased lifespan suggest that the ectopic 5-HT₇ couples to Gs and stimulates IPCs and insulin signaling

lead to increased storage of both lipid and carbohydrate [11] and knockdown of the GABA_B receptor on IPCs altered the lipid storage profile over time at starvation [23]. We tested flies with 5-HT_{1A} knocked down in IPCs for whole-body lipid levels over 0, 12 and 24 h starvation. Compared to controls, the 5-HT_{1A}-RNAi flies display significantly reduced lipid levels at 12 and 24 h of starvation (Fig. 10a). The heterozygous 5-HT_{1A} mutants also displayed a significant reduction of lipid levels compared to control (wild-type) flies at 12 and 24 h starvation (Fig. 10b). Thus, knockdown of 5-HT_{1A} in IPCs, or globally, produces a lipid storage phenotype consistent with increased insulin signaling, similar to findings in the GABA study [23].

from cold coma induced by exposure to 0°C for 4 h (Fig. 9c, d).

Knockdown of 5-HT_{1A} has no effect on growth

From these stress experiments, it can be suggested that increased DILP signaling, due to diminished 5-HT_{1A} leads to decreased temperature tolerance. These findings are contrary to those in an earlier paper where ablation of IPCs, and presumably reduced insulin signaling, leads to decreased heat and cold tolerance in *Drosophila* [11]. Our results are, however, in accordance with findings in *C. elegans* [49].

Finally, we tested whether knockdown of 5-HT_{1A} has any effect on growth. Adult flies (4–6 days old) with the receptor knocked down in IPCs by *Dilp2*-driven 5-HT_{1A}-RNAi or globally in the mutant were weighed. There was no significant difference between controls and flies with receptor knockdown (S Fig. 5), suggesting that 5-HT_{1A}-mediated signaling plays no major role in control of organismal growth as determined by body weight. Similar results were obtained in flies where the GABA_B receptor was knocked down in IPCs [23].

Discussion

Knockdown of 5-HT_{1A} alters lipid storage in flies

We have shown here that the inhibitory serotonin 5-HT_{1A} receptor is expressed in IPCs of adult *Drosophila* and affects IPC activity and probably insulin-like signaling as monitored by different stress responses, DILP levels and lipid storage. On the other hand, we did not obtain evidence

DILP signaling is important in regulation of carbohydrate and lipid metabolism in flies [1, 11, 16]. Ablation of IPCs




Fig. 8 Feeding flies a 5-HT_{1A} antagonist increases sensitivity to starvation. The 5-HT_{1A} antagonist WAY100635 was fed to the flies via aqueous agarose at a concentration of 0.18 mM (0.1 g/L) and the flies were kept on this agarose for the duration of the starvation experiment. **a** Survival of wild-type (w¹¹¹⁸) flies kept on agarose with or without the antagonist. Antagonist-fed flies displayed a significant reduction in lifespan ($p = 0.001$, Log rank test; $n = 101$ and 165 for the two test groups; run in two replicates). **b** Survival of heterozygous 5-HT_{1A} mutant (*mut*) flies fed agarose with or without antagonist. The antagonist further reduces lifespan in the mutant flies ($p < 0.0001$; $n = 166$ and 147 for the two test groups; run in two replicates). **c** Comparison of median survival (lifespan) of the four groups of flies tested in (a) and (b). It can be seen that the antagonist action is additive to the heterozygous mutation of the receptor. Thus, the shortest median life span is seen for mutant flies fed the antagonist, which is significantly shorter than both wild-type flies fed antagonist and mutants fed agarose alone ($*p < 0.05$, $**p < 0.01$; one-way Anova with Tukey comparison)

insulin signaling pathway [22, 50, 51]. Therefore, activation or inhibition of the IPCs could cause actions that are non-insulin mediated. These could be either via other messengers released by the IPCs or by indirect action of DILPs on other circuits or endocrine cells. As an example, the reduced 5-HT_{1A} inhibition of the IPCs could cause an increased starvation-induced hyperlocomotion (see [50]). This could in turn result in increased rates of carbohydrate and lipid consumption, leading to reduced survival at starvation. Thus, it cannot be excluded that some of the phenotypes discussed below are not mediated by typical insulin signaling to the fat body.

Our assays indicated that temperature tolerance and starvation resistance were impaired in flies where the 5-HT_{1A} receptor was knocked down in IPCs as well as in 5-HT_{1A} mutants. Furthermore, the 5-HT_{1A} ablated flies and mutants displayed a higher lipid consumption rate, likely because insulin signaling was increased. However, no clear effect on growth (body weight) was observed after 5-HT_{1A} knockdown, similar to findings in a study of GABA_B receptor knockdown on IPCs [23]. Possibly, this lack of effect on growth is due to the absence of 5-HT_{1A} expression on larval IPCs suggested by our expression data. Thus, although we could induce increased insulin signaling in the adult fly by knockdown of the 5-HT_{1A} receptor, our experiments with mutants and targeted RNAi seem not to affect larval IPCs. Other studies have demonstrated effects on growth after ablation of IPCs or interference with DILP signaling in the larva [2, 6, 18, 24, 52]. Also, a paper by Kaplan and coworkers [24] indicated that serotonergic neurons modulate IPC activity and DILP signaling in larvae and that this signaling affects growth. However, that paper did not provide evidence for direct action of serotonin on the IPCs. Certainly, it is possible that one of the other 5-HT receptors is mediating effects on growth and/or

for expression or action of the 5-HT_{1B} or 5-HT₇ receptors on IPCs, and an earlier report provided no support for 5-HT₂ expression in these neurosecretory cells [31]. Thus, it is likely that serotonin signaling to the brain IPCs in adult flies is mainly mediated by the 5-HT_{1A} receptor.

When discussing the role of 5-HT_{1A} in inhibiting IPCs, it should be noted that the IPCs are also known to regulate locomotor activity, sleep-wakefulness and sensitivity to ethanol, and this regulation may be independent of the




Fig. 9 Responses to temperature stress are influenced by knockdown of 5-HT_{1A} receptor in IPCs or globally. Flies with the 5-HT_{1A} receptor knocked down in IPCs by the transgene *Dilp2-Gal4/UAS-5-HT_{1A}-RNAi* or globally in the mutant were tested for responses to temperature stress. All experiments were run in two replicates. **a, b** Response to heat was tested by exposing flies to 39°C and monitoring time to knockdown (given as percent survival with heat). **a** The flies with the 5-HT_{1A} receptor diminished in IPCs (*Dilp2-5-HT_{1A}Ri*) displayed a faster knockdown at 39°C ($p = 0.0005$ and $p < 0.0001$ to parental controls; Log rank test, Mantel-Cox; $n = 48-55$

for the three genotypes). **b** The 5-HT_{1A} mutant (*mut*) flies displayed a similar increased sensitivity to heat compared to wild-type flies (*w¹¹¹⁸*) ($p < 0.0001$; $n = 50$ and 54). **c, d** Recovery from cold knockdown (coma) was monitored in the same genotypes. Flies were kept at 0°C for 4 h and the time to recovery was monitored (given as percent in chill coma). **c** The flies with receptor knockdown in IPCs were slower in their recovery from cold ($p = 0.0009$ and $p = 0.0017$ to controls; $n = 52-58$ for the three genotypes). **d** The receptor mutant flies also display a longer recovery time ($p < 0.0001$ to wild-type control; $n = 56$ and 58)




Fig. 10 Knockdown of 5-HT_{1A} receptor in IPCs or globally affects lipid storage at starvation. Whole-body lipid was measured, as given in “Materials and methods”, in fed flies (0 h) and flies exposed to starvation for 12 and 24 h. **a** Flies with the 5-HT_{1A} receptor diminished in IPCs (*Dilp2-5-HT_{1A}Ri*) displayed a significantly lower amount of lipid than control flies both in fed and starved flies (***) ($p < 0.001$; one-way Anova with Tukey’s comparison; $n = 120$ for each genotype; experiment run in three replicates). The decrease in lipid

over time was also significant for both genotypes ($p < 0.001$; two-way ANOVA). **b** In the 5-HT_{1A} mutant flies, the lipid levels were lower than in wild-type flies at 12 and 24 h of starvation. (***) ($p < 0.001$, ns not significant; one-way Anova, $n = 120$ for each genotype; experiment run in three replicates). The decrease in lipid over time was also significant for both genotypes ($p < 0.001$; two-way ANOVA)

that the signaling to the IPCs in growth regulation is indirect.

Since the 5-HT_{1A} knockdown flies display a reduced stress resistance and altered lipid consumption, it is suggestive that DILP release from IPCs increases at stress due to the manipulation. However, the DILP signaling from IPCs is complex, since the different DILPs in the IPCs seem to be individually regulated transcriptionally [13, 17, 18]. If production of DILPs can be individually regulated, it is likely that the ratio of DILP2, 3 and 5 stored in vesicles in the IPC axon terminations also vary. Since the calcium-dependent release of the different DILPs is unlikely to be individually controlled, the release at any given point may result in different ratios of circulating DILPs. Then, if the different DILPs mediate different actions on physiology, as suggested [9], one would expect that the composition of the DILP cocktail could be functionally important.

Several studies have assayed DILP-immunofluorescence levels in cell bodies of IPCs after various manipulations to monitor “insulin signaling” [13, 19, 23, 24]. Our experiments here show increased DILP2 immunolabeling in the IPC cell bodies after 5-HT_{1A} knockdown. This would suggest an increase of DILP production, but it is likely that DILP release is visible only in the axon terminals of the IPCs in the corpora cardiaca and aorta (not monitored here). Thus, the IPC cell bodies are not optimal for monitoring release. Previous studies showed similar changes in DILP2 immunolabeling after knockdown of the inhibitory GABA_B receptor on IPCs [23] and in NS3 mutants [24]. It may be that the increase in DILP2 immunolabeling seen in the cell bodies of the IPCs is due to an increased production of DILP as a consequence of increased release into the circulation from axon terminals. A sensitive assay is, however, required for monitoring DILP levels in the hemolymph of *Drosophila* to obtain a measure of bona fide release.

Due to the extensive arborizations of serotonergic neuron processes in the small and compact brain of *Drosophila*, we could not identify the individual neurons that innervate the IPCs. In a study of the much larger blowfly brain, two pairs of candidate serotonergic neurons with axon terminations in the pars intercerebralis were shown: one pair with cell bodies in posterior protocerebrum and one pair in the subesophageal ganglion [53]. Also in the larva, in spite of a simpler neuronal organization, these serotonergic neurons could not be individually identified [24, 47]. However, it was suggested that in the larva serotonergic neurons in the subesophageal ganglion may have branches that impinge on the IPCs [47]. Such a location would make sense since the subesophageal ganglion is known to receive chemosensory inputs and contain neurons regulating aspects of feeding and neuroendocrine function [54–57].

The changing expression pattern of 5-HT_{1A} in IPCs during development may indicate that this receptor is involved in a behavioral transition from feeding to non-feeding stage. Neuropeptide F (NPF), an ortholog of NPY in mammals, was shown to be a molecular switch in the transition from feeding to non-feeding (wandering) stage of larvae [56]. These authors showed that NPF is expressed at higher levels in certain brain neurons in feeding third instar larvae and is downregulated in the late third instar non-feeding stage. Maybe the 5-HT_{1A} expression on IPCs starts during the same behavioral transition, and perhaps there is a link between NPF and serotonin receptor expression. There is another relevant example of a change in the larval brain at transition to wandering stage. At this transition, the larvae become positively phototactic, and simultaneously a set of serotonergic neuron branches grow into the larval optic center [37, 58, 59]. Importantly, serotonin signaling, probably mediated by 5-HT_{1A}, is required in the larval photic response [37].

Our findings here that brain IPCs can be directly inactivated by serotonergic signaling suggests that these neurosecretory cells are under complex stimulatory and inhibitory regulation. Production and/or release of DILPs in IPCs is induced by a circulating factor released from the fat body [19] and possibly by the peptide sNPF and the monoamine octopamine, both released by brain neurons [21, 22] and is inhibited by GABA [23] as well as serotonin, as shown here. Such a control by multiple neuronal systems and hormonal factors may serve to fine-tune the activity of the IPCs in the production and release of the physiologically very important DILPs. Similarly, the insulin release from mammalian pancreatic beta cells is under control of circulating glucose levels as well as several neuromediators, such as serotonin, GABA, glucagon-like peptide and other peptides [60–62].

It is not clear whether the two inhibitory signals to the IPCs, GABA and serotonin, are mediated by different core intracellular pathways. The postsynaptic GABA_B receptor commonly couples to G-protein-coupled inwardly rectifying potassium channels (GIRKs), but can also inhibit adenylate cyclase via G_{α_i}/G_{α_o}, or even inhibit voltage-dependent Ca²⁺ channels, all leading to hyperpolarization [63–67]. In *Drosophila*, knockdown of a putative GIRK subunit in the IPCs phenocopied GABA_B receptor knockdown in starvation assays [23], but other pathways were not tested. The 5-HT_{1A} can also act on GIRKs, as well as couple negatively to adenylate cyclase [25, 32]. Interestingly, the stimulatory action of octopamine on IPCs was shown to be mediated by the OAMB receptor through activation of adenylate cyclase, increased cyclic AMP and PKA activation [22]. For the future, it would be interesting to investigate whether octopamine and serotonin act antagonistically on IPCs by

converging on adenylate cyclase. Perhaps these biogenic amines play antagonistic roles in tuning insulin signaling during stress, as well as in regulation of sleep wakefulness.

Acknowledgments We thank the persons and organizations listed in “Materials and methods” for flies and reagents. This study was supported by the Swedish Research Council (VR).

References

- Baker KD, Thummel CS (2007) Diabetic larvae and obese flies—emerging studies of metabolism in *Drosophila*. *Cell Metab* 6:257–266
- Brogiolo W, Stocker H, Ikeya T, Rintelen F, Fernandez R et al (2001) An evolutionarily conserved function of the *Drosophila* insulin receptor and insulin-like peptides in growth control. *Curr Biol* 11:213–221
- Giannakou ME, Partridge L (2007) Role of insulin-like signalling in *Drosophila* lifespan. *Trends Biochem Sci* 32:180–188
- Géminard G, Arquier N, Layalle S, Bourouis M, Slaidina M et al (2006) Control of metabolism and growth through insulin-like peptides in *Drosophila*. *Diabetes* 55:S5–S8
- Kenyon C (2010) A pathway that links reproductive status to lifespan in *Caenorhabditis elegans*. *Ann N Y Acad Sci* 1204:156–162
- Teleman AA (2010) Molecular mechanisms of metabolic regulation by insulin in *Drosophila*. *Biochem J* 425:13–26
- Wu Q, Brown MR (2006) Signaling and function of insulin-like peptides in insects. *Annu Rev Entomol* 51:1–24
- Fernandez R, Tabarini D, Azpiazu N, Frasch M, Schlessinger J (1995) The *Drosophila* insulin receptor homolog: a gene essential for embryonic development encodes two receptor isoforms with different signaling potential. *EMBO J* 14:3373–3384
- Grönke S, Clarke DF, Broughton S, Andrews TD, Partridge L (2010) Molecular evolution and functional characterization of *Drosophila* insulin-like peptides. *PLoS Genet* 6:e1000857
- Slaidina M, Delanoue R, Grönke S, Partridge L, Leopold P (2009) A *Drosophila* insulin-like peptide promotes growth during nonfeeding states. *Dev Cell* 17:874–884
- Broughton SJ, Piper MD, Ikeya T, Bass TM, Jacobson J et al (2005) Longer lifespan, altered metabolism, and stress resistance in *Drosophila* from ablation of cells making insulin-like ligands. *Proc Natl Acad Sci USA* 102:3105–3110
- Tatar M, Kopelman A, Epstein D, Tu MP, Yin CM et al (2001) A mutant *Drosophila* insulin receptor homolog that extends lifespan and impairs neuroendocrine function. *Science* 292:107–110
- Broughton SJ, Slack C, Alic N, Metaxakis A, Bass TM et al (2010) DILP-producing median neurosecretory cells in the *Drosophila* brain mediate the response of lifespan to nutrition. *Aging Cell* 9:336–346
- Zhang H, Liu J, Li CR, Momen B, Kohanski RA et al (2009) Deletion of *Drosophila* insulin-like peptides causes growth defects and metabolic abnormalities. *Proc Natl Acad Sci USA* 106:19617–19622
- Cao C, Brown MR (2001) Localization of an insulin-like peptide in brains of two flies. *Cell Tissue Res* 304:317–321
- Rulifson EJ, Kim SK, Nusse R (2002) Ablation of insulin-producing neurons in flies: growth and diabetic phenotypes. *Science* 296:1118–1120
- Broughton S, Alic N, Slack C, Bass T, Ikeya T et al (2008) Reduction of DILP2 in *Drosophila* triages a metabolic phenotype from lifespan revealing redundancy and compensation among DILPs. *PLoS ONE* 3:e3721
- Ikeya T, Galic M, Belawat P, Nairz K, Hafen E (2002) Nutrient-dependent expression of insulin-like peptides from neuroendocrine cells in the CNS contributes to growth regulation in *Drosophila*. *Curr Biol* 12:1293–1300
- Geminard C, Rulifson EJ, Leopold P (2009) Remote control of insulin secretion by fat cells in *Drosophila*. *Cell Metab* 10:199–207
- Lee KS, You KH, Choo JK, Han YM, Yu K (2004) *Drosophila* short neuropeptide F regulates food intake and body size. *J Biol Chem* 279:50781–50789
- Lee KS, Kwon OY, Lee JH, Kwon K, Min KJ et al (2008) *Drosophila* short neuropeptide F signalling regulates growth by ERK-mediated insulin signalling. *Nat Cell Biol* 10:468–475
- Crocker A, Shahidullah M, Levitan IB, Sehgal A (2010) Identification of a neural circuit that underlies the effects of octopamine on sleep:wake behavior. *Neuron* 65:670–681
- Enell LE, Kapan N, Söderberg JA, Kabsai L, Nässel DR (2010) Insulin signaling, lifespan and stress resistance are modulated by metabotropic GABA receptors on insulin producing cells in the brain of *Drosophila*. *PLoS ONE* 5:e15780
- Kaplan DD, Zimmermann G, Suyama K, Meyer T, Scott MP (2008) A nucleostemin family GTPase, NS3, acts in serotonergic neurons to regulate insulin signaling and control body size. *Genes Dev* 22:1877–1893
- Nichols DE, Nichols CD (2008) Serotonin receptors. *Chem Rev* 108:1614–1641
- Witz P, Amlaiky N, Plassat JL, Maroteaux L, Borrelli E et al (1990) Cloning and characterization of a *Drosophila* serotonin receptor that activates adenylate cyclase. *Proc Natl Acad Sci USA* 87:8940–8944
- Saudou F, Boschert U, Amlaiky N, Plassat JL, Hen R (1992) A family of *Drosophila* serotonin receptors with distinct intracellular signalling properties and expression patterns. *EMBO J* 11:7–17
- Blenau W, Thamm M (2011) Distribution of serotonin (5-HT) and its receptors in the insect brain with focus on the mushroom bodies. Lessons from *Drosophila melanogaster* and *Apis mellifera*. *Arthropod Struct Dev* (in press)
- Yuan Q, Lin F, Zheng X, Sehgal A (2005) Serotonin modulates circadian entrainment in *Drosophila*. *Neuron* 47:115–127
- Yuan Q, Joiner WJ, Sehgal A (2006) A sleep-promoting role for the *Drosophila* serotonin receptor 1A. *Curr Biol* 16:1051–1062
- Nichols CD (2007) 5-HT₂ receptors in *Drosophila* are expressed in the brain and modulate aspects of circadian behaviors. *Dev Neurobiol* 67:752–763
- Johnson O, Becnel J, Nichols CD (2009) Serotonin 5-HT₂ and 5-HT_{1A}-like receptors differentially modulate aggressive behaviors in *Drosophila melanogaster*. *Neuroscience* 158:1292–1300
- Becnel J, Johnson O, Luo J, Nässel DR, Nichols CD (2011) The serotonin 5-HT₇Dro receptor is expressed in the brain of *Drosophila*, and is essential for normal courtship and mating. *PLoS ONE* 6(6):e20800
- Alekseyenko OV, Lee C, Kravitz EA (2010) Targeted manipulation of serotonergic neurotransmission affects the escalation of aggression in adult male *Drosophila melanogaster*. *PLoS ONE* 5:e10806
- Daubert EA, Condron BG (2010) Serotonin: a regulator of neuronal morphology and circuitry. *Trends Neurosci* 33:424–434
- Dierick HA, Greenspan RJ (2007) Serotonin and neuropeptide F have opposite modulatory effects on fly aggression. *Nat Genet* 39:678–682

37. Rodriguez Moncalvo VG, Campos AR (2009) Role of serotonergic neurons in the *Drosophila* larval response to light. *BMC Neurosci* 10:66
38. Dacks AM, Green DS, Root CM, Nighorn AJ, Wang JW (2009) Serotonin modulates olfactory processing in the antennal lobe of *Drosophila*. *J Neurogenet* 23:366–377
39. Sitaraman D, Zars M, Laferriere H, Chen YC, Sable-Smith A et al (2008) Serotonin is necessary for place memory in *Drosophila*. *Proc Natl Acad Sci USA* 105:5579–5584
40. Wu Q, Zhang Y, Xu J, Shen P (2005) Regulation of hunger-driven behaviors by neural ribosomal S6 kinase in *Drosophila*. *Proc Natl Acad Sci USA* 102:13289–13294
41. Wang Y, Guo HF, Pologruto TA, Hannan F, Hakker I et al (2004) Stereotyped odor-evoked activity in the mushroom body of *Drosophila* revealed by green fluorescent protein-based Ca²⁺ imaging. *J Neurosci* 24:6507–6514
42. Kerr M, Davies SA, Dow JA (2004) Cell-specific manipulation of second messengers; a toolbox for integrative physiology in *Drosophila*. *Curr Biol* 14:1468–1474
43. Thamm M, Balfanz S, Scheiner R, Baumann A, Blenau W (2010) Characterization of the 5-HT1A receptor of the honeybee (*Apis mellifera*) and involvement of serotonin in phototactic behavior. *Cell Mol Life Sci* 67:2467–2479
44. Service FJ, O'Brien PC, Rizza RA (1987) Measurements of glucose control. *Diabetes Care* 10:225–237
45. Cognigni P, Bailey AP, Miguel-Aliaga I (2011) Enteric neurons and systemic signals couple nutritional and reproductive status with intestinal homeostasis. *Cell Metab* 13:92–104
46. Troppmann B, Balfanz S, Baumann A, Blenau W (2010) Inverse agonist and neutral antagonist actions of synthetic compounds at an insect 5-HT1 receptor. *British J Pharmacol* 159:1450–1462
47. Agrawal N, Padmanabhan N, Hasan G (2009) Inositol 1, 4, 5-trisphosphate receptor function in *Drosophila* insulin producing cells. *PLoS ONE* 4:e6652
48. Fornal CA, Metzler CW, Gallegos RA, Veasey SC, McCreary AC et al (1996) WAY-100635, a potent and selective 5-hydroxytryptamine_{1A} antagonist, increases serotonergic neuronal activity in behaving cats: comparison with (S)-WAY-100135. *J Pharmacol Exp Ther* 278:752–762
49. Walker GA, Lithgow GJ (2003) Lifespan extension in *C. elegans* by a molecular chaperone dependent upon insulin-like signals. *Aging Cell* 2:131–139
50. Mattaliano MD, Montana ES, Parisky KM, Littleton JT, Griffith LC (2007) The *Drosophila* ARC homolog regulates behavioral responses to starvation. *Mol Cell Neurosci* 36:211–221
51. Corl AB, Rodan AR, Heberlein U (2005) Insulin signaling in the nervous system regulates ethanol intoxication in *Drosophila melanogaster*. *Nat Neurosci* 8:18–19
52. Walkiewicz MA, Stern M (2009) Increased insulin/insulin growth factor signaling advances the onset of metamorphosis in *Drosophila*. *PLoS ONE* 4:e5072
53. Nässel DR (1988) Serotonin and serotonin-immunoreactive neurons in the nervous system of insects. *Prog Neurobiol* 30:1–85
54. Melcher C, Pankratz MJ (2005) Candidate gustatory interneurons modulating feeding behavior in the *Drosophila* brain. *PLoS Biol* 3:e305
55. Bader R, Colomb J, Pankratz B, Schrock A, Stocker RF et al (2007) Genetic dissection of neural circuit anatomy underlying feeding behavior in *Drosophila*: distinct classes of hugin-expressing neurons. *J Comp Neurol* 502:848–856
56. Wu Q, Wen T, Lee G, Park JH, Cai HN et al (2003) Developmental control of foraging and social behavior by the *Drosophila* neuropeptide Y-like system. *Neuron* 39:147–161
57. Miyazaki T, Ito K (2010) Neural architecture of the primary gustatory center of *Drosophila melanogaster* visualized with GAL4 and LexA enhancer-trap systems. *J Comp Neurol* 518:4147–4181
58. Campos AR, Lee KJ, Steller H (1995) Establishment of neuronal connectivity during development of the *Drosophila* larval visual system. *J Neurobiol* 28:313–329
59. Sawin-McCormack EP, Sokolowski MB, Campos AR (1995) Characterization and genetic analysis of *Drosophila melanogaster* photobehavior during larval development. *J Neurogenet* 10:119–135
60. Paulmann N, Grohmann M, Voigt JP, Bert B, Vowinkel J et al (2009) Intracellular serotonin modulates insulin secretion from pancreatic beta-cells by protein serotonylation. *PLoS Biol* 7:e1000229
61. Adeghate E, Ponery AS, Pallot DJ, Singh J (2001) Distribution of vasoactive intestinal polypeptide, neuropeptide-Y and substance P and their effects on insulin secretion from the in vitro pancreas of normal and diabetic rats. *Peptides* 22:99–107
62. Adeghate E, Ponery AS (2002) GABA in the endocrine pancreas: cellular localization and function in normal and diabetic rats. *Tissue Cell* 34:1–6
63. Mezler M, Muller T, Raming K (2001) Cloning and functional expression of GABA(B) receptors from *Drosophila*. *Eur J Neurosci* 13:477–486
64. Kaupmann K, Schuler V, Mosbacher J, Bischoff S, Bittiger H et al (1998) Human gamma-aminobutyric acid type B receptors are differentially expressed and regulate inwardly rectifying K⁺ channels. *Proc Natl Acad Sci USA* 95:14991–14996
65. Bettler B, Kaupmann K, Mosbacher J, Gassmann M (2004) Molecular structure and physiological functions of GABA(B) receptors. *Physiol Rev* 84:835–867
66. Hamasaka Y, Wegener C, Nässel DR (2005) GABA modulates *Drosophila* circadian clock neurons via GABAB receptors and decreases in calcium. *J Neurobiol* 65:225–240
67. Kolaj M, Bai D, Renaud LP (2004) GABAB receptor modulation of rapid inhibitory and excitatory neurotransmission from sub-fornical organ and other afferents to median preoptic nucleus neurons. *J Neurophysiol* 92:111–122