

Randomized Trial of a Smartphone Mobile Application Compared to Text Messaging to Support Smoking Cessation

David B. Buller, PhD,¹ Ron Borland, PhD,² Erwin P. Bettinghaus, PhD,¹ James H. Shane,¹ and Donald E. Zimmerman, PhD³

¹Klein Buendel, Inc., Golden, Colorado.

²Cancer Council Victoria, Carlton South, Victoria, Australia.

³Colorado State University, Ft. Collins, Colorado.

The findings, views, and opinions expressed here are those of the authors and not necessarily those of the National Institutes of Health.

Abstract

Background: Text messaging has successfully supported smoking cessation. This study compares a mobile application with text messaging to support smoking cessation. **Materials and Methods:** Young adult smokers 18–30 years old ($n=102$) participated in a randomized pretest–posttest trial. Smokers received a smartphone application (REQ-Mobile) with short messages and interactive tools or a text messaging system (onQ), managed by an expert system. Self-reported usability of REQ-Mobile and quitting behavior (quit attempts, point-prevalence, 30-day point-prevalence, and continued abstinence) were assessed in posttests. **Results:** Overall, 60% of smokers used mobile services (REQ-Mobile, 61%, mean of 128.5 messages received; onQ, 59%, mean of 107.8 messages), and 75% evaluated REQ-Mobile as user-friendly. A majority of smokers reported being abstinent at posttest (6 weeks, 53% of completers; 12 weeks, 66% of completers [44% of all cases]). Also, 37% (25% of all cases) reported 30-day point-prevalence abstinence, and 32% (22% of all cases) reported continuous abstinence at 12 weeks. OnQ produced more abstinence ($p < 0.05$) than REQ-Mobile. Use of both services predicted increased 30-day abstinence at 12 weeks (used, 47%; not used, 20%; $p = 0.03$). **Conclusions:** REQ-Mobile was feasible for delivering cessation support but appeared to not move smokers to quit as quickly as text messaging. Text messaging may work better because it is simple, well known, and delivered to a primary inbox. These advantages may disappear as smokers become more experienced with new handsets. Mobile phones may be promising delivery platforms for cessation services using either smartphone applications or text messaging.

Key words: e-health, technology, telecommunications

Introduction

Approximately one in five U.S. adults continue to smoke.¹ Fortunately, many try to quit,^{2–4} and community-based brief cessation counseling is effective.^{5–7} Cessation programs delivered over the Internet and on cell phones have

been successful.^{8–16} Text messages tailored to smoking habits and barriers to quitting have achieved high use (>50%),^{17,18} encouraged quit attempts, and helped smokers cope with craving^{11–14} and relapse.¹⁵

Multimedia, data-rich smartphones hold promise for delivery of cessation counseling. By proactively, unobtrusively, and confidentially reaching out, mobile services should gain smokers' attention quickly, deliver meaningful in-the-moment help,^{19–22} create a sense of urgency to respond,^{19,20,23,24} and increase involvement in quitting.²⁵ Mobile applications also have the potential to mimic autonomous social support instrumental to quitting^{26–29} by creating a virtual relationship with accountability and emotional support.²⁰ However, in 2009, only 11% of iPhone® (Apple®, Cupertino, CA) cessation applications followed the U.S. Public Health Service's 2008 Clinical Practice Guideline for Treating Tobacco Use and Dependence; few directed smokers to effective treatments.³⁰ Research is needed on mobile applications that provide direct help to smokers, follow recommended guidelines, and do not simply refer to cessation services.³¹

A small-scale randomized trial tested delivering cessation support over a smartphone mobile application, before embarking on a large community-based trial. The research questions explored in this pilot study were the following: (1) Would smokers use the application and rate it high on usability? (2) How many smokers receiving the application engaged in quitting, and how did use affect quitting? (3) How would the application compare with a text messaging program in usage and quitting? The mobile application was compared with a text messaging program because text messaging has been efficacious, is simpler to use, and is potentially less expensive and complicated to deploy to determine whether it is worth investing in a mobile application.

Materials and Methods

SAMPLE

Adult U.S. smokers ($n=102$) 18–30 years old were recruited on Google Adwords, Facebook, and Adbrite advertising systems in 2010 (advertising cost, \$47 per enrollee). Young adults should be most adept at using smartphones. Of smokers expressing interest ($n=357$), 187 screened by telephone were eligible (i.e., current smokers, interested in quitting, U.S. resident, and English proficient); 126 completed the consent form and pretest online, and 102 (28% of interested smokers; 54% of eligible smokers) signed a cell phone contract and were randomized (Fig. 1). Treatment effects were of primary interest, so we focused on confidence intervals (CIs): a sample of 100 would yield a two-sided 90% CI of ± 0.163 for a difference in proportions of 0.40 versus 0.50 and ± 0.141 for 0.20 versus 0.30.




Fig. 1. CONSORT diagram for randomized trial.

ONQ TEXT MESSAGING SYSTEM

The onQ text messaging system^{32,33} had 1,126 messages managed by the Quit Coach expert system³⁴⁻³⁶ and sent to the text inbox. Messages were grounded in social cognitive theory³⁷ and a modified version of the transtheoretical model³⁸ and designed to increase/sustain motivation and enhance self-efficacy across four phases of quitting (without a quit date, with a quit date, quit for <1 week, and quit for 1-4 weeks). Messages suggested tasks to plan, set, and maintain a quit date, cope with cravings and relapse, and consolidate a nonsmoking lifestyle (see Borland et al.³² and linked files on the journal's Web site for more details).

REAL E QUIT MOBILE SMOKING CESSATION APPLICATION

The Real e Quit mobile application (REQ-Mobile) ran on Windows Mobile 6.1 or higher smartphones. When the application was produced, Android® (Google, Mountain View, CA) handsets did not exist, and Apple's policy limited the number of phones that could be used in a study. Smokers received short messages from onQ slightly revised for REQ-Mobile in an e-mail-like inbox. Audio phase-transition messages were delivered to enhance comprehension.³⁹ Additional features were designed to perform tasks advised in onQ messages, based on the Quit Coach theoretical framework. Smokers could create lists on (1) reasons for and (2) benefits of quitting and plans for coping with (3) challenging situations and (4) stressful circumstances, listen to 13 audio testimonials from former smokers,

and read 16 short support documents (e.g., benefits of quitting, strategies for stopping, using nicotine replacement therapy [NRT], coping with withdrawal). They could set a quit date and update the expert system. The short messages directed smokers to quitting tools, and 56 messages encouraged them to use the study Web site to view additional resources on smoking cessation. Once smokers quit for more than 1 week, they were shifted to onQ because maintenance counseling was not programmed into REQ-Mobile owing to limited funds.

Two rounds of formative testing of REQ-Mobile with smokers 18-30 years old (59% female) confirmed usability. In Round 1 (n = 17), 76-100% of smokers completed 11 tasks.

After REQ-Mobile was revised, 90% of smokers in Round 2 (n = 10) completed seven of nine tasks, and completion times decreased. In both rounds, smokers rated REQ-Mobile favorably.

PROCEDURES

A randomized pretest-posttest two-group design was conducted in 2010-2011. After consenting online, participants completed the pretest online at the study Web site. Half were then assigned to REQ-Mobile, and half were assigned to onQ, using an online randomization program. The uniform random number generation function in SAS software (SAS Institute, Cary, NC) was used to generate random pairs assigned to treatment or control (i.e., block size of 2). Pairs were loaded into the study Web site without revealing them to the investigators and used sequentially to assign participants who completed the pretest. A smartphone with 3 months of unlimited phone, text, and data service was mailed to all study participants. If study requirements were met, smokers could keep the phone and receive a fourth month of service. Project staff explained how to use the assigned service. All participants first had to log on to Quit Coach, located on the study Web site, and complete a brief assessment, separate from the pretest, to initialize the mobile programs. Participants in both conditions could return to the study Web site to read longer tailored advice from Quit Coach and read longer support documents than were available on either onQ or REQ-Mobile. Smokers could request a free 2-week course of nicotine patches. All smokers were invited by e-mail to complete follow-up questionnaires online at 6 and 12 weeks.

MEASURES

Questionnaires. At pretest, 6-week posttest, and 12-week posttest, smokers reported smoking status (i.e., smoked every day or some days and 30-day prevalence), readiness to quit, nicotine dependence (i.e., cigarettes/day, time to first cigarette, and difficulty not smoking around other smokers), quit attempts in the past year, quitting self-efficacy, smoking by significant other and five closest people, and home smoking rules. At 6 and 12 weeks, continued smokers reported their likelihood of quitting in the next 6 weeks, whether they had set a quit date, and if they had tried to quit since the pretest. Participants not smoking reported how long ago they quit, if they had smoked since quitting (and if so, 7-day smoking prevalence), likelihood of smoking again, self-efficacy for staying abstinent, and use of cessation aids. Smokers assigned to REQ-Mobile and who had used it evaluated its usability at 6 weeks.⁴⁰ Nonusers were not asked about usability for they had no experience with the mobile application. Smokers provided demographics, Internet experience, use of mobile phones, and health status.

Smoking abstinence was assessed in four ways. At both posttests, point-prevalence abstinence on the day interviewed was indicated by a declaration of quitting. At 6 weeks, 7-day point-prevalence abstinence was defined as not smoking since quitting or in the past 7 days (one smoker who quit < 1 week ago and not smoked since was considered abstinent). At 12 weeks, 30-day point-prevalence abstinence was defined as not smoking in the past 30 days and declaring a quit, and continuous abstinence was defined as also not smoking since quitting.

Usage. Use of REQ-Mobile and onQ was obtained from the Web servers, which recorded the number of messages sent and opened, the use of REQ-Mobile quitting tools through the first week after quitting, and number of messages sent and text responses returned to onQ.

STATISTICAL ANALYSIS PROCEDURES

Comparisons between REQ-Mobile and onQ were performed by chi-squared and logistic regression in SAS software. Spearman correlations estimated the relationship of program use to quitting. Controlling for mobile phone experience and use of NRT did not change the outcomes. Effects of loss to follow-up were probed by analyzing abstinence by those completing posttests and then with all cases assuming those lost still smoked. The alpha criterion was $p=0.05$ (two-tailed).

Results

PROFILE OF THE SAMPLE

Table 1 gives the sample profile. Randomization balanced groups on all demographics and pretest smoking status except income (Table 1). Most participants were successfully followed up (65% at 6 weeks; 67% at 12 weeks; 56% completed both posttests) (Table 2), with no differences in demographics ($p>0.05$), condition ($p>0.05$), or use of interventions ($p>0.05$). Follow-up in the REQ-Mobile group was marginally less than in the onQ group.

Table 1. Demographic Characteristics and Smoking Status by Experimental Condition

DEMOGRAPHIC CHARACTERISTIC	REQ-MOBILE GROUP (N= 51)	ONQ GROUP (N= 51)	P	OVERALL (N= 102)
Sex				
Male	55%	43%	0.23	49%
Female	45%	57%		51%
Hispanic ethnicity	4%	8%	0.68	6%
Race				
White	70%	76%	0.79	74%
African American	16%	12%		14%
Other	14%	12%		12%
Education				
High school diploma or less	41%	57%	0.28	49%
Education beyond high school	45%	33%		39%
College or postgraduate degree	14%	10%		12%
Annual household income				
Less than \$25,000	65%	41%	0.02	53%
\$25,000 or more	35%	59%		47%
Marital status				
Married	33%	51%	0.17	42%
Divorced/widowed	10%	10%		10%
Never married	57%	39%		48%
Age (mean years)	25.5	24.3	0.07	24.9
Smoking status				
Cigarettes smoked per day (mean number)	16.8	17.1	0.84	17.0
Attempted to quit in past year	66%	71%	0.62	69%
Readiness to quit				
Open to quitting	12%	10%	0.82	11%
Like to quit	18%	14%		16%
Planning to quit	58%	66%		62%
Have set a quit date	10%	6%		8%
Have quit	2%	10%		3%

Table 2. Number of Participants Followed Up by Use of Mobile Smoking Cessation Services

	REQ-MOBILE			ONQ			GRAND TOTAL (N)
	USED (N)	DID NOT USE (N)	TOTAL (N)	USED (N)	DID NOT USE (N)	TOTAL (N)	
6-week posttest							
Completed	19	14	33	23	10	33	66
Not completed	12	6	18	7	11	18	36
12-week posttest							
Completed	19	14	33	24	11	35	68
Not completed	12	6	18	6	10	16	34
Both posttests							
Completed	13	12	25	22	10	32	57
Not completed	18	8	26	8	11	19	45
Grand total	31	20	51	30	21	51	102

However, 9% were struggling to maintain abstinence at 6 weeks because only 44% reported 7-day point-prevalence abstinence. At 12 weeks, 37% (25% of all cases) of smokers reported being abstinent when assessed by 30-day point-prevalence abstinence, but continuous abstinence was lower (32% completers; 22% all cases). No participant still smoking at 12 weeks reported a quit attempt (i.e., no relapsers).

Use of both services was positively correlated with quitting among completers at 12 weeks ($n = 68$). More smokers who used either program reported 30-day point-prevalence abstinence (47%) than nonusers (20%; $p = 0.03$; odds ratio = 3.30; 95% CI 1.03, 10.56; $p = 0.04$; use by type of service interaction $p > 0.05$) but not continuous abstinence (used, 40%; not used, 20%; $p = 0.10$; odds ratio = 2.47; 95% CI 0.77, 7.94; $p = 0.13$; use by type of service interaction $p > 0.05$).

**RESEARCH QUESTION 1:
USE AND USABILITY OF CESSATION SERVICES**

Overall, 60% ($n = 61$ of 102) of smokers used the allocated service. In the REQ-Mobile group, 61% of smokers ($n = 31$) opened the inbox, and these smokers did so an average of 26.0 times (standard deviation [SD], 30.3), received 128.5 short messages (SD, 61.9), and opened just over 76 short messages (Table 3). Short support documents were used most, testimonials and list-making tool less, and quit date screen least. Use was highly intercorrelated (Table 3). With onQ, 59% of smokers ($n = 30$) used the service, receiving on average 107.8 text messages (SD, 71.8; mean, 5.2 text messages sent to onQ during [SD, 3.6]). The Web server also recorded that many smokers revisited the Quit Coach program on the study Web site: revisits to the Web site, 73% ($n = 55$ of 75; $n = 27$ REQ-Mobile, $n = 28$ onQ), mean of 5.2 revisits; revisits to Quit Coach, 26% ($n = 18$ of 68; 8 REQ-Mobile, 10 onQ), mean of 2.7 revisits.

Smokers who used REQ-Mobile and were followed up at 6 weeks ($n = 16$) evaluated its usability favorably. They considered it simple (75%), reliable (75%), useful (75%), easy (69%), designed for them (69%), satisfying (69%), and easy to learn (69%). Most felt confident using it (63%), did not need help (69%), and would likely use it frequently (94%). Evaluations were less positive on REQ-Mobile being straightforward (63%), consistent (56%), and well integrated (56%). Most smokers (75%) rated it overall as user-friendly.

QUITTING BEHAVIOR

Of the 63 smokers who initialized REQ-Mobile or onQ, 97% set a quit date. A majority of posttest completers reported a quit attempt at 6 weeks (53%) and 12 weeks (99%; 66% of all cases [i.e., lost = no attempt]) and declared they were abstinent (point-prevalence abstinence) at 6 weeks (53%) and 12 weeks (66%; 44% of all cases).

QUITTING BEHAVIOR BY EXPERIMENTAL CONDITION

Significantly more onQ smokers had quit at 6 weeks than REQ-Mobile smokers (Table 4). OnQ smokers were at significantly later readiness to quit phases, remained abstinent for longer, and were using NRT more compared with REQ-Mobile smokers, but these differences were less apparent at 12 weeks (Table 4).

More frequent use of REQ-Mobile was positively associated with greater abstinence at 12 weeks (number of REQ-Mobile sessions, 30-day point-prevalence abstinence Spearman rank $r = 0.40$, $p = 0.02$ and continuous abstinence $r = 0.35$, $p = 0.04$; number of onQ text messages sent, 30-day point-prevalence abstinence $r = 0.28$, $p = 0.14$ and continuous abstinence $r = 0.31$, $p = 0.09$).

The number of times each REQ-Mobile quitting tool was used, assessed individually, was positively correlated with 12-week abstinence (Table 3). Both absolute and relative use (i.e., proportion each feature was used) showed that the list-making tool was positively correlated with success. Spending relatively more occasions listening to audio testimonials was associated with less abstinence; the positive correlation with opening short messages may be an artifact of taking longer to quit.

Discussion

This study confirmed that a smartphone mobile application could be successfully deployed and that most smokers had considerable interest in its advice. Although the sample may have been highly motivated, smokers provided highly favorable evaluations of REQ-Mobile's usability (although the subsample that used it and completed follow-up may be more favorable than nonusers or non-completers). It was as popular as text messaging by usage. Smokers received slightly more messages from REQ-Mobile than onQ, perhaps

Table 3. Mean Use of REQ-Mobile Application Tools, Intercorrelation (Spearman *r*) of Use of Various Tools, and Correlation of Tool Use with Smoking Abstinence by Smokers Allocated to the REQ-Mobile Group Who Used the Mobile Application and Completed the 12-Week Posttest (*n*= 18)

	LIST MAKER	SHORT SUPPORT DOCUMENTS	AUDIO TESTIMONIALS	SET QUIT DATE	READ A MESSAGE IN INBOX
Mean					
Number of times used	8.00	36.50	9.11	2.00	76.33
Percentage of use	6.16	23.86	6.57	3.38	60.03
Intercorrelation of use of tools and features					
List maker					
Short support documents					
Number of times used	0.82 ^a				
Percentage of use	0.15				
Audio testimonials					
Number of times used	0.79 ^a	0.91 ^a			
Percentage of use	-0.02	-0.25			
Set quit date					
Number of times used	0.59 ^b	0.46	0.53 ^a		
Percentage of use	0.07	-0.68 ^a	-0.01		
Opened message in inbox					
Number of times used	0.81 ^a	0.93 ^a	0.92 ^a	0.50 ^b	
Percentage of use	-0.43	-0.88 ^a	0.22	0.37	
12-week 30-day point-prevalence abstinence					
Number of times used	0.49 ^b	0.19	0.03	0.38	0.12
Percentage of use	0.56 ^b	0.41	-0.31	-0.08	-0.57 ^b
12-week continuous abstinence					
Number of times used	0.45	0.21	0.06	0.35	0.15
Percentage of use	0.54 ^b	0.27	-0.24	-0.09	-0.40

^a*p*<0.05.

^b*p*<0.01.

because they took longer to quit and fewer were abstinent at 6 weeks, a finding of concern. REQ-Mobile's short messages were the most popular feature probably because they were the gateway to other features; short support documents were popular, too, perhaps because of simplicity. The less-popular list-making tool required more effort. Audio testimonials were used less, as in another study.⁴¹

Mobile cessation services can encourage quitting. Nearly all smokers set a quit date during the study. Although the sample's selectivity may have influenced the quitting base rate, more than one in five smokers had continuously quit at 12 weeks, more than achieved in other studies,^{8-10,42,43} including text messaging.^{11-13,16}

It must be pointed out that many smokers returned to use the online Quit Coach, so some of the apparent success of mobile cessation services could be due to the online content. Smokers may expect flexibility in interfaces as wireless connections and cloud computing expands mobility across devices, eliminating barriers of availability,⁴⁴ location,⁴⁵ function,⁴⁶ connection speed, and marketplace fragmentation.⁴⁷

REQ-Mobile, with added interactive features, seemed to move smokers less quickly to initiate quitting than onQ. Use of it was associated with quitting, but both could be a consequence of motivation. Unlike many existing iPhone cessation applications,³⁰ REQ-Mobile was built on theoretical models³⁴⁻³⁶ and directed smokers to use effective treatments.⁴⁸⁻⁵⁵ Its continuous, real-time, mobile nature^{19,56} should have integrated cessation support into smokers' communication environments more than other cessation services (e.g., group sessions, telephone, Web site) when and where they need it. This could have increased involvement with the counseling and quitting,^{19,20,23-25} and provided social support for quitting.^{20,26-29}

Even with the seemingly apparent advantages of mobile applications, the onQ program appeared to produce more successful quitting, a finding that needs further exploration. The most plausible explanation is simplicity. Smokers are often ambivalent about their smoking and quitting,⁵⁷ so are easily discouraged or distracted from making quit attempts. Simplicity predicts adopting new products⁵⁸ and complexity interfered with another mobile cessation program.¹⁵ It should be noted that REQ-Mobile smokers were switched to onQ messages after having quit for 1 week (because of

Table 4. Differences in Percentage of Smokers Quitting (95% Confidence Intervals), Readiness to Quit, Length of Time Since Quitting, and Use of Nicotine Replacement Therapy Between REQ-Mobile Application and onQ Text Messaging Conditions

	REQ-MOBILE APPLICATION	ONQ TEXT MESSAGING	ESTIMATED EFFECT SIZE ^a	P
6-week posttest				
Made a quit attempt (completers <i>n</i> =66)	33% (16–50%)	73% (57–89%)	0.82	<0.01
Point-prevalence abstinence (on day of interview; completers <i>n</i> =66) ^b	33% (16–50%)	73% (57–89%)	0.82	<0.01
7-day point prevalence abstinence (completers <i>n</i> =66)	30% (14–47%)	58% (40–75%)	0.57	0.03
Readiness to quit (completers <i>n</i> =66)				
Open to quitting	0%	0%	0.89 ^c	0.01
Like to quit	12%	6%		
Planning to quit	15%	9%		
Have set a quit date	40%	12%		
Have quit	33%	73%		
How long ago quit (<i>n</i> =35 who reported point-prevalence abstinence)				
Less than 1 month	45%	29%	0.33	0.35
1 month or more	55%	71%		
Use of NRT (completers <i>n</i> =66)				
None	31%	9%	0.26 ^d	0.02
1 week or less	38%	21%		
2 weeks	12%	40%		
More than 2 weeks	19%	30%		
12-week posttest				
Made a quit attempt				
Intent-to-treat (<i>n</i> =102)	63% (49–76%)	69% (55–82%)	0.13	0.53
Completers (<i>n</i> =68)	97% (91–100%)	100%	0.35	0.49
Point-prevalence abstinence (on day of interview) ^b				
Intent-to-treat (<i>n</i> =102)	33% (20–47%)	55% (41–69%)	0.45	0.03
Completers (<i>n</i> =68)	52% (34–70%)	80% (66–94%)	0.60	0.01
30-day point prevalence abstinence				
Intent-to-treat (<i>n</i> =102)	18% (7–28%)	31% (18–45%)	0.30	0.11
Completers (<i>n</i> =68)	27% (11–43%)	46% (28–63%)	0.40	0.12
Continuous abstinence				
Intent-to-treat (<i>n</i> =102)	16% (5–26%)	27% (15–40%)	0.27	0.15
Completers (<i>n</i> =68)	24% (9–40%)	40% (23–57%)	0.35	0.17

continued →

Table 4. Differences in Percentage of Smokers Quitting (95% Confidence Intervals), Readiness to Quit, Length of Time Since Quitting, and Use of Nicotine Replacement Therapy Between REQ-Mobile Application and onQ Text Messaging Conditions *continued*

	REQ-MOBILE APPLICATION	ONQ TEXT MESSAGING	ESTIMATED EFFECT SIZE ^a	P
Readiness to quit (completers <i>n</i> =68)				
Open to quitting	6%	5%	0.60 ^c	0.04
Like to quit	3%	6%		
Planning to quit	15%	3%		
Have set a quit date	24%	6%		
Have quit (point-prevalence abstinence)	52%	80%		
How long ago quit (<i>n</i> =45 who reported point-prevalence abstinence)				
Less than 1 month	29%	14%	0.37	0.27
1 month or more	71%	86%		
Use of NRT (completers <i>n</i> =68)				
None	25%	6%	0.06 ^d	0.07
1 week or less	22%	17%		
2 weeks	19%	40%		
More than 2 weeks	34%	37%		

^aEffect size was computed as $h = 2 \arcsin(\sqrt{p_1}) - 2 \arcsin(\sqrt{p_2})$.

^bIn some cases, the 95% confidence intervals for the quitting measures overlap, but the difference between conditions is statistically significant. This can occur because confidence intervals are calculated based on the subsample within a condition and are conservative, whereas the statistical test is calculated based on the larger entire sample.⁷²

^cEffect size on readiness to quit variable was estimated by converting responses to two categories: "have quit" versus all other responses combined.

^dEffect size on nicotine replacement therapy (NRT) variables was estimated by converting response to two categories: "more than 2 weeks" versus all other responses combined.

lack of project funds for programming this phase), which may explain why this superiority of onQ declined from the 6-week to 12-week follow-up. It is also possible that smokers were new to smartphones and mastered them over time, improving REQ-Mobile's impact at 12 weeks. Furthermore, smokers may have monitored their text inbox, where onQ messages were sent, more routinely than REQ-Mobile's private inbox. The free phone provided to all participants may have been a second phone for some smokers. Thus, smokers needed to monitor two inboxes, an extra step that may have allowed them to ignore messages and read them in batches (76 messages were opened in 26 REQ-Mobile sessions). However, REQ-Mobile (but not onQ) displayed a notice on the main screen every 15 min when its inbox contained unread messages to prompt more immediate reading, but integrating REQ-Mobile's inbox with the normal text messaging inbox might improve it. Thus, quitting trajectories should be tracked when studying mobile technologies. Less plausible is that the difference could have arisen because randomization failed or groups were nonequivalent on income.

Some components of REQ-Mobile seemed to support quitting more than others. Theoretically, the list-making tool should have enhanced motivation and self-efficacy for quitting, but more motivated smokers

may have been most willing to use it. What combination of content, involvement, and motivation makes any tool effective needs further exploration. Setting a quit date can be an important step toward initiating a quit.⁵⁹ Some smokers spontaneously, successfully quit,^{60,61} but this may be less common among younger smokers.⁶² Audio testimonials did not appear useful, perhaps because their content was inappropriate or they distracted smokers from other important tasks. They might be eliminated to simplify the mobile application. This suggests caution when considering using social media, which is dominated by user-generated content, for they may divert smokers from more useful cessation content.

This study had several strengths and limitations. Strengths included recruiting smokers throughout the United States and balancing differences with randomization. However, the small sample limited statistical power. Self-reported abstinence is over-reported,⁶³⁻⁶⁵ but randomization to two active interventions should have equalized this bias. Prescreening and free phones may have recruited highly motivated smokers, limiting generalizability to populations such as quit line callers, or for free stuff (although the high quit rates suggest this is unlikely).

Mobile applications may be effective for brief community-based smoking cessation, as smartphone ownership increases (half of U.S. mobile phone users use applications^{66,67}). Future efforts are needed on how to direct smokers to use applications,³¹ a challenge for all cessation counseling.^{5,7,15,68-71}

Acknowledgments

This project was supported by grant CA107444 from the National Cancer Institute.

Disclosure Statement

D.B.B. is an employee of Klein Buendel, Inc, a for-profit company. D.B.B.'s spouse is the sole owner and President of Klein Buendel. D.B.B. is a member of Wedge Communications LLC, a technology transfer company that has licensed the REQ-Mobile mobile application. R.B. was responsible for developing the onQ and Quit Coach programs, but he holds no commercial interest in them. E.P.B. is an employee of Klein Buendel, Inc., and a member of Wedge Communications LLC. J.H.S. is an employee of Klein Buendel, Inc. D.E.Z. has no competing financial interests.

REFERENCES

1. Centers for Disease Control and Prevention. Fast facts. 2012. Available at www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/ (last accessed May 3, 2013).
2. American Cancer Society. Guide to quitting smoking. 2012. Available at www.cancer.org/Healthy/StayAwayfromTobacco/GuidetoQuittingSmoking/index (last accessed May 3, 2013).
3. Centers for Disease Control and Prevention. Cigarette smoking among adults and trends in smoking cessation—United States, 2008. *MMWR Morb Mortal Wkly Rep* 2009;58:1227-1232.
4. Centers for Disease Control and Prevention. Smoking cessation. 2013. Available at www.cdc.gov/tobacco/data_statistics/fact_sheets/cessation/quitting/index.htm (last accessed May 3, 2013).
5. Murthy P, Subodh BN. Current developments in behavioral interventions for tobacco cessation. *Curr Opin Psychiatry* 2010;23:151-156.
6. Gerbert B, Berg-Smith S, Mancuso M, et al. Using innovative video doctor technology in primary care to deliver brief smoking and alcohol intervention. *Health Promot Pract* 2003;4:249-261.
7. Gaglio B, Smith TL, Estabrooks PA, Ritzwoller DP, Ferro EF, Glasgow RE. Using theory and technology to design a practical and generalizable smoking reduction intervention. *Health Promot Pract* 2010;11:675-684.
8. Cijvljak M, Sheikh A, Stead LF, Car J. Internet-based interventions for smoking cessation. *Cochrane Database Syst Rev* 2010;9:CD007078.
9. Myung SK, McDonnell DD, Kazinets G, Seo HG, Moskowitz JM. Effects of Web- and computer-based smoking cessation programs: Meta-analysis of randomized controlled trials. *Arch Intern Med* 2009;169:929-937.
10. Shahab L, McEwen A. Online support for smoking cessation: A systematic review of the literature. *Addiction* 2009;104:1792-1804.
11. Obermayer JL, Riley WT, Asif O, Jean-Mary J. College smoking-cessation using cell phone text messaging. *J Am Coll Health* 2004;53:71-78.
12. Rodgers A, Corbett T, Bramley D, et al. Do u smoke after txt? Results of a randomised trial of smoking cessation using mobile phone text messaging. *Tob Control* 2005;14:255-261.

13. Bramley D, Riddell T, Whittaker R, et al. Smoking cessation using mobile phone text messaging is as effective in Maori as non-Maori. *N Z Med J* 2005;118(1216):U1494.
14. Free C, Knight R, Robertson S, et al. Smoking cessation support delivered via mobile phone text messaging (txt2stop): A single-blind, randomised trial. *Lancet* 2011;378:49-55.
15. Wetter DW, McClure JB, Cofta-Woerpel L, et al. A randomized clinical trial of a palmtop computer-delivered treatment for smoking relapse prevention among women. *Psychol Addict Behav* 2011;25:365-371.
16. Whittaker R, Borland R, Bullen C, Lin RB, McRobbie H, Rodgers A. Mobile phone-based interventions for smoking cessation. *Cochrane Database Syst Rev* 2009;7:CD006611.
17. Whittaker R, Maddison R, McRobbie H, et al. A multimedia mobile phone-based youth smoking cessation intervention: Findings from content development and piloting studies. *J Med Internet Res* 2008;10:e49.
18. Berkman ET, Dickenson J, Falk EB, Lieberman MD. Using SMS text messaging to assess moderators of smoking reduction: Validating a new tool for ecological measurement of health behaviors. *Health Psychol* 2011;30:186-194.
19. Abroms L, Padmanabhan P, Evans W. Mobile phones for health communication to promote behavior change. In: Noar S, Harrington N, eds. *E-health applications: Promising strategies for behavior change*. New York: Routledge, 2012:147-166.
20. Fjeldsoe B, Miller Y, Marshall A. Text messaging interventions for chronic disease management and health promotion. In: Noar S, Harrington N, eds. *E-health applications: Promising strategies for behavior change*. New York: Routledge, 2012:167-186.
21. Heron KE, Smyth JM. Ecological momentary interventions: Incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010;15:1-39.
22. Riley W, Rivera D, Atienza A, Nilsen W, Allison S, Mermelstein R. Health behavior models in the age of mobile interventions: Are our theories up to the task? *Trans Behav Med* 2011;1:53-71.
23. Gerber BS, Stolley MR, Thompson AL, Sharp LK, Fitzgibbon ML. Mobile phone text messaging to promote healthy behaviors and weight loss maintenance: A feasibility study. *Health Inform J* 2009;15:17-25.
24. Suffoletto B, Callaway C, Kristan J, Kraemer K, Clark DB. Text-message-based drinking assessments and brief interventions for young adults discharged from the emergency department. *Alcohol Clin Exp Res* 2012;36:552-560.
25. Parrott R. Motivation to attend to health messages: Presentation of content and linguistic considerations. In: Maibach E, Parrott R, eds. *Designing health messages: Approaches from communication theory and public health practice*. Thousand Oaks, CA: Sage, 1995:7-23.
26. Westmaas JL, Bontemps-Jones J, Bauer JE. Social support in smoking cessation: Reconciling theory and evidence. *Nicotine Tob Res* 2010;12:695-707.
27. Mermelstein R, Cohen S, Lichtenstein E, Baer JS, Kamarck T. Social support and smoking cessation and maintenance. *J Consult Clin Psychol* 1986;54:447-453.
28. Cohen S, Lichtenstein E. Partner behaviors that support quitting smoking. *J Consult Clin Psychol* 1990;58:304-309.
29. Graham AL, Papanonatos GD, Kang H, Moreno JL, Abrams DB. Development and validation of the online social support for smokers scale. *J Med Internet Res* 2011;13:e69.
30. Abroms LC, Padmanabhan N, Thaweethai L, Phillips T. iPhone apps for smoking cessation: A content analysis. *Am J Prev Med* 2011;40:279-285.
31. Backinger CL, Augustson EM. Where there's an app, there's a way? *Am J Prev Med* 2011;40:390-391.
32. Borland R, Balmford J, Benda P. Population-level effects of automated smoking cessation help programs: A randomized controlled trial. *Addiction* 2012;108:618-628.
33. Balmford J, Borland R, Benda P, Howard S. Factors associated with use of automated smoking cessation interventions: Findings from the eQuit study. *Health Educ Res* 2012;28:288-299.

34. Borland R, Balmford J, Hunt D. The effectiveness of personally tailored computer-generated advice letters for smoking cessation. *Addiction* **2004**;99:369–377.
35. Balmford J, Borland R, Benda P. Patterns of use of an automated interactive personalized coaching program for smoking cessation. *J Med Internet Res* **2008**;10:e54.
36. Balmford J, Borland R, Li L, Ferretter I. Usage of an Internet smoking cessation resource: The Australian QuitCoach. *Drug Alcohol Rev* **2009**;28:66–72.
37. Bandura A. *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs, NJ: Prentice Hall, **1986**.
38. Prochaska JO, Redding CA, Evers KE. The transtheoretical model and stages of change. In: Glanz K, Rimer BK, Lewis FM, eds. *Health behavior and health education*. San Francisco: Jossey-Bass, **2005**:60–66.
39. Mayer RE, Moreno R. Nine ways to reduce cognitive load in multimedia learning. *Educ Psychol* **2003**;38:43–52.
40. Bangor A, Kortum P, Miller J. Determining what individual SUS scores mean: Adding an adjective rating scale. *J Usabil Stud* **2009**;4:114–123.
41. Whittaker R, Dorey E, Bramley D, et al. A theory-based video messaging mobile phone intervention for smoking cessation: Randomized controlled trial. *J Med Internet Res* **2011**;13:e10.
42. Stead LF, Lancaster T. Group behavior therapy programmes for smoking cessation. *Cochrane Database Syst Rev* **2005**;2:CD0001007.
43. Baillie AJ, Mattick RP, Hall W, Webster P. Meta-analytic review of the efficacy of smoking cessation interventions. *Drug Alcohol Rev* **1994**;13:157–170.
44. Nielsen Wire. Connected devices: How we use tablets in the U.S. **2011**. Available at www.nielsen.com/us/en/newswire/2011/connected-devices-how-we-use-tablets-in-the-u-s.html (last accessed May 3, 2013).
45. Bosker B. The top use for tablets: Gaming (STATS). **2011**. Available at www.huffingtonpost.com/2011/04/08/the-top-use-for-tablets-stats_n_846565.html (last accessed May 3, 2013).
46. ComScore. ComScore reports March 2011 U.S. mobile subscriber market share. **2011**. Available at www.comscore.com/Press_Events/Press_Releases/2011/5/comScore_Reports_March_2011_U.S._Mobile_Subscriber_Market_Share (last accessed May 3, 2013).
47. Linsalata D, Slawsby A. White paper: Addressing growing handset complexity with software solutions. **2005**. Available at <http://mobiledevelopers.pbworks.com/f/Challenges+along+mobile+supply+chain+players+by+IDC.pdf> (last accessed May 3, 2013).
48. Miller N, Frieden TR, Liu SY, et al. Effectiveness of a large-scale distribution programme of free nicotine patches: A prospective evaluation. *Lancet* **2005**;365:1849–1854.
49. Cummings KM, Hyland A, Fix B, et al. Free nicotine patch giveaway program 12-month follow-up of participants. *Am J Prev Med* **2006**;31:181–184.
50. Cummings KM, Hyland A, Carlin-Menter S, Mahoney MC, Willett J, Juster HR. Costs of giving out free nicotine patches through a telephone quit line. *J Public Health Manag Pract* **2011**;17:E16–E23.
51. Cummings KM, Fix B, Celestino P, Carlin-Menter S, O'Connor R, Hyland A. Reach, efficacy, and cost-effectiveness of free nicotine medication giveaway programs. *J Public Health Manag Pract* **2006**;12:37–43.
52. Fellows JL, Bush T, McAfee T, Dickerson J. Cost effectiveness of the Oregon quitline “free patch initiative.” *Tob Control* **2007**;16(Suppl 1):i47–i52.
53. Bush TM, McAfee T, Deprey M, et al. The impact of a free nicotine patch starter kit on quit rates in a state quit line. *Nicotine Tob Res* **2008**;10:1511–1516.
54. Tinkelman D, Wilson SM, Willett J, Sweeney CT. Offering free NRT through a tobacco quitline: Impact on utilisation and quit rates. *Tob Control* **2007**;16(Suppl 1):i42–i46.
55. An LC, Schillo BA, Kavanaugh AM, et al. Increased reach and effectiveness of a statewide tobacco quitline after the addition of access to free nicotine replacement therapy. *Tob Control* **2006**;15:286–293.
56. Fjeldsoe BS, Marshall AL, Miller YD. Behavior change interventions delivered by mobile telephone short-message service. *Am J Prev Med* **2009**;36:165–173.
57. Balmford J, Borland R. What does it mean to want to quit? *Drug Alcohol Rev* **2008**;27:21–27.
58. Rogers EM. *Diffusion of innovations*, 5th ed. New York: Free Press, **2003**.
59. Prochaska JO, Velicer WF. The transtheoretical model of health behavior change. *Am J Health Promot* **1997**;12:38–48.
60. West R, Sohal T. “Catastrophic” pathways to smoking cessation: Findings from national survey. *BMJ* **2006**;332:458–460.
61. Ferguson SG, Shiffman S, Gitchell JG, Sembower MA, West R. Unplanned quit attempts—Results from a U.S. sample of smokers and ex-smokers. *Nicotine Tob Res* **2009**;11:827–832.
62. Cooper J, Borland R, Yong HH, et al. To what extent do smokers make spontaneous quit attempts and what are the implications for smoking cessation maintenance? Findings from the International Tobacco Control four country survey. *Nicotine Tob Res* **2010**;12(Suppl):S51–S57.
63. Lynch B, Bonnie R; Institute of Medicine Committee on Preventing Nicotine Addiction in Children and Youths. *Growing up tobacco free: Preventing nicotine addiction in children and youths*. Washington, DC: National Academy Press, **1993**.
64. Lando HA, Pirie PL, McGovern PG, Pechacek TF, Swim J, Loken B. A comparison of self-help approaches to smoking cessation. *Addict Behav* **1991**;16:183–193.
65. Orleans CT, Schoenbach VJ, Wagner EH, et al. Self-help quit smoking interventions: Effects of self-help materials, social support instructions, and telephone counseling. *J Consult Clin Psychol* **1991**;59:439–448.
66. Smith A. Mobile access 2010. **2010**. Available at www.pewinternet.org/Reports/2010/Mobile-Access-2010/Summary-of-Findings.aspx%20on%20October%202013 (last accessed March 21, 2013).
67. Purcell K, Rainie L, Rosenstiel T, Mitchell A. How mobile devices are changing community information environments. **2011**. Available at www.pewinternet.org/Reports/2011/Local-mobile-news.aspx (last accessed May 3, 2013).
68. Zhu SH, Gardiner P, Cummins S, et al. Quitline utilization rates of African-American and white smokers: The California experience. *Am J Health Promot* **2011**;25(5 Suppl):S51–S58.
69. Brendryen H, Kraft P. Happy ending: A randomized controlled trial of a digital multi-media smoking cessation intervention. *Addiction* **2008**;103:478–484.
70. An LC, Betzner A, Schillo B, et al. The comparative effectiveness of clinic, work-site, phone, and Web-based tobacco treatment programs. *Nicotine Tob Res* **2010**;12:989–996.
71. Zbikowski SM, Jack LM, McClure JB, et al. Utilization of services in a randomized trial testing phone- and web-based interventions for smoking cessation. *Nicotine Tob Res* **2011**;13:319–327.
72. Washington State Department of Health. Guidelines for using confidence intervals for public health assessment. **2002**. Available at http://www.doh.wa.gov/data/guidelines/WordDocs/CI_guidelines.pdf (last accessed January 17, 2012).

Address correspondence to:

David B. Buller, PhD

Klein Buendel, Inc.

1667 Cole Boulevard, Suite 225

Golden, CO 80401

E-mail: dbuller@kleinbuendel.com

Received: May 9, 2013

Revised: June 14, 2013

Accepted: June 20, 2013