



# Intravitreal Bevacizumab as Anti-Vascular Endothelial Growth Factor in the Management of Complications of Proliferative Diabetic Retinopathy

J. Fernando Arevalo, MD, FACS

Retina Division, Wilmer Eye Institute, Johns Hopkins University School of Medicine, Baltimore, USA & The Vitreoretinal Division, King Khaled Eye Specialist Hospital, Riyadh, Kingdom of Saudi Arabia & Faculty of Health Sciences, Stellenbosch University, South Africa

## ABSTRACT

Vascular endothelial growth factor (VEGF) has been shown to be an endothelial cell-specific mitogen and an angiogenic inducer in a variety of in vitro and in vivo models. Furthermore, it has been demonstrated to increase retinal vessel permeability by increasing the phosphorylation of tight junction proteins. Recent work has found elevated levels of VEGF in ocular fluids of patients with proliferative diabetic retinopathy (PDR). Thus, it makes sense to consider anti-VEGF treatments in the management of PDR. The purpose of the current research is to determine if intravitreal bevacizumab as anti-VEGF is helpful in the management of complications of PDR.

## KEY WORDS

Bevacizumab; Anti-Vascular Endothelial Growth Factor; Diabetic Retinopathy

©2013, Medical Hypothesis, Discovery & Innovation (MEHDI) Ophthalmology Journal

All rights reserved.

---

## Correspondence to:

Dr J. Fernando Arevalo, Chief of Vitreoretinal Division, The King Khaled Eye Specialist Hospital, Al-Oruba Street, PO Box 7191, Riyadh 11462, Kingdom of Saudi Arabia. Phone: +966 (1) 482-1234 ext. 3860. Fax: +966 (1) 482-1234 ext. 3727, E-mail: [arevalo@jhmi.edu](mailto:arevalo@jhmi.edu)

---

## INTRODUCTION

Dietary habits and an increase in sedentarism are the main culprits of a worldwide epidemic in diabetes mellitus (DM). In the year 2010, more than 285 million people suffered from DM. The World Health Organization has estimated that by the year 2030 there will be 370 million persons affected with DM in the world, and every one of them will be at risk of developing diabetic retinopathy (DR) [1]. Other studies predict that

the number will be even higher at 439 million persons affected with DM in the world [2]. Prior to the advent of panretinal photocoagulation (PRP), proliferative diabetic retinopathy (PDR) was the main cause of diabetic blindness.

More than 60 years ago, Michaelson speculated on the presence of a Factor X that was capable of inducing retinal angiogenesis or neovascularization [3]. Vascular endothelial growth factor (VEGF) appears to be the most likely candidate to be Michaelson's factor X and the main



molecular mediator in diabetic retinopathy. The underlying problem in diabetic patients is the progressive retinal ischemia caused by the metabolic disarray of chronic hyperglycemia. Hypoxia is a major inducer of VEGF gene transcription. VEGF has been shown to be an endothelial cell-specific mitogen and an angiogenic inducer in a variety of in vitro and in vivo models [4]. Furthermore, it has been demonstrated to increase retinal vessel permeability by increasing the phosphorylation of tight junction proteins. Recent work has found elevated levels of VEGF in ocular fluids of patients with PDR [5-10]. Thus it makes sense to consider anti-VEGF treatments in the management of diabetic macular edema (DME) and PDR.

Several anti-VEGF agents are currently available in clinical practice. Pegaptanib sodium (Macugen®, Eyetech, NY, USA) is an aptamer against the VEGF-165 isoform [11]. Ranibizumab (Lucentis®, Genentech, San Francisco, CA, USA) is a fragment of a humanized monoclonal antibody against all VEGF isoforms [12]. Bevacizumab (Avastin®, Genentech, San Francisco, CA, USA) is a humanized, recombinant monoclonal IgG antibody that binds and inhibits all VEGF isoforms [13-15]. Aflibercept, previously known as VEGF-Trap eye (Eylea®, Regeneron Pharmaceuticals Inc., Tarrytown, NY) is a recombinant fusion protein that consists of portions of human VEGF receptors 1 and 2 that allows it to bind to VEGF-A, VEGF-B and placental growth factor [16]. All of these agents have been used to differing extents in the management of DME and PDR [17-27]. However, there is a tremendous difference in price between these drugs. Bevacizumab costs only a small fraction of Ranibizumab, Pegaptanib or Aflibercept. If Bevacizumab were not available and research for its off-label use not performed, many diabetics in developing countries would not have enjoyed the benefit of this remarkable medication.

We have long been interested in studying the effects of intravitreal bevacizumab in several vitreoretinal conditions including DME and PDR [20-27]. Given the off-label nature of intravitreal bevacizumab, its pharmacokinetics and safety have not been as thoroughly studied as other approved similar drugs. We have previously shown in an open label uncontrolled clinical study that intravitreal bevacizumab appears to be safe and well tolerated [28]. A recent randomized clinical

trial has shown no differences with regards to systemic side effects between intravitreal ranibizumab and intravitreal bevacizumab [29].

### HYPOTHESIS

Intravitreal bevacizumab (IVB) may decrease retinal neovascularization in patients with PDR after six months of follow-up. However, the effect may decrease at 24 months of follow up due to tachyphylaxis, and pan-retinal photocoagulation and/or vitrectomy will be necessary.

### DISCUSSION

The underlying problem in diabetic retinopathy is progressive retinal ischemia caused by the metabolic disarray of chronic hyperglycemia. If the hyperglycemia is not controlled, up-regulation of growth factors and cytokines occur. VEGF is an important, if not the most important cytokine involved. Activation of the VEGF-receptor pathway triggers a network of signaling processes that promotes endothelial cell growth, migration, survival from pre-existing vessels, differentiation and mobilization of endothelial progenitor cells from the bone marrow into the peripheral circulation. An increase in vascular permeability followed by angiogenesis leads to retinal, optic disc and even anterior segment neovascularization [30].

PRP has been the mainstay of treatment for PDR, and its suppressive effect on ocular neovascularization has been well documented [31]. However, substantial regression of new vessels may take weeks after completion of PRP, and in up to one-third of cases, new vessels continue to grow despite initial PRP [32]. In these cases, vitreous hemorrhage may induce visual loss and prevent the need for complete laser. Moreover, macular edema may increase after PRP and cause transient or persistent visual loss [33]. Neovascularization on and around the optic disc (NVD) and vitreous hemorrhage were found to be more frequently associated with severe visual loss despite PRP in the Diabetic Retinopathy Study (DRS) and ETDRS [34-35]. Long intervals between PRP sessions and the variable amount of time required for a favorable response may increase the incidence of complications due to the progression of PDR [35]. In fact, a single



episode of PRP or shorter intervals between PRP episodes, although desirable in severe PDR, and when the patient must travel long distances for treatment, are often associated with acute visual disturbances due to exudative choroidal detachment, retinal detachment, and macular edema [32,36-38]. Several reports document the regression of retinal neovascularization following the administration of intravitreal anti-VEGF compounds in PDR [39-41].

We previously conducted a retrospective study in 43 eyes of 39 patients with PDR that had persistent retinal neovascularization and concomitant DME [24]. Of the total of 43 eyes, 31 (72.1%) eyes had been previously treated with scatter photocoagulation at least six months before intravitreal bevacizumab, no eyes had prior focal/grid laser photocoagulation, and no eyes had a previous intravitreal triamcinolone injection. Twenty-four (55.8%) eyes were treated with an intravitreal injection of 1.25 mg and 19 (44.2%) eyes were treated with 2.5 mg of bevacizumab.

At the end of the two year follow-up (Restrepo et al. The Association for Research in Vision and Ophthalmology Annual Meeting 2010. Fort Lauderdale, FL. Unpublished data), the mean baseline BCVA of logMAR =  $0.94 \pm 0.38$  ETDRS (20/176) improved to a mean BCVA of logMAR =  $0.67 \pm 0.39$  ETDRS (20/94) ( $p < 0.0005$ ). Final BCVA analysis by sub-groups demonstrated that 35 (81.4%) eyes remained stable, five (11.6%) eyes improved two or more ETDRS lines of BCVA, and three (7%) eyes decreased two or more ETDRS lines of BCVA. These visual improvements were accompanied by a statistically significant decrease in OCT CMT [23-24].

Of the 43 eyes, 17 (39.5%) eyes showed total regression of retinal neovascularization. Fifteen (34.9%) eyes demonstrated partial regression of retinal neovascularization, and 11 (25.6) eyes showed no regression of retinal neovascularization. When divided by type of retinal neovascularization, of the total of 43 eyes, 15 (34.9%) eyes showed total regression of NVE, 10 (23.3%) eyes demonstrated partial regression of NVE, and 12 (27.9%) eyes showed no regression of NVE. Thirteen (30.2%) eyes showed total regression of NVD, 14 (32.5%) eyes demonstrated partial regression of NVD, and 16 (37.2%) eyes showed no regression of NVD [24]. Interestingly, the above 24 month neovascularization regression data compares unfavorably with our

previously published six-month data [26]. This suggests that anti-VEGF treatment of PDR is not a permanent solution and that consolidation with either PRP and vitrectomy is needed.

## CONCLUSION

Given the off-label cost effective nature of intravitreal bevacizumab, its clinical effectiveness and safety have not been as thoroughly studied as other approved similar drugs. Our results may add evidence that IVB should be used as an adjuvant to PRP in patients with PDR, and that PRP and/or vitrectomy is necessary to control PDR in the long-run in a larger number of patients.

## DISCLOSURE

The author reports no conflicts of interest in this work. This manuscript is part of PhD thesis on "Intravitreal Bevacizumab as Anti-Vascular Endothelial Growth Factor in the Management of Complications of Diabetic Retinopathy". This study partially supported by the Arevalo-Coutinho Foundation for Research in Ophthalmology, Caracas, Venezuela.

## REFERENCES

1. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care*. 2004 May;27(5):1047-53. PMID: 15111519.
2. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res Clin Pract*. 2010 Jan;87(1):4-14. PMID: 19896746.
3. Michaelson IC. The Mode of Development of the Vascular System of the Retina, with some Observations on the Significance for Certain Retinal Diseases. *Trans Ophthalmol Soc UK*. 1948;68:137-80.
4. Ferrara N. Vascular endothelial growth factor: basic science and clinical progress. *Endocr Rev*. 2004 Aug;25(4):581-611. PMID: 15294883.
5. Adamis AP, Miller JW, Bernal MT, D'Amico DJ, Folkman J, Yeo TK, Yeo KT. Increased vascular endothelial growth factor levels in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol*. 1994 Oct 15;118(4):445-50. PMID: 7943121.
6. Pe'er J, Folberg R, Itin A, Gnessin H, Hemo I, Keshet E. Upregulated expression of vascular endothelial growth factor in proliferative diabetic retinopathy. *Br J Ophthalmol*. 1996 Mar;80(3):241-5. PMID: 8703862.
7. Aiello LP, Avery RL, Arrigg PG, Keyt BA, Jampel HD, Shah ST, Pasquale LR, Thieme H, Iwamoto MA, Park JE, et al. Vascular endothelial growth



- factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *N Engl J Med.* 1994 Dec 1;331(22):1480-7. PMID: 7526212.
8. Malecaze F, Clamens S, Simorre-Pinatel V, Mathis A, Chollet P, Favard C, Bayard F, Plouet J. Detection of vascular endothelial growth factor messenger RNA and vascular endothelial growth factor-like activity in proliferative diabetic retinopathy. *Arch Ophthalmol.* 1994 Nov;112(11):1476-82. PMID: 7980139.
9. Tolentino MJ, McLeod DS, Taomoto M, Otsuji T, Adamis AP, Luty GA. Pathologic features of vascular endothelial growth factor-induced retinopathy in the nonhuman primate. *Am J Ophthalmol.* 2002 Mar;133(3):373-85. PMID: 11860975.
10. Tolentino MJ, Miller JW, Gragoudas ES, Jakobiec FA, Flynn E, Chatzistefanou K, Ferrara N, Adamis AP. Intravitreal injections of vascular endothelial growth factor produce retinal ischemia and microangiopathy in an adult primate. *Ophthalmology.* 1996 Nov;103(11):1820-8. PMID: 8942877.
11. Fine SL, Martin DF, Kirkpatrick P. Pegaptanib sodium. *Nat Rev Drug Discov.* 2005 Mar;4(3):187-8. PMID: 15770779.
12. Dugel PU. Ranibizumab treatment of patients with ocular diseases. *Int Ophthalmol Clin.* 2006 Fall;46(4):131-40. PMID: 17060799.
13. Bevacizumab. Anti-VEGF monoclonal antibody, avastin, rhumab-VEGF. *Drugs R D.* 2002;3(1):28-30. PMID: 11881526.
14. Rosenfeld PJ, Fung AE, Puliafito CA. Optical coherence tomography findings after an intravitreal injection of bevacizumab (avastin) for macular edema from central retinal vein occlusion. *Ophthalmic Surg Lasers Imaging.* 2005 Jul-Aug;36(4):336-9. PMID: 16156153.
15. Rosenfeld PJ, Moshfeghi AA, Puliafito CA. Optical coherence tomography findings after an intravitreal injection of bevacizumab (avastin) for neovascular age-related macular degeneration. *Ophthalmic Surg Lasers Imaging.* 2005 Jul-Aug;36(4):331-5. PMID: 16156152.
16. Aflibercept: AVE 0005, AVE 005, AVE0005, VEGF Trap - Regeneron, VEGF Trap (R1R2), VEGF Trap-Eye. *Drugs R D.* 2008;9(4):261-9. PMID: 18588357.
17. Nguyen QD, Brown DM, Marcus DM, Boyer DS, Patel S, Feiner L, Gibson A, Sy J, Rundle AC, Hopkins JJ, Rubio RG, Ehrlich JS; RISE and RIDE Research Group. Ranibizumab for diabetic macular edema: results from 2 phase III randomized trials: RISE and RIDE. *Ophthalmology.* 2012 Apr;119(4):789-801. PMID: 22330964.
18. Cunningham ET Jr, Adamis AP, Altaweel M, Aiello LP, Bressler NM, D'Amico DJ, Goldbaum M, Guyer DR, Katz B, Patel M, Schwartz SD; Macugen Diabetic Retinopathy Study Group. A phase II randomized double-masked trial of pegaptanib, an anti-vascular endothelial growth factor aptamer, for diabetic macular edema. *Ophthalmology.* 2005 Oct;112(10):1747-57. PMID: 16154196.
19. Sultan MB, Zhou D, Loftus J, Dombi T, Ice KS; Macugen 1013 Study Group. A phase 2/3, multicenter, randomized, double-masked, 2-year trial of pegaptanib sodium for the treatment of diabetic macular edema. *Ophthalmology.* 2011 Jun;118(6):1107-18. PMID: 21529957.
20. Arevalo JF, Garcia-Amaris RA. Intravitreal bevacizumab for diabetic retinopathy. *Curr Diabetes Rev.* 2009 Feb;5(1):39-46. Review. PMID: 19199897.
21. Arevalo JF, Fromow-Guerra J, Quiroz-Mercado H, Sanchez JG, Wu L, Maia M, Berrocal MH, Solis-Vivanco A, Farah ME; Pan-American Collaborative Retina Study Group. Primary intravitreal bevacizumab (Avastin) for diabetic macular edema: results from the Pan-American Collaborative Retina Study Group at 6-month follow-up. *Ophthalmology.* 2007 Apr;114(4):743-50. PMID: 17398322.
22. Arevalo JF, Sanchez JG, Fromow-Guerra J, Wu L, Berrocal MH, Farah ME, Cardillo J, Rodríguez FJ; Pan-American Collaborative Retina Study Group (PACORES). Comparison of two doses of primary intravitreal bevacizumab (Avastin) for diffuse diabetic macular edema: results from the Pan-American Collaborative Retina Study Group (PACORES) at 12-month follow-up. *Graefes Arch Clin Exp Ophthalmol.* 2009 Jun;247(6):735-43. PMID: 19189118.
23. Arevalo JF, Sanchez JG, Lasave AF, Wu L, Maia M, Bonafonte S, Brito M, Alezzandrini AA, Restrepo N, Berrocal MH, Saravia M, Farah ME, Fromow-Guerra J, Morales-Canton V; Pan-American Collaborative Retina Study Group (PACORES). Intravitreal Bevacizumab (Avastin®) for Diabetic Retinopathy at 24-months: The 2008 Juan Verdaguera-Planas Lecture. *Curr Diabetes Rev.* 2010 Sep;6(5):313-22. PMID: 20594160.
24. Arevalo JF, Sanchez JG, Lasave AF, Wu L, Maia M, Bonafonte S, Brito M, Alezzandrini AA, Restrepo N, Berrocal MH, Saravia M, Farah ME, Fromow-Guerra J, Morales-Canton V. Intravitreal Bevacizumab (Avastin) for Diabetic Retinopathy: The 2010 GLADAOF Lecture. *J Ophthalmol.* 2011;2011:584238. PMID: 21584260.
25. Arevalo JF, Sanchez JG, Wu L, Maia M, Alezzandrini AA, Brito M, Bonafonte S, Lujan S, Diaz-Llopis M, Restrepo N, Rodríguez FJ, Udaondo-Mirete P; Pan-American Collaborative Retina Study Group. Primary intravitreal bevacizumab for diffuse diabetic macular edema: the Pan-American Collaborative Retina Study Group at 24 months. *Ophthalmology.* 2009 Aug;116(8):1488-97, 1497.e1. PMID: 19545900.
26. Arevalo JF, Wu L, Sanchez JG, Maia M, Saravia MJ, Fernandez CF, Evans T. Intravitreal bevacizumab (Avastin) for proliferative diabetic retinopathy: 6-months follow-up. *Eye (Lond).* 2009 Jan;23(1):117-23. PMID: 17891058.
27. Do DV, Schmidt-Erfurth U, Gonzalez VH, Gordon CM, Tolentino M, Berliner AJ, Vittori R, Rückert R, Sandbrink R, Stein D, Yang K, Beckmann K, Heier JS. The DA VINCI Study: phase 2 primary results of VEGF Trap-Eye in patients with diabetic macular edema. *Ophthalmology.* 2011 Sep;118(9):1819-26. PMID: 21546089.
28. Wu L, Martínez-Castellanos MA, Quiroz-Mercado H, Arevalo JF, Berrocal MH, Farah ME, Maia M, Roca JA, Rodríguez FJ; Pan American Collaborative Retina Study Group (PACORES). Twelve-month safety of intravitreal injections of bevacizumab (Avastin): results of the Pan-American Collaborative Retina Study Group (PACORES). *Graefes Arch Clin Exp Ophthalmol.* 2008 Jan;246(1):81-7. PMID: 17674014.
29. CATT Research Group, Martin DF, Maguire MG, Ying GS, Grunwald JE, Fine SL, Jaffe GJ. Ranibizumab and bevacizumab for neovascular age-related macular degeneration. *N Engl J Med.* 2011 May 19;364(20):1897-908. PMID: 21526923.
30. Grant MB, Afzal A, Spoerri P, Pan H, Shaw LC, Mames RN. The role of growth factors in the pathogenesis of diabetic retinopathy. *Expert Opin Investig Drugs.* 2004 Oct;13(10):1275-93. PMID: 15461557.



31. Photocoagulation treatment of proliferative diabetic retinopathy. Clinical application of Diabetic Retinopathy Study (DRS) findings, DRS Report Number 8. The Diabetic Retinopathy Study Research Group. *Ophthalmology*. 1981 Jul;88(7):583-600. PMID: 7196564.
32. Doft BH, Blankenship GW. Single versus multiple treatment sessions of argon laser panretinal photocoagulation for proliferative diabetic retinopathy. *Ophthalmology*. 1982 Jul;89(7):772-9. PMID: 6181452.
33. Tsujikawa A, Kiryu J, Dong J, Yasukawa T, Suzuma I, Takagi H, Ogura Y. Quantitative analysis of diabetic macular edema after scatter laser photocoagulation with the scanning retinal thickness analyzer. *Retina*. 1999;19(1):59-64. PMID: 10048375.
34. Fong DS, Ferris FL 3rd, Davis MD, Chew EY. Causes of severe visual loss in the early treatment diabetic retinopathy study: ETDRS report no. 24. Early Treatment Diabetic Retinopathy Study Research Group. *Am J Ophthalmol*. 1999 Feb;127(2):137-41. PMID: 10030553.
35. Kaufman SC, Ferris FL 3rd, Seigel DG, Davis MD, DeMets DL. Factors associated with visual outcome after photocoagulation for diabetic retinopathy. Diabetic Retinopathy Study Report #13. *Invest Ophthalmol Vis Sci*. 1989 Jan;30(1):23-8. PMID: 2912911.
36. McDonald HR, Schatz H. Visual loss following panretinal photocoagulation for proliferative diabetic retinopathy. *Ophthalmology*. 1985 Mar;92(3):388-93. PMID: 4039432.
37. McDonald HR, Schatz H. Macular edema following panretinal photocoagulation. *Retina*. 1985 Winter-Spring;5(1):5-10. PMID: 4001591.
38. Shimura M, Yasuda K, Nakazawa T, Kano T, Ohta S, Tamai M. Quantifying alterations of macular thickness before and after panretinal photocoagulation in patients with severe diabetic retinopathy and good vision. *Ophthalmology*. 2003 Dec;110(12):2386-94. PMID: 14644723.
39. Avery RL, Pearlman J, Pieramici DJ, Rabena MD, Castellarin AA, Nasir MA, Giust MJ, Wendel R, Patel A. Intravitreal bevacizumab (Avastin) in the treatment of proliferative diabetic retinopathy. *Ophthalmology*. 2006 Oct;113(10):1695.e1-15. PMID: 17011951.
40. Avery RL. Regression of retinal and iris neovascularization after intravitreal bevacizumab (Avastin) treatment. *Retina*. 2006 Mar;26(3):352-4. PMID: 16508438.
41. Spaide RF, Fisher YL. Intravitreal bevacizumab (Avastin) treatment of proliferative diabetic retinopathy complicated by vitreous hemorrhage. *Retina*. 2006 Mar;26(3):275-8. PMID: 16508426.