



Published in final edited form as:

J Nippon Med Sch. 2013 ; 80(2): 90–94.

Role of the Placenta in Adverse Perinatal Outcomes Among HIV-1 Seropositive Women

William Ackerman IV^{1,‡} and Jesse J. Kwiek²

¹Laboratory of Perinatal Research, Department of Obstetrics and Gynecology, College of Medicine, The Ohio State University, Columbus OH, USA

²The Division of Infectious Diseases, Department of Medicine, The Center for Microbial Interface Biology, and Public Health Preparedness for Infectious Diseases Program, The Ohio State University, Columbus, OH, USA

Abstract

Women seropositive for human immunodeficiency virus type 1 (HIV-1) are at an increased risk for a number of adverse perinatal outcomes. Although efforts to reduce mother-to-child transmission of HIV (MTCT) remain a priority in resource-limited countries, HIV testing and treatment have led to steep declines in MTCT in well-resourced countries. Even so, HIV seropositive pregnant women in the United States continue to deliver a disproportionately high number of preterm and low birth weight infants. In this mini-review, we address the role of the placenta in such HIV-related perinatal sequelae. We posit that adverse perinatal outcomes may result from two mutually non-exclusive routes: (1) HIV infection of the placenta proper, potentially leading to impaired maternal-fetal exchange; and (2) infection of the maternal decidual microenvironment, possibly disrupting normal placental implantation and development. Further research into the relationship between HIV-1 infection and placental pathology may lead to the development of novel strategies to improve birth outcomes among HIV-1 seropositive parturients.

Keywords

Human immunodeficiency virus type 1; placenta; placenta diseases; low birth weight; preterm birth

Women of child-bearing age comprise approximately half of the estimated 33.3 million people currently living with human immunodeficiency virus type 1 (HIV-1)¹. It is well-established that HIV-1 seropositive pregnant women are at increased risk for a number of adverse perinatal outcomes, including spontaneous abortion, stillbirth, intrauterine growth restriction (IUGR), low birth weight (LBW), preterm birth (PTB), infant mortality, and mother-to-child transmission of HIV (MTCT)^{2–13}. In well-resourced countries, access to HIV testing, antiretroviral therapy, and other interventions has resulted in precipitous reductions in MTCT rates^{14,15}; however, similarly pronounced reductions in the risks of PTB and LBW among HIV seropositive women in the United States have not been realized¹⁶. Here, we briefly review possible reasons for these observations.

In normal pregnancy, the placenta facilitates maternal-fetal nutrient and gas exchange, serves as a tolerant immunological barrier, and produces hormones required to maintain pregnancy. Deficiencies in placental function are causally linked to obstetrical complications

[‡]Corresponding Author: William E. Ackerman IV, N525 Doan Hall, 410 West 10th Avenue, Columbus, OH 43210, william.ackerman@osumc.edu, T: (614) 293-4438, F: (614) 293-5728.

and adverse birth outcomes¹⁷. Nevertheless, in the setting of maternal HIV-1 seropositivity, the relationship between placental HIV-1 infection and adverse perinatal outcomes has been difficult to establish¹⁸. At issue has been whether cells of the placenta are susceptible to HIV infection, and by extension, whether such infections could be sufficient to account for perinatal sequelae. Evidence that viral replication may occur locally within the placenta is supported by phylogenetic studies demonstrating genetic divergence between placental and circulating maternal HIV-1 quasispecies^{19–21}. However, efforts to characterize HIV-1 infection within villous placental have proved challenging, resulting in conflicting estimates of the nature and extent of infection depending upon clinical context and the sensitivity and specificity of given means of detection^{18,20,22–30}. Most authors have observed that that *ex vivo* cultures of placental villi support HIV-1 infection^{18,30–35}, yet primary trophoblasts or trophoblast-derived cell lines are generally resistant to infection^{36–42}. It is possible that these discordant results indicate that HIV-1 does not preferentially infect trophoblastic cells, but rather, leukocytes present within the villous parenchyma of *ex vivo* cultures^{24,36,43}. Collectively, these data suggest that while the villous placenta may be infected by HIV in some instances, the exact nature of viral replication within this “placental compartment” requires clarification.

Assuming that placental HIV infection occurs in an appreciable number of clinical cases, it is striking that most histopathological studies have failed to demonstrate specific placental lesions^{18,23,44–47}, which is in contrast to many other infectious pathogens⁴⁷. Among the non-specific pathologies, villitis has been an inconsistent finding^{23,29,44,47,48}, and some (but not all) studies have reported alterations in villous maturity^{45,48,49}. A fairly consistent finding among these studies has been an increase in the proportion of cases with chorioamnionitis among placentas delivered of seropositive women relative to uninfected controls^{23,44,50–52}. Chorioamnionitis is strongly associated with PTB and may also increase the risk for MTCT⁵³; as such, HIV-associated chorioamnionitis might contribute to adverse outcomes among seropositive pregnant women.

In addition to chorioamnionitis, Goldenberg et al. observed in a large study that a high proportion of placentas delivered of HIV seropositive women exhibited deciduitis⁵⁴. Moreover, these authors found that marked leukocyte infiltration into the decidua basalis was significantly associated with adverse outcomes, including LBW and PTB. Such observations highlight the clinical importance of the decidual microenvironment, and particularly the decidua basalis, which is the portion of the uterine mucosa into which the placenta implants. Proper implantation is essential for normal placental function and fetal growth, and requires that placental trophoblastic cells interact with cells of the decidua (decidualized fibroblasts, uterine natural killer cells, and maternal macrophages, among others) in a highly orchestrated manner. Studies of feline immunodeficiency virus (FIV) infection showed that experimental inoculation of pregnant cats triggered decidual inflammation, resulting in reproductive failures^{55,56}. The high incidence of early spontaneous abortions among HIV-infected women⁵⁷ has invited speculation that similar immune dysregulation may occur in HIV-infected human decidua⁵⁸. That HIV can infect maternal immune cells in the decidua is supported by the *ex vivo* experiments of Menu and colleagues, who showed in first trimester decidual histocultures that macrophage-like CD14⁺ cells could be productively infected with some HIV strains⁵⁹. In addition, HIV has been detected in lymphocytes and macrophages of the decidua and chorion *in situ*²⁶. Based on these observations in aggregate, we put forth the hypothesis that HIV infection of decidual macrophages (and possibly lymphocytes) prior to or early in pregnancy might serve to create an inflammatory intrauterine environment that is unfavorable for normal implantation; this may explain a portion of the adverse perinatal outcomes associated with maternal HIV infection.

In summary, infection of the placenta and/or decidual microenvironment represent two mutually non-exclusive routes through which HIV-1 might result in adverse perinatal outcomes. Further research into the relationship between HIV-1 infection and placental pathology may help to improve birth outcomes among HIV-1 seropositive parturients.

References

1. UN Joint Programme on HIV/AIDS. [accessed 30 October 2012] Global Report: UNAIDS Report on the Global AIDS Epidemic: 2010. 2010. Available at: <http://www.unhcr.org/refworld/docid/4cfca9c62.html>
2. Leroy V, Ladner J, Nyiraziraje M, et al. Effect of HIV-1 infection on pregnancy outcome in women in Kigali, Rwanda, 1992–1994. Pregnancy and HIV Study Group. *AIDS*. 1998; 12:643–650. [PubMed: 9583605]
3. Goldstein PJ, Smit R, Stevens M, Sever JL. Association between HIV in pregnancy and antiretroviral therapy, including protease inhibitors and low birth weight infants. *Infect Dis Obstet Gynecol*. 2000; 8:94–98. [PubMed: 10805364]
4. Brocklehurst P, French R. The association between maternal HIV infection and perinatal outcome: a systematic review of the literature and meta-analysis. *Br J Obstet Gynaecol*. 1998; 105:836–848. [PubMed: 9746375]
5. Dreyfuss ML, Msamanga GI, Spiegelman D, et al. Determinants of low birth weight among HIV-infected pregnant women in Tanzania. *Am J Clin Nutr*. 2001; 74:814–826. [PubMed: 11722965]
6. Villamor E, Msamanga G, Spiegelman D, Peterson KE, Antelman G, Fawzi WW. Pattern and predictors of weight gain during pregnancy among HIV-1-infected women from Tanzania. *J Acquir Immune Defic Syndr*. 2003; 32:560–569. [PubMed: 12679710]
7. van Eijk AM, De Cock KM, Ayisi JG, et al. Pregnancy interval and delivery outcome among HIV-seropositive and HIV-seronegative women in Kisumu, Kenya. *Trop Med Int Health*. 2004; 9:15–24. [PubMed: 14728603]
8. Boer K, Nellen JF, Patel D, et al. The AmRo study: pregnancy outcome in HIV-1-infected women under effective highly active antiretroviral therapy and a policy of vaginal delivery. *BJOG*. 2007; 114:148–155. [PubMed: 17305888]
9. Mitgitti R, Seanchaisuriya P, Schelp FP, Marui E, Yanai H. Low birth weight infants born to HIV-seropositive mothers and HIV-seronegative mothers in Chiang Rai, Thailand. *Southeast Asian J Trop Med Public Health*. 2008; 39:273–278. [PubMed: 18564713]
10. Olagbuji BN, Ezeanochie MC, Ande AB, Oboro VO. Obstetric and perinatal outcome in HIV positive women receiving HAART in urban Nigeria. *Arch Gynecol Obstet*. 2010; 281:991–994. [PubMed: 19629507]
11. Parekh N, Ribaudo H, Souda S, et al. Risk factors for very preterm delivery and delivery of very-small-for-gestational-age infants among HIV-exposed and HIV-unexposed infants in Botswana. *Int J Gynaecol Obstet*. 2011; 115:20–25. [PubMed: 21767835]
12. Asavapiryanont S, Kasiwat S. Prevalence of low birthweight infants in HIV-infected women delivered in Rajavithi Hospital. *J Med Assoc Thai*. 2011; 94 (Suppl 2):S66–70. [PubMed: 21717881]
13. Nkhoma ET, Kalilani-Phiri L, Mwapasa V, Rogerson SJ, Meshnick SR. Effect of HIV infection and *Plasmodium falciparum* parasitemia on pregnancy outcomes in Malawi. *Am J Trop Med Hyg*. 2012; 87:29–34. [PubMed: 22764288]
14. Fowler MG, Lampe MA, Jamieson DJ, Kourtis AP, Rogers MF. Reducing the risk of mother-to-child human immunodeficiency virus transmission: past successes, current progress and challenges, and future directions. *Am J Obstet Gynecol*. 2007; 197:S3–9. [PubMed: 17825648]
15. EC Study. The mother-to-child HIV transmission epidemic in Europe: evolving in the East and established in the West. *AIDS*. 2006; 20:1419–1427. [PubMed: 16791017]
16. Schulte J, Dominguez K, Sukalac T, Bohannon B, Fowler MG. Declines in low birth weight and preterm birth among infants who were born to HIV-infected women during an era of increased use of maternal antiretroviral drugs: Pediatric Spectrum of HIV Disease, 1989–2004. *Pediatrics*. 2007; 119:e900–906. [PubMed: 17353299]

17. Jauniaux E, Van Oppenraaij RH, Burton GJ. Obstetric outcome after early placental complications. *Curr Opin Obstet Gynecol.* 2010; 22:452–457. [PubMed: 20930630]
18. Miller RK, Polliotti BM, Laughlin T, et al. Role of the placenta in fetal HIV infection. *Teratology.* 2000; 61:391–394. [PubMed: 10777837]
19. Zachar V, Zacharova V, Fink T, et al. Genetic analysis reveals ongoing HIV type 1 evolution in infected human placental trophoblast. *AIDS Res Hum Retroviruses.* 1999; 15:1673–1683. [PubMed: 10606090]
20. Menu E, Mbopi-Keou FX, Lagaye S, et al. Selection of maternal human immunodeficiency virus type 1 variants in human placenta. European Network for In Utero Transmission of HIV-1. *J Infect Dis.* 1999; 179:44–51. [PubMed: 9841821]
21. Kumar SB, Handelmann SK, Voronkin I, et al. Different regions of HIV-1 subtype C env are associated with placental localization and in utero mother-to-child transmission. *J Virol.* 2011; 85:7142–7152. [PubMed: 21543508]
22. Peuchmaur M, Delfraissy JF, Pons JC, et al. HIV proteins absent from placentas of 75 HIV-1-positive women studied by immunohistochemistry. *AIDS.* 1991; 5:741–745. [PubMed: 1679334]
23. Chandwani S, Greco MA, Mittal K, Antoine C, Krasinski K, Borkowsky W. Pathology and human immunodeficiency virus expression in placentas of seropositive women. *J Infect Dis.* 1991; 163:1134–1138. [PubMed: 1902246]
24. Mattern CF, Murray K, Jensen A, Farzadegan H, Pang J, Modlin JF. Localization of human immunodeficiency virus core antigen in term human placentas. *Pediatrics.* 1992; 89:207–209. [PubMed: 1734385]
25. Martin AW, Brady K, Smith SI, et al. Immunohistochemical localization of human immunodeficiency virus p24 antigen in placental tissue. *Hum Pathol.* 1992; 23:411–414. [PubMed: 1563742]
26. Backe E, Jimenez E, Unger M, Schafer A, Jauniaux E, Vogel M. Demonstration of HIV-1 infected cells in human placenta by in situ hybridisation and immunostaining. *J Clin Pathol.* 1992; 45:871–874. [PubMed: 1430256]
27. Tscherning-Casper C, Papadogiannakis N, Anvret M, et al. The trophoblastic epithelial barrier is not infected in full-term placentae of human immunodeficiency virus-seropositive mothers undergoing antiretroviral therapy. *J Virol.* 1999; 73:9673–9678. [PubMed: 10516083]
28. Dictor M, Lindgren S, Bont J, et al. HIV-1 in placentas of untreated HIV-1-infected women in relation to viral transmission, infectious HIV-1 and RNA load in plasma. *Scand J Infect Dis.* 2001; 33:27–32. [PubMed: 11234974]
29. Katz JM, Fox CH, Eglinton GS, Meyers WA 3rd, Queenan JT. Relationship between human immunodeficiency virus-1 RNA identification in placenta and perinatal transmission. *J Perinatol.* 1997; 17:119–124. [PubMed: 9134510]
30. Sheikh AU, Polliotti BM, Miller RK. Human immunodeficiency virus infection: in situ polymerase chain reaction localization in human placentas after in utero and in vitro infection. *Am J Obstet Gynecol.* 2000; 182:207–213. [PubMed: 10649180]
31. Maury W, Potts BJ, Rabson AB. HIV-1 infection of first-trimester and term human placental tissue: a possible mode of maternal-fetal transmission. *J Infect Dis.* 1989; 160:583–588. [PubMed: 2477467]
32. Amirhessami-Aghili N, Spector SA. Human immunodeficiency virus type 1 infection of human placenta: potential route for fetal infection. *J Virol.* 1991; 65:2231–2236. [PubMed: 2016757]
33. Polliotti BM, Sheikh AU, Subbarao S, et al. HIV-1 infection of human placental villous tissue *in vitro*. *Trophoblast Research.* 1998; 12:205–223.
34. Polliotti BM, Gnall-Sazenski S, Laughlin TS, Miller RK. Inhibitory effects of human chorionic gonadotropin (hCG) preparations on HIV infection of human placenta in vitro. *Placenta.* 2002; 23 (Suppl A):S102–106. [PubMed: 11978066]
35. Kfutwah AK, Mary JY, Nicola MA, et al. Tumour necrosis factor-alpha stimulates HIV-1 replication in single-cycle infection of human term placental villi fragments in a time, viral dose and envelope dependent manner. *Retrovirology.* 2006; 3:36. [PubMed: 16796744]
36. Mano H, Chermann JC. Replication of human immunodeficiency virus type 1 in primary cultured placental cells. *Res Virol.* 1991; 142:95–104. [PubMed: 1896650]

37. Douglas GC, Fry GN, Thirkill T, et al. Cell-mediated infection of human placental trophoblast with HIV in vitro. *AIDS Res Hum Retroviruses*. 1991; 7:735–740. [PubMed: 1742080]
38. Bourinbaier AS, Nagorny R. Human immunodeficiency virus type 1 infection of choriocarcinoma-derived trophoblasts. *Acta Virol*. 1993; 37:21–28. [PubMed: 8105649]
39. Kilani RT, Chang LJ, Garcia-Lloret MI, Hemmings D, Winkler-Lowen B, Guilbert LJ. Placental trophoblasts resist infection by multiple human immunodeficiency virus (HIV) type 1 variants even with cytomegalovirus coinfection but support HIV replication after provirus transfection. *J Virol*. 1997; 71:6359–6372. [PubMed: 9261353]
40. Dolcini G, Derrien M, Chaouat G, Barre-Sinoussi F, Menu E. Cell-free HIV type 1 infection is restricted in the human trophoblast choriocarcinoma BeWo cell line, even with expression of CD4, CXCR4 and CCR5. *AIDS Res Hum Retroviruses*. 2003; 19:857–864. [PubMed: 14585217]
41. Vidricaire G, Tardif MR, Tremblay MJ. The low viral production in trophoblastic cells is due to a high endocytic internalization of the human immunodeficiency virus type 1 and can be overcome by the pro-inflammatory cytokines tumor necrosis factor-alpha and interleukin-1. *J Biol Chem*. 2003; 278:15832–15841. [PubMed: 12604606]
42. Ross AL, Cannou C, Barre-Sinoussi F, Menu E. Proteasome-independent degradation of HIV-1 in naturally non-permissive human placental trophoblast cells. *Retrovirology*. 2009; 6:46. [PubMed: 19445667]
43. McGann KA, Collman R, Kolson DL, et al. Human immunodeficiency virus type 1 causes productive infection of macrophages in primary placental cell cultures. *J Infect Dis*. 1994; 169:746–753. [PubMed: 8133087]
44. Jauniaux E, Nessmann C, Imbert MC, Meuris S, Puissant F, Hustin J. Morphological aspects of the placenta in HIV pregnancies. *Placenta*. 1988; 9:633–642. [PubMed: 3257096]
45. Backe E, Jimenez E, Unger M, et al. Vertical human immunodeficiency virus transmission: a study of placental pathology in relation to maternal risk factors. *Am J Perinatol*. 1994; 11:326–330. [PubMed: 7993509]
46. Temmerman M, Chomba EN, Ndinya-Achola J, Plummer FA, Coppens M, Piot P. Maternal human immunodeficiency virus-1 infection and pregnancy outcome. *Obstet Gynecol*. 1994; 83:495–501. [PubMed: 7907777]
47. Benirschke, KB.; Burton, GJ.; Baergen, RN. Pathology of the Human Placenta. Springer-Verlag; Berlin Heidelberg: 2012. Infectious diseases; p. 557-655.
48. Baurakiades E, Martins AP, Victor Moreschi N, et al. Histomorphometric and immunohistochemical analysis of infectious agents, T-cell subpopulations and inflammatory adhesion molecules in placentas from HIV-seropositive pregnant women. *Diagn Pathol*. 2011; 6:101. [PubMed: 22024147]
49. Kumar SB, Rice CE, Milner DA Jr, et al. Elevated cytokine and chemokine levels in the placenta are associated with in-utero HIV-1 mother-to-child transmission. *AIDS*. 2012; 26:685–694. [PubMed: 22301415]
50. Gichangi PB, Nyongo AO, Temmerman M. Pregnancy outcome and placental weights: their relationship to HIV-1 infection. *East Afr Med J*. 1993; 70:85–89. [PubMed: 8513748]
51. D'Costa GF, Khadke K, Patil YV. Pathology of placenta in HIV infection. *Indian J Pathol Microbiol*. 2007; 50:515–519. [PubMed: 17883121]
52. Schuetz AN, Guarner J, Packard MM, Zaki SR, Shehata BM, Opreas-Ilie G. Infectious disease immunohistochemistry in placentas from HIV-positive and HIV-negative patients. *Pediatr Dev Pathol*. 2011; 14:180–188. [PubMed: 21054157]
53. Goldenberg RL, Vermund SH, Goepfert AR, Andrews WW. Choriodecidual inflammation: a potentially preventable cause of perinatal HIV-1 transmission? *Lancet*. 1998; 352:1927–1930. [PubMed: 9863804]
54. Goldenberg RL, Mudenda V, Read JS, et al. HPTN 024 study: histologic chorioamnionitis, antibiotics and adverse infant outcomes in a predominantly HIV-1-infected African population. *Am J Obstet Gynecol*. 2006; 195:1065–1074. [PubMed: 16875654]
55. Coats KS, Boudreaux CE, Clay BT, Lockett NN, Scott VL. Placental immunopathology in the FIV-infected cat: a role for inflammation in compromised pregnancy? *Vet Immunol Immunopathol*. 2010; 134:39–47. [PubMed: 19896219]

56. Scott VL, Boudreaux CE, Lockett NN, Clay BT, Coats KS. Cytokine dysregulation in early- and late-term placentas from feline immunodeficiency virus (FIV)-infected cats. *Am J Reprod Immunol.* 2011; 65:480–491. [PubMed: 20825375]
57. Shearer WT, Langston C, Lewis DE, et al. Early spontaneous abortions and fetal thymic abnormalities in maternal-to-fetal HIV infection. *Acta Paediatr Suppl.* 1997; 421:60–64. [PubMed: 9240860]
58. Patterson BK, Behbahani H, Kabat WJ, et al. Leukemia inhibitory factor inhibits HIV-1 replication and is upregulated in placentae from nontransmitting women. *J Clin Invest.* 2001; 107:287–294. [PubMed: 11160152]
59. Marlin R, Nugeyre MT, de Truchis C, et al. Antigen-presenting cells represent targets for R5 HIV-1 infection in the first trimester pregnancy uterine mucosa. *PLoS One.* 2009; 4:e5971. [PubMed: 19543402]