



Published in final edited form as:

Cell Microbiol. 2014 March ; 16(3): 396–410. doi:10.1111/cmi.12224.

Identification of a type IV secretion substrate of *Brucella abortus* that participates in the early stages of intracellular survival

Peter H. Döhmer¹, Ezequiel Valguarnera, Cecilia Czibener^{1,*}, and Juan E. Ugalde^{1,*}

¹Instituto de Investigaciones Biotecnológicas “Dr. Rodolfo A. Ugalde”, Instituto Tecnológico de Chascomús, IIB-INTECH, CONICET, Universidad Nacional de San Martín, San Martín, Buenos Aires, Argentina

SUMMARY

Brucella abortus, the etiological agent of bovine brucellosis, is an intracellular pathogen whose virulence is completely dependent on a type IV secretion system. This secretion system translocates effector proteins into the host cell to modulate the intracellular fate of the bacterium in order to establish a secure niche where it actively replicates. Although much has been done in understanding how this secretion system participates in the virulence process, few effector proteins have been identified to date. We describe here the identification of a type IV secretion substrate (SepA) that is only present in *Brucella* spp. and has no detectable homology to known proteins. This protein is secreted in a *virB* dependent manner in a two-step process involving a periplasmic intermediate and secretion is necessary for its function. The deletion mutant showed a defect in the early stages of intracellular replication in professional and non-professional phagocytes although it invades the cells more efficiently than the wild type parental strain. Our results indicate that, even though the mutant was more invasive, it had a defect in excluding the lysosomal marker Lamp-1 and was inactivated more efficiently during the early phases of the intracellular life cycle.

INTRODUCTION

Intracellular pathogens have evolved a variety of strategies to survive and multiply within the host cells. These mechanisms are very versatile and in many cases involve specialized secretion systems devoted to the secretion and translocation of effector proteins to the host cell that interfere or modulate the cellular response to favor the infectious process (Cossart *et al.*, 2010, Ninio *et al.*, 2007). *Brucella* spp are gram-negative facultative intracellular bacteria that cause brucellosis, a wide spread zoonosis that affects mainly domestic animals and humans, causing important economical and sanitary problems in many areas of the world (Corbel, 1997, Godfroid *et al.*, 2011). One of the hallmarks of the infectious cycle of *Brucella* is its ability to survive and multiply in professional phagocytes like macrophages (Atluri *et al.*, 2011), a process that requires the *virB* system, a type IV secretion system homologous to the *Agrobacterium tumefaciens* conjugation-like system that translocates the T-DNA into the plant cell (Alvarez-Martinez *et al.*, 2009). This secretion system is necessary to modulate the intracellular trafficking of the bacteria in the host cell. In the early phases this system contributes to avoid the fusion of the *Brucella* containing vacuole (BCV) with the lysosomes and, in a second phase, redirects its fate to a calnexin-positive

*Corresponding authors Mailing address Instituto de Investigaciones Biotecnológicas “Dr. Rodolfo A. Ugalde”, IIB-INTECH Universidad Nacional de San Martín. Avda 25 de Mayo y Francia, Campus Miguelete, UNSAM, San Martín (1650), Buenos Aires, Argentina. Phone: (54)11-4006-1500 (2129 or 2135) Fax: (54)11-4006-1559 jugalde@iibintech.com.ar or czibener@iibintech.com.ar.

The authors declare no conflict of interest.

endoplasmic reticulum derived membrane niche where it can actively replicate (Celli *et al.*, 2003, Comerci *et al.*, 2001). The intracellular traffic of *Brucella* in non-professional and professional phagocytes has been extensively studied (Celli *et al.*, 2003, Celli *et al.*, 2005, Comerci *et al.*, 2001, Pizarro-Cerda *et al.*, 1999, Pizarro-Cerda *et al.*, 1998). Shortly after its internalization, the BCV transiently acquires the lysosomal marker Lamp-1, which is subsequently excluded as the BCV is directed to its final replicative organelle. The BCV actively excludes the mature lysosomal marker Cathepsin-D, avoids maturation into an active lysosome and acquires endoplasmic reticulum derived membranes creating a suitable organelle for the replication of the bacteria.

Although much has been done on how the *virB* system affects the intracellular trafficking, the virulence process, and how this complex secretion system is transcriptionally regulated (Arocena *et al.*, 2010, Arocena *et al.*, 2012, Delrue *et al.*, 2005, Sieira *et al.*, 2010, Sieira *et al.*, 2012, Sieira *et al.*, 2004), very little is known on the identity of the secreted/translocated proteins as well as the molecular mechanisms they use to modulate the intracellular trafficking. To date eleven effectors have been identified and, of them, the activity of only one has been elucidated.

RicA was identified as a *Brucella* protein that interacts with the GTPase Rab2 and is translocated into the host cell in a *virB* dependent manner. This effector protein binds preferentially the GDP-bound Rab2, avoids the recruitment of this Rab to the BCV and affects the normal intracellular trafficking (de Barsy *et al.*, 2011). An interesting observation made by the authors of this report is that, even though RicA is translocated into the host in a *virB* dependent manner, the protein is also secreted into the supernatant in bacteriological cultures in a *virB* independent manner, suggesting that there could be a different secretion system involved in this process. This raises the interesting point if the *virB* dependent translocation of effector proteins involves a two-step process with other secretion systems also participating.

Two effector proteins that were isolated identifying promoter regions similar to the *virB* promoter box are VceA and VceC. The authors of this report demonstrated, using a TEM1 β -lactamase reporter, that these proteins are translocated into the host cells in a *virB* dependent manner and, in one of these proteins (VceE), a C-terminal type four secretion signal was identified that is also functional in *Legionella* (de Jong *et al.*, 2008).

Recently, Marchesini *et al.* performed a bioinformatic and biochemical screening searching for novel *virB* effector proteins (Marchesini *et al.*, 2011). They identified, using as a reporter the adenylate cyclase (Cya) from *Bordetella pertussis*, four new *virB* dependent translocated proteins. One of these proteins, Bab2_0123, was also detected as secreted in a *virB* dependent manner by immunofluorescence and biochemical fractionation. The authors demonstrated that the first 25 amino acids of the protein are necessary for translocation into the host cell and showed that the mutant had no detectable phenotype.

Very recently Myeni *et al.* (Myeni *et al.*, 2013) informed the identification of five new type IV effector proteins in *B. abortus*. The authors identified these proteins through a bioinformatic search and verified that they are translocated into the host cells using β -lactamase and CyaA adenylate cyclase reporter assays. The ectopic expression of several of them as well as single and combined deletion mutant strains indicate that *Brucella* modulates the secretory pathway using multiple effector proteins (Myeni *et al.*, 2013).

We describe here the identification of a *virB* secretion substrate present in all *Brucella* species sequenced to date. We demonstrate by immunofluorescence that this protein is secreted in a *virB* dependent manner during the infection of host cells and that this secretion

requires a periplasmic intermediate. We also show that the mutant strain has a significant defect in the early stages of the intracellular multiplication process and that is less effective in excluding the lysosomal marker Lamp-1. Surprisingly, even though the mutant strain is more efficiently inactivated during the first stages of the intracellular replication process, it invades the host cell more efficiently than the wild type suggesting that this gene might play a role in "selecting" the phagocytosis route and, thus, modulating the final destination of the resulting phagosome. This hypothesis is discussed.

RESULTS

Identification of the gene product of Bab1_1492 as a secreted protein in *Brucella abortus*

In order to identify novel virulence factors in *B. abortus* we performed a bioinformatic search seeking for potential horizontally transmitted regions identifying transposases or recombinases close to transfer tRNAs. This rationale is based on the observation that, in many bacteria, horizontally transmitted pathogenic islands are located adjacent to tRNAs and encode enzymes involved in DNA processing. For this we used the program ARTEMIS (Rutherford *et al.*, 2000) and searched both *Brucella abortus* 2308 chromosomes for putative transposases or recombinases that were adjacent to genes coding for transfer tRNAs. Using this approach we identified several potentially horizontally transmitted regions and analyzed each one of them individually identifying the open reading frames and determining, for each, conservation among *Brucella* species, presence in other bacteria and domain structure. This methodology allowed us to select a group of potential proteins whose genes were cloned upstream of a 3xFLAG, subcloned in a *Brucella* replicative vector (pBBR1MCS-4) and introduced into a wild type *B. abortus* 23208 strain. The resulting clones were used to infect J774 A.1 cells and, at different times post-infection, the infected cells fixed and stained to detect the FLAG epitope and the bacteria. The objective of this approach was to identify gene products secreted by the bacteria during the infection of host cells. Figure 1A shows a schematic representation of the screening method utilized. Out of 5 candidates analyzed one was detected on the surface of the intracellular bacteria. Figure 1B shows that the protein product of the gene Bab1_1492 was detected on the bacterial surface during the infection of J774 A.1 cells but not the protein product of gene Bab1_0740 (Figure 1C) although both genes were expressed (Figure 1D). To further examine the secretion of the protein product of Bab1_1492, confocal images were obtained at 4 hrs post-infection and as a positive control a strain expressing a 3xFLAG fusion to the type IV effector Bpe123 was used (Marchesini *et al.*, 2011). Figure 2A shows that at 4 hrs surface exposed Bab1_1492 was observed in most of the bacteria analyzed, while only 15 to 20 % of bacteria were positive for Bpe123 as has been informed (Del Giudice *et al.*, 2013, Marchesini *et al.*, 2011) (Figure 2B). A scanning of the fluorescence in both channels through a linear path indicated that both marks showed the same pattern (Figure 2D). These results demonstrate that the protein product of Bab1_1492 is secreted during infection, for this reason we named the gene *sepA* for Secreted Effector Protein A.

Secretion of SepA is a two-step *virB* dependent process

Because the type IV secretion system (*virB*) is one of the main virulence traits required for intracellular survival we asked if secretion of SepA was dependent on this system. For this we introduced the plasmid carrying the FLAG-tagged version of *sepA* in the *virB10* and *virB11* mutant strains and determined the number of intracellular bacteria displaying the protein on the surface. As can be observed in Figure 3A and 3B, secretion of SepA was significantly reduced in the *virB* mutant backgrounds measured as the number of bacteria showing surface exposed FLAG staining, although expression of the tagged protein was equivalent between strains (Figure 3C). To determine if SepA is secreted immediately after internalization we analyzed by immunofluorescence surface exposed SepA in the wild type

and the *virB10* mutant at 30 min post-infection. The results shown in Supplementary Figure 1 showed that SepA is secreted very early during infection and in a *virB* dependent manner. Altogether, these results demonstrate that secretion of SepA is dependent on a functional *virB* system.

It has been informed that some type IV secretion substrates possess a canonical sec-dependent secretion signal that targets them to the periplasm or the inner membrane as a first secretion step (Burns, 2003, Nagai *et al.*, 2001, Pantoja *et al.*, 2002). Since the analysis of the protein product of *sepA* using different prediction programs for the detection of putative sec-dependent secretion signals were not conclusive, we decided to determine if this protein is targeted to the periplasm and, if so, if this intermediate state is necessary for the *virB* secretion. To test we generated a double-tagged strain carrying a chromosomal FLAG-tagged version of the gene Bab2_0448 coding for a cytoplasmic tellurite resistant protein that we are currently studying in the laboratory (unpublished results) in which we introduced the plasmid coding the *sepA*-3xFLAG. This strain was used to produce periplasmic and cytoplasmic fractions (see Experimental procedures) and the presence of SepA was determined in these fractions. As control proteins we used the presence of the protein product of Bab2_0448 and GroEL for cytoplasm and OMP1 for periplasm. As can be seen in Figure 4A, SepA was found partitioned in both the periplasm and the cytoplasm as was OMP1, while the cytoplasmic controls were only detected in the corresponding fraction (cytoplasmic). This result demonstrates that SepA is exported to the periplasm. In order to determine if this periplasmic intermediate is required for the *virB* secretion we analyzed the bacterial surface display of the construct lacking the first 25 amino acids of the protein (Δ N-SepA). As shown in Figure 4B, deletion of the amino terminal secretion signal abolished the surface display of SepA as did the absence of the *virB* system. Loading controls indicated that the deleted protein was expressed with lower levels than the wild type and that it migrated slightly less (Figure 4C). We do not have an explanation for this puzzling observation but speculate that maybe the truncated protein, because it cannot be transported to the periplasm, aggregates or associates with other proteins resistant in our gel conditions. In order to determine if in the *virB* mutant background SepA is still present in the periplasm we performed a periplasmic extraction with the *virB10* and *virB11* strains expressing the SepA-3xFLAG fusion protein and determined the fractionation pattern of SepA in these strains. As can be observed in Figure 4D, SepA was found in the periplasm of these strains indicating that, even though it was not secreted to the bacterial surface, it retained a periplasmic localization. This result also demonstrates that the bacterial surface display of SepA is not the consequence of “leakage” of the periplasm of the extracellular space. Altogether these results indicate that SepA is targeted to the periplasm and that this intermediate state is required for the *virB* secretion.

SepA participates in the early stages of intracellular replication

To determine if *sepA* plays a role during the intracellular life cycle we generated a clean deletion strain and analyzed the intracellular multiplication in HeLa, J774 A.1 cells and bone marrow derived macrophages (BMDM) with an antibiotic protection assay as described in Experimental procedures. As can be observed in Figure 5A, 5B and 5C the strain Δ *sepA* showed a significant reduction in the intracellular CFU at 4 and 24 hrs post-infection in both professional and non-professional phagocytes. At 48 hrs, even though the mutant had less viable counts than the wild type strain, the differences were less dramatic in HeLa and J774 A.1 cells indicating that the mutant had a defect in the early stages of the intracellular replication process but was able to replicate. In BMDM the mutant was unable to replicate at 48 hr post-infection indicating that the phenotype was more pronounced in these cells. The defects observed were not the consequence of a general replication phenotype as the mutant had an *in vitro* growth identical to the wild type parental strain (Figure 5D). To determine if

the phenotype observed at 4 hrs post-infection was dependent on the multiplicity of infection (m.o.i.) we carried out an antibiotic protection assay on J774 A.1 cells with different multiplicity of infections and determined intracellular CFUs at that time point. The results indicated that the $\Delta sepA$ phenotype was not m.o.i. dependent (Figure 5E). As expected, we were unable to complement the $\Delta sepA$ mutant with the construction lacking the N-terminal periplasmic localization region indicating that the phenotype is dependent on the secretion of the protein (Figure 5F). The lower viable bacteria at 4 hrs post-infection could be result of a defect in the invasion or early intracellular survival. To discern between these possibilities, we initially determined the invasion capacity as described in Experimental procedures. The results, shown in Figure 6 indicated that invasion was not decreased in the mutant strongly suggesting that the defect was probably due a defect in the capacity to avoid the fusion of the BCV with the lysosomes. Moreover, an interesting observation was that, even though at 4 hrs post-infection the $\Delta sepA$ mutant showed a 10-fold reduction in the intracellular bacteria, it actually invaded the host cells more efficiently than the wild type (Figure 6C). To determine if at early time points the internalized bacteria were viable, we performed a gentamicin protection assay and recovered intracellular bacteria at 1, 2, 3 and 4 hrs post-infection and found that at all times tested the mutant had less viable counts (Figure 6D) that decreased over time in comparison to the wild type strain. These results strongly suggested that the mutant probably had a defect in the early stages of the intracellular trafficking. To test this we determined the co-localization of the lysosomal marker Lamp-1 with the BCV at 4 and 24 hrs post-infection in HeLa cells as described in Experimental procedures. As shown in Figure 7A and 7B, the BCV containing the $\Delta sepA$ mutant acquired the lysosomal marker Lamp-1 as the wild type at 4 hrs post-infection but was less efficient in excluding it at 24 hrs suggesting a defect in avoiding the maturation into an active lysosome. Even though the strain $\Delta sepA$ showed a defect in the initial phases of the intracellular life cycle the bacteria that were able to bypass this initial attack and replicated in an ER-like calnexin positive niche confirming that the defect was restricted to the initial phases of the intracellular cycle (data not shown).

$\Delta sepA$ has no impact in the virulence in mice but over-expression of the gene causes a severe defect

To determine if *sepA* plays a role in the virulence in the mouse model of infection, we infected intraperitoneally groups of five mice with the wild type 2308, the $\Delta sepA$ and the complemented strains and measured, at 15 days post-infection, the bacterial load in the spleens. As can be seen in Figure 8A, the mutant strain did not display any differences in the CFUs per spleen in comparison to the parental strain. An interesting observation was that the complemented strain colonized the spleens significantly less than the mutant and the parental strains suggesting that over-expression of the gene might cause a negative effect. To confirm this we introduced plasmid pBBR1-MCS4/F2-3xF over-expressing SepA in the wild type 2308 strain and evaluated its virulence in mice in comparison with the non-transformed wild type and the deletion strains. As can be observed in Figure 8B, over-expression of SepA dramatically reduced the bacterial load in the spleens of infected mice at 15 days post-infection and the mutant, as indicated before, was as virulent as the wild type parental strain. This effect was not the consequence of a defect in the viability of the strain as the strain grew as the parental strain (Supplementary Figure 2) and did not show any major membrane alterations evidenced by several indicators as for example LPS integrity (Figure 8C). These results strongly suggest that the expression of *sepA* is probably highly regulated and that an excess of the protein is detrimental for the virulence process although more work should be performed in order to understand the molecular basis of this defect.

DISCUSSION

Many intracellular pathogens have, as virulence strategies, the capacity to establish their multiplication niches inside professional phagocytes. To achieve this they have evolved complex secretion systems devoted to secrete and translocate into the host cells virulence factors that modulate the biological functions in their one benefit. Understanding at the molecular level this biochemical crosstalk has been one of the main focuses of the microbial pathogenesis field of the last decade. *Brucella*, the causative agent of Brucellosis, is an intracellular pathogen that replicates in macrophages and dendritic cells. The *virB* operon encodes one of the central virulence traits required for the intracellular lifestyle of the bacterium. This operon codes for a type IV secretion system responsible for the translocation of effector proteins that actively avoid the fusion of the *Brucella* containing vacuole with the lysosomes and recruit membranes from the endoplasmic reticulum to establish a competent replication niche (Celli *et al.*, 2003, Comerci *et al.*, 2001, O'Callaghan *et al.*, 1999, Sieira *et al.*, 2000). This macromolecular apparatus is essential for virulence and, to date, few secretion substrates have been identified (de Bary *et al.*, 2011, de Jong *et al.*, 2008, Marchesini *et al.*, 2011). We identify here a new virulence factor in *Brucella abortus* that is secreted in a *virB* dependent manner and we have named this gene *sepA*. We demonstrate by immunofluorescence that secretion of SepA occurs during the early phases of the intracellular life cycle of *B. abortus*, that this secretion is *virB* dependent and show that it involves a periplasmic intermediate. To date it is not clear the exact mechanism in type IV secretion systems. Even though it is highly accepted that secretion is a process in which the substrate is engaged in the cytoplasm, there have been reports of periplasmic and membrane proteins that are secreted in a type IV dependent process in some pathogens (Burns, 2003, Nagai *et al.*, 2001, Pantoja *et al.*, 2002). One of the canonical examples of these is the pertussis toxin, that assembles in the periplasm and is secreted in a type IV dependent manner (Burns, 2003). In the case of SepA we have clearly established that secretion to the surface of the bacteria is a process that depends on the *virB* system and on an N-terminal signal peptide that targets the protein to the periplasm indicating that its secretion is a two-step process. With the purpose of determining the role of *sepA* in the pathogenesis of *B. abortus* we constructed a deletion mutant and evaluated the intracellular replication capacity of the strain. Our results showed that the mutant had a defect in the early stages of the intracellular replication curve and that this alteration was the consequence of a reduced capacity to avoid the fusion of the BCV with the lysosomes. Interestingly, even though the mutant did not display any defects in its invasion it was significantly higher than that of the parental wild type strain. Consistently, the same observation was seen for the *virB10* polar mutant. These results are, apparently, inconsistent with the reduced viable bacteria at 1, 2, 3 and 4 hrs post-infection. One hypothesis could be proposed here. Because it has been established that the route of entry in phagocytes can influence the final destination of the phagosome generated (Aderem *et al.*, 1999), it could be possible that SepA is involved in the initial steps of the intracellular life cycle facilitating certain routes that will ultimately influence the initial composition and final destination of the BCV. In this model the type IV secretion system and some effectors would be involved in the very early stages of the intracellular life cycle selecting the internalization route of the bacterium. Even though the mutant is more invasive than the parental wild type strain, the BCV generated in the process are less efficient for evading the early phagosome-lysosome fusion. Mechanistically, this process could be either through a positive or a negative selection, this is SepA could facilitate phagocytosis through certain routes or inhibit others. We speculate that *sepA* participates in this early selection process and that its biological activity modifies the final destination of the BCV. At this stage the molecular mechanism used by SepA to promote infection is not known and the fact that the protein has no significant homology to any known proteins complicates this task significantly. Albeit the fact that a molecular mechanism is still lacking, to our knowledge this is the first Type IV secretion substrate in

Brucella that shows a phenotype in cells altering the kinetics of Lamp-1 acquisition and exclusion.

EXPERIMENTAL PROCEDURES

Bacterial strains and growth conditions

B. melitensis biovar abortus 2308 was used as a wild-type strain. *B. abortus* strains were cultured in Tryptic Soy Agar (TSA; Difco/Becton-Dickinson, Sparks, MD), or in Tryptic Soy Broth (TSB) at 37°C on a rotary shaker for 16–24 hrs. When needed, media were supplemented with 50 µg/ml Kanamycin (Km^r), 5 µg/ml Nalidixic Acid (Nal^r), 100 µg/ml Ampicillin (Amp^r) or 3 µg/ml Gentamicin (Gm^r). Work with *B. abortus* was performed at the Biosafety Level 3 laboratory facility at the University of San Martín. *E. coli* strains were grown on Luria-Bertani (LB) agar and broth at 37°C overnight. Antibiotics, when required, were added at the following concentrations: 50 µg/ml Kanamycin or 100 µg/ml Ampicillin.

Recombinant DNA techniques

To generate plasmid pBBR1-MCS4-3xFlag, a fragment containing a 3xFLAG epitope was amplified from plasmid pBAD24-3xFLAG (Spano *et al.*, 2008) using primers CC7-Flag (5'-CGCGGATCCACTCTAGAGATCGTCATCCTTGTA-3') and pbad/HindIII (5'-CCCAAGCTTGTTCCTCATAACCCGTTT-3'). The product was digested with BamHI-HindIII, and ligated into the same sites of plasmid pBBR1-MCS4 (Kovach *et al.*, 1995).

For the construction of vectors expressing a C-terminal 3xFLAG-tagged version of genes Bab1_1492, Bab1_0296, Bab1_0752, Bab1_0740, Bab1_0745, the plasmid pBBR1-MCS4-3xFLAG was used. DNA fragments were amplified by PCR from *B. abortus* 2308 genomic DNA using primers F1492 (5'-CGGAATTCTTGAGCGGAGAGAGCC-3') and R1492 (5'-CGCCATGGCGGCGGACGCCGGGCC-3') for the Bab1_1492 gene; F0296 (5'-CGGAATTCTGAGACATGCGAATTAAG-3') and R0296 (5'-CGCCATGGCAAGCTCCAAGCATCTAAT-3') for the Bab1_0296 gene; F0752 (5'-TAGCTAGCTGTGCGGAAGTCAGCC-3') and R0752 (5'-GCCATGGCTAGGCGTCCAGACATTC-3') for the Bab1_0752 gene; F0740 (5'-CGGAATTCTTTGGTCGCTACGCGT-3') and R0740 (5'-CGCCATGGCGAATTTGTCTAGCAGGT-3') for the Bab1_0740 gene and F0745 (5'-CGGAATTCAAATCAACGTTCAAATCTATC-3') and R0745 (5'-CGCCATGGCATATTCCAAAATATTCCTTTG-3') for the Bab1_0745 gene. The truncated version (amino acids 25 to 190) of Bab1_1492 was generated, using primers FsNH2-1492 (5'-GGAATTCAGAGAGGGATGACCCGAGCGAAACCATT-3') and R1492 (5'-CGCCATGGCGGCGGACGCCGGGCC-3'). All the PCR products were digested with EcoRI-NcoI, except the fragment 0752 that was digested with EcoRI-NheI. The resulting fragments were cloned in pBBR1-MCS4-3xFlag in the same sites generating in frame fusions to the 3xFLAG epitope. The resulting plasmids were named pBBR1-MCS4/ F2-3xF (Bab1_1492), pBBR1-MCS4/ F2NH2-3xF (truncated version of Bab1_1492), pBBR1-MCS4/ 0296-3xF (Bab1_0296), pBBR1-MCS4/ 0752-3xF (Bab1_0752), pBBR1-MCS4/ 0740-3xF (Bab1_0740) and pBBR1-MCS4/ 0745-3xF (Bab1_0745). Plasmids expressing 3xFLAG tagged fusion proteins were introduced in *B. abortus* strains by biparental mating. The integrity of all constructs was confirmed by sequence analysis.

A *B. abortus* $\Delta sepA$ strain with a clean deletion of Bab1_1492 was constructed by standard recombinant DNA and allelic exchange procedures as previously described (Czibener *et al.*, 2012). The flanking regions of Bab1_1492 were amplified by PCR using primers Del1492-1 (5'-CGGGATCCAGTTTCGCAGACGCGGTCCA-3'), Del1492-2 (5'-

TATTCTCTCGTTGACCGCAA-3'), Del1492-3 (5'-GTCAACGAGAGAATACTTCCTGTTTCGAAGCCGTTT-3') and Del1492-4 (5'-CCCAAGCTTTGAGCGGCCATTGCGTCAGG-3'). The PCR products were ligated using the recombinant PCR technique (Czibener *et al.*, 2012), and the resulting DNA fragment digested with BamHI-HindIII and ligated to the same sites of suicide vector pK18mobsacB, giving rise to plasmid pK18mobsacB/ Δ sepA. The plasmid was introduced in *B. abortus* strain by biparental mating, and the mutants Km^S and Suc^R were confirmed by colony PCR with primers Del1492-1 and Del1492-4.

Intracellular replication curves

Gentamicin protection assays were performed as previously described (Ugalde *et al.*, 2000). All tissue cultures were maintained at 37°C in a humidified atmosphere containing 5% CO₂. HeLa and J774.A1 macrophage-like cell lines were maintained in Dulbecco modified Eagle medium (Gibco Life Technologies) and RPMI 1640 (Gibco Life technologies), respectively, supplemented with 5% fetal bovine serum (PAA lab's GmbH, Pasching, Austria) and Streptomycin (50 µg/ml)/Penicillin (50 U/ml) (Gibco Life technologies). Cells were trypsinized, counted and seeded on 24-well plates in media, at a density of 5×10^4 cells per well, and were grown for 24 hrs. *B. abortus* infections were carried out at a multiplicity of infection (MOI) of 500:1 for HeLa cells and 100:1 for J774.A1 cells. Bacteria were centrifuged onto cells at $400 \times g$ for 10 min. After 60 min wells were gently washed three times with phosphate-buffered saline (PBS) and incubated for 3 hrs with fresh medium containing 50 µg/ml Gentamicin to kill non-internalized bacteria. At 4 hrs post-infection, cells were washed three times with PBS, and fresh medium containing 10 µg/ml Gentamicin was added. Bone marrow derived macrophages (BMDM) were prepared as described in (Czibener *et al.*, 2006). At the indicated times, infected cells were either washed three times with PBS and lysed with 500 µl 0.1% Triton X-100 in H₂O (Sigma-Aldrich) or processed for immunofluorescence staining as described below. The intracellular CFU counts were determined by plating serial dilutions on TSA with the appropriate antibiotics.

Antibodies and reagents

All the antibodies were diluted in 10% horse serum (Gibco), 5% BSA (Sigma) and 0.1% saponin (Sigma), except for the anti-FLAG M2 monoclonal antibody (Sigma), that was diluted in 0.2 % Triton X-100 instead of 0.1% saponin for cell permeabilization. The primary antibodies used were rabbit anti-*Brucella* polyclonal antibody (1:1,500), mouse immunoglobulin G anti-O-antigen monoclonal antibody M-84 (1:1000) (Nielsen *et al.*, 1995), mouse anti-human LAMP-1 H4A3 (1:400) (Developmental Studies Hybridoma Bank, Department of Biological Sciences, University of Iowa), and mouse anti-FLAG M2 monoclonal antibody (1:4000). The secondary antibodies used were Alexa Fluor 488 goat anti-rabbit or anti-mouse and Alexa Fluor 568 goat anti-rabbit or anti-mouse (Molecular Probes, Invitrogen Co.). For DNA staining, DAPI dye at 2 µg/ml (final concentration) was used.

Immunofluorescence microscopy

Eukaryotic cells were plated on glass coverslips (Marienfeld GmbH & co. KG) and infected as described above. For HeLa cells the MOI was 1000:1, and for J774.A1 cells the MOI was 500:1. At the indicated times, the coverslips were washed with PBS and the cells were fixed for 15 min in 4% paraformaldehyde (pH 7.4) at room temperature. After fixation, cells were washed three times with PBS and incubated for 10 min with PBS-50 mM NH₄CL in order to quench free aldehyde groups. For the In/Out assay, the coverslips were incubated with rabbit anti-*Brucella* antibody for 45 min at room temperature, washed in PBS and incubated for 30 min with PBS-10% horse serum-5% BSA-0.1% saponin permeabilization solution.

Then washed in PBS, and incubated for 45 min with mouse anti-O-antigen antibody M-84. After the second antibody the cells were washed with PBS and incubated for 45 min with Alexa Fluor 568 goat anti-rabbit, Alexa Fluor 488 goat anti-mouse secondary antibodies and the DAPI dye.

For the Lamp-1 co-localization assay, coverslips were incubated for 30 min with PBS–10% horse serum–5% BSA–0.1% saponin permeabilization solution, washed with PBS, and incubated for 45 min with mouse anti-human Lamp-1 H4A3 antibody. After 5 washes with PBS, they were incubated for 45 min with rabbit anti-*Brucella* antibody, washed again with PBS, and then incubated for 45 min with Alexa Fluor 568 goat anti-rabbit and Alexa Fluor 488 goat anti-mouse secondary antibodies.

For the Screening assays, the coverslips were incubated for 30 min with PBS–10% horse serum–5% BSA–0.1% saponin permeabilization solution, washed with PBS, and incubated for 45 min with a rabbit anti-*Brucella* antibody. After 5 washes with PBS, cells were incubated for 30 min with PBS–10% horse serum–5% BSA– 0.2% Triton X-100 permeabilization solution, washed again with PBS and incubated for 45 min with mouse anti-FLAG M2. Then washed with PBS, and finally incubated for 45 min with Alexa Fluor 568 goat anti-rabbit, Alexa Fluor 488 goat anti-mouse secondary antibodies and DAPI dye.

After immunofluorescence staining, all the coverslips were mounted onto slides with FluorSave (Calbiochem). Samples were examined on a Nikon microscope (Eclipse TE 2000) at a magnification of X600. The software MBF ImageJ v1.43m (Wayne Rasband, NIH, USA) was used to merge the microscopic images. Con-Focal images were acquired in an Olympus FV 1000 inverted microscope and images, as well as Z stacks, processed with the Fluoview FV100 software provided with the microscope.

Mouse infections

Mice infections were performed as previously described (Ugalde *et al.*, 2003). Groups of 5 eight to nine week-old female BALB/c mice were intraperitoneally inoculated with 1×10^5 CFU of *B. abortus* 2308 wild type, $\Delta sepA$, complemented or SepA overexpressing strains in PBS. At 2 weeks post-infection, spleens from infected mice were removed and homogenized in 2 ml of PBS. Serial dilutions from individualized spleens were plated on TSA with the appropriate antibiotics to quantify recovered CFU.

Periplasmic and cytoplasmic localization assay

For the periplasmic localization assays the *B. abortus* strains were cultivated in TSB for 16h – 24h at 37°C until an OD_{600nm} 1 was reached. 2.5×10^{10} bacterial cells were centrifuged 10 min at 3300×g. The pellets were washed with physiologic solution, centrifuged for 10 min at 3300×g and resuspended in 1 ml of 0.2M Tris-HCl pH 7.6. 1 ml of 0.2M Tris-HCl pH 7.6, 1M Sucrose and 0.25% Zwitterion 3–16 solution was added to the cell suspension and incubated for 10 min at room temperature. The samples were centrifuged for 30 min at 8000×g, the pellets separated from the supernatants and stored at –20°C until used. The pellets and supernatants were processed for Western-Blot, using an anti-FLAG M2 monoclonal antibody (1:5000), anti GroEL (1:2000) and anti OMP-1 (1:2000) kindly provided by Dr. Axel Cloeckert as primary antibodies, and goat anti-mouse HRP (1:2000) (Dako) as secondary antibody. All antibodies were diluted in Tris-buffered saline (TBS)- 5% non-fat milk- 0.05% Tween solution.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Pablo Dosantos and Francisco Guaimas for confocal microscopy assistance. This work was supported by grants PICT-PRH08-160 and PICT-PRH08-230 to C.C. and J.E.U. respectively, NIAID grant AI078891 to J.E.U., and grants from the University of San Martín to C.C. and J.E.U. P.H.D and E.V. were supported by fellowships of the National Research Council of Argentina (CONICET). CC and J.E.U. are members of the CONICET.

REFERENCES

- Aderem A, Underhill DM. Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol.* 1999; 17:593–623. [PubMed: 10358769]
- Alvarez-Martinez CE, Christie PJ. Biological diversity of prokaryotic type IV secretion systems. *Microbiol Mol Biol Rev.* 2009; 73:775–808. [PubMed: 19946141]
- Arocena GM, Sieira R, Comerci DJ, Ugalde RA. Identification of the quorum-sensing target DNA sequence and N-Acyl homoserine lactone responsiveness of the *Brucella abortus* virB promoter. *J Bacteriol.* 2010; 192:3434–3440. [PubMed: 20400542]
- Arocena GM, Zorreguieta A, Sieira R. Expression of VjbR under nutrient limitation conditions is regulated at the post-transcriptional level by specific acidic pH values and urocanic acid. *PLoS One.* 2012; 7:e35394. [PubMed: 22530016]
- Atluri VL, Xavier MN, de Jong MF, den Hartigh AB, Tsolis RE. Interactions of the human pathogenic *Brucella* species with their hosts. *Annu Rev Microbiol.* 2011; 65:523–541. [PubMed: 21939378]
- Burns DL. Type IV transporters of pathogenic bacteria. *Curr Opin Microbiol.* 2003; 6:29–34. [PubMed: 12615216]
- Celli J, de Chastellier C, Franchini DM, Pizarro-Cerda J, Moreno E, Gorvel JP. *Brucella* evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic reticulum. *J Exp Med.* 2003; 198:545–556. [PubMed: 12925673]
- Celli J, Salcedo SP, Gorvel JP. *Brucella* coopts the small GTPase Sar1 for intracellular replication. *Proc Natl Acad Sci U S A.* 2005; 102:1673–1678. [PubMed: 15632218]
- Comerci DJ, Martinez-Lorenzo MJ, Sieira R, Gorvel JP, Ugalde RA. Essential role of the VirB machinery in the maturation of the *Brucella abortus*-containing vacuole. *Cell Microbiol.* 2001; 3:159–168. [PubMed: 11260139]
- Corbel MJ. Brucellosis: an overview. *Emerg Infect Dis.* 1997; 3:213–221. [PubMed: 9204307]
- Cossart P, Roy CR. Manipulation of host membrane machinery by bacterial pathogens. *Curr Opin Cell Biol.* 2010; 22:547–554. [PubMed: 20542678]
- Czibener C, Sherer NM, Becker SM, Pypaert M, Hui E, Chapman ER, et al. Ca²⁺ and synaptotagmin VII-dependent delivery of lysosomal membrane to nascent phagosomes. *J Cell Biol.* 2006; 174:997–1007. [PubMed: 16982801]
- Czibener C, Ugalde JE. Identification of a unique gene cluster of *Brucella* spp. that mediates adhesion to host cells. *Microbes Infect.* 2012; 14:79–85. [PubMed: 21911075]
- de Barsey M, Jamet A, Filopon D, Nicolas C, Laloux G, Rual JF, et al. Identification of a *Brucella* spp. secreted effector specifically interacting with human small GTPase Rab2. *Cell Microbiol.* 2011; 13:1044–1058. [PubMed: 21501366]
- de Jong MF, Sun YH, den Hartigh AB, van Dijl JM, Tsolis RM. Identification of VceA and VceC, two members of the VjbR regulon that are translocated into macrophages by the *Brucella* type IV secretion system. *Mol Microbiol.* 2008; 70:1378–1396. [PubMed: 19019140]
- Del Giudice MG, Ugalde JE, Czibener C. A lysozyme-like protein in *Brucella abortus* is involved in the early stages of intracellular replication. *Infect Immun.* 2013; 81:956–964. [PubMed: 23319555]
- Delrue RM, Deschamps C, Leonard S, Nijskens C, Danese I, Schaus JM, et al. A quorum-sensing regulator controls expression of both the type IV secretion system and the flagellar apparatus of *Brucella melitensis*. *Cell Microbiol.* 2005; 7:1151–1161. [PubMed: 16008582]
- Godfroid J, Scholz HC, Barbier T, Nicolas C, Wattiau P, Fretin D, et al. Brucellosis at the animal/ecosystem/human interface at the beginning of the 21st century. *Prev Vet Med.* 2011; 102:118–131. [PubMed: 21571380]

- Kovach ME, Elzer PH, Hill DS, Robertson GT, Farris MA, Roop RM 2nd, Peterson KM. Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene*. 1995; 166:175–176. [PubMed: 8529885]
- Marchesini MI, Herrmann CK, Salcedo SP, Gorvel JP, Comerci DJ. In search of *Brucella abortus* type IV secretion substrates: screening and identification of four proteins translocated into host cells through VirB system. *Cell Microbiol*. 2011; 13:1261–1274. [PubMed: 21707904]
- Myeni S, Child R, Ng TW, Kupko JJ 3rd, Wehrly TD, Porcella SF, et al. *Brucella* Modulates Secretory Trafficking via Multiple Type IV Secretion Effector Proteins. *PLoS Pathog*. 2013; 9:e1003556. [PubMed: 23950720]
- Nagai H, Roy CR. The DotA protein from *Legionella pneumophila* is secreted by a novel process that requires the Dot/Icm transporter. *EMBO J*. 2001; 20:5962–5970. [PubMed: 11689436]
- Nielsen KH, Kelly L, Gall D, Nicoletti P, Kelly W. Improved competitive enzyme immunoassay for the diagnosis of bovine brucellosis. *Vet Immunol Immunopathol*. 1995; 46:285–291. [PubMed: 7502488]
- Ninio S, Roy CR. Effector proteins translocated by *Legionella pneumophila*: strength in numbers. *Trends Microbiol*. 2007; 15:372–380. [PubMed: 17632005]
- O'Callaghan D, Cazevieuille C, Allardet-Servent A, Boschirolu ML, Bourg G, Foulongne V, et al. A homologue of the *Agrobacterium tumefaciens* VirB and *Bordetella pertussis* Ptl type IV secretion systems is essential for intracellular survival of *Brucella suis*. *Mol Microbiol*. 1999; 33:1210–1220. [PubMed: 10510235]
- Pantoja M, Chen L, Chen Y, Nester EW. *Agrobacterium* type IV secretion is a two-step process in which export substrates associate with the virulence protein VirJ in the periplasm. *Mol Microbiol*. 2002; 45:1325–1335. [PubMed: 12207700]
- Pizarro-Cerda J, Desjardins M, Moreno E, Akira S, Gorvel JP. Modulation of endocytosis in nuclear factor IL-6(–/–) macrophages is responsible for a high susceptibility to intracellular bacterial infection. *J Immunol*. 1999; 162:3519–3526. [PubMed: 10092809]
- Pizarro-Cerda J, Meresse S, Parton RG, van der Goot G, Sola-Landa A, Lopez-Goni I, et al. *Brucella abortus* transits through the autophagic pathway and replicates in the endoplasmic reticulum of nonprofessional phagocytes. *Infect Immun*. 1998; 66:5711–5724. [PubMed: 9826346]
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. Artemis: sequence visualization and annotation. *Bioinformatics*. 2000; 16:944–945. [PubMed: 11120685]
- Sieira R, Arocena GM, Bukata L, Comerci DJ, Ugalde RA. Metabolic control of virulence genes in *Brucella abortus*: HutC coordinates virB expression and the histidine utilization pathway by direct binding to both promoters. *J Bacteriol*. 2010; 192:217–224. [PubMed: 19854911]
- Sieira R, Arocena GM, Zorreguieta A, Comerci DJ, Ugalde RA. A MarR-Type Regulator Directly Activates Transcription from the *Brucella abortus* virB Promoter by Sharing a Redundant Role with HutC. *J Bacteriol*. 2012; 194:6431–6440. [PubMed: 23002224]
- Sieira R, Comerci DJ, Pietrasanta LI, Ugalde RA. Integration host factor is involved in transcriptional regulation of the *Brucella abortus* virB operon. *Mol Microbiol*. 2004; 54:808–822. [PubMed: 15491369]
- Sieira R, Comerci DJ, Sanchez DO, Ugalde RA. A homologue of an operon required for DNA transfer in *Agrobacterium* is required in *Brucella abortus* for virulence and intracellular multiplication. *J Bacteriol*. 2000; 182:4849–4855. [PubMed: 10940027]
- Spano S, Ugalde JE, Galan JE. Delivery of a *Salmonella Typhi* exotoxin from a host intracellular compartment. *Cell Host and Microbe*. 2008 In press.
- Ugalde JE, Comerci DJ, Leguizamón MS, Ugalde RA. Evaluation of *Brucella abortus* phosphoglucomutase (pgm) mutant as a new live rough-phenotype vaccine. *Infect Immun*. 2003; 71:6264–6269. [PubMed: 14573645]
- Ugalde JE, Czibener C, Feldman MF, Ugalde RA. Identification and characterization of the *Brucella abortus* phosphoglucomutase gene: role of lipopolysaccharide in virulence and intracellular multiplication. *Infect Immun*. 2000; 68:5716–5723. [PubMed: 10992476]




Figure 1. The protein product of Bab1_1492 is secreted during the infection of host cells
A. Schematic representation of the screening method used to identify novel secreted virulence effectors in *Brucella abortus*. **B.** Immunofluorescence of J774 A.1 cells infected with a *B. abortus* strain coding for Bab1_1492-3xFLAG in a replicative vector at 4 hrs post-infection. Red, *Brucella*; green, Flag. **C.** Immunofluorescence of J774 A.1 cells infected with a *B. abortus* strain coding for Bab1_0740-3xFLAG in a replicative vector at 4 hrs post-infection. Red, *Brucella*; green, Flag. **D.** Western-blot with an anti-FLAG of the strains shown in panels B and C.




Figure 2. Confocal analysis of the secretion of the protein product of Bab1_1492
Confocal images of J774 A.1 cells infected with the *B. abortus* strains coding for Bab1_1492-3xFLAG (**A**) or Bpe123-3xFLAG (**B**) in replicative vectors at 4 hrs post-infection. **C**. A 45° rotation view of the line traced in the inset of panel A. **D**. Quantification of the red and green fluorescence intensity over the line traced in the inset of panel A showing a correlation between both markers. Red, *Brucella*; green, FLAG.




Figure 3. Secretion of SepA is a *virB* dependent process

A. Representative Immunofluorescence of J774 A.1 cells infected with *B. abortus* wild type, *virB10* and Δ *virB11* mutant strains coding for SepA-3xFLAG in a replicative vector at 4 hrs post-infection. Red, *Brucella*; green, FLAG. **B.** Quantification of the percent of SepA secreting bacteria of the representative images shown in A. * $P < 0.05$. **C.** Western-blot with an anti-FLAG of the strains showing equivalent levels of expression of SepA.




Figure 4. Secretion of SepA involves a periplasmic intermediate

A. Western-blot with an anti-FLAG, anti-GroEL and anti OMP1 monoclonal antibodies of periplasmic or cytoplasmic fractions of a periplasmic fractionation assay performed with *B. abortus* chromosomal Bab2_0448-3xFLAG tagged strain carrying the plasmid coding for SepA-3xFLAG. Schematic representation of the Δ N-SepA-3xFLAG (Δ N-SepA) construction carrying a N-terminal 25 amino acid deletion. **B.** Immunofluorescence of J774 A.1 cells infected with *B. abortus* strains carrying plasmids coding for SepA-3xFLAG (SepA) or Δ N-SepA-3xFLAG (Δ N-SepA) at 4 hrs post-infection. Red, *Brucella*; green, FLAG. **C.** Western-blot showing the expression levels of SepA and Δ N-SepA in Wild-type

B. abortus. Loading control, OMP1. **D.** Western-blot with an anti-FLAG, anti-GroEL and anti OMP1 monoclonal antibodies of periplasmic or cytoplasmic fractions of a periplasmic fractionation assay performed with *B. abortus virB10* and *virB11* mutant strains carrying the plasmid coding for SepA-3xFLAG.




Figure 5. *SepA* participates in the early stages of intracellular replication

A. Intracellular replication of *B. abortus* wild type and $\Delta sepA$ deletion strains in HeLa cells determined by the gentamicin protection assay. **B.** Intracellular replication of *B. abortus* wild type and $\Delta sepA$ deletion strains in J774 A.1 cells determined by the gentamicin protection assay. **C.** Intracellular replication of *B. abortus* wild type and $\Delta sepA$ deletion strains in bone marrow derived macrophages determined by the gentamicin protection assay **D.** Viable bacteria of the wild type (*Ba wt*), *Ba* $\Delta sepA$ mutant and complemented strains at 4 hrs post-infection measured by an antibiotic protection assay in J774 A.1 cells with different multiplicity of infection ratios (50, 100 and 500). **E.** Growth curve of the wild type and the $\Delta sepA$ deletion strain in TSB. **F.** Complementation of the 4 hrs phenotype with the construction lacking the N-terminal periplasmic localization signal. ** $P < 0.002$ and *** $P < 0.001$.




Figure 6. The Δ sepA mutant strain invades cells more efficiently

A and B. In/out staining of the wild type strain (*Ba wt*) and the Δ sepA (*Ba Δ sepA*) mutant strain respectively at 1 hr post-infection in J774 A.1 cells. **C.** Quantification of the immunofluorescence shown in panels A and B. *Ba virB10*, assay performed with the *B. abortus virB10* mutant. ** $P < 0.002$. **D.** Intracellular viable bacteria of the wild type mutant and complemented strains at 1, 2, 3 and 4 hrs post-infection determined by an antibiotic protection assay. * $P < 0.001$.





Figure 7. The $\Delta sepA$ mutant strain is affected in its capacity to exclude the lysosomal marker Lamp-1

A. Representative immunofluorescence images of Lamp-1 positive and negative BCVs of the wild type (*Ba wt*) and mutant (*Ba ΔsepA*) strains in infected HeLa cells at 24 hours post-infection. Red, *Brucella*; green, Lamp-1. **B.** Quantification of the percent of Lamp-1 positive BCVs at 4 and 24 hrs post-infection with the wild type and the $\Delta sepA$ mutant strains.




Figure 8. Over-expression of SepA affects virulence in mice

A and B. Spleen colonization at 15 days post-infection of mice intraperitoneally infected with the wild type (*Ba wt*), *sepA* mutant (*Ba Δ sepA*), complemented (*Ba Δ sepA(sepA)*) or SepA over-expressing (*Ba wt(sepA)*) strains. Bacterial load was determined as indicated in Experimental procedures. * $P < 0.002$. **C.** Western-blot with a monoclonal anti-O-antigen antibody (M84) against whole bacteria showing that neither the mutant nor the over-expressing strains are rough.