



Optimal Combination of VNTR Typing for Discrimination of Isolated *Mycobacterium tuberculosis* in Korea

Jihye Lee, M.Sc., Heeyoon Kang, M.Sc., Sarang Kim, M.Sc., Heekyung Yoo, M.Sc., Hee Jin Kim, M.D. and Young Kil Park, Ph.D.

Korean Institute of Tuberculosis, Cheongwon, Korea

Background: Variable-number tandem repeat (VNTR) typing is a promising method to discriminate the *Mycobacterium tuberculosis* isolates in molecular epidemiology. The purpose of this study is to determine the optimal VNTR combinations for discriminating isolated *M. tuberculosis* strains in Korea.

Methods: A total of 317 clinical isolates collected throughout Korea were genotyped by using the IS6110 restriction fragment length polymorphism (RFLP), and then analysed for the number of VNTR copies from 32 VNTR loci.

Results: The results of discriminatory power according to diverse combinations were as follows: 25 clusters in 83 strains were yielded from the internationally standardized 15 VNTR loci (Hunter-Gaston discriminatory index [HGDI], 0.9958), 25 clusters in 65 strains by using IS6110 RFLP (HGDI, 0.9977), 14 clusters in 32 strains in 12 hyper-variable VNTR loci (HGDI, 0.9995), 6 clusters in 13 strains in 32 VNTR loci (HDGI, 0.9998), and 7 clusters in 14 strains of both the 12 hyper-variable VNTR and IS6110 RFLP (HDGI, 0.9999).

Conclusion: The combination of 12 hyper-variable VNTR typing can be an effective tool for genotyping Korean *M. tuberculosis* isolates where the Beijing strains are predominant.

Keywords: *Mycobacterium tuberculosis*; Molecular Epidemiology; Minisatellite Repeats

Introduction

Tuberculosis (TB) is a contagious disease that develops from infection with *Mycobacterium tuberculosis* bacilli in droplets projected by coughing of active TB patients. DNA

typing of *M. tuberculosis* is an efficient tool to show the transmission link of TB¹. In particular, IS6110 restriction fragment length polymorphism (RFLP) has been used for discrimination of Korean *M. tuberculosis* isolates because of high IS6110 copies and diverse patterns²⁻⁴. However, IS6110 RFLP has some disadvantages such as a lengthy turnaround time, difficulty in comparison between laboratory DNA typing data, and uncertain discrimination of isolates containing similar fragments sizes for inter- or intra-strains⁵.

Among more than 40 variable-number tandem repeat (VNTR) loci scattered on the *M. tuberculosis* chromosome⁶, 15 and 24 VNTR loci have been proposed as the international standard⁷. However, the discriminatory power is not enough in countries that have a high proportion of Beijing type *M. tuberculosis*⁸⁻¹². K strains that have 10 IS6110 RFLP copies are most dominant strain and often found in TB outbreak in Korea. K strains occupy about 4–5% out of any group of *M. tuberculosis* isolates isolated in Korea^{3,13}. K strains causes difficulty in discerning the direct transmission link by IS6110

Address for correspondence: Young Kil Park, Ph.D.

Korean Institute of Tuberculosis, 168-5 Osongsangmyung 4 ro, Osong-up, Cheongwon 363-954, Korea

Phone: 82-43-249-4960, **Fax:** 82-43-249-4965

E-mail: ypark7@empas.com

Received: Oct. 10, 2013

Revised: Nov. 4, 2013

Accepted: Nov. 7, 2013

©It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>).

Copyright © 2014

The Korean Academy of Tuberculosis and Respiratory Diseases.

All rights reserved.

RFLP typing only.

Therefore, we attempted to optimize a combination of VNTR loci and IS6110 RFLP for discrimination of Korean *M. tuberculosis* isolates.

Materials and Methods

1. Strains

A total of 317 strains were randomly selected among 2,400 strains isolated from smear and culture-positive primary pulmonary TB patients that registered at the Public Health Center (PHC) of Korea between 2006 and 2011. When the strains were clustered in IS6110 RFLP or VNTR typing, we collected

Table 1. Primer sequences and labeled dyes of 32 VNTR loci for multiplex PCR

Multiplex	Locus	Alias	Primer sequence	Unit (bp)	Dye
Mix 1	2163a	QUB-11a	CGTGATGTTGATCGGGATGT/ACCCCTGGAGTCTGGCATC	69	PET
	2996	MIRU-26	TAGGTCTACCGTCGAAATCTGTGAC/CATAGGCGACCAGGCGAATAG	51	VIC
	3192	MIRU-31	ACTGATTGGCTTCATACGGCTTTA/GTGCCGACGTGGTCTTGAT	53	NED
Mix 2	3820		TGCGCGGTGAATGAGACG/ACCTTCATCCTTGCGCAG	57	PET
	2163b	QUB-11b	CCGATGTAGCCCGTGAAGA/AGGGTCTGATTGGCTACTCA	69	FAM
	4156		ACCGCAAGGCTGATGATCC/GTGCATCTCGTCTGACTTCC	59	NED
Mix 3	3232		CCCCAGCCTTACGACTGA/GTCGGGCTTGGTGAAGG	56	FAM
	3336		ATCCCCGCGGTACCCATC/GCCAGCGGTGTCGACTATCC	59	PET
	4052	QUB-26	GTGCCGGCCAGGTCCTTCC/CACCGCGTGTGACCCGAAC	111	NED
Mix 4	1955	Mtub21	AGACGTCAGATCCCAGTT/ACCCGACAACAAGCCCA	57	VIC
	4120		GTTCACCGGAGCCAACC/GAGGTGGTTTCGTGGTTCG	57	PET
	0424	Mtub04	CTTGGCCGGCATCAAGCGCATTATT/GGCAGCAGAGCCCGGGATTCTTC	51	FAM
Mix 5	0580	MIRU-4	GCGCGAGAGCCCGAACTGC/GCGCAGCAGAAACGTCAGC	77	FAM
	2165	ETR-A	AAATCGGTCCCATCACCTTCTTAT/CGAAGCCTGGGGTGCCCGGATTT	75	NED
	1644	MIRU-16	TCGGTGATCGGGTCCAGTCCAAGTA/CCCCTCGTGCAGCCCTGGTAC	53	VIC
Mix 6	0960	MIRU-10	GTTCTTGACCAACTGCAGTCGTCC/GCCACCTTGGTGATCAGCTACCT	53	FAM
	3690	Mtub39	CGGTGGAGGCGATGAACGTCTTC/TAGAGCGGCACGGGGGAAAGCTTAG	58	VIC
	2074	Mtub24	TGTGTCACCTGACGATTTCAAGG/TGGCCGGCAAATAATGGATGC	56	PET
Mix 7	2347	Mtub29	GCCAGCCGCCGTGCATAAACCT/AGCCACCCGGTGTGCCTTGATGAC	57	FAM
	3007	MIRU-27	TCGAAAGCCTCTGCGTGCCAGTAA/GCGATGTGAGCGTGCCACTCAA	53	NED
	2461	ETR-B	GCGAACACCAGGACAGCATCATG/GGCATGCCGGTGATCGAGTGG	57	VIC
Mix 8	2531	MIRU-23	CTGTGATGGCCGCAACAAAACG/AGCTCAACGGGTTGCCCCCTTTTGTC	53	VIC
	4348	MIRU-39	CGCATCGACAAACTGGAGCCAAAC/CGGAAACGTCTACGCCCCACACAT	53	NED
	3155	QUB-15	GCCAGCCGTAACCCGACCAG/GGGCCGAAATTCGAGTGG	54	PET
Mix 9	2687	MIRU-24	CGACCAAGATGTGCAGGAATACAT/GGGCGAGTTGAGCTCACAGAA	54	VIC
	0802	MIRU-40	GGGTTGCTGGATGACAACGTGT/GGGTGATCTCGGCGAAATCAGATA	54	NED
	2372		ACCTCCGTTCGATAATC/CAGCTTTCAGCCTCCACA	57	PET
Mix 10	2401	Mtub30	CGTCGTCGCCGAGCTGGATT/CACCGGGGCTGGCAGCTAAG	58	PET
	3171	Mtub34	GGTGCGCACCTGCTCCAGATAA/GGCTCTCATTGCTGGAGGGTTGTAC	54	NED
	0577	ETR-C	GTGAGTCGCTGCAGAACCTGCAG/GGCGTCTTGACCTCCACGAGTG	58	VIC
Single	0154	MIRU-2	TGGACTTGCAGCAATGGACCAACT/TACTCGGACGCCGGCTCAAAAT	53	FAM
Single	2059	MIRU-20	TCGAGAGATGCCCTTCGAGTTAG/GGAGACCGCGACCAGGTACTTGTA	77	FAM

VNTR: variable-number tandem repeat; PCR: polymerase chain reaction.

the epidemiological information of the strains from the PHCs and each person who had the clustered strain by documents and telephone calls.

2. IS6110 DNA fingerprinting

For all 317 isolates, DNA isolation and IS6110 RFLP typing were performed as described previously². An RFLP cluster was defined by completely identical patterns after analysis of BioNumerics version 5.1 software (Applied Maths, Kortrijk, Belgium) within two or more isolates. K strains were identified according to the previously reported definition^{4,13}.

3. Variable number of tandem repeats

We selected 32 VNTR loci for this study^{7,9}. Primer sets for polymerase chain reaction (PCR) were prepared for the 32 VNTR loci. PCR was conducted as described previously^{14,16} and VNTR 4052, 3155, and 2074 were designated by the Research Institute of Tuberculosis in Japan (Table 1). Each primer set was labelled with 4 kinds of fluorescent dye for capillary sequencer analysis. The capillary sequencer, a 3500 Genetic analyzer (Applied Biosystems, Foster City, CA, USA), was used for measuring the precise size of PCR products for each VNTR loci. However, in cases of large fragment sizes over 12,000 bp or an ambiguous size in the capillary sequencer, we measured the size by 1% agarose gel electrophoresis. Clusters were found as the result of comparison of genotyping data by BioNumerics version 5.1 software (Applied Maths). For PCR amplification, we mainly used *Ex-Taq* polymerase (Takara, Tokyo, Japan), except for Mix4, Mix7, and Mix9, which used *KOD FX Taq* polymerase (TOYOBO, Tokyo, Japan). PCR conditions were as follows: pre-denaturation for 5 minutes at 94°C, denaturation for 30 seconds at 94°C, annealing for 30 seconds at 63°C, extension for 1 minute at 72°C, and post-extension for 10 minutes at 72°C, except for Mix9 that had an annealing temperature of 60°C.

4. Comparison of discriminatory power

The allelic diversity (h) of the 32 VNTR loci was calculated by the following formula: $h=1-\sum x_i^2$, where x_i is the frequency of the i th allele at the locus¹⁷. The discriminatory power of each method was calculated by the Hunter-Gaston discriminatory index (HGDI)¹⁸.

5. Ethical considerations

Ethical clearance was obtained from the Ethics Review Committee of Korean Institute of Tuberculosis.

Results

1. Analysis of the 32 VNTR loci

As a result of VNTR typing of the 317 *M. tuberculosis* strains isolated from Korea, VNTR loci showing a high h value of over 0.6 were VNTR 3232, 3820, 4120, 3336, 2163b, 0424, 1955, 4052, 3192, 4156, 2996, and 2163a. Interestingly, VNTR 0802, 3690, 2165, and 0960 revealed a high h value (over 0.6) only in non-Beijing *M. tuberculosis* isolates (Table 2).

Two PCR products amplified from VNTR 3802 and 0580 (MIRU04) were an unusual size, and were regarded as uncountable repeated numbers (Table 2). The imprecise PCR product of the VNTR 3802 loci was located between 8 and 9 copies, and that of VNTR 0580 was located between 3 and 4 copies of the repeated unit. Four PCR products in VNTR 3820, 4052 (QUB26), and 2372 had multiple bands. Non-amplified PCR products were found in VNTR 3232, 4120, and 2163b (QUB11b), 3192 (MIRU31), 2163a (QUB11a), 2165 (ETR-A), and 2074 (Mtub24).

2. Epidemiological linkage of the VNTR clusters

Six clusters in 13 patients were found after VNTR typing of the 32 loci, and only 3 clusters in 6 patients were found when additionally analysed by IS6110 RFLP typing (Table 3). Out of the 3 clusters, we found only one definite epidemiological linkage through personal contact. Patients with the V32C6 cluster were brother and sister. Patient 06-1731 with the cluster was the brother who developed TB in 2006, and was cured completely in 2007. Patient 08-148, who was the sister of patient 06-1731, had developed TB, and was cured in 2008. Even though random selection of PHC strains, we found the strains isolated from brother and sister TB-developed in different year by accident.

Two patients with V32C3 and IS6110 clusters had lived in Jeju province, even though they were not acquainted with each other. The two strains (06-890 and 09-1184) with the C32C4 cluster and identical IS6110 RFLP types were the K strain that is the most frequent endemic strain in Korea (4–5% in any TB population). The other strain (10-335) with the C32C4 cluster was the K family that exhibited a difference of only one band. Interestingly, another patient (10-179) with the V32C1 cluster also had the K strain. However, we found different copies in VNTR 2163a, 3232, and 3820 compared with that in V32C4 (data not shown).

3. Discriminatory power of VNTR and IS6110 RFLP typing

When the internationally standardized 15 VNTR loci were applied to the 317 strains, we found 25 clusters in 83 strains (HGDI, 0.9958) (Table 4), and at least 7 VNTR loci (3232,

Table 2. Allelic diversity and *h* values of 32 *Mycobacterium tuberculosis* VNTR loci determined by 317 Korean isolates

VNTR locus (alias)	Allele																															<i>h</i> value		
	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	>20	Imprecision	Multi-band	Non-PCR	Total (n=317)	Beijing (n=258)	Non-Beijing (n=59)						
3232	-	-	2	5	1	13	7	13	8	11	34	28	64	29	51	12	18	2	7	3	1	7	-	-	1	0.894	0.866	0.866						
3820	-	-	-	6	1	43	2	21	14	67	16	19	26	7	51	9	8	7	2	7	2	6	1	2	-	0.887	0.865	0.541						
4120	-	-	4	23	32	18	11	31	85	28	48	11	8	7	1	1	3	3	1	-	1	-	-	1	0.864	0.827	0.678							
3336	-	-	1	2	7	19	13	112	11	18	49	18	8	9	43	1	1	-	3	1	1	-	-	-	-	0.818	0.752	0.846						
2163b (QUB11b)	-	3	27	29	42	99	79	29	4	1	-	-	1	-	-	-	-	-	-	1	-	-	-	2	0.796	0.751	0.750							
0424 (Mtrub 04)	-	5	102	100	105	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.687	0.665	0.510							
1955 (Mtrub 21)	3	9	62	112	126	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.678	0.588	0.450							
4052 (QUB 26)	-	1	9	1	1	13	18	38	162	66	6	1	-	-	-	-	-	-	-	-	-	-	1	-	0.673	0.627	0.782							
3192 (MIRU 31)	-	-	18	46	70	157	20	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	0.669	0.561	0.507							
4156 (QUB4156)	-	1	8	126	130	52	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.646	0.640	0.298							
2996 (MIRU26)	-	1	2	34	20	40	13	185	16	6	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.623	0.482	0.622							
2163a (QUB11a)	-	1	3	-	15	68	7	12	182	9	1	2	-	-	-	-	-	-	-	-	-	11	-	6	0.604	0.483	0.606							
3155 (QUB15)	-	-	128	6	173	10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.538	0.539	0.448						
0802 (MIRU40)	-	9	80	202	22	3	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.525	0.488	0.642						
3690 (Mtrub39)	-	5	26	217	34	29	2	3	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.504	0.426	0.742						
2372	-	10	49	232	17	7	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	0.433	0.280	0.474						
4348 (MIRU39)	-	11	70	230	4	-	1	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.423	0.250	0.390						
2165 (ETR-A)	-	2	17	43	251	-	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	0.339	0.154	0.665						
0960 (MIRU10)	-	8	40	255	10	2	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.335	0.175	0.606						
2401 (Mtrub30)	-	4	56	-	255	1	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.322	0.103	0.429						
2074 (Mtrub24)	-	4	43	259	10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	0.309	0.173	0.591						
1644 (MIRU16)	-	5	10	288	14	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.171	0.139	0.299						
3007 (MIRU27)	-	2	12	292	11	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.149	0.111	0.298						
0580 (MIRU04)	5	5	297	5	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	-	-	0.121	0.083	0.275						
2347 (Mtrub29)	-	-	12	4	298	1	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.115	0.104	0.159						
2531 (MIRU23)	-	3	3	-	2	300	8	-	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.104	0.061	0.270						
2461 (ETR-B)	-	16	301	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.096	0.023	0.344						
0577 (ETR-C)	-	-	8	4	302	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.092	0.038	0.297						
2059 (MIRU20)	-	8	305	4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.073	0.075	0.065						
3171 (Mtrub34)	-	-	7	307	3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.062	0.046	0.128						
0154 (MIRU02)	-	5	310	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.043	0.031	0.097						
2687 (MIRU24)	-	317	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0	0	0						

VNTR: variable-number tandem repeat; PCR: polymerase chain reaction.

Table 3. Characteristics of 6 clusters isolated from VNTR typing analysis in 32 VNTR loci

VNTR cluster	Year	Region	ID	No. of IS6110 copies	IS6110 RFLP pattern
V32C1	2006	Ansan	06-681	9	
	2010	Jungseong	10-179	10	
V32C2	2007	Pyongtaek	07-230	12	
	2007	Pyongtaek	07-977	13	
V32C3	2006	Jeju	06-481	17	
	2006	Jeju	06-571	17	
V32C4	2006	Ansan	06-890	10	
	2009	Gosung	09-1184	10	
	2010	Pohang	10-335	11	
V32C5	2007	Anyang	07-2256	18	
	2007	Tongyoung	07-1645	19	
V32C6	2006	Busan	06-1731	16	
	2008	Busan	08-148	16	

VNTR: variable-number tandem repeat; RFLP: restriction fragment length polymorphism.

Table 4. Comparison of discriminatory power according to DNA typing method alone or combination

Typing method	No. of DNA types	No. of unique types	No. of clusters	Total isolates belonging to clusters	Maximum isolates belonging to a cluster	HGDI
IS6110 RFLP	277	252	25	65	13	0.9977
12 VNTR	299	285	14	32	4	0.9995
15 VNTR loci	257	232	25	83	16	0.9958
32 VNTR	310	304	6	13	3	0.9998
32 VNTR+IS6110 RFLP	314	311	3	6	2	0.9999
12 VNTR+IS6110 RFLP	310	303	7	14	2	0.9999

Fifteen variable-number tandem repeat (VNTR) loci are of international standards, and 12 VNTR are hyper-variables. RFLP: restriction fragment length polymorphism.

3336, 3820, 4120, 2372, 4348, and 3155) were needed to be equivalent to the discriminatory power of the 32 VNTR loci (data not shown). However, in cases of the hyper-variable 12 VNTR loci (3232, 3820, 4120, 3336, 2163b, 0424, 1955, 4052, 3192, 4156, 2996, and 2163a), 14 clusters in 32 strains (HGDI, 0.9995) were found, and we needed only 4 VNTR loci (4348, 2165, 0577, and 2372) to be equal to the discriminatory power of the 32 VNTR loci (data not shown). Furthermore, in the IS6110 RFLP typing, 25 clusters in 65 strains (HGDI, 0.9977) (Table 4) were obtained, and we needed at least 14 VNTR loci (4120, 3820, 2372, 1955, 0424, 4052, 3155, 2163b, 2163a, 0577, 0802, 4348, 3192, and 2996) for 3 clusters as a result of both the 32 VNTR loci and IS6110 RFLP typing (data not shown). When the 12 hyper-variable VNTR loci were applied to the 317 strains, followed by IS6110 RFLP typing, the discrimina-

tory power was almost identical to the result of both the 32 VNTR loci and IS6110 RFLP typing (Table 4).

Discussion

VNTR typing has been proposed as an alternative to IS6110 RFLP typing that bears intrinsic drawbacks for *M. tuberculosis* DNA typing. However, the disadvantage of VNTR typing is low discriminatory power with few combinations of VNTR loci. Therefore, we tried to find an optimal combination of DNA typing methods to discriminate *M. tuberculosis* isolated from Korea, which is accompanied by a high proportion of Beijing type^{3,4}. Furthermore, compared with the surrounding countries, there is a higher proportion of RD181 among the

ancient Beijing types⁴. These peculiar characteristics of Korean *M. tuberculosis* strains lead to an unclear distinction of *M. tuberculosis* isolates with only the 15 international VNTR standard VNTR loci.

Recently, the utility of hyper-variable VNTR loci has been used to compensate for the lack of discriminatory power of the 15 VNTR loci^{10,11,16}. Hyper-variable VNTR loci, including 3232, 3820, 4120, 3336, and 2163a, also revealed a high *h* value in this study, which was similar to that in other studies^{9-11,16}. In the report Murase et al.⁹, more than 4% of these 5 loci had 15 or more copies, leading to difficulty in interpreting the exact copy number. We also found that 2–4% had 15 or more copies in VNTR 2163b, 3336, and 4120, and 13–16% had 15 or more copies in VNTR 3820, and 3232. These hyper-variable VNTR loci are excluded in the international standard 15 VNTR loci⁷ and JATA 12⁹ because of the absence of PCR products, PCR products that are difficult to interpret with 15 or more copies, and amplification of multiple alleles. Non-amplification of PCR products not only occurred for hyper-variable VNTR loci but also general VNTR loci such as 2165 (ETR-A), and 2074 (Mtub24) in this study (Table 2). The major problem of hyper-variable VNTR loci was a high copy number of the repeats, which required additional analysis such as agarose gel electrophoresis. However, the copies in these hyper-variable VNTR loci are so diverse that they have a higher discriminatory power that is too valuable to exclude. Iwamoto et al.¹⁶ also recommended that these hyper-variable VNTR loci for second-line typing of clustering following the international standard 15 loci.

MIRU 40, Mtub 21, VNTR 4156, Mtub 04, QUB26, and QUB1b also revealed high *h* values in a previous study of South Korean *M. tuberculosis*¹⁹. In this study, we obtained a good result using hyper-variable VNTR loci, and IS6110 RFLP was useful as a secondary tool to discriminate the clusters.

An intriguing characteristic was found after comparing the mode of VNTR loci among Korea, Taiwan, and Japan^{9,20}. Between Taiwan and Korea, there were some differences in the mode of copies of VNTR loci 0154, 3192, 2163b, 4052, 0424, 1955, 2347, and 2401. In particular, 5 copies of VNTR 0154, 0424, and 1955, and 3 copies of VNTR 2347 were modes in Taiwan but rare in Korea. Between Japan and Korea, there were some differences in VNTR 2163b, 1955, 3155, 3232, 3820, and 4120. Notably, 5 copies of VNTR 3155 and 4 copies of VNTR 3336 were more frequent in Japan but rare in Korea. These differences may be a clue for differentiation among the three countries. Compared with Beijing strains, we found that VNTR 2165, 0960, and 2074 loci had excellent higher *h* values for the non-Beijing strains, indicating that these VNTR loci may be useful to discriminate *M. tuberculosis* strains in countries with a high proportion of non-Beijing strains. The optimal combination of VNTR loci may be different depending on the proportion of Beijing strains or non-Beijing strains in each country.

We could not find any cluster consisted of only multi-drug resistant (MDR) strains in this study. The strains included in this study were selected randomly among strains collected from PHCs. Therefore, most of them (281 strains) were pan-susceptible, only a few of strains (13 strains) were MDR, and the remains (23 strains) were any drug resistant with non-MDR.

Even though we analysed 32 VNTR loci for discrimination of 317 Korean *M. tuberculosis* isolates, we found 6 clusters. Among them, 3 clusters were not found to be clusters when additionally analysed by the IS6110 RFLP typing method, indicating that IS6110 RFLP typing is very useful for sub-classifying VNTR clusters. In terms of cost efficiency, it is difficult to use 32 VNTR loci for discrimination of *M. tuberculosis*, and the 15 international standard VNTR loci do not have satisfactory discrimination power for Korean strains. Inevitably, we need hyper-variable VNTR loci and the additional IS6110 RFLP typing method for effective discrimination of Korean *M. tuberculosis* strains.

The combination of 12 hyper-variable VNTR typing can be an effective tool for genotyping Korean *M. tuberculosis* isolates in which Beijing strains are predominant.

References

1. van Embden JD, Cave MD, Crawford JT, Dale JW, Eisenach KD, Gicquel B, et al. Strain identification of *Mycobacterium tuberculosis* by DNA fingerprinting: recommendations for a standardized methodology. *J Clin Microbiol* 1993;31:406-9.
2. Park YK, Bai GH, Kim SJ. Restriction fragment length polymorphism analysis of *Mycobacterium tuberculosis* isolated from countries in the western pacific region. *J Clin Microbiol* 2000;38:191-7.
3. Park YK, Shin S, Ryu S, Cho SN, Koh WJ, Kwon OJ, et al. Comparison of drug resistance genotypes between Beijing and non-Beijing family strains of *Mycobacterium tuberculosis* in Korea. *J Microbiol Methods* 2005;63:165-72.
4. Kang HY, Wada T, Iwamoto T, Maeda S, Murase Y, Kato S, et al. Phylogeographical particularity of the *Mycobacterium tuberculosis* Beijing family in South Korea based on international comparison with surrounding countries. *J Med Microbiol* 2010;59(Pt 10):1191-7.
5. Cowan LS, Diem L, Monson T, Wand P, Temporado D, Oemig TV, et al. Evaluation of a two-step approach for large-scale, prospective genotyping of *Mycobacterium tuberculosis* isolates in the United States. *J Clin Microbiol* 2005;43:688-95.
6. Smittipat N, Palittapongarnpim P. Identification of possible loci of variable number of tandem repeats in *Mycobacterium tuberculosis*. *Tuber Lung Dis* 2000;80:69-74.
7. Supply P, Allix C, Lesjean S, Cardoso-Oelemann M, Rusch-Gerdes S, Willery E, et al. Proposal for standardization of optimized mycobacterial interspersed repetitive unit-

- variable-number tandem repeat typing of *Mycobacterium tuberculosis*. J Clin Microbiol 2006;44:4498-510.
8. Jiao WW, Mokrousov I, Sun GZ, Guo YJ, Vyazovaya A, Narvskaya O, et al. Evaluation of new variable-number tandem-repeat systems for typing *Mycobacterium tuberculosis* with Beijing genotype isolates from Beijing, China. J Clin Microbiol 2008;46:1045-9.
 9. Murase Y, Mitarai S, Sugawara I, Kato S, Maeda S. Promising loci of variable numbers of tandem repeats for typing Beijing family *Mycobacterium tuberculosis*. J Med Microbiol 2008;57(Pt 7):873-80.
 10. Mokrousov I, Narvskaya O, Vyazovaya A, Millet J, Otten T, Vishnevsky B, et al. *Mycobacterium tuberculosis* Beijing genotype in Russia: in search of informative variable-number tandem-repeat loci. J Clin Microbiol 2008;46:3576-84.
 11. Mokrousov I, Vyazovaya A, Otten T, Zhuravlev V, Pavlova E, Tarashkevich L, et al. *Mycobacterium tuberculosis* population in northwestern Russia: an update from Russian-EU/Latvian border region. PLoS One 2012;7:e41318.
 12. Iwamoto T, Grandjean L, Arikawa K, Nakanishi N, Caviedes L, Coronel J, et al. Genetic diversity and transmission characteristics of Beijing family strains of *Mycobacterium tuberculosis* in Peru. PLoS One 2012;7:e49651.
 13. Kim SJ, Bai GH, Lee H, Kim HJ, Lew WJ, Park YK, et al. Transmission of *Mycobacterium tuberculosis* among high school students in Korea. Int J Tuberc Lung Dis 2001;5:824-30.
 14. Roring S, Scott A, Brittain D, Walker I, Hewinson G, Neill S, et al. Development of variable-number tandem repeat typing of *Mycobacterium bovis*: comparison of results with those obtained by using existing exact tandem repeats and spoligo-typing. J Clin Microbiol 2002;40:2126-33.
 15. Smittipat N, Billamas P, Palittapongarnpim M, Thong-On A, Temu MM, Thanakijcharoen P, et al. Polymorphism of variable-number tandem repeats at multiple loci in *Mycobacterium tuberculosis*. J Clin Microbiol 2005;43:5034-43.
 16. Iwamoto T, Yoshida S, Suzuki K, Tomita M, Fujiyama R, Tanaka N, et al. Hypervariable loci that enhance the discriminatory ability of newly proposed 15-loci and 24-loci variable-number tandem repeat typing method on *Mycobacterium tuberculosis* strains predominated by the Beijing family. FEMS Microbiol Lett 2007;270:67-74.
 17. Sun YJ, Bellamy R, Lee AS, Ng ST, Ravindran S, Wong SY, et al. Use of mycobacterial interspersed repetitive unit-variable-number tandem repeat typing to examine genetic diversity of *Mycobacterium tuberculosis* in Singapore. J Clin Microbiol 2004;42:1986-93.
 18. Hunter PR, Gaston MA. Numerical index of the discriminatory ability of typing systems: an application of Simpson's index of diversity. J Clin Microbiol 1988;26:2465-6.
 19. Shamputa IC, Lee J, Allix-Beguec C, Cho EJ, Lee JI, Rajan V, et al. Genetic diversity of *Mycobacterium tuberculosis* isolates from a tertiary care tuberculosis hospital in South Korea. J Clin Microbiol 2010;48:387-94.
 20. Chen YY, Chang JR, Huang WF, Kuo SC, Su IJ, Sun JR, et al. Genetic diversity of the *Mycobacterium tuberculosis* Beijing family based on SNP and VNTR typing profiles in Asian countries. PLoS One 2012;7:e39792.