



CrossMark  
click for updates

## Review

**Cite this article:** Demmig-Adams B, Stewart JJ, Adams III WW. 2014 Multiple feedbacks between chloroplast and whole plant in the context of plant adaptation and acclimation to the environment. *Phil. Trans. R. Soc. B* **369**: 20130244.

<http://dx.doi.org/10.1098/rstb.2013.0244>

One contribution of 20 to a Theme Issue 'Changing the light environment: chloroplast signalling and response mechanisms'.

### Subject Areas:

plant science

### Keywords:

phloem loading, photosynthesis, PsbS, source–sink, tocopherol, zeaxanthin

### Author for correspondence:

Barbara Demmig-Adams

e-mail: [barbara.demmig-adams@colorado.edu](mailto:barbara.demmig-adams@colorado.edu)

# Multiple feedbacks between chloroplast and whole plant in the context of plant adaptation and acclimation to the environment

Barbara Demmig-Adams, Jared J. Stewart and William W. Adams III

Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309-0334, USA


This review focuses on feedback pathways that serve to match plant energy acquisition with plant energy utilization, and thereby aid in the optimization of chloroplast and whole-plant function in a given environment. First, the role of source–sink signalling in adjusting photosynthetic capacity (light harvesting, photochemistry and carbon fixation) to meet whole-plant carbohydrate demand is briefly reviewed. Contrasting overall outcomes, i.e. increased plant growth versus plant growth arrest, are described and related to respective contrasting environments that either do or do not present opportunities for plant growth. Next, new insights into chloroplast-generated oxidative signals, and their modulation by specific components of the chloroplast's photoprotective network, are reviewed with respect to their ability to block foliar phloem-loading complexes, and, thereby, affect both plant growth and plant biotic defences. Lastly, carbon export capacity is described as a newly identified tuning point that has been subjected to the evolution of differential responses in plant varieties (ecotypes) and species from different geographical origins with contrasting environmental challenges.

## 1. Introduction

Multiple pathways of communication, over both short and long distances, between the various functional parts of the plant are being elucidated. It appears self-evident that such communication pathways should enable the plant to respond appropriately to opportunities as well as challenges in its physical and biological environment. We place the communication between chloroplast and the whole plant into a context of the adjustment in form and function via physiological *acclimation* of individual plants to their growth environment as well as genetic *adaptation* of species and varieties (ecotypes) to the climate and other features of the habitats in which the respective lines evolved.

## 2. Matching energy acquisition with energy utilization: established and emerging signalling pathways serving to optimize chloroplast and whole-plant function in a given environment

Here, we review the signalling pathways serving to match light availability, and energy acquisition in photosynthesis, with energy utilization for plant growth, development and defences—in the context of plant genetic make-up, developmental status and a wide range of environmental conditions. Figure 1 presents the principal functional sequence from solar energy acquisition and sugar production to sugar distribution throughout the plant (solid arrows), starting with light absorption in the light-harvesting complexes (light harvesting) and primary photochemistry (in photosystem II (PSII) and the oxygen-evolving complex (OEC)) supporting photosynthetic electron transport and CO<sub>2</sub> fixation



**Figure 1.** Schematic depiction of principal steps involved in photosynthesis (light harvesting, photochemistry, involving photosystem II (PSII) and the oxygen-evolving complex (OEC) and carbon fixation in the Calvin cycle (CC) and in sugar export (phloem loading and sugar transport) to the plant's sinks (solid arrows) as well as (dashed arrows) feedback loops from sugar export and the plant's sinks to photosynthesis and (dotted arrows) feed-forward and feedback loops related to the generation of reactive oxygen species (ROS) in light harvesting and photochemistry. Heat = heat released via thermal dissipation of excess excitation energy in light-harvesting antennae. (Online version in colour.)

(in the Calvin cycle (CC)), followed by the loading of sugars into the phloem conduits and the transport of sugars to the plant's sinks (sugar-consuming and sugar-storing tissues). In addition, figure 1 features a host of feedback loops falling into two principal categories of either (i) (dashed arrows along the top) providing feedback from the plant's sugar-transport system and its sinks, via modulation of foliar sugar/starch content, back to light harvesting, photochemistry and CO<sub>2</sub> fixation or (ii) (dotted arrows along the bottom) providing both feed-forward and feedback loops (dotted arrows along the bottom) associated with chloroplast redox balance (determined by the ratio, and action of, oxidants and antioxidants) as influenced by the environment and by the chloroplast's photoprotective network. In the following, we will examine examples of source- and sink-driven signalling networks, and will argue that both source- and sink-driven signals modulate phloem loading/transport, and thereby impact plant response to environmental stresses.


### (a) Sink-to-source signalling in photosynthetic adjustment

It has long been recognized that the leaf's capacity for photosynthesis is adjusted in response to the level of demand generated by the plant's sinks for the products of photosynthesis [1–3]. A plant's inherent growth rate, developmental status and the opportunity (or lack thereof) for plant growth presented in the environment all contribute to setting this 'sink demand' as the level of plant sink strength. For example, (i) short-lived annual species tend to have higher inherent growth rates than slowly growing evergreens, (ii) a plant in its reproductive state tends to have greater sink strength than a vegetative plant and (iii) soil conditions (e.g. water and nutrient availability, temperature and soil

texture/structure), together with climatic features (e.g. light availability, temperature, humidity), cumulatively generate either plant growth-promoting or plant growth-impeding environmental conditions. It has been shown that regulation of foliar photosynthetic capacity occurs via modulation of the expression of photosynthetic genes involved in e.g. light harvesting, photochemistry and other thylakoid processes, and CO<sub>2</sub> fixation (e.g. [4–6]). The negative feedback via limiting carbon export was established by studying photosynthetic repression in response to inhibition of carbon export from leaves via e.g. girdling of branches or trees (e.g. [7–9]) or cold-girdling of stems [1,5]; the negative feedback from limiting sink strength was demonstrated e.g. via photosynthetic repression in response to removal of fruit and other sinks (e.g. [7,10–12]). Foliar sugar and/or starch accumulation are typically observed under either limiting carbon export at the leaf level or limiting plant sink strength [13].

### (b) Contrasting overall outcomes in environments with contrasting opportunities for plant growth: increased plant growth versus plant growth arrest

Figure 2 illustrates that two principal scenarios with opposite outcomes can be distinguished by examining the level of 'opportunity' for plant growth presented by the environment. We will illustrate these scenarios for the example of overall plant growth response to resource availability in the environment, while acknowledging that the same principles also apply to other more complex features, including temporal dynamics in the environment and in plant-organ-specific growth patterns. Plant response to opportunity presented by the environment presumably involves the feedback loops detailed in §2a and some of those discussed in §2c.



**Figure 2.** Schematic depiction of contrasting scenarios, representing increased, limiting or no opportunity for growth in the environment, and their respective effects on photosynthesis, carbon export, plant growth and the flexibility of thermal dissipation. NPQ, non-photochemical quenching on chlorophyll fluorescence, as an indicator for thermal dissipation of excess excitation energy; OEC, oxygen-evolving complex; PSII, photosystem II.

On the one hand (figure 2, left-hand side), there is the scenario of an increase in the opportunity for plant growth presented by the environment. Such a scenario can arise via increased availability of previously limiting resources such as light, water and/or nutrients, a shift to a more favourable temperature regime, etc., experienced by a species with the appropriate genetic adaptations and developmental state allowing it to upregulate photosynthetic capacity, carbon export capacity (see §3) and growth rate. As an additional response to increased light availability, any plant not increasing its growth and photosynthesis rates enough to use *all* of the additional available light will probably upregulate its capacity for thermal dissipation of any excess of absorbed light [14,15] and/or employ a variety of mechanisms to decrease light interception, e.g. via changes in leaf angle or leaf reflectance [16–19], or decrease light absorption, via decreases in chlorophyll content and light-harvesting capacity (e.g. [20,21]).

On the other hand (figure 2, right-hand side), limiting or no opportunity for plant growth is presented in environments with either limiting or excessive resources (like water, nutrients) or unfavourable temperatures. Such environmental conditions typically result in reduced growth or even growth arrest and a resulting decrease in plant sink strength [13]. Furthermore, in the latter environments, photosynthetic depression, with decreased photosynthetic capacity and light-harvesting capacity and/or inactivation of photochemistry ('photoinhibition'), is seen [13,22,23].

While decreased photosynthetic capacity and light-harvesting capacity are features that have long been associated with photosynthetic repression under sink limitation (see above), the phenomenon of the 'photoinhibition' of photosynthesis—involving an inactivation of PSII and/or OEC—is commonly instead interpreted as a sign of 'damage' (e.g. [24–31]). We have recently summarized evidence showing that, in all cases in which photoinhibitory inactivation of photochemistry and foliar carbohydrate levels were both assessed, photoinhibition

was found to be associated with greater accumulation of foliar sugar and/or starch [13]. Viewing such responses in the context of the whole plant in its environment thus offers opportunities for a more holistic interpretation of responses like the phenomenon of 'photoinhibition'. The invariable association of photoinhibition with foliar carbohydrate accumulation in plants under field conditions suggests that the tacit assumption that photoinhibition limits plant productivity (e.g. [32–40]) should be re-assessed for the possibility that limiting plant productivity instead triggers photoinhibition [13]. Based on the above-described data, the possibility that inactivation of PSII and/or OEC is, in fact, a principal component (together with decreased light harvesting and CO<sub>2</sub> fixation) of photosynthetic repression under certain sink limiting conditions should be seriously considered.

There is a third scenario (figure 2, middle), where light availability is suddenly increased for a plant that is not genetically adapted for, and/or finds itself in an environment not suitable for, increasing carbon export and/or plant sink strength. Such a scenario is not common in natural settings, but is a frequently used experimental approach to study photoinhibition in response to sudden transfer of shade-grown plants to high light. We have shown that sudden transfer of evergreens grown under non-fluctuating low light levels to highly excessive light results in photoinhibition of PSII and of photosynthetic capacity accompanied by foliar carbohydrate accumulation [13,41]. Especially in evergreens, which, as their name implies, do not commonly lower their chlorophyll content (or their light-harvesting capacity), photoinhibition is associated with a conversion of the otherwise rapidly reversible, flexible thermal dissipation of excess light (assessed from non-photochemical quenching (NPQ) of chlorophyll fluorescence) to a less flexible form of thermal dissipation (sustained NPQ) remaining continuously engaged for prolonged time periods [14,15,22,41–43]. An analogous scenario for annuals, as species presumably possessing sufficient carbon

**Table 1.** Summary of evidence for a role of the antioxidant tocopherol, the photosystem II protein PsbS and the xanthophyll zeaxanthin in modulating cell wall ingrowths in phloem-loading complexes, herbivore resistance and oxylipin levels. While the *npq1* mutant exhibited a trend for increased cell wall ingrowths that was not significant, double mutant *npq1 lut2* exhibited significantly greater wall ingrowths than wild-type when transferred from low to high growth light. The PsbS-deficient *npq4* mutant exhibited significantly greater oxylipin levels than wild-type under herbivore attack in the field but not in the absence of herbivores.

mutant or treatment	resulting effect	sources
<i>vte</i> mutant (tocopherol-/vitamin E'-deficient)	callose deposition resulting in enhanced wall ingrowths in phloem-loading complexes	[57]
oxylipin (methyl jasmonate) treatment	enhanced wall ingrowths in phloem-loading complexes	[58]
<i>npq1 lut2</i> mutants (zeaxanthin- and lutein-deficient)	enhanced wall ingrowths in phloem-loading complexes	[56]
<i>npq4</i> mutant (PsbS-deficient)	enhanced herbivore resistance	[59]
<i>vte</i> mutant (tocopherol-/vitamin E'-deficient)	enhanced oxylipin levels and enhanced ROS levels	[60–63]
<i>npq1</i> mutant (zeaxanthin-deficient)	enhanced oxylipin levels	[56]
<i>npq4</i> mutant (PsbS-deficient)	enhanced oxylipin levels and enhanced ROS levels	[59,64]

export capacity and sink strength to respond to increased light availability, is a sudden transfer from low to high growth light environment under *limiting soil nitrogen* supply, which results in photoinhibition as well [44].


### (c) Oxidative signals and phloem blocking/occlusion as modulated by chloroplast photoprotection

In the lower portion of figure 1 (dotted arrows), selected examples are featured for the generation of oxidative signals associated with light-harvesting and primary photochemistry. In plants adapted and acclimated to environments where leaves absorb a greater level of light than they are able to use through electron transport, a large fraction of the resulting excess excitation energy is safely dissipated (as heat; figure 1) via thermal dissipation of excess excitation energy [23]. The low remaining levels of excitation energy in excess of what is used in electron transport lead to the formation of reactive oxygen species (ROS; figure 1). At low concentrations, ROS trigger formation of oxidative signals (figure 1), while massive ROS formation presumably leads to cellular damage [45,46]. In light-harvesting complexes, excess excitation energy promotes the formation of (electronically excited) singlet oxygen as a form of ROS, while charge separation in primary photochemistry can give rise to the formation of singly reduced superoxide as the second major form of ROS arising in the light reactions [47]. Chloroplast antioxidant defences (figure 1) de-excite, and thereby detoxify, both singlet oxygen and superoxide, but highly excessive light levels presumably allow remaining ROS to trigger multiple redox-associated signalling pathways. For selected reviews of various examples of the many different components of the redox-signalling network originating in the chloroplast, see the studies of Munné-Bosch *et al.* [48], Baginsky & Link [49], Dietz *et al.* [50], Foyer *et al.* [51] and Mullineaux *et al.* [52]. We will highlight a selected example of one such signalling pathway, leading to the lipid-peroxidation-derived oxylipin messengers (figure 1), that had previously been shown to be involved in the regulation of the biosynthesis of chloroplast antioxidants and facilitators of thermal dissipation [53–55] and are now emerging as themselves being subjected to modulation by some of these same chloroplast antioxidants and facilitators of thermal dissipation [56] (table 1). We are highlighting this

specific example among the many redox-signalling networks because of the connections that have been made between this particular signalling pathway and phloem loading/transport.

Table 1 summarizes evidence for modulation of the formation of oxylipin messengers by the antioxidant vitamin E (tocopherols) and two factors directly associated with thermal energy dissipation in the light-harvesting complexes, the xanthophylls zeaxanthin and lutein and the PSII protein PsbS. Mutants deficient in the antioxidant vitamin E, in the xanthophylls zeaxanthin and lutein or in the PsbS protein generated greater levels of ROS [62,64] and greater levels of oxylipin messengers compared with wild-type (table 1). The key oxylipin studied in this work (figure 1 and table 1) was the stress hormone jasmonic acid, known to regulate plant growth and development as well as (biotic) defences against pathogens and pests (e.g. [65]). In addition, the latter studies also gave attention to actual evidence for altered biotic defences; it was found that the PsbS-deficient mutant (with lower levels of flexible NPQ and enhanced production of ROS and jasmonic acid) was more resistant to *herbivore* attack than the wild-type under field conditions (table 1). Similarly, evidence was also accumulated for an alteration of plant responses associated with defences against *pathogens*. An occlusion of the leaf's phloem-loading complexes and its long-distance phloem conduits presumably blocks the internal spreading of pathogens (like fungi and viruses that use these transport channels) within the plant [66–68]. Mutants deficient in the antioxidant vitamin E, and thus forming increased levels of jasmonic acid (table 1), were shown to be impaired in phloem loading, carbon export, and plant growth and accumulated foliar carbohydrates (for a review, see [48] and also [57,63,69–72]). Similarly, mutants deficient in zeaxanthin exhibited not only enhanced oxylipin production but also increased cell wall deposition in phloem-loading cells (table 1).

It has been recognized that cell wall architecture is a component of plant biotic defence: 'A rigid cell wall can fend off pathogen attack by forming an impenetrable, physical barrier' [73, p. 549]. The specific occlusion of the gateways involved in phloem loading and carbon export from leaves is a defence mechanism that presumably comes at a cost to plant growth and thus has far-reaching implications. It is increasingly recognized that combinations of abiotic and biotic stresses in a plant's environment can either exacerbate or ameliorate each other [74].




**Figure 3.** Differences in total cross-sectional sieve tube area per foliar sugar-loading vein induced by growth temperature (temp.) and growth light environment in *Arabidopsis* ecotypes from Sweden (filled squares) and Italy (filled circles) (as well as in the model line Col-0 (open triangles)). Sieve tubes are also referred to as sieve elements in the literature on plant vasculature. \* $p < 0.05$ ; \*\*\* $p < 0.001$ ; light intensity is given in  $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ .

Likewise, multiple plant-hormone-based signalling pathways exhibit both *antagonistic* and *cooperative* interactions [75–77]. It is feasible that trade-offs may exist not only between plant growth versus defence and thus plant abiotic and biotic stress resistance, but also between plant resistance to pests versus pathogens, and/or between additional specific features. For example, *greater* vitamin E content of certain pepper varieties was associated with *inhibition* of the development of an *insect pest* of pepper [78]—whereas *decreased* vitamin E content of *Arabidopsis* mutants was associated with *enhanced* putative *pathogen* defence via phloem occlusion (table 1). Future research should examine combinations of individual plant species and specific pests or pathogens in specific abiotic environments.

### 3. Adaptation to contrasting environmental challenges with respect to carbon export-related features

In addition to being the site of the above-described occlusion (table 1 and figure 1) in the possible service of defence against pathogens that spread through the phloem, phloem-loading complexes were recently proposed to be the target point for overall adjustment of foliar carbon export capacity in response to environmental change [79,80]. Differences in the adjustability of phloem loading and carbon export were, furthermore, shown to be excellent predictors for differences in *photosynthetic acclimation* to the environment between plant varieties (ecotypes) and species from different geographical regions [81]. The latter work indicates that control points and within-plant signalling networks are subjected to genetic adaptation of



**Figure 4.** (a) Light- and CO<sub>2</sub>-saturated photosynthetic rate (measured at 25°C), (b) sieve element number per foliar minor vein and (c) foliar vein density, in the winter annual *Arabidopsis thaliana* and the summer annual squash, *Cucurbita pepo* L. cv. Italian Zucchini Romanesco, both grown at 25°C under a 9-h photoperiod of 1000  $\mu\text{mol photon m}^{-2} \text{s}^{-1}$ . \*\*\* $p < 0.001$ . For further details on methods and growing conditions, see Cohu *et al.* [80,81].

plants to their environment over evolutionary time. These responses are featured in figures 1 and 2 as (increased, decreased or genetically limited) carbon export capacity (phloem loading/sugar transport) and photosynthetic capacity.

Firstly, it was shown that number and size of the cells loading sugars into the phloem's sugar-transporting sieve tubes as well as number and size of the sieve tubes are adjusted to growth temperature in the winter annual species *Arabidopsis* and spinach [79–81]. The characterized winter annuals typically germinate in the autumn, overwinter as rosettes, and complete their life cycle in the following spring. These winter annuals exhibited larger and more numerous phloem cells in foliar minor veins when grown under cool temperatures, which was suggested to serve to maintain a high level of sugar

export—and a high photosynthetic capacity—at cool growth temperatures despite an increased viscosity of the phloem sap (and other limitations to sugar transport) under these cool temperatures. The latter concomitant upregulation of photosynthetic capacity and phloem-loading capacity also co-occurs with upregulation of other leaf anatomical features, e.g. increased leaf thickness and, especially, an increase the number of the chloroplast-packed palisade layers of the leaf [82]. Future research should elucidate which signalling networks are involved in this concerted upregulation. Transcription factors, such as CBF/DREB (C-repeat binding factor/dehydration-responsive-element-binding protein), involved in plant adaptation/acclimation to cold stress [83,84], are attractive candidates as key components of such networks.

Furthermore, local varieties (ecotypes) of *Arabidopsis* adapted to the latitudinal extremes (Sweden versus Italy) of this species' geographical range exhibited different sensitivities to environmental cues. Only the Swedish ecotype responded to a decrease in growth temperature (without an increase in growth light intensity) with an increase in the size of the sieve tubes of the leaves' minor veins (figure 3; see also [80,81]). A combination of both low temperature and high light during plant growth was required to elicit a similar response in the Italian ecotype (figure 3; see also [80,81]). These findings suggest that the two ecotypes of this winter annual species do not differ in the principal ability to adjust leaf anatomy to cool growth temperature, but do differ in the *intensity of cues required* to elicit such a response. The greater responsiveness of the Swedish ecotype to cool temperature makes intuitive sense. These responses are consistent with putative genetic differences in the operation of the signalling networks involved.

In addition to differential adaptations to the environment between members (ecotypes) of a given winter annual

species, different strategies are also seen between winter annual and summer annual species. In contrast to winter annuals, summer annuals typically germinate in the spring, grow over the summer and complete their life cycle in the autumn. Figure 4 shows similar photosynthetic capacities in both the winter annual *Arabidopsis* and the summer annual squash grown under warm temperatures and high light. Anatomical adjustments to phloem-loading cells are made by plants using either of the two major mechanisms for active sugar loading into the sugar-transporting phloem (sieve) tubes (see [85]), i.e. apoplastic loading via membrane transporters (as the phloem-loading mode used by *Arabidopsis*) and symplastic loading through plasmodesmatal openings in the cell wall (as the phloem-loading mode used by squash). However, while carbon export in *Arabidopsis* apparently relies on a *few large* veins per area of leaf (large number of sieve tubes per given sugar-loading vein and a low vein density per leaf area), carbon export in squash instead apparently relies on *many small* veins per area of leaf (small number of sieve tubes per given sugar-loading vein and a high vein density per leaf area; figure 4). Future research should address the implications of this difference in number (density) and size of sugar-loading and -exporting leaf veins between winter and summer annuals (for data on additional species, see [79]) for sugar export and phloem loading in source leaves (as well as, possibly, for sugar import and phloem unloading in sink tissues), and e.g. tolerance of cold versus hot temperatures as well as for abiotic versus biotic stress tolerance.

**Acknowledgements.** We thank Onno Muller for the characterization of squash.

**Funding statement.** This work was supported by the National Science Foundation (Award nos. IOS-0841546 and DEB-1022236 to B.D.-A. and W.W.A.) and the University of Colorado at Boulder.

## References

- Krapp A, Stitt M. 1995 An evaluation of direct and indirect mechanisms for the 'sink-regulation' of photosynthesis in spinach: changes in gas exchange, carbohydrates, metabolites, enzyme activities and steady-state transcript levels after cold-girdling source leaves. *Planta* **195**, 313–323. (doi:10.1007/BF00202587)
- Paul MJ, Driscoll SP. 1997 Sugar repression of photosynthesis: the role of carbohydrates in signalling nitrogen deficiency through source:sink imbalance. *Plant Cell Environ.* **20**, 110–116. (doi:10.1046/j.1365-3040.1997.d01-17.x)
- Paul MJ, Foyer CH. 2001 Sink regulation of photosynthesis. *J. Exp. Bot.* **52**, 1383–1400. (doi:10.1093/jexbot/52.360.1383)
- Krapp A, Quick WP, Stitt M. 1991 Ribulose-1,5-bisphosphate carboxylase-oxygenase, other Calvin-cycle enzymes, and chlorophyll decrease when glucose is supplied to mature spinach leaves via the transpiration stream. *Planta* **186**, 58–69. (doi:10.1007/BF00201498)
- Krapp A, Hofmann B, Schäfer C, Stitt M. 1993 Regulation of the expression of *rbcS* and other photosynthetic genes by carbohydrates: a mechanism for the 'sink regulation' of photosynthesis? *Plant J.* **3**, 817–828. (doi:10.1111/j.1365-3113X.1993.00817.x)
- Kilb B, Wietoska H, Godde D. 1996 Changes in the expression of photosynthetic genes precede loss of photosynthetic activities and chlorophyll when glucose is supplied to mature spinach leaves. *Plant Sci.* **115**, 225–235. (doi:10.1016/0168-9452(96)04362-2)
- Myers DA, Thomas RB, DeLucia EH. 1999 Photosynthetic responses of loblolly pine (*Pinus taeda*) needles to experimental reduction in sink demand. *Tree Physiol.* **19**, 235–242. (doi:10.1093/treephys/19.4-5.235)
- Rivas F, Gravina A, Agusti M. 2007 Girdling effects on fruit set and quantum yield efficiency of PSII in two *Citrus* cultivars. *Tree Physiol.* **27**, 527–535. (doi:10.1093/treephys/27.4.527)
- Urban L, Alphonsout L. 2007 Girdling decreases photosynthetic electron fluxes and induces sustained photoprotection in mango leaves. *Tree Physiol.* **27**, 345–352. (doi:10.1093/treephys/27.3.345)
- Mondal MH, Brun WA, Brenner ML. 1978 Effects of sink removal on photosynthesis and senescence in leaves of soybean (*Glycine max* L.) plants. *Plant Physiol.* **61**, 394–397. (doi:10.1104/pp.61.3.394)
- Wünsche JN, Greer DH, Laing WA, Palmer JW. 2005 Physiological and biochemical leaf and tree responses to crop load in apple. *Tree Physiol.* **25**, 1253–1263. (doi:10.1093/treephys/25.10.1253)
- Duan W, Fan PG, Wan LJ, Li WD, Yan ST, Li SH. 2008 Photosynthetic response to low sink demand after fruit removal in relation to photoinhibition and photoprotection in peach trees. *Tree Physiol.* **28**, 123–132. (doi:10.1093/treephys/28.1.123)
- Adams III WW, Muller O, Cohu CM, Demmig-Adams B. 2013 May photoinhibition be a consequence, rather than a cause, of limited plant productivity? *Photosynth. Res.* **117**, 31–44. (doi:10.1007/s11120-013-9849-7)
- Demmig-Adams B, Moeller DL, Logan BA, Adams III WW. 1998 Positive correlation between levels of retained zeaxanthin + antheraxanthin and degree of photoinhibition in shade leaves of *Schefflera arboricola*. *Planta* **205**, 367–374. (doi:10.1007/s004250050332)
- Demmig-Adams B *et al.* 2006 Modulation of PsbS and flexible versus sustained energy dissipation by

- light environment in different species. *Physiol. Plant.* **127**, 670–680. (doi:10.1111/j.1399-3054.2006.00698.x)
16. Mooney HA, Ehleringer J, Björkman O. 1977 The leaf energy balance of leaves of the evergreen desert shrub *Atriplex hymenelytra*. *Oecologia* **29**, 301–310. (doi:10.1007/BF00345804)
  17. Ehleringer JR, Björkman O. 1978 Comparison of photosynthetic characteristics of *Encelia* species possessing glabrous and pubescent leaves. *Plant Physiol.* **62**, 185–190. (doi:10.1104/pp.62.2.185)
  18. McMillen GG, McClendon JH. 1979 Leaf angle: an adaptive feature of sun and shade leaves. *Bot. Gaz.* **140**, 437–442. (doi:10.1086/337110)
  19. Ludlow MM, Björkman O. 1984 Paraheliotropic leaf movement in *Siratro* as a protective mechanism against drought-induced damage to primary photosynthetic reactions: damage by excessive light and heat. *Planta* **161**, 505–518. (doi:10.1007/BF00407082)
  20. Adams III WW, Barker DH. 1998 Seasonal changes in xanthophyll cycle-dependent energy dissipation in *Yucca glauca* Nuttall. *Plant Cell Environ.* **21**, 501–511. (doi:10.1046/j.1365-3040.1998.00283.x)
  21. Verhoeven AS, Demmig-Adams B, Adams III WW. 1997 Enhanced employment of the xanthophyll cycle and thermal energy dissipation in spinach exposed to high light and nitrogen stress. *Plant Physiol.* **113**, 817–824.
  22. Demmig-Adams B, Adams III WW. 2006 Photoprotection in an ecological context: the remarkable complexity of thermal energy dissipation. *New Phytol.* **172**, 11–21. (doi:10.1111/j.1469-8137.2006.01835.x)
  23. Demmig-Adams B, CoHu CM, Muller O, Adams III WW. 2012 Modulation of photosynthetic energy conversion efficiency in nature: from seconds to seasons. *Photosynth. Res.* **113**, 75–88. (doi:10.1007/s11120-012-9761-6)
  24. Adir N, Zer H, Shochat S, Ohad I. 2003 Photoinhibition—a historical perspective. *Photosynth. Res.* **76**, 343–370. (doi:10.1023/A:1024969518145)
  25. Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI. 2007 Photoinhibition of photosystem II under environmental stress. *Biochim. Biophys. Acta* **1767**, 414–421. (doi:10.1016/j.bbabi.2006.11.019)
  26. Murata N, Allakhverdiev SI, Nishiyama Y. 2012 The mechanism of photoinhibition *in vivo*: Re-evaluation of the roles of catalase,  $\alpha$ -tocopherol, non-photochemical quenching, and electron transport. *Biochim. Biophys. Acta* **1817**, 1127–1133. (doi:10.1016/j.bbabi.2012.02.020)
  27. Tyystjärvi E. 2008 Photoinhibition of photosystem II and photodamage of the oxygen evolving manganese cluster. *Coord. Chem. Rev.* **252**, 361–376. (doi:10.1016/j.ccr.2007.08.021)
  28. Tyystjärvi E. 2013 Photoinhibition of photosystem II. *Intl. Rev. Cell Mol. Biol.* **300**, 243–303.
  29. Keren N, Krieger-Liszskay A. 2011 Photoinhibition: molecular mechanisms and physiological significance. *Physiol. Plant.* **142**, 1–5. (doi:10.1111/j.1399-3054.2011.01467.x)
  30. Takahashi S, Badger MR. 2011 Photoprotection in plants: a new light on photosystem II damage. *Trends Plant Sci.* **16**, 53–60. (doi:10.1016/j.tplants.2010.10.001)
  31. Vass I. 2012 Molecular mechanisms of photodamage in the Photosystem II complex. *Biochim. Biophys. Acta* **1817**, 209–217. (doi:10.1016/j.bbabi.2011.04.014)
  32. Farage PR, Long SP. 1991 The occurrence of photoinhibition in an over-wintering crop of oil-seed rape (*Brassica napus* L.) and its correlation with changes in crop growth. *Planta* **185**, 279–286. (doi:10.1007/BF00194071)
  33. Baker NR, Farage PK, Stirling CM, Long SP. 1994 Photoinhibition of crop photosynthesis in the field at low temperatures. In *Photoinhibition of photosynthesis; from molecular mechanisms to the field* (eds NR Baker, JR Bowyer), pp. 349–363. Oxford, UK: Bios Scientific Publishers.
  34. Long SP, Humphries S, Falkowski PG. 1994 Photoinhibition of photosynthesis in nature. *Annu. Rev. Plant. Physiol. Plant Mol. Biol.* **45**, 633–662. (doi:10.1146/annurev.pp.45.060194.003221)
  35. Raven JA. 1994 The cost of photoinhibition to plant communities. In *Photoinhibition of photosynthesis; from molecular mechanisms to the field* (eds NR Baker, JR Bowyer), pp. 449–464. Oxford, UK: Bios Scientific Publishers.
  36. Raven JA. 2011 The cost of photoinhibition. *Physiol. Plant.* **142**, 87–104. (doi:10.1111/j.1399-3054.2011.01465.x)
  37. Losciale P, Chow WS, Grappadelli LC. 2010 Modulating the light environment with the peach ‘asymmetric orchard’: effects on gas exchange performances, photoprotection, and photoinhibition. *J. Exp. Bot.* **61**, 1177–1192. (doi:10.1093/jxb/erp387)
  38. Reynolds M, Foulkes J, Furbank R, Griffiths S, King J, Murchie E, Parry M, Slafer G. 2012 Achieving yield gains in wheat. *Plant Cell Environ.* **35**, 1799–1823. (doi:10.1111/j.1365-3040.2012.02588.x)
  39. Zhu XG, Ort DR, Whitmarsh J, Long SP. 2004 The slow reversibility of photosystem II thermal energy dissipation on transfer from high to low light may cause large losses in carbon gain by crop canopies: a theoretical analysis. *J. Exp. Bot.* **55**, 1167–1175. (doi:10.1093/jxb/erh141)
  40. Zhu X-G, Long SP, Ort DR. 2010 Improving photosynthetic efficiency for greater yield. *Annu. Rev. Plant Biol.* **61**, 235–261. (doi:10.1146/annurev-arplant-042809-112206)
  41. Adams III WW, Watson AM, Mueh KE, Amiard V, Turgeon R, Ebbert V, Logan BA, Combs AF, Demmig-Adams B. 2007 Photosynthetic acclimation in the context of structural constraints to carbon export from leaves. *Photosynth. Res.* **94**, 455–466. (doi:10.1007/s11120-006-9123-3)
  42. Adams III WW, Demmig-Adams B, Verhoeven AS, Barker DH. 1995 ‘Photoinhibition’ during winter stress: involvement of sustained xanthophyll cycle-dependent energy dissipation. *Aust. J. Plant Physiol.* **22**, 261–276. (doi:10.1071/PP9950261)
  43. Adams III WW, Zarter CR, Mueh KE, Amiard V, Demmig-Adams B. 2006 Energy dissipation and photoinhibition: a continuum of photoprotection. In *Photoprotection, photoinhibition, gene regulation, and environment; advances in photosynthesis and respiration*, vol. 21 (eds B Demmig-Adams, WW Adams, AK Mattoo), pp. 49–64. Dordrecht, The Netherlands: Springer.
  44. Ferrar PJ, Osmond CB. 1986 Nitrogen supply as a factor influencing photoinhibition and photosynthetic acclimation after transfer of shade-grown *Solanum dulcamara* to bright light. *Planta* **168**, 563–570. (doi:10.1007/BF00392277)
  45. Dat J, Vandenabeele S, Vranová E, Van Montagu M, Inzé D, Van Breusegem F. 2000 Dual action of the active oxygen species during plant stress responses. *Cell. Mol. Life Sci.* **57**, 779–795. (doi:10.1007/s000180050041)
  46. Nanda AK, Andrio E, Marino D, Pauly N, Dunand C. 2010 Reactive oxygen species during plant–microorganism early interactions. *J. Integr. Plant Biol.* **52**, 195–204. (doi:10.1111/j.1744-7909.2010.00933.x)
  47. Ledford HK, Niyogi KK. 2005 Singlet oxygen and photo-oxidative stress management in plants and algae. *Plant Cell Environ.* **28**, 1037–1045. (doi:10.1111/j.1365-3040.2005.01374.x)
  48. Munné-Bosch S, Queval G, Foyer CH. 2013 The impact of global change factors on redox signaling underpinning stress tolerance. *Plant Physiol.* **161**, 5–19. (doi:10.1104/pp.112.205690)
  49. Baginsky S, Link G. 2006 Redox regulation of chloroplast gene expression. In *Photoprotection, photoinhibition, gene regulation, and environment. Advances in photosynthesis and respiration*, vol. 21 (eds B Demmig-Adams, WW Adams, AK Mattoo), pp. 269–287. Dordrecht, The Netherlands: Springer.
  50. Dietz K-J, Stork T, Finkemeier I, Lamkemeyer P, Li W-X, El-Tayeb MA, Michel K-P, Pistorius E, Baier M. 2006 The role of peroxiredoxins in oxygenic photosynthesis of cyanobacteria and higher plants: Peroxide detoxification or redox sensing? In *Photoprotection, photoinhibition, gene regulation, and environment. Advances in photosynthesis and respiration*, vol. 21 (eds B Demmig-Adams, WW Adams, AK Mattoo), pp. 303–319. Dordrecht, The Netherlands: Springer.
  51. Foyer CH, Trebst A, Noctor G. 2006 Signaling and integration of defense functions of tocopherol, ascorbate and glutathione. In *Photoprotection, photoinhibition, gene regulation, and environment. Advances in photosynthesis and respiration*, vol. 21 (eds B Demmig-Adams, WW Adams, AK Mattoo), pp. 241–268. Dordrecht, The Netherlands: Springer.
  52. Mullineaux PM, Karpinski S, Creissen GP. 2006 Integration of signaling in antioxidant defenses. In *Photoprotection, photoinhibition, gene regulation, and environment. Advances in photosynthesis and respiration*, vol. 21 (eds B Demmig-Adams, WW Adams, AK Mattoo), pp. 223–239. Dordrecht, The Netherlands: Springer.
  53. Pérez AG, Sanz C, Richardson DG, Ollás JM. 1993 Methyl jasmonate vapor promotes  $\beta$ -carotene synthesis and chlorophyll degradation in Golden Delicious apple peel. *J. Plant Growth Regul.* **12**, 163–167. (doi:10.1007/BF00189648)

54. Sandorf I, Holländer-Czytko H. 2002 Jasmonate is involved in the induction of tyrosine aminotransferase and tocopherol biosynthesis in *Arabidopsis thaliana*. *Planta* **216**, 173–179. (doi:10.1007/s00425-002-0888-0)
55. Wolucka BA, Goossens A, Inzé D. 2005 Methyl jasmonate stimulates the *de novo* biosynthesis of vitamin C in plant cell suspensions. *J. Exp. Bot.* **56**, 2527–2538. (doi:10.1093/jxb/eri246)
56. Demmig-Adams B, Cochu CM, Amiard V, van Zadelhoff G, Veldink GA, Muller O, Adams III WW. 2013 Emerging trade-offs—impact of photoprotectants (PsbS, xanthophylls, and vitamin E) on oxylipins and biotic defense. *New Phytol.* **197**, 720–729. (doi:10.1111/nph.12100)
57. Maeda H, Song W, Sage TL, DellaPenna D. 2006 Tocopherols play a crucial role in low-temperature adaptation and phloem loading in *Arabidopsis*. *Plant Cell* **18**, 2710–2732. (doi:10.1105/tpc.105.039404)
58. Amiard V, Demmig-Adams B, Mueh KE, Turgeon R, Combs AF, Adams III WW. 2007 Role of light and jasmonic signaling in regulating foliar phloem cell wall ingrowth development. *New Phytol.* **173**, 722–731. (doi:10.1111/j.1469-8137.2006.01954.x)
59. Frenkel M *et al.* 2009 Improper excess light energy dissipation in *Arabidopsis* results in a metabolic reprogramming. *BMC Plant Biol.* **9**, 12. (doi:10.1186/1471-2229-9-12)
60. Munné-Bosch S, Weiler EW, Alegre L, Muller M, Duchting P, Falk J. 2007  $\alpha$ -Tocopherol may influence cellular signaling by modulating jasmonic acid levels in plants. *Planta* **225**, 681–691. (doi:10.1007/s00425-006-0375-0)
61. Semchuk NM, Lushchak OV, Falk J, Krupinska K, Lushchak VI. 2009 Inactivation of genes, encoding tocopherol biosynthetic pathway enzymes, results in oxidative stress in outdoor grown *Arabidopsis thaliana*. *Plant Physiol. Biochem.* **47**, 384–390. (doi:10.1016/j.plaphy.2009.01.009)
62. Havaux M, Eymery F, Profirova S, Rey P, Dorman P. 2005 Vitamin E protects against photoinhibition and photooxidative stress in *Arabidopsis thaliana*. *Plant Cell* **17**, 3451–3469. (doi:10.1105/tpc.105.037036)
63. Sattler SE, Mene-Saffrane L, Farmer EE, Kruschke M, Mueller MJ, DellaPenna D. 2006 Nonenzymatic lipid peroxidation reprograms gene expression and activates defense markers in *Arabidopsis* tocopherol-deficient mutants. *Plant Cell* **18**, 3706–3720. (doi:10.1105/tpc.106.044065)
64. Jänkänpää HJ *et al.* 2013 Non-photochemical quenching capacity in *Arabidopsis thaliana* affects herbivore behaviour. *PLoS ONE* **8**, e53232. (doi:10.1371/journal.pone.0053232)
65. Rohwer CL, Erwin JE. 2008 Horticultural application of jasmonates: a review. *J. Hort. Sci.* **83**, 283–304.
66. Gilbertson RL, Lucas WJ. 1996 How do viruses traffic on the 'Vascular highway'? *Trends Plant Sci.* **1**, 260–268. (doi:10.1016/1360-1385(96)10029-7)
67. Waignan E, Ueki S, Trutnyeva K, Citovsky V. 2004 The ins and outs of nondestructive cell-to-cell systemic movement of plant viruses. *Crit. Rev. Plant Sci.* **23**, 195–250. (doi:10.1080/07352680490452807)
68. Scholthof HB. 2005 Plant virus transport: motions of functional equivalence. *Trends Plant Sci.* **10**, 376–382. (doi:10.1016/j.tplants.2005.07.002)
69. Sattler SE, Cajon EB, Coughlin SJ, DellaPenna D. 2003 Characterization of tocopherol cyclases from higher plants and cyanobacteria: evolutionary implications for tocopherol synthesis and function. *Plant Physiol.* **132**, 2184–2195. (doi:10.1104/pp.103.024257)
70. Sattler SE, Gilliland LU, Magallanes-Lundback M, Pollard M, DellaPenna D. 2004 Vitamin E is essential for seed longevity and for preventing lipid peroxidation during germination. *Plant Cell* **16**, 1419–1432. (doi:10.1105/tpc.021360)
71. Hofius D, Hajirezaei MR, Geiger M, Tschiersch H, Melzer M, Sonnewald U. 2004 RNAi-mediated tocopherol deficiency impairs photoassimilate export in transgenic potato plants. *Plant Physiol.* **135**, 1256–1268. (doi:10.1104/pp.104.043927)
72. Maeda H, Sage TL, Isaac G, Welti R, DellaPenna D. 2008 Tocopherols modulate extraplastidic polyunsaturated fatty acid metabolism in *Arabidopsis* at low temperature. *Plant Cell* **20**, 452–470. (doi:10.1105/tpc.107.054718)
73. Sasidharan R, Voesenek LACJ, Pierik R. 2011 Cell wall modifying proteins mediate plant acclimatization to biotic and abiotic stresses. *Crit. Rev. Plant Sci.* **30**, 548–562. (doi:10.1080/07352689.2011.615706)
74. Atkinson NJ, Urwin PE. 2012 The interaction of plant biotic and abiotic stresses: from genes to the field. *J. Exp. Bot.* **63**, 3523–3543. (doi:10.1093/jxb/ers100)
75. Bari R, Jones JGD. 2008 Role of plant hormones in plant defence responses. *Plant Mol. Biol.* **69**, 473–488. (doi:10.1007/s11103-008-9435-0)
76. Kim TH. 2012 Plant stress surveillance monitored by ABA and disease signaling interactions. *Mol. Cells* **33**, 1–7. (doi:10.1007/s10059-012-2299-9)
77. Derksen H, Rampitsch C, Daayf F. 2013 Signaling cross-talk in plant disease resistance. *Plant Sci.* **207**, 79–87. (doi:10.1016/j.plantsci.2013.03.004)
78. Maharijaya A, Vosman B, Verstappen F, Steenhuis-Broers G, Mumm R, Purwito A, Visser RGF, Voorrips RE. 2012 Resistance factors in pepper inhibit larval development of thrips (*Frankliniella occidentalis*). *Entomol. Exp. Appl.* **145**, 62–71. (doi:10.1111/j.1570-7458.2012.01304.x)
79. Adams III WW, Cochu CM, Muller O, Demmig-Adams B. 2013 Foliar phloem infrastructure in support of photosynthesis. *Front. Plant Sci.* **4**, 194. (doi:10.3389/fpls.2013.00194)
80. Cochu CM, Muller O, Demmig-Adams B, Adams III WW. 2013 Minor loading vein acclimation for three *Arabidopsis thaliana* ecotypes in response to growth under different temperature and light regimes. *Front. Plant Sci.* **4**, 240. (doi:10.3389/fpls.2013.00240)
81. Cochu CM, Muller O, Stewart JJ, Demmig-Adams B, Adams III WW. 2013 Association between minor loading vein architecture and light- and CO<sub>2</sub>-saturated rates of photosynthetic oxygen evolution among *Arabidopsis thaliana* ecotypes from different latitudes. *Front. Plant Sci.* **4**, 264. (doi:10.3389/fpls.2013.00264)
82. Dumlaoui MR, Darehshouri A, Cochu CM, Muller O, Mathias J, Adams III WW, Demmig-Adams B. 2012 Low temperature acclimation of photosynthetic capacity and leaf morphology in the context of phloem loading type. *Photosynth. Res.* **113**, 181–189. (doi:10.1007/s11120-012-9762-5)
83. Thomashow MF. 2010 Molecular basis of plant cold acclimation: insights gained from studying the CBF cold response pathway. *Plant Phys.* **154**, 571–577. (doi:10.1104/pp.110.161794)
84. Akhtar M, Jaiswal A, Taj G, Jaiswal JP, Qureshi MI, Singh NK. 2012 DREB1/CBF transcription factors: their structure, function and role in abiotic stress tolerance in plants. *J. Genet.* **91**, 385–395. (doi:10.1007/s12041-012-0201-3)
85. Muller O, Cochu CM, Stewart JJ, Protheroe JA, Demmig-Adams B, Adams III WW. In press. Association between photosynthesis and contrasting features of minor veins in leaves of summer annuals loading phloem via symplastic versus apoplastic routes. *Physiologia Plantarum*. (doi:10.1111/ppl.12155).