

Factors influencing physical activity in older adults

Chae-Hee Park¹, Steriani Elavsky², Kyo-Man Koo^{3*}

¹Department of Sport and Healthy Aging, Korea National Sport University, Seoul, Korea

²Department of Kinesiology, Penn State University, Pennsylvania, USA

³Department of Adapted Physical Education, Baekseok University, Cheonan, Korea

The purpose of this study was to investigate the extent to which Korean seniors report employing different motivational and social cognitive strategies related to physical activity, and to evaluate which motivational and social cognitive strategies were related to physical activity, and which motivational and social cognitive strategies differentiate between high active and low-active Korean seniors. Community-dwelling older adults (N = 187) participated in the study and completed questionnaires assessing self-reported physical activity and a range of motivational and social cognitive variables. The results showed that physical activity was predicted by quality goal-setting practices, self-efficacy, social support from family and physical activity self-regulation subscales of

social support and exercise planning and scheduling. Between the groups of highly physically active and low-active participants, we observed differences in satisfaction with life, self-efficacy, quality goal-setting practices, and self-regulatory strategies related to self-monitoring, goal-setting, social support and time management. In conclusion, these findings indicate that physical activity promotion efforts among Korean older adults should focus on enhancing self-efficacy, social support, and self-regulation skills.

Keywords: Korean older adults, Physical activity, Social cognitive factors

INTRODUCTION

By 2026, people aged 65 and over will account for 20 percent of the South Korean population. These numbers indicate that South Korea (hereafter Korea) will become the most aged society among advanced economies in 2050, with nearly four out of every 10 Koreans being aged 65 or over due to low birthrates and the rapidly aging population. Korea also boasts life expectancy (LE) one year higher than the OECD average of 80 yr, raising the important question of whether people are spending these extra years of life in good health and with good quality of life. To this end, the concept of healthy life expectancy (HALE) has been proposed, estimating the equivalent years in full health that a person can expect to live on the basis of the current mortality rates and prevalence distribution of health states in the population (OECD, 2009). HALE at birth in Korea is 71 yr (female -74 yr and male -68 yr). Difference years between LE and HALE are about 10 yr.

Sufficient and regular physical activity is one of the most widely

recommended health promotion strategies for managing chronic illnesses and is known to have various health benefits (Braith and Stewart, 2006; Laaksonen et al., 2005; Lam et al., 2004; Yeom et al., 2011). Over the past 30 yr, an extensive body of evidence has accumulated regarding the benefits that accrue to older adults who participate in regular physical activity. The important role of physical activity in promoting functional health, delaying or preventing non-communicable disease such as osteoporosis, coronary artery disease, non-insulin-dependent diabetes mellitus and disability, and reducing mortality has been established throughout years (Christ and Ross, 2010; Ferrucci et al., 1999; Hubert et al., 2002; Jonker et al., 2006; Leveille et al., 1999). In addition, physical activities decrease the risk of falling, improve sleep, enhance mood and general well-being, and improve blood pressure and decrease relative abdominal fat (Alessi et al., 1999; Resnick, 2001).

In spite of the established benefits of physical activity, physical activity participation remains insufficient. Approximately 28-34% of adults aged 65-74 yr and 35-44% of seniors aged 75 yr or older

*Corresponding author: Kyo-Man Koo

Department of Adapted Physical Education, Baekseok University, 76 Munam-ro, Dongnam-gu, Cheonan 330-704, Korea

Tel: +82-41-550-2819, Fax: +82-2-410-1877, E-mail: nicekm@bu.ac.kr

Received: February 8, 2014/ Revised: February 15, 2014/ Accepted: February 17, 2014

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

spend no time engaging in physical activity and inactivity is more common in women than men in the United States (USDHHS, 2006). Compared to the US, participation in regular physical activity is even less common in seniors in Korea. Approximately 80% of Korean older adults aged 60-70 yr do not engaged in moderate physical activity and 90% of older adults over 71 yr of age are inactive (Korea Ministry of Health and Welfare, 2007).

Participation in physical activity in older adults is influenced by a number of variables including demographic factors such as gender, education, and marital status. For example, physical activity participation is lower among older females (Janke et al., 2006; Weiss et al., 2007) and less educated older seniors (Droomerset al., 2001; Janke et al., 2006; Weiss et al., 2007). Interestingly, active men are more likely to have an active spouse and some studies suggest that higher levels of physical activity for older married persons have been observed (Janke et al., 2006; Pettee et al., 2006).

Additionally, choices of older adults to be regularly physically active are influenced by social support from family members or friends, availability of facilities for exercise and/or recreational activities, personal determinants especially one's motivation, self-efficacy (i.e., a belief a person has in his or her capacity to perform a course of action), and self-regulation skills (e.g., feasible goal-setting, regular tracking of physical activity) (e.g., McAuley, et al., 2007; King and King, 2010).

In the Korean context, few studies explored the reasons why older adults engage in physical activity in Korea. Studies on physical activity with seniors have mostly focused on programs, mental health including depression and anxiety, and demographic factors but do not broadly examine social cognitive factors related to older adults' physical activity (Cha, 2009; Lee et al., 2011; You and Won, 2010). In this study, motivational and social cognitive constructs related to Korean seniors' participation in physical activity were examined. The main objectives were to examine 1) the extent to which Korean seniors report employing different motivational and social cognitive strategies; 2) which motivational and social cognitive strategies were related to physical activity; and 3) which motivational and social cognitive strategies differentiate between high- active and low-active Korean seniors.

MATERIALS AND METHODS

Subjects

Community-dwelling older adults (N = 199) partook in a survey assessing self-reported physical activity and a range of motivational and social cognitive variables. One hundred eighty seven older

adults provided sufficient data (Range_{age} = 57-96 yr; Mean_{age} = 71.62 ± 5.894 yr) to be included in the analysis after listwise deletion for missing and unreliable values. The respondents lived in a metropolitan area and a medium sized city. Researchers contacted the participants through institutions frequented by older adults such as senior centers offering various social and educational activities for seniors. Most of the participants were women (70.1%), married (71.1%), had above average education (31.6% finished high school, 30% finished undergraduate and graduate school), and more than half of the participants (52.4%) reported that they did not have any healthy problems.

Methods

Several questionnaires were used to acquire self-reported estimates of physical activity and assess the participants' motivation to physical activity represented by a mixture of self-regulatory constructs, perceived self-efficacy and perceived social support. The Lifestyle Physical Activity Self-Efficacy Scale (LSE) (Elavsky, McAuley, 2007) and Barriers Self-Efficacy Scale (BASE) (McAuley, 1993) were used to assess the level of confidence that one can perform sufficient physical activity as part of one's lifestyle during the following six months and in the face of barriers, respectively. The Exercise Planning and Scheduling Scale (EPS) and Exercise Goal-Setting Scale (EGS) (Rovniak et al., 2002) were used to assess strategies such as planning and goal setting. The Physical activity Self-Regulation (PASR) measured a range of motivational constructs related to self-regulation (Umstatted et al., 2009) and the Social Support for Exercise (SSE) (Sallis et al., 1987) scale was used to capture perceived social support in physical activity from friends (including acquaintances and co-workers) and family (referring to anyone living in the household). Physical Activity Survey for the Elderly (PASE) (Washburn et al., 1993) was used to measure physical activity in older adults. The 12-Item Short-Form Health Survey (SF-12) was used to assess physical and mental health status (Ware et al., 1996) and the Satisfaction with Life Scale (SWLS) measured global quality of life (i.e., satisfaction with the respondent's life as a whole; Diener et al., 1985).

All used questionnaires were translated from English for the purpose of the study and supplemented by back-translation to ensure the accuracy of the translation; they also showed acceptable internal consistency (Cronbach's alpha = 0.61-0.97) in the study. Most scales were 5 point Likert scales with 1 representing "never" and 5 representing "very often." The self efficacy scales (LSE and BASE) were 100 points scales with responses indicated in 10-point increments.

Analysis

All data obtained from the measures were processed by the SPSS (ver. 21.0) statistical software. To address goal 1, descriptive statistics were generated to describe the relative use of a variety of motivational strategies in the participants. To address goal 2, bivariate correlations were computed between the motivational constructs and self-reported measures of physical activity and conducted multiple regression analysis was used to evaluate unique contribution of each factor to explained variance in physical activity. To address goal 3, we compared motivation strategies implemented by the physically high active and physically low-active groups using independent two-sample t-test. We used 150min of moderate and/or 75 min of vigorous activity per week to divide the sample into highly physically active/low-active. That is, if a person did 150 min of moderate PA or more they would be considered highly active. If they did 75 min of vigorous they would be considered active (even if they did not report 150 of moderate).

RESULTS

As can be seen in Table 1, older adults in this study strongly believed that they would be able to participate in regular physical activity (5 or more days per week for at least 30+ minutes of accumulated activities per day in the future) during the next six months (for an average LSE score of 74%). Additionally, they did believe

Table 1. Motivational strategies for physical activity (Descriptive statistics)

Motivation for physical activity (self-efficacy scales in percent, all other on Likert-type scales ranging from 1 [never] to 5 [very often])	Physical activity and social cognitive variables-Spearman correlations		
	Mean	Median	St. deviation
Lifestyle physical activity self-efficacy (LSE)	74.5968	80.0000	23.02621
Exercise as displaced priority (EPS)	3.6952	3.7500	0.75402
Exercise planning and scheduling (EPS)	3.3717	3.5000	0.88078
Quality goal-setting practices (EGS)	3.1881	3.1667	0.83652
Barriers self-efficacy (BASE)	73.4329	78.4615	22.27381
Self-monitoring (PASR)	3.4465	3.5000	0.96711
Goal-setting (PASR)	3.3690	3.5000	0.96174
Social support (PASR)	2.7513	3.0000	1.16671
Reinforcement (PASR)	3.7701	4.0000	0.91761
Time management (PASR)	3.5481	3.5000	0.97569
Relapse prevention (PASR)	3.2406	3.5000	1.08551
Social support for exercise (SSE) - family	2.5705	2.3571	0.93407
Social support for exercise (SSE) - friends	2.6914	2.5714	0.99302
Social support (SSE) - overall	2.6401	2.5357	0.77727
Physical health Score	38.6390	38.5693	6.21324
Mental health Score	53.7999	56.0904	10.57845
Satisfaction with Life	5.0075	5.2000	1.24837

that they would continue sufficient physical activity when they encountered diverse barriers (for an average BASE score of 73%).

Participants mostly reported that they did set exercise goals (they scored high on the PASR subscale “Goal-setting”) and it was consistent with the quality of the goal setting practices such as setting both short term and long term exercise goals or analyzing the progress towards goals on the EGS scale. According to the scores on PASR “Social support” subscale, respondents in the study only rarely sought social support. They also perceived receiving low levels of support from family and friends, scoring on average 2.6 on the 5-point SSE scales.

As reflected in PASR “Reinforcement” subscale, the most prevalent motivational strategy in our samples was reinforcement seeking, such as focusing on positive emotions or health benefits of exercise. Also commonly used strategies were time management for example, reserving specific times for physical activity) and self-monitoring (i.e. participants focused on things that helped them to be active).

The total PASE scores were moderately correlated with a number of the motivational and social cognitive variables (Table 2), except for scores on the exercise displayed priority, exercise planning and scheduling, reinforcement, social support from family and friends, physical health and mental health scales.

Table 2. Motivational and social cognitive influences on physical activity

Physical activity and social cognitive variables-Spearman correlations	
	PASE
Lifestyle physical activity self-efficacy	0.305***
Exercise as Displaced Priority	0.097
Exercise Planning and Scheduling	0.070
Quality Goal-Setting Practices	0.315***
Barriers Self-Efficacy	0.227**
Self-monitoring PASR	0.212**
Goal-setting PASR	0.182*
Social support PASR	0.245**
Reinforcement PASR	0.123
Time management PASR	0.206**
Relapse prevention PASR	0.210**
Social Support for Exercise - FAMILY	-0.045
Social Support for Exercise - FRIENDS	0.131
Social Support Overall	0.049
Physical Health Score	0.073
Mental Health Score	0.007
Satisfaction with Life	0.163*

Spearman correlations.
*Result significant on 0.05 level; **Result significant on 0.01 level; ***Result significant on 0.001 level.

Table 3. Motivational and social cognitive influences on physical activity

Physical activity and social cognitive variables-Multiple regression		
Model	PASE	
	Beta	t
Quality goal-setting practices	0.321	3.619***
Lifestyle physical activity self-efficacy	0.254	3.595***
Social support for exercise - FAMILY	-0.220	-3.107**
Social support (PASR)	0.197	2.686**
Exercise planning and scheduling	-0.187	-2.335*
	MR=0.467	R ² =0.218

Multiple Regression.

*Result significant on 0.05 level; **Result significant on 0.01 level; ***Result significant on 0.001 level.

Next, we conducted multiple regression analysis regressing separately each of the physical activity scores on the motivational and social cognitive variables. The regression coefficients and unique contributions of all variables to variance in physical activity are presented in Table 3. As can be seen, physical activity was predicted by Quality Goal-Setting Practices ($\beta = 0.321, P < 0.001$), Lifestyle physical activity self-efficacy ($\beta = 0.254, P < 0.001$), Social support exercise-family ($\beta = -0.220, P < 0.01$), and physical activity self-regulation (PASR) subscales of Social support ($\beta = 0.197, P < 0.01$) and Exercise Planning and Scheduling ($\beta = -0.187, P < 0.05$). Overall, the motivational and social cognitive variables predicted significantly PASE score ($F = 10.075, P = 0.000$; explaining 21.8% of variance in PASE).

Table 4. Comparison of physically high-active and low-active older adults

Comparison of physically active and low-active seniors					
Low-active/High-active		N	Mean	St. deviation	t
Lifestyle physical activity self-efficacy	Low-active	94	69.7404	24.03479	-2.959**
	High-active	93	79.5054	20.96510	
Exercise as displaced priority	Low-active	94	3.6011	0.77799	-1.725
	High-active	93	3.7903	0.72071	
Exercise planning and scheduling	Low-active	94	3.2952	0.88822	-1.195
	High-active	93	3.4489	0.87116	
Quality goal-setting practices	Low-active	94	2.9638	0.76874	-3.817***
	High-active	93	3.4147	0.84503	
Barriers self-efficacy	Low-active	94	70.2823	22.45376	-1.959
	High-active	93	76.6170	21.74830	
Self-monitoring PASR	Low-active	94	3.2819	0.98253	-2.369*
	High-active	93	3.6129	0.92704	
Goal-setting PASR	Low-active	94	3.1915	0.96750	-2.575*
	High-active	93	3.5484	0.92685	
Social support PASR	Low-active	94	2.5160	1.22134	-2.827**
	High-active	93	2.9892	1.06316	
Reinforcement PASR	Low-active	94	3.6915	1.01892	-1.180*
	High-active	93	3.8495	0.80008	
Time management PASR	Low-active	94	3.3936	1.02881	-2.200*
	High-active	93	3.7043	0.89757	
Relapse prevention PASR	Low-active	94	3.1064	1.19333	-1.711
	High-active	93	3.3763	0.95170	
Social support for exercise - Family	Low-active	94	2.5893	0.91201	0.277
	High-active	93	2.5514	0.96042	
Social support for exercise - Friends	Low-active	94	2.5540	1.00736	-1.916
	High-active	93	2.8303	0.96385	
Social support overall	Low-active	94	2.5899	0.82018	-0.888
	High-active	93	2.6908	0.73229	
Physical health score	Low-active	94	38.3932	6.35295	-0.543
	High-active	93	38.8875	6.09304	
Mental health score	Low-active	94	53.4390	10.78296	-0.468
	High-active	93	54.1643	10.41329	
Satisfaction with life	Low-active	94	4.7872	1.23247	-2.458*
	High-active	93	5.2301	1.23110	

Independent two sample t-test.

*Result significant on 0.05 level; **Result significant on 0.01 level; ***Result significant on 0.001 level.

Between the groups of highly physically active and low-active participants (Table 4), we observed differences in Lifestyle physical activity self-efficacy ($\tau = -2.959$, $P < 0.01$), Quality Goal-Setting Practices ($\tau = -3.817$, $P < 0.001$), PASR subscales of Self-monitoring ($\tau = -2.369$, $P < 0.05$), Goal-setting ($\tau = -2.575$, $P < 0.05$), Social support ($\tau = -2.827$, $P < 0.01$), Reinforcement ($\tau = -1.180$, $P < 0.05$), Time management ($\tau = -2.200$, $P < 0.05$), Satisfaction of life ($\tau = -2.485$, $P < 0.05$). Highly active older adults had significantly stronger beliefs that they would be able to maintain sufficient physical activity during the next six months. In addition, highly active participants did set exercise goals, monitored their physical activity more and participated more in physical activity when they received advises from exercise and/or health professionals. On the other hand, the high-active and low-active did not significantly differ in their scores on the EPS subscale of giving priority to exercise and planning and scheduling for exercise, BASE, PASR subscales of relapse prevention, and SSE. In both groups, barriers self efficacy scores were rated quite highly meaning that they would be able to participate in physical activity when they face various environmental, social and motivational obstacles. It possibly indicated that both high-active and low-active participants recognized the importance of participating in physical activity. In contrast, both group scored low on the social support especially from family indicating that the participants' approach to physical activity was predominantly individualistic, although the high-active group perceived more support from friends rather than family.

DISCUSSION

Despite the increasing need for physical activity promotion efforts targeted at seniors, very little is known about motivations of Korean older adults to participate in physical activity. The aim of this study was to examine which motivational and social cognitive strategies were used by Korean older adults, which were related to their physical activity; and which differentiated between high and low active older adults.

The results showed that lifestyle physical activity self-efficacy and quality goal-setting practices were important motivational factors related to physical activity in our participants and consistent with other literature supporting the importance of these constructs. Self-efficacy is an important determinant of physical activity participation in older adults (Choi, 2004) and has predicted the maintenance of physical activity in seniors for up to 5 yr (McAuley et al., 2003, 2011). Additionally, Elavsky (2005) showed that self-efficacy often mediates that relationship between physical activity

and satisfaction with life or measures of health-related quality of life in seniors. It indicates that self-efficacy plays a significant role between physical activity and some of its outcomes.

Interestingly, the participants in the study did not seem to consider physical activity as the first priority in their everyday lives. However, older adults in our study claimed that they set their own exercise goals and tried to achieve them. We found that a significant difference between high- and low-active group in this variable. This suggest that highly active Korean older adults seem to employ self-selected goal setting strategies for example setting multiple goals, monitoring the progress toward goals, setting short- and long term goals, and analyzing their goals. Using self-selected goals rather than assigned ones is associated with greater dedication and often is more valued by the individual (Hall et al., 2010; Locke and Latham, 2002).

In general we observed a weak relationship between self-regulation strategies except social support and physical activity. On the other hand, we found significant differences between high- and low-active group in some of the self-regulation strategies including self-monitoring, goal-setting, social support, reinforcement, and time management. It is possible those self-regulation strategies may be indirectly associated with physical activity or that Korean older adults participate more in lifestyle (habitual) physical activity as opposed to planful or structured physical activity that may benefit from self-regulation more directly. Mudrak et al. (2012) observed a similar set of relationship in Czech older adults. Whereas there was a weak relationship between physical activity and self-regulation strategies, self-monitoring and relapse prevention significantly differentiated between active and inactive older adults. These results suggest that the effectiveness of motivational strategies such as self-regulation may vary depending on the position that physical activity occupies in older adults' lives, which is partly culturally dependent. In Korea, not many studies have conducted the relationship between the self-regulatory skills such as realistic goal-setting, self-monitoring and physical activity among older adults even though those strategies are an important influence of behavior changes. Future researches should examine how the self-regulatory strategies play a role on behavior changes related to physical activity in the older adult population.

A number of studies demonstrated that the importance of social support to exercise behavior for older adults. Orsega-Smith et al. (2007) suggested that social support is an active and cost-effective approach to increase physical activity, and can be offered at an individual level by family, friends, or others who provide encouragement to strengthen an individual's motives to be physically active.

Indeed, social support has been shown to be an important predictor of exercise adherence among older adults (Oka et al., 1995) and to be among the most influential forces for older women to participate in active types of activities (O'Brien Cousins, 1995). In this study, the most useful form of social support appeared to stem from others such as participating in a program with friends, receiving advices from health professionals and getting demonstrations from exercise experts as opposed to from family members. The high-active group reported more support from friends and others as compared to family, a finding that is consistent with other studies suggesting that social support provided by friends rather than family domain of perceived physical ability was significantly related to leisure time physical activity (Orsega-Smith et al., 2007). More studies are needed to explore the most useful sources of social support in older adults to determine the most effective ways in which social support could be enhanced in this population.

The ambiguous role of social support from family may be indirectly related to other social phenomena such as marital status. Crespo et al. (2000) found that currently or formerly married men were more likely to be more physically active than never-married men, whereas physical activity participation did not seem to vary by marital status for women. In this study most of the participants were women (70.1%) and married (71.1%). It seems that Korean older adults, women especially, may not be receiving enough support regarding physical activity from anyone living in the household. This effect may be more pronounced for individuals who experience a divorce in later life. A study (Kim, 2009) indicated that total divorce rate has decreased in Korea but a divorce rate among older adults 65 yr and older has been increasing steadily posing additional challenges for older adults' social support.

Further studies should be systematically conducted about different types of social support in influencing physical activity behaviors and which resources are important elements of promoting physical activity for older adults. And a variety of types of social support can be created or enhanced via social network and policy interventions to promote physical activity for seniors. Even though, self-report measures of physical activity characterize a convenient and realistic approach, and all measures used in the study have been validated for use in seniors, it may be influenced by recall bias and social desirability and differ from other objective measures for example pedometers or accelerometers. The results from this study provide a more powerful and sustainable influence on physical activity among older adults.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

The study was supported by 2012 faculty research fund by the Korea National Sport University.

REFERENCES

- Alessi CA, Yoon EJ, Schnelle JF, Al-Samarrai NR, Cruise PA. A randomized trial of a combined physical activity and environmental intervention in nursing home residents: Do sleep and agitation improve? *J Am Geriatr Soc* 1999;47:784-791.
- Braith RW, Stewart K J. Resistance exercise training: its role in the prevention of cardiovascular disease. *Circulation* 2006;113:2642-2650.
- Cha JW. Meta-regression analysis of variables related to effects of exercise program applied to the elderly. *J Korean Physic Educ Girls Women* 2009;23(3):203-220.
- Choi JH, Lee GM, Kim HJ, Seo JW. The contributions of gender and physical activity levels on physical function, psychological function, and health-related quality of life in the elderly. *Korean J Physic Educ* 2004; 43(6):975-983.
- Christ H, Ross A. Correlates of physical activity participation in community-dwelling older adults. *J Aging Phys Act* 2010;18:375-389.
- Crespo CJ, Smit E, Andersen RE, Carter-Pokras O, Ainsworth BE. Race/ethnicity, social class and their relation to physical inactivity during leisure time: Results from the third national health and nutrition examination survey, 1988-1994. *Am J Prev Med* 2000;18(1):46-53.
- Diener E, Emmons RA, Larsen RJ, Griffin S. The Satisfaction with Life Scale. *J Pers Assess* 1985;49:71-75.
- Droomers M, Schrijvers CTM, Mackenbach JP. Educational level and decreases in leisure time physical activity: Predictors from the longitudinal GLOBE study. *J Epidemiol Community Health* 2001;55:562-568.
- Elavsky S. Physical activity enhances long-term quality of life in older adults: efficacy, esteem, and affective influences. *Ann Behav Med* 2005; 30(2):138-145.
- Elavasky S, McAuley E. Physical activity and mental health outcomes during menopause: A randomized controlled trial. *Ann Behav Med* 2007;33(2): 132-142.
- Ferrucci L, Izmirlian G, Leveille S, Phillips, CL, Corti MC, Brock DB, Guralnik JM. Smoking, physical activity, and active life expectancy. *Am J Epidemiol* 1999;149:645-653.

- Hall KS, Crowley GM, Bosworth HB, Howard TA, Morey MC. Individual progress toward self-selected goals among older adults enrolled in physical activity counseling intervention. *J Aging Phys Act* 2010;18:439-450.
- Hubert HB, Bloch DA, Oehlert JW, Fries JF. Lifestyle habits and compression of morbidity. *J Gerontol A Biol Sci Med Sci* 2002;57A(6):M347-M351.
- Janke M, Davey A, Kleiber D. Modeling change in older adults' leisure activities. *Leisure Sci* 2006;28:285-303.
- Jonker JT, De Laet C, Franco OH, Peeters A, Mackenbach J, Nusselder WJ. Physical activity and life expectancy with and without diabetes. *Diabetes Care* 2006;29(1):38-43.
- Kim SJ. A life history study on the elderly women who have divorced. *J Korean Gerontol Soc* 2009;29(3):1087-1105.
- King AC, King DK. Physical activity for an aging population. *Public Health Rev* 2010;32(2):401-426.
- Korea Ministry of Health and Welfare. Guidelines for healthy life projects. Retrieved April 1, 2011, from http://mchp.hp.go.kr/hpMchp/board.dia?method=downFile&FI_FID=884, 2007.
- Laaksonen DE, Lindstrom J, Lakka TA, Eriksson JG, Niskanen L, Wikström K. Physical activity in the prevention of type 2 diabetes: the Finnish diabetes prevention study. *Diabetes* 2005;54:158-165.
- Lam TH, Ho SY, Hedley AJ, Mak KH, Leung GM. Leisure time physical activity and mortality in Hong Kong: case-control study of all adult deaths in 1998. *Ann Epidemiol* 2004;14:391-398.
- Lee CW, Lee DH, Yoon JH. The effects of circuit exercise programs on body composition, blood lipid and liver function variables in elderly people with obesity. *J Sport Leisure Stud* 2011;45:913-922.
- Leverille SG, Guralnik JM, Ferrucci L, Langlois JA. Aging successfully until death in old age: Opportunities for increasing active life expectancy. *Am J Epidemiol* 1999;149(7):654-664.
- Locke EA, Latham GP. Building a practically useful theory of goal setting and task motivation. *Am Psychol* 2002;57(9):705-717.
- McAuley E, Morris KS, Motl RW, Hu L, Konopack JF, Elavsky S. Long-term follow-up of physical activity behavior in older adults. *Health Psychol* 2007;26(3):375-380.
- McAuley E. Self-efficacy and the maintenance of exercise participation in older adults. *J Behav Med* 1993;16(1):103-113.
- McAuley E, Jerome GJ, Elavsky S, Marquez DX, Ramsey SN. Predicting long-term maintenance of physical activity in older adults. *Prev Med* 2003;37:110-118.
- McAuley E, Mullen SP, Szabo AN, White SM, Wójcicki TR, Mailey EL, Gothe NP, Olson EA, Voss M, Erickson K, Prakash R, Kramer AF. Self-regulatory processes and exercise adherence in older adults: Executive function and self-efficacy effects. *Am J Prev Med* 2011;41(3):284-290.
- Mudrak J, Slepicka P, Elavsky S. Motivation for physical activity in Czech seniors. *Acta Universitatis Carolinae-Kinanthropologica* 2012;47(2):7-18.
- O'Brien Cousins S. Social support for exercise among elderly women in Canada. *Health Promot Int* 1995;10:273-282.
- OECD Better Life Index (Korea) available at <http://www.oecdbetterlifeindex.org/countries/korea/>
- OECD (2009), *Health at a Glance 2010: OECD Indicators*, OECD Publishing, Paris. available at <http://www.oecd.org/berlin/47570143.pdf>
- Oka RK, King AC, Young DR. Sources of social support as predictors of exercise adherence in women and men ages 50 to 65 years. *Women's Health: Res Gender Behav Policy* 1995;1:161-175.
- Orsega-Smith EM, Payne LL, Mowen AJ, Ho CH, Godbey GC. The role of social support and self-efficacy in shaping the leisure time physical activity of older adults. *J Leisure Res* 2007;39(4):705-727.
- Pettee KK, Brach JS, Kriska AM, Boudreau R, Richardson CR, Colbert LH, Newman AB. Influence of marital status on physical activity levels among older adults. *Med Sci Sports Exerc* 2006;38(3):541-546.
- Resnick B. Testing a model of overall activity in older adults. *J Aging Phys Act* 2001;9:142-160.
- Rovniak LS, Anderson ES, Winett RA, Stephens RS. Social cognitive determinants of physical activity in young adults: A prospective structural equation analysis. *Ann Behav Med* 2002; 24(2):149-156.
- Sallis JF, Grossman RM, Pinski RB, Patterson TL, Nader PR. The development of scales to measure social support for diet and exercise behaviors. *Prev Med* 1987;16:825-836.
- Yang S, Khang YH, Harper S, Smith GD, Leon DA, Lynch J. Understanding the rapid increase in life expectancy in South Korea. *Am J Public Health* 2010;100(5):896-903.
- Umstattd MR, Motl R, Wilcox S, Saunders R, Watford M. Measuring physical activity self-regulation strategies in older adults. *J Phys Act Health* 2009;6(1):105-112.
- UN data- Healthy life expectancy (HALE) at birth (years) Available at http://data.un.org/Data.aspx?q=life+expectancy&d=WHO&f=MEASURE_CODE%3AWHOSIS_000002#WHO
- United States Department of Health and Human Services. *Health, United States: Health and aging chartbook*. Washington, D. C: Author, 2006.
- Yeom HA, Jung D, Mona C. Adherence to physical activity among older adults using a geographic information system: Korean national health and nutrition examinations survey IV. *Asian Nurs Res* 2011;5(2):118-127.
- You KU, Won YB. The correlation between physical activity, negative emotion, economic level and mental health condition of elderly people. *Korean J Sport Psychol* 2010;21(4):197-205.
- Ware JE, Kosinski M, Keller SD. A 12-Item Short-Form Health Survey:

- Construction of scales and preliminary tests of reliability and validity. *Medical Care* 1996;34(3):220-233.
- Washburn R, Smith K, Jette A, Jenney, C. The Physical Activity Scale for the Elderly (PASE): Development and evaluation. *J Clin Epidemiol* 1993;46:153-162.
- Weiss DR, O'Loughlin JL, Platt RW, Paradis G. Five-year predictors of physical activity decline among adults in low-income communities: A prospective study. *Int J Behav Nutr Phys Act* 2007;4(2):1-7.