

Single-step protocol for the differentiation of human-induced pluripotent stem cells into hepatic progenitor-like cells

MINORU TOMIZAWA¹, FUMINOBU SHINOZAKI², TAKAO SUGIYAMA³,
SHIGENORI YAMAMOTO⁴, MAKOTO SUEISHI³ and TAKANOBU YOSHIDA⁵

Departments of ¹Gastroenterology, ²Radiology, ³Rheumatology, ⁴Pediatrics and ⁵Internal Medicine,
National Hospital Organization Shimoshizu Hospital, Yotsukaido, Chiba 284-0003, Japan

Received May 28, 2012; Accepted July 31, 2012

DOI: 10.3892/br.2012.2

Abstract. Induced pluripotent stem (iPS) cells are ideal sources of hepatocyte for transplantation into patients experiencing hepatic failure. Growth and transcription factors were analyzed to design a single-step protocol for the differentiation of iPS cells into hepatocytes. The expression of transcription factors was analyzed using reverse transcription-polymerase chain reaction (RT-PCR) and compared among iPS cells, as well as fetal and adult liver cells. iPS cells were cultured with growth factors and RT-PCR was performed to analyze the expression of transcription factors. iPS cells were introduced with transcription factors, cultured with growth factors and subjected to real-time quantitative PCR. Indocyanine green (ICG) was added to the medium as a hepatocyte marker. Sox17, GATA4, GATA6, FoxA2, HEX, HNF4 α and C/EBP α were expressed in fetal and adult liver cells, but not in iPS cells. Sox17, GATA6 and HNF4 α were expressed after exposure a combination of oncostatin M, epidermal growth factor, retinoic acid, dexamethasone and ITS (OERDITS). When iPS cells were introduced with FoxA2, GATA4, HEX and C/EBP α and cultured with OERDITS for 8 days, the cells expressed α -fetoprotein, δ -like (Dlk)-1 and γ -glutamyl transpeptidase (GTP), and ICG uptake was observed. Exposure to FoxA2, GATA4, HEX and C/EBP α and culturing with OERDITS supplementation potentially serves as a single-step inducer for the differentiation of iPS cells into hepatic progenitor-like cells within 8 days.

Introduction

Fulminant hepatitis is characterized by sudden, severe liver dysfunction leading to coagulopathy and hepatic encephalopathy

(1). In their study, Alam *et al* (2) reported the natural course of fulminant hepatitis with most of the patients developing fulminant hepatic failure within 2 weeks after the onset of jaundice, resulting in a 73.1% rate in mortality in the study population. Hepatocyte transplantation offers a potential therapeutic option for the treatment of fulminant hepatitis. Induced pluripotent stem (iPS) cells are an ideal source of autologous hepatocytes as they potentially prevent the need for immunosuppression prior to cell engraftment (3). However, the prompt differentiation of iPS cells into hepatocytes is essential.

Strategies used for the direct differentiation of iPS cells into hepatocytes have indicated the sequential supplementation of cytokines and growth factors involved in the embryonic development of the mammalian liver (3). Current strategies to generate hepatocytes from iPS cells that mimic liver embryogenesis by adding essential growth factors and simulators to serum-free culture medium have yielded more hepatocytes and hepatocyte-like cells and an increased homogeneity in the final cell population (3). Endodermal markers were previously analyzed in human embryonic stem (ES) cells, cultured with growth factors (4). Retinoic acid (RA), nerve growth factor (NGF), hepatocyte growth factor (HGF) and endodermal growth factor (EGF) are known to increase the expression of α -fetoprotein (AFP), while dexamethasone (Dex) and ITS are used to maintain the *in vitro* functions of hepatocytes (5). The use of activin A for the differentiation of iPS cells into endodermal cells is controversial. Activin A at a concentration of 100 ng/ml promotes the differentiation of ES cells into endoderm, whereas activin A maintains pluripotency, regulating the expression of Nanog (6,7).

Transcription factors play an important role in hepatocyte differentiation. Mice deficient in GATA-binding protein (GATA) 6 died between embryonic Days 6.5 and 7.5 and exhibited a specific defect in endodermal differentiation (8). The expression of Sry-related HMG box (Sox) 17 is restricted to the nascent primitive endodermal epithelium (9). The absence of Sox17 leads to the premature delamination and migration of the parietal endoderm. Forkhead box protein (Fox) A2 and GATA4 are the first known proteins that bind to the albumin gene enhancer in liver precursor cells in embryos (10). Disruption of the hematopoietically expressed homeobox (HEX) resulted in embryonic lethality attributable to insubstantial liver formation, making it essential for liver

Correspondence to: Dr Minoru Tomizawa, Department of Gastroenterology, National Hospital Organization Shimoshizu Hospital, 934-5 Shikawatashi, Yotsukaido, Chiba 284-0003, Japan
E-mail: nihminor-cib@umin.ac.jp

Key words: growth factor, transcription factor, hepatocyte

organogenesis (11). The CCAAT/enhancer binding protein (C/EBP) α promotes the differentiation of hepatoblasts into mature hepatocytes (12,13). Clones expressing the activating albumin promoter express C/EBP β (14). HNF4 α is essential for the hepatic specification of iPS cells (15). These data suggest that Sox17, GATA6, FoxA2, GATA4, HEX, C/EBP α , C/EBP β and HNF4 α are expected to promote the differentiation of iPS cells into hepatocytes.

Thus, the aim of this study was to analyze various growth and transcription factors involved in hepatocyte differentiation from iPS cells within 8 days.

Materials and methods

Cell culture. The iPS cell line (RIKEN Cell Bank, Tsukuba, Japan) 201B7 was cultured in ReproFF media (ReproCELL, Inc., Yokohama, Japan) as a feeder-free culture (ReproCELL, Inc., Tokyo, Japan) in dishes (Asahi Techno Glass, Funabashi, Japan) coated with Matrigel (Becton-Dickinson, Franklin Lakes, NJ) and were kept in 5% CO₂ at 37°C in a humidified chamber. The cells were collected with Accutase (Innovative Cell Technologies, Inc., San Diego, CA, USA) for the experiments. 201B7 cells were cultured in Dulbecco's modified Eagle's medium/F12-medium (Sigma-Aldrich Corp., St. Louis, MO, USA) supplemented with 20% of knockout serum replacement (Life Technologies, Grand Island, NY, USA), 10% of minimum essential amino acids (Life Technologies), 2 mM of L-glutamine (Life Technologies), and 1 mM of 2-mercaptoethanol (iPSm(-); Sigma-Aldrich Corp.). Basic fibroblast growth factor (bFGF) (5 ng/ml; Wako Pure Chemical Industries, Ltd., Osaka, Japan), bone morphogenic protein (BMP) 4 (20 ng/ml; Wako Pure Chemical Industries, Ltd.), oncostatin M (20 ng/ml; Wako Pure Chemical Industries), EGF (20 ng/ml; Wako Pure Chemical Industries, Ltd.), NGF (100 ng/ml; R&D Systems, Inc., Minneapolis, MN), transforming growth factor (TGF)- β (1 ng/ml; R&D Systems, Inc.), RA (1 μ M; Sigma-Aldrich Corp.), HGF (10 ng/ml; Sigma-Aldrich Corp.), Dex (10⁻⁷ M; Wako Pure Chemical Industries, Ltd.), and insulin, transferrin, selenium (ITS) (Wako Pure Chemical Industries, Ltd.) were added in iPSm(-).

Reverse transcriptase and real-time quantitative polymerase chain reaction. Total RNA (5 μ g), isolated using Isogen (Nippon Gene, Tokyo, Japan), was used for first-strand cDNA synthesis using SuperScript III and oligo(dT), according to the manufacturer's instructions (Life Technologies). Polymerase chain reaction (PCR) was performed using the GeneAmp[®] PCR System 9700 (Life Technologies) and subjected to gel electrophoresis. Real-time quantitative PCR was performed with Fast SYBR-Green Master mix (Life Technologies) and analyzed using the MiniOpticon (Bio-Rad, Hercules, CA, USA). The primer pairs used for RT-PCR were: GATA4, 5'-GAAAACGGAAGCCCAAGAACC and 5'-AGACA TCGCACTGACTGAGAACG (NM_002052, 56°C, 218 bp); GATA6, 5'-TTCATCACGGCGGCTTGGATTGTC and 5'-GTGTTGTGGGGGAAGTATTTTTGTC (NM_005257, 56°C, 299 bp); Sox17, 5'-CGCTTTCATGGTGTGGGCTAA GGACG and 5'-TAGTTGGGGTGGTCTGCATGTGCTG (NM_022454, 63°C, 186 bp); FoxA2, 5'-CCACCACCAACC CCACAAAATG and 5'-TGCAACACCGTCTCCCCAAAGT




Figure 1. The expression of transcription factors in iPS and liver cells. RT-PCR analysis indicated that GATA4, GATA6, Sox17, FoxA2, HEX and C/EBP α were not expressed in iPS cells, whereas their expression was detected in fetal or adult liver cells. C/EBP β was expressed in iPS, fetal and adult liver cells. Lanes: 1, H₂O; 2, iPS cells; 3, fetal liver and 4, adult liver.

(NM_021784, 60°C, 294 bp); HEX, 5'-TTCTCCAACGACC AGACCATCG and 5'-TTTTATCGCCCTCAATGTCCAC (NM_002729, 56°C, 364 bp); C/EBP α , 5'-TGGAGACGCA GCAGAAGGTG and 5'-TCGGGAAGGAGGCAGGAAAC (U34070, 69°C, 538 bp); and C/EBP β , 5'-AGACGCAGCAC AAGGTCCTG and 5'-GAGAGGGGCAGAGGGGAGAGC (X52560, 60°C, 421 bp). RT-PCR was performed with 30 cycles of 1 min of denaturation, 1 min of annealing and 1 min of extension. The primer pairs for real-time quantitative PCR of AFP, ribosomal protein L19 (RPL19), δ -like (Dlk)-1, GATA6, HNF4 α and γ -glutamyl transpeptidase (GTP) were 5'-ACA CAAAAGCCCACTCCAG and 5'-GGTGCATACAGGAA GGGATG (NM_005618, 147 bp), 5'-CGAATGCCAGA GAAGGTCAC and 5'-CCATGAGAATCCGCTTGTTT (157 bp), 5'-GGATGAGTGCGTCATAGCAA and 5'-CCT CCTCTTCAGCAGCATTC (121 bp), 5'-CCACTCGTGTC TGCTTTTGTGC and 5'-CCCTTCCCTTCCATCTTCT CTCAC (139 bp), 5'-CAACGGACAGATGTGTGAGTGG and 5'-ATAACTTCCTGCTTGGTGTATGGTC (NM_000457, 183 bp), and 5'-CCTCATCCTCAACATCCTCAAAGG and 5'-CACCTCAGTCACATCCACAAACTTG (J04131, 163 bp), respectively. The primer pairs for the quantitative PCR of Sox17 were the same as those for RT-PCR. Real-time quantitative PCR was performed for 40 cycles, using 5 sec for denaturation and 5 sec for annealing-extension.

Indocyanine green uptake study. Indocyanine green (ICG, 25 mg; Dai-ichi Pharmaceutical, Co., Ltd., Tokyo, Japan) was dissolved in 5 ml of water in a sterile vial and 20 ml were added to each medium at a final concentration of 1 mg/ml (16). The ICG solution was added to the cell culture and incubated at 37°C for 15 min (5). After the dish was rinsed with phosphate-buffered saline (PBS), the ICG uptake was observed through microscopy.

Results

Expression of transcription factors in iPS and liver cells. RT-PCR was performed to identify the transcription factors involved in hepatocyte differentiation (Fig. 1). GATA4,




Figure 2. RT-PCR analysis of transcription and growth factors. Sox17 and GATA6 were strongly expressed in the presence of oncostatin M and EGF, respectively. GATA4, FoxA2, HEX and C/EBP α were not sufficiently expressed with either growth factor. Lanes: 1, H₂O; 2, ReproFF; 3, no growth factors; 4, basic fibroblast growth factor; 5, BMP4; 6, oncostatin M; 7, epidermal growth factor; 8, nerve growth factor; 9, transforming growth factor β ; 10, retinoic acid and 11, hepatocyte growth factor.




Figure 3. The expression of GATA6, Sox17 and HNF4 α in the presence of growth factors. iPS cells were cultured using a single or a combination of growth factors. The expression of GATA6, Sox17 and HNF4 α were analyzed. (A and B) The relative expression levels were normalized against iPS cells in ReproFF media. (C) The expression of HNF4 α was analyzed using gel electrophoresis as it was too low to be normalized in iPS cells. (A and B) Columns and (C) lanes 1-8 show the following: 1, ReproFF; 2, oncostatin M; 3, epidermal growth factor; 4, retinoic acid; 5, dexamethasone; 6, ITS; 7, oncostatin M + epidermal growth factor + retinoic acid and 8, oncostatin M + epidermal growth factor + retinoic acid + dexamethasone + ITS. Error bar, standard error.

GATA6, Sox17, FoxA2, HEX and C/EBP α were expressed in fetal or adult liver cells, but not in iPS cells, whereas C/EBP β was expressed in iPS cells as well as in fetal and adult liver cells.

RT-PCR analysis of transcription and growth factors. Using RNA isolated from cells cultured with each growth factor for 8 days, RT-PCR was performed to detect the growth factors




Figure 4. The expression of α -fetoprotein (AFP) and δ -like (Dlk)-1 in the presence of growth factors. The expression of AFP and Dlk-1 in iPS cells cultured with growth factors was analyzed. The relative expression levels were normalized against iPS cells using ReproFF media. Columns: 1, ReproFF; 2, oncostatin M; 3, epidermal growth factor; 4, retinoic acid; 5, dexamethasone; 6, ITS; 7, oncostatin M + epidermal growth factor + retinoic acid and 8, oncostatin M + epidermal growth factor + retinoic acid + dexamethasone + ITS. Error bar, standard error.

promoting iPS cells to express GATA4, GATA6, Sox17, FoxA2, HEX and C/EBP α (Fig. 2). C/EBP β was not included in this experiment as it was expressed in iPS cells. GATA6 and Sox17 were strongly expressed in iPS cells cultured with EGF and oncostatin M, respectively. Sox7 and GATA6 were expressed in cells cultured with RA. C/EBP α was weakly expressed with oncostatin M. GATA4, FoxA2 and HEX were not expressed in iPS cells using any of the growth factors. These results suggest that oncostatin M, EGF and RA stimulated the expression of GATA6 and Sox17 in iPS cells.

Expression of GATA6, Sox17 and HNF4 α in the presence of growth factors. Real-time quantitative PCR conducted in cell cultures with a combination of oncostatin M, EGF and RA showed an increase in the expression of GATA6, Sox17 and HNF4 α (Fig. 3). The expression of GATA6 and Sox17 showed the greatest increase during culture with a combination of oncostatin M, EGF, RA, dexamethasone and ITS. HNF4 α was expressed in iPS cells cultured with oncostatin M, dexamethasone, ITS and their combinations.

Expression of α -fetoprotein (AFP) and δ -like (Dlk)-1 in the presence of growth factors. The expression of AFP and Dlk-1, markers of hepatic progenitor cells, was then analyzed in iPS cells with single or a combination of growth factors (Fig. 4). The combination of oncostatin M, EGF, RA, dexamethasone and ITS (OERDITS) resulted in the greatest increase in the expression of AFP. Dlk-1 expression was stimulated with either oncostatin M or OERDITS. Thus, OERDITS was regarded as the most suitable combination of growth factors to promote the differentiation of iPS cells into hepatocytes.




Figure 5. Expression of (A) AFP, (B) Dlk-1, (C) γ -GTP and (D) Nanog in the presence of growth factors. iPS cells were introduced with a combination of 3 or 4 growth factors from FoxA2, GATA4, HEX and C/EBP α and cultured in oncostatin M, EGF, RA, dexamethasone and ITS. The expression of AFP, Dlk-1, γ -GTP and Nanog was analyzed. The relative expression levels were normalized against iPS cells in ReproFF media. Columns: 1, ReproFF; 2, GHA; 3, FHA; 4, FGA; 5, FGH; 6, FGHA and 7, fetal liver. FGHA: F, FoxA2; G, GATA4; H, HEX; A, C/EBP α . Error bar, standard error.




Figure 6. Indocyanine green (ICG) uptake in iPS cells cultured at varying conditions is shown. Indocyanine green (ICG) was added in the medium of iPS cells introduced with FoxA2, GATA4, HEX and C/EBP α and cultured with oncostatin M, EGF, RA, dexamethasone and ITS. (A) iPS cells before the addition of ICG in the medium and (B) 15 min after the addition of ICG. Original magnification, x200; bar, 25 μ m.

Expression of AFP, Dlk-1, γ -GTP and Nanog in the presence of growth factors. FoxA2, GATA4, HEX and C/EBP α were introduced with iPS cells and cultured with OERDITS (Fig. 5). The expression of AFP and Dlk-1 was comparable to that observed in fetal liver cells with a combination of FoxA2, GATA4, HEX and C/EBP α (FGHA). The expression of γ -GTP with FGHA was the strongest, although it was lower than that shown by fetal liver cells. All the conditions significantly suppressed the expression of Nanog.

Indocyanine green (ICG) uptake in iPS cells cultured at varying conditions. ICG was added in the iPS cell culture medium 8 days after the introduction of FGHA and cultured in OERDITS. ICG was taken-up by iPS cells (Fig. 6).

Discussion

Stepwise differentiation protocols are currently employed to promote the differentiation of iPS cells into hepatocytes (15,17-19). Current protocols involve the sequential application of growth factors. DeLaForest *et al* (15) used LY294002, an inhibitor of phosphatidylinositol 3 kinase. Si-Tayeb *et al* (17) changed the oxygen concentration to 4 and 20%. Song *et al* (20) added supplements (N2 and B27). Takayama *et al* (19) transduced Sox17, HEX and HNF4 α . However, these protocols require more than 20 days for iPS cells to differentiate into hepatocytes. Shorter induction periods are necessary for the transplantation of cells to patients diagnosed with hepatic failure. Thus, we developed a new

protocol that promoted the differentiation of iPS cells into hepatocytes within 8 days. This study has also investigated growth factors that increased the expression of transcription factors. The OERDITS medium was found to stimulate the expression of GATA6, Sox17 and HNF4 α . Sox17 is essential for the differentiation of iPS cells into endodermal cells, whereas HNF4 α promotes the maturation of hepatocytes (21). In addition, the introduction of GATA6, Sox17 and HNF4 α was not necessary with the OERDITS medium. However, the OERDITS medium did not increase the expression of FoxA2, GATA4 and HEX. Therefore, the introduction of FoxA2, GATA4 and HEX was necessary for the induction of cell differentiation.

AFP expression was lowest in iPS cells introduced with FoxA2, GATA4 and HEX (Fig. 4), suggesting that C/EBP α was necessary for iPS cells to differentiate into hepatocytes. Thus, the combination of GATA4, FoxA2, HEX and C/EBP α was essential for the generation of hepatocytes from iPS cells.

The role of activin A in the endodermal differentiation of iPS cells depends on its concentration. The concentration of 100 ng/ml of activin A was shown to promote differentiation of human embryonic stem (ES) cells into endoderm cells (6). Activin A maintains pluripotency, regulating the Nanog expression (7). No endodermal markers were expressed with activin A at 20 ng/ml. Xiao *et al* (22) succeeded in the long-term feeder-free culture of human ES cells (H1) for >150 days and 20 passages using 5 ng/ml of activin A. However, the stepwise protocols employed high concentrations of activin A. Activin A was not analyzed in the present study as we had succeeded in the passage culture of 201B7 cells using a medium with activin A at 10 ng/ml (in preparation for submission).

Albumin was not expressed in iPS cells cultured with FGHA and OERDITS (data not shown). Future investigations should therefore be performed on the application of the extracellular matrix (23). Moreover, exposure to FoxA2, GATA4, HEX and C/EBP α and culturing with OERDITS supplementation potentially serves as a single-step inducer for the differentiation of iPS cells into hepatic progenitor-like cells.

Acknowledgements

This study was supported by the Grant-in-Aid for Encouragement of Scientists (no. 22931047) and the Research Grant-in-Aid for Scientific Research (C) (no. 23591002) from the Japan Society for the Promotion of Science (JSPS).

References

- Inoue K, Watanabe T, Maruoka N, Kuroki Y, Takahashi H and Yoshida M: Japanese-style intensive medical care improves prognosis for acute liver failure and the perioperative management of liver transplantation. *Transplant Proc* 42: 4109-4112, 2010.
- Alam S, Azam G, Mustafa G, *et al*: Natural course of fulminant hepatic failure: the scenario in Bangladesh and the differences from the west. *Saudi J Gastroenterol* 15: 229-233, 2009.
- Chistiakov DA and Chistiakov PA: Strategies to produce hepatocytes and hepatocyte-like cells from pluripotent stem cells. *Hepatol Res* 42: 111-119, 2012.
- Schuldiner M, Yanuka O, Itskovitz-Eldor J, Melton DA and Benvenisty N: Effects of eight growth factors on the differentiation of cells derived from human embryonic stem cells. *Proc Natl Acad Sci USA* 97: 11307-11312, 2000.
- Tomizawa M, Toyama Y, Ito C, *et al*: Hepatoblast-like cells enriched from mouse embryonic stem cells in medium without glucose, pyruvate, arginine, and tyrosine. *Cell Tissue Res* 333: 17-27, 2008.
- Yao S, Chen S, Clark J, *et al*: Long-term self-renewal and directed differentiation of human embryonic stem cells in chemically defined conditions. *Proc Natl Acad Sci USA* 103: 6907-6912, 2006.
- Shin M, Alev C, Wu Y, Nagai H and Sheng G: Activin/TGF-beta signaling regulates *Nanog* expression in the epiblast during gastrulation. *Mech Dev* 128: 268-278, 2011.
- Morrissey EE, Tang Z, Sigrist K, *et al*: GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev* 12: 3579-3590, 1998.
- Artus J, Piliszek A and Hadjantonakis AK: The primitive endoderm lineage of the mouse blastocyst: sequential transcription factor activation and regulation of differentiation by Sox17. *Dev Biol* 350: 393-404, 2011.
- Cirillo LA, Lin FR, Cuesta I, Friedman D, Jarnik M and Zaret KS: Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4. *Mol Cell* 9: 279-289, 2002.
- Keng VW, Yagi H, Ikawa M, *et al*: Homeobox gene HEX is essential for onset of mouse embryonic liver development and differentiation of the monocyte lineage. *Biochem Biophys Res Commun* 276: 1155-1161, 2000.
- Tomizawa M, Garfield S, Factor V and Xanthopoulos KG: Hepatocytes deficient in CCAAT/enhancer binding protein α (C/EBP α) exhibit both hepatocyte and biliary epithelial cell character. *Biochem Biophys Res Commun* 249: 1-5, 1998.
- Yamasaki H, Sada A, Iwata T, *et al*: Suppression of C/EBP α expression in periportal hepatoblasts may stimulate biliary cell differentiation through increased *Hnf6* and *Hnf1b* expression. *Development* 133: 4233-4243, 2006.
- Kheolamai P and Dickson AJ: Liver-enriched transcription factors are critical for the expression of hepatocyte marker genes in mES-derived hepatocyte-lineage cells. *BMC Mol Biol* 10: 35, 2009.
- DeLaForest A, Nagaoka M, Si-Tayeb K, *et al*: HNF4A is essential for specification of hepatic progenitors from human pluripotent stem cells. *Development* 138: 4143-4153, 2011.
- Yamada T, Yoshikawa M, Kanda S, *et al*: In vitro differentiation of embryonic stem cells into hepatocyte-like cells identified by cellular uptake of indocyanine green. *Stem Cells* 20: 146-154, 2002.
- Si-Tayeb K, Noto FK, Nagaoka M, *et al*: Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. *Hepatology* 51: 297-305, 2010.
- Song Z, Cai J, Liu Y, *et al*: Efficient generation of hepatocyte-like cells from human induced pluripotent stem cells. *Cell Res* 19: 1233-1242, 2009.
- Takayama K, Inamura M, Kawabata K, *et al*: Efficient generation of functional hepatocytes from human embryonic stem cells and induced pluripotent stem cells by HNF4 α transduction. *Mol Ther* 20: 127-137, 2012.
- Song H, Hollstein M and Xu Y: p53 gain-of-function cancer mutants induce genetic instability by inactivating ATM. *Nat Cell Biol* 9: 573-580, 2007.
- Takayama K, Inamura M, Kawabata K, *et al*: Efficient and directive generation of two distinct endoderm lineages from human ESCs and iPSCs by differentiation stage-specific SOX17 transduction. *PLoS One* 6: e21780, 2011.
- Xiao L, Yuan X and Sharkis SJ: Activin A maintains self-renewal and regulates fibroblast growth factor, Wnt, and bone morphogenic protein pathways in human embryonic stem cells. *Stem Cells* 24: 1476-1486, 2006.
- Shiraki N, Yamazoe T, Qin Z, *et al*: Efficient differentiation of embryonic stem cells into hepatic cells in vitro using a feeder-free basement membrane substratum. *PLoS One* 6: e24228, 2011.