

Green Fluorescent Diamidines as Diagnostic Probes for Trypanosomes

Federica Giordani,^c Manoj Munde,^a W. David Wilson,^a Mohamed A. Ismail,^{a,b} Arvind Kumar,^a David W. Boykin,^a Michael P. Barrett^c

Department of Chemistry, Georgia State University, Atlanta, Georgia, USA^a; Department of Chemistry, College of Science, King Faisal University, Hofuf, Saudi Arabia^b; Wellcome Trust Centre for Molecular Parasitology, Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom^c

Light-emitting diode (LED) fluorescence microscopy offers potential benefits in the diagnosis of human African trypanosomiasis and in other aspects of diseases management, such as detection of drug-resistant strains. To advance such approaches, reliable and specific fluorescent markers to stain parasites in human fluids are needed. Here we describe a series of novel green fluorescent diamidines and their suitability as probes with which to stain trypanosomes.

Early diagnosis of human African trypanosomiasis (HAT) (1) is paramount for favorable prognosis, but it is difficult to achieve in the field (2). Conventional light microscopy, in particular, is insensitive and time-consuming. Deployment of better diagnostics, together with more effective drugs, will be essential to eliminate the disease as a public health problem.

The development of portable and efficient light-emitting diode (LED)-illuminated microscopes (3–5) allows fluorescence microscopy to offer improvements in detection. Its use in the diagnosis of other tropical diseases has already been shown to increase the sensitivity and rapidity of sample preparation and screening (6, 7). A similar approach would also be useful in diagnosis of animal trypanosomiasis, which suffers limitations similar to those of diagnosis of HAT. The diagnostic fluorescence tests developed for HAT (8, 9) use acridine orange to stain trypanosomes. This is a cheap but aspecific fluorophore, which can label the nucleic acids of any cell in the specimen, and as a DNA intercalator, it poses potential risks to laboratory technicians (although, based on available data, the IARC inserts it in group 3, as not classifiable as to its carcinogenicity for humans).

The aim of this work was to synthesize and evaluate a series of fluorescent diamidines as targeted diagnostic probes for trypanosomes. Diamidines (10, 11) appeared promising since they are rapidly internalized by *Trypanosoma brucei* via specific surface transporters, including the aminopurine P2/TbAT1 transporter (12–14), and several of them are fluorescent (11, 15, 16). The UV emission of DB99 was previously exploited in our laboratory to develop a simple assay for detection of drug-resistant strains lacking P2 (17). Unfortunately, although LED fluorescence microscopes are being increasingly used, UV LED-illuminated instruments are not yet widely available, most fluorescence biological work being carried out at longer wavelengths.

(Biological aspects of this work were conducted by F. Giordani in partial fulfillment of the requirements for a Ph.D. from the University of Glasgow, Glasgow, Scotland, 2011.)

By addition of a series of ultraconjugated ring structures in the linker between the two amidine groups (containing the recognition motif for the parasite's P2 transporter), we generated a series of new diamidines with longer emission wavelengths than the parental UV-emitting DB75 (chemistry data are presented in the supplemental material). Spectral measurements confirmed that all novel dications emitted in the green wavelength range (Table 1). As expected for diamidines (11), the new molecules had a high affinity for DNA. Values of DNA binding by thermal melting

(ΔT_m , measured as described in reference 15) revealed the binding affinity to be greater than that of pentamidine (12.6°C) but lower than that of DB75 (25°C) (Table 1).

The dependence of the compounds on P2 transport for uptake into trypanosomes was confirmed by an alamarBlue assay (18) performed on three *Trypanosoma brucei brucei* strains: a wild-type line (bloodstream Lister 427), a line not expressing P2 (*tbat1*^{-/-}) (19), and a derived clone, in which high-affinity pentamidine transporter (HAPT1, now identified as an aquaglyceroporin [20]) activity is also lost (B48) (21). Compound DB1692 showed the greatest dependency on P2 for transport (50% inhibitory concentration [IC₅₀] 25-fold higher for the *tbat1*^{-/-} line than for the wild type), followed by DB1645 (16-fold higher), DB1465 (10-fold), and DB1464 (8-fold) (Table 2). Data using the B48 line showed that HAPT1 activity also contributed to the accumulation of DB1645 and DB1465. In contrast, the addition of a single furan ring in the linker of DB1680 highly affected the rate of uptake of this compound through the P2 transporter, and further loss of HAPT1 did not affect its toxicity, suggesting other routes of internalization (e.g., other carriers or endocytosis). Equally noteworthy is the potent *in vitro* activity shown by these new compounds against the wild-type line, especially by DB1464, which had an IC₅₀ close to that of pentamidine. This indicates that these molecules may be attractive candidates to pursue as possible leads against HAT, provided that ADME (absorption, distribution, metabolism, and excretion) characterization demonstrates good pharmacokinetic properties and rules out potential toxicity issues previously observed for some molecules belonging to this class.

All five diamidines stained live *T. b. brucei* in infected rat blood films using a standard fluorescein isothiocyanate (FITC) filter set (excitation, band-pass, 450 to 490 nm; beam splitter, 510 nm; and emission, band-pass, 515 to 565 nm) and a Zeiss Axioplan fluo-

Received 16 September 2013 Returned for modification 6 December 2013

Accepted 17 December 2013

Published ahead of print 23 December 2013

Address correspondence to Michael P. Barrett, michael.barrett@glasgow.ac.uk.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.02024-13>.

Copyright © 2014 Giordani et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

doi:10.1128/AAC.02024-13

TABLE 1 Structures and physical properties of the fluorescent diamidines

Compound	Structure	λ_{EX} (nm) ^a	λ_{EM} (nm) ^b	Fluorescence intensity at max ^c	ΔT_m values ^d with poly(dA) · poly(dT)
DB1464		405	542	73	9.4
DB1465		417	513	123	18.5
DB1645		424	576	36	14.6
DB1680		418	546	8	15.0
DB1692		400	546	40	17.0

^a Excitation wavelength.^b Emission wavelength.^c max, maximum.^d Melting temperature of the compound-DNA complexes as compared to values for free DNA. T_m values are in degrees Celsius and have an error of $\pm 0.5^\circ\text{C}$.

rescence microscope (Fig. 1A). Under our experimental conditions (*ex vivo* incubation for 15 min at 37°C, with 50 μM [each] fluorophore), all five diamidines provided specific staining of trypanosomes without labeling erythrocytes or white blood cells. Staining of the parasite nucleus was not observed, but all com-

TABLE 2 *In vitro* trypanocidal activities of the five diamidines against *T. b. brucei* S427 wild-type and derived cell lines^a

Compound	IC ₅₀ [nM] (\pm SEM) for:				
	Wild type	<i>tbat1</i> ^{-/-} line	RF ^b	IC ₅₀ [nM] (\pm SEM) for B48	RF
DB1464	0.9 \pm 0.3	6.9 \pm 2.1	8	7.3 \pm 0.7	8
DB1465	6.9 \pm 1.9	71.4 \pm 4.4	10	334 \pm 11	48
DB1645	14.5 \pm 2.8	233 \pm 43	16	610 \pm 60	42
DB1680	266 \pm 21	591 \pm 43	2	341 \pm 11	1
DB1692	17.6 \pm 4.4	436 \pm 41	25	344 \pm 14	20
Pentamidine	0.6 \pm 0.3	0.9 \pm 0.2	1.5	167 \pm 10	278

^a $n = 3$.^b RF, resistance factor (ratio of the IC₅₀ measured against the *tbat1*^{-/-} and B48 cell lines to its value obtained for the wild type).

pounds selectively accumulated inside the kinetoplast (the parasite's mitochondrial genome, at the posterior of the cell) and other cytoplasmic organelles, spread throughout the cell body and also visualized at longer wavelengths (Fig. 1B). These were likely to be acidocalcisomes, previously observed to be stained by other diamidines (15). For DB1692, a prominent red fluorescent spot, putatively identified as the lysosome, also appeared in a region between the nucleus and the kinetoplast after 1 h of treatment. Under the FITC filter, parasites were readily detectable among blood cells based on their movement and fluorescent dots (especially the bright kinetoplast). Cell labeling was achieved only by incubating at room temperature or 37°C but not at 4°C. This temperature sensitivity could lead to variability in field tests using ambient temperature in different sites; hence, the use of temperature-controlled devices (e.g., water baths or heating blocks) might be required in such settings. Although the fluorescence intensity increased with the time of incubation for all probes, the overall yields remained below those identified with DB75 and could not be improved by increasing the fluorophore concentration. The low emission intensity of the new compounds could be




FIG 1 (A) Fluorescence images of infected rat blood incubated *ex vivo* with the five green fluorescent diamidines (note that the parasite position can change between the bright-field and corresponding fluorescence micrographs), 40 \times objective. (B) *In vitro* trypanosomes treated with DB1680 and DB1692 (50 μ M, 1 h, 37 $^{\circ}$ C), 100 \times objective; 4',6-diamidino-2-phenylindole (DAPI) (20 μ M) was used as a DNA counterstain. Asterisk, kinetoplast; arrow, cytoplasmic corpuscles, possibly acidocalcisomes; arrowhead, nucleus; hash mark, possibly lysosome. Bar, 10 μ m.

due to their intrinsically low fluorescence yields (Table 1) but also to the discrepancy between their excitation maxima, all close to 400 nm, and the standard FITC filter wavelength utilized for microscopy. Hence, the use of these specific fluorophores with these particular filters may not be directly applicable to HAT diagnosis due to the risk of false negatives. However, these same compounds with customized filters or further derivatives with higher fluorescence yields could be considered as probes for the specific staining of trypanosomes using existing LED-based fluorescence microscopes.

Despite the compounds' partial dependence on the P2 transporter for uptake, they did not distinguish wild-type and *tbat1*^{-/-} knockout line by fluorescence microscopy (50 μ M for incubations ranging from 15 min to up to 2 h), indicating that uptake routes independent of P2 play important roles in their internalization when used under these conditions.

Implementation of fluorescence tests for trypanosome detection in human fluids must rely on affordable, efficient and preferably specific markers. This work shows the feasibility of generating cheap fluorophores, emitting at differential wavelengths, by simple addition of cyclic groups within the chemical structure of diamidines, the precursor of which, DB75, is fluorescent only in the UV region. The exploitation of surface transporters peculiar to trypanosomes is a straightforward way to specifically target fluorophores to these parasites, and improved compounds may be developed following this approach to target both human and animal infective trypanosomes.

ACKNOWLEDGMENTS

This work was supported by an award from the Bill and Melinda Gates Foundation through the Consortium for Parasitic Drug Development (CPDD) (to M.P.B., W.D.W., and D.W.B.), an NIH grant (AI064200) (to W.D.W. and

D.W.B.), the Sir Halley Stewart Trust (to M.P.B. and F.G.), and the Wellcome Trust. The Wellcome Trust Centre for Molecular Parasitology is supported by core funding from the Wellcome Trust (085349).

REFERENCES

- Barrett MP, Burchmore RJ, Stich A, Lazzari JO, Frasch AC, Cazzulo JJ, Krishna S. 2003. The trypanosomiasis. *Lancet* 362:1469–1480. [http://dx.doi.org/10.1016/S0140-6736\(03\)14694-6](http://dx.doi.org/10.1016/S0140-6736(03)14694-6).
- Chappuis F, Loutan L, Simarro P, Lejon V, Buscher P. 2005. Options for field diagnosis of human African trypanosomiasis. *Clin. Microbiol. Rev.* 18:133–146. <http://dx.doi.org/10.1128/CMR.18.1.133-146.2005>.
- Hanscheid T. 2008. The future looks bright: low-cost fluorescent microscopes for detection of *Mycobacterium tuberculosis* and *Coccidiae*. *Trans. R. Soc. Trop. Med. Hyg.* 102:520–521. <http://dx.doi.org/10.1016/j.trstmh.2008.02.020>.
- Jones D, Nyalwidhe J, Tetley L, Barrett MP. 2007. McArthur revisited: fluorescence microscopes for field diagnostics. *Trends Parasitol.* 23:468–469. <http://dx.doi.org/10.1016/j.pt.2007.08.003>.
- Minion J, Sohn H, Pai M. 2009. Light-emitting diode technologies for TB diagnosis: what is on the market? *Expert. Rev. Med. Devices* 6:341–345. <http://dx.doi.org/10.1586/erd.09.26>.
- Keiser J, Utzinger J, Premji Z, Yamagata Y, Singer BH. 2002. Acridine orange for malaria diagnosis: its diagnostic performance, its promotion and implementation in Tanzania, and the implications for malaria control. *Ann. Trop. Med. Parasitol.* 96:643–654. <http://dx.doi.org/10.1179/000349802125001834>.
- Steingart KR, Henry M, Ng V, Hopewell PC, Ramsay A, Cunningham J, Urbanczik R, Perkins M, Aziz MA, Pai M. 2006. Fluorescence versus conventional sputum smear microscopy for tuberculosis: a systematic review. *Lancet Infect. Dis.* 6:570–581. [http://dx.doi.org/10.1016/S1473-3099\(06\)70578-3](http://dx.doi.org/10.1016/S1473-3099(06)70578-3).
- Bailey JW, Smith DH. 1992. The use of the acridine orange QBC technique in the diagnosis of African trypanosomiasis. *Trans. R. Soc. Trop. Med. Hyg.* 86:630. [http://dx.doi.org/10.1016/0035-9203\(92\)90160-E](http://dx.doi.org/10.1016/0035-9203(92)90160-E).
- Bieler S, Matovu E, Mitashi P, Ssewanyana E, Bi Shamamba SK, Bessell PR, Ndung'u JM. 2012. Improved detection of *Trypanosoma brucei* by lysis of red blood cells, concentration and LED fluorescence microscopy. *Acta Trop.* 121:135–140. <http://dx.doi.org/10.1016/j.actatropica.2011.10.016>.
- Paine MF, Wang MZ, Generaux CN, Boykin DW, Wilson WD, de Koning HP, Olson CA, Pohlig G, Burri C, Brun R, Murilla GA, Thuita JK, Barrett MP, Tidwell RR. 2010. Diamidines for human African trypanosomiasis. *Curr. Opin. Investig. Drugs* 11:876–883.
- Wilson WD, Nguyen B, Tanious FA, Mathis A, Hall JE, Stephens CE, Boykin DW. 2005. Dications that target the DNA minor groove: compound design and preparation, DNA interactions, cellular distribution and biological activity. *Curr. Med. Chem. Anticancer Agents* 5:389–408. <http://dx.doi.org/10.2174/1568011054222319>.
- Carter NS, Berger BJ, Fairlamb AH. 1995. Uptake of diamidine drugs by the P2 nucleoside transporter in melarsen-sensitive and -resistant *Trypanosoma brucei brucei*. *J. Biol. Chem.* 270:28153–28157. <http://dx.doi.org/10.1074/jbc.270.47.28153>.
- de Koning HP, Anderson LF, Stewart M, Burchmore RJ, Wallace LJ, Barrett MP. 2004. The trypanocide diminazene aceturate is accumulated predominantly through the TbAT1 purine transporter: additional insights on diamidine resistance in African trypanosomes. *Antimicrob. Agents Chemother.* 48:1515–1519. <http://dx.doi.org/10.1128/AAC.48.5.1515-1519.2004>.
- Lanteri CA, Stewart ML, Brock JM, Alibu VP, Meshnick SR, Tidwell RR, Barrett MP. 2006. Roles for the *Trypanosoma brucei* P2 transporter in DB75 uptake and resistance. *Mol. Pharmacol.* 70:1585–1592. <http://dx.doi.org/10.1124/mol.106.024653>.
- Mathis AM, Bridges AS, Ismail MA, Kumar A, Francesconi I, Anbazhagan M, Hu Q, Tanious FA, Wenzler T, Saulter J, Wilson WD, Brun R, Boykin DW, Tidwell RR, Hall JE. 2007. Diphenyl furans and aza analogs: effects of structural modification on *in vitro* activity, DNA binding, and accumulation and distribution in trypanosomes. *Antimicrob. Agents Chemother.* 51:2801–2810. <http://dx.doi.org/10.1128/AAC.00005-07>.
- Mathis AM, Holman JL, Sturk LM, Ismail MA, Boykin DW, Tidwell RR, Hall JE. 2006. Accumulation and intracellular distribution of antitrypanosomal diamidine compounds DB75 and DB820 in African trypanosomes. *Antimicrob. Agents Chemother.* 50:2185–2191. <http://dx.doi.org/10.1128/AAC.00192-06>.
- Stewart ML, Krishna S, Burchmore RJ, Brun R, de Koning HP, Boykin DW, Tidwell RR, Hall JE, Barrett MP. 2005. Detection of arsenical drug resistance in *Trypanosoma brucei* with a simple fluorescence test. *Lancet* 366:486–487. [http://dx.doi.org/10.1016/S0140-6736\(05\)66793-1](http://dx.doi.org/10.1016/S0140-6736(05)66793-1).
- Vincent IM, Creek D, Watson DG, Kamleh MA, Woods DJ, Wong PE, Burchmore RJ, Barrett MP. 2010. A molecular mechanism for eflornithine resistance in African trypanosomes. *PLoS Pathog.* 6:e1001204. <http://dx.doi.org/10.1371/journal.ppat.1001204>.
- Matovu E, Stewart ML, Geiser F, Brun R, Maser P, Wallace LJ, Burchmore RJ, Enyaru JC, Barrett MP, Kaminsky R, Seebeck T, de Koning HP. 2003. Mechanisms of arsenical and diamidine uptake and resistance in *Trypanosoma brucei*. *Eukaryot. Cell* 2:1003–1008. <http://dx.doi.org/10.1128/EC.2.5.1003-1008.2003>.
- Baker N, Glover L, Munday JC, Aguinaga AD, Barrett MP, de Koning HP, Horn D. 2012. Aquaglyceroporin 2 controls susceptibility to melarsoprol and pentamidine in African trypanosomes. *Proc. Natl. Acad. Sci. U. S. A.* 109:10996–11001. <http://dx.doi.org/10.1073/pnas.1202885109>.
- Bridges DJ, Gould MK, Nerima B, Maser P, Burchmore RJ, de Koning HP. 2007. Loss of the high-affinity pentamidine transporter is responsible for high levels of cross-resistance between arsenical and diamidine drugs in African trypanosomes. *Mol. Pharmacol.* 71:1098–1108. <http://dx.doi.org/10.1124/mol.106.031351>.