

Antileishmanial Activity of Semisynthetic Lupane Triterpenoids Betulin and Betulinic Acid Derivatives: Synergistic Effects with Miltefosine

Maria C. Sousa^{1,2*}, Raquel Varandas¹, Rita C. Santos¹, Manuel Santos-Rosa⁴, Vera Alves⁴, Jorge A. R. Salvador^{1,3*}

1 Faculty of Pharmacy, University of Coimbra, Pólo das Ciências da Saúde, Azinhaga de Santa Comba, Coimbra, Portugal, **2** Centre of Pharmaceutical Studies, Faculty of Pharmacy of University of Coimbra (CEF/FFUC), Pólo das Ciências da Saúde, Azinhaga de Santa Comba, Coimbra, Portugal, **3** CNC- Center for Neurosciences and Cell Biology, University of Coimbra, Coimbra, Portugal, **4** Institute of Immunology, Faculty of Medicine, University of Coimbra, Pólo I, Rua Larga, Coimbra, Portugal

Abstract

Leishmaniasis is a neglected tropical disease (NTDs), endemic in 88 countries, affecting more than 12 million people. The treatment consists in pentavalent antimony compounds, amphotericin B, pentamidine and miltefosine, among others. However, these current drugs are limited due to their toxicity, development of biological resistance, length of treatment and high cost. Thus, it is important to continue the search for new effective and less toxic treatments. The anti-*Leishmania* activity of sixteen semisynthetic lupane triterpenoids derivatives of betulin (BT01 to BT09) and betulinic acid (AB10 to AB16) were evaluated. Drug interactions between the active compounds and one current antileishmanial drug, miltefosine, were assessed using the fixed ratio isobologram method. In addition, effects on the cell cycle, apoptosis/necrosis events, morphology and DNA integrity were studied. The derivatives BT06 (3 β -Hydroxy-(20R)-lupan-29-oxo-28-yl-1H-imidazole-1-carboxylate) and AB13 (28-(1H-imidazole-1-yl)-3,28-dioxo-lup-1,20(29)-dien-2-yl-1H-imidazole-1-carboxylate) were found to be the most active, with IC₅₀ values of 50.8 μ M and 25.8 μ M, respectively. Interactions between these two compounds and miltefosine were classified as synergistic, with the most effective association being between AB13 and miltefosine, where decreases of IC₅₀ values to 6 μ M were observed, similar to the miltefosine activity alone. AB13 induced significant morphological changes, while both derivatives produced anti-proliferative activity through cell cycle arrest at the G0/G1 phase. Neither of these derivatives induced significant apoptosis/necrosis, as indicated by phosphatidylserine externalization and DNA fragmentation assays. In addition, neither of the derivatives induced death in macrophage cell lines. Thus, they do not present any potential risk of toxicity for the host cells. This study has identified the betulin derivative BT06 and the betulinic acid derivative AB13 as promising molecules in the development of new alternative therapies for leishmaniasis, including those involving combined-therapy with miltefosine.

Citation: Sousa MC, Varandas R, Santos RC, Santos-Rosa M, Alves V, et al. (2014) Antileishmanial Activity of Semisynthetic Lupane Triterpenoids Betulin and Betulinic Acid Derivatives: Synergistic Effects with Miltefosine. PLoS ONE 9(3): e89939. doi:10.1371/journal.pone.0089939

Editor: Andrea Cavalli, University of Bologna & Italian Institute of Technology, Italy

Received: October 24, 2013; **Accepted:** January 23, 2014; **Published:** March 18, 2014

Copyright: © 2014 Sousa et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by “(POCI)/FEDER” da Fundação para a Ciência e Tecnologia (FCT). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: mcsousa@ci.uc.pt (MCS); salvador@ci.uc.pt (JARS)

Introduction

Over one billion people are infected by one or more neglected tropical diseases (NTDs). These diseases comprise a group of parasitic, viral and bacterial infections that affect some of the poorest and most marginalized populations globally. Leishmaniasis is one such NTD that is endemic in 88 countries, affecting more than 12 million people and threatening 350 million people worldwide. The disease is associated with an incidence of 1.5 to 2 million cases per year, and an annual mortality rate of over 59,000 deaths [1]. It is caused by the *Leishmania* species, and the disease is broadly classified into three different clinical forms: visceral leishmaniasis (VL), cutaneous leishmaniasis (CL) and mucocutaneous leishmaniasis (MCL), which differ in the pattern and clinical manifestations of infection. VL can be fatal if left untreated, CL is localized and frequently self-heals within 3–18 months, while MCL leaves disfiguring scars. The parasite growth occurs through different morphological stages: the flagellated promastigotes

develop in the gut of the phlebotomine sandfly female (vector), while the non-flagellated amastigotes develop in mammalian host macrophages.

Currently, the treatment consists of chemotherapeutic agents, such as the pentavalent antimony compounds (sodium stibogluconate or meglumine antimoniate), polyene amphotericin B (as the deoxycholate salt or a liposomal formulation, AmBisome), the alkylphosphocholine miltefosine, aminoglycoside paromomycin and pentamidine [2]. All of these are limited due to their high toxicity, life-threatening side-effects, cost, length of treatment and emergence of resistance [1,3]. Because of the limited viable treatment options, with few alternatives available in the pipeline, it is important to continue the search for new effective chemotherapeutics and less toxic treatments.

The development of drug resistance is associated with monotherapy regimes [3,4]. Combination therapy is one interesting approach to decrease this development, to reduce the duration




Figure 1. Structures of betulin derivatives and IUPAC nomenclature.

doi:10.1371/journal.pone.0089939.g001

and cost of the treatment, and to increase the lifetime of old and new drugs [5,6]. Several combined treatments for visceral leishmaniasis have been tested with positive results, leading to a

reduction of adverse symptoms and a shorter duration of therapy [7].




Figure 2. Structures of betulinic acid derivatives and IUPAC nomenclature.

doi:10.1371/journal.pone.0089939.g002

Recent pharmacological studies elicited interest in several molecules with activities that trigger apoptotic death in cancerous cells as potential antiparasitic agents [8]. In trypanosomatids, features suggesting programmed cell death (PCA), commonly named as apoptosis, have been reported in response to a wide range of stimuli such as heat shock, reactive oxygen species, antiparasitic drugs, prostaglandins, and antimicrobial peptides [9–11]. There are several reports showing that *Leishmania* apoptosis occur in response to antileishmanial drugs. The treatment with the pentavalent antimony Sb(V) shows a significant induction of caspase-like activity resulting in DNA fragmentation [12]. Inhibitors of respiratory chain complexes were able to induce apoptotic cell death on the blood stream form of *L. donovani* [13]. Topoisomerase I poison camptothecin, promotes protein-DNA cleavable complex formation leading to apoptosis-like cell death in *Leishmania donovani* [14,15]. Miltefosine, the latest antileishmanial

drug introduced in the market and the first effective oral treatment of VL, was initially developed as an anticancer drug [16] and induce apoptosis-like cell death in *Leishmania donovani* [17]. All these findings point the importance of testing known anticancer agents against *Leishmania* due of its potential to inducing parasite death.

Triterpenoids in general, and particularly betulin and its derivative betulinic acid, show antitumor [18–20], anti-inflammatory [21,22], antiviral [19,23,24], antibacterial and antimalarial activity [25,26]. There are few studies in literature reporting the activity of betulin and betulinic acid derivatives in *Leishmania*. Alakurtti and collaborators [27] determined the activity of heterocyclic betulin derivatives on *Leishmania donovani* amastigotes and Dominguez-Carmona and collaborators [28] found activity against promastigotes of *L. amazonensis* of betulinic acid acetate and of betulinic acid methyl ester.

Previously, a series of new imidazole carboxylic esters (carbamates) and N-acylimidazole derivatives of betulin and betulinic acid have been synthesized and were shown to have cytotoxicity activity against human cancer cell lines HepG2, Jurkat and HeLa [29]. This cytotoxicity was related with apoptosis mechanisms associated with caspases signalling and DNA topoisomerases inhibition [30,31]. These results prompt us to evaluate their potential activity on *Leishmania*. We studied the susceptibility of *Leishmania infantum* to sixteen betulin and betulinic acid derivatives, together with the interactions between the most active compounds and miltefosine. Additionally, we have undertaken other assays to demonstrate the safety of the triterpenoid compounds for mammalian cells, and to elucidate the mechanisms of action involved in the leishmanicidal activity.

Materials and Methods

Chemicals and reagents

RPMI-1640 Medium, Phosphate Buffered Saline (PBS), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), dimethylsulfoxide (DMSO) were obtained from Sigma–Aldrich Co. Fetal bovine serum (FBS) was purchased from Gibco-Invitrogen.

Drugs and inhibitors

Miltefosine was obtained from Sigma Chemical Co.(St. Louis,USA). The nine derivatives of betulin (Figure 1) and the seven betulinic acid derivatives (Figure 2) used in this study were previously synthesized as part of a large library of compounds for potential antitumor evaluation [29]. The stock solutions (10 mM) derivatives of betulin and betulinic acid were prepared in dimethylsulfoxide (DMSO) and stored at -20°C . A stock solution of miltefosine (10 mM) was prepared in deionizer water and stored at 4°C . Final DMSO concentration (1%) had no effect on parasite clearance.

Parasites and cultures

Promastigote forms of *Leishmania infantum* Nicolle (zymodeme MON-1) were maintained at 26°C with weekly transfers in HEPES (25 mM)-buffered RPMI 1640 medium enriched with 10% inactivated fetal bovine serum (FBS). Log phase promastigotes were used to study the effects of the betulin and betulinic acid derivatives, miltefosine and drug combinations.

Viability assay

In drug susceptibility assays, stock solutions of betulin and betulinic acid derivatives were diluted in culture medium (RPMI 1640) in order to get a range of concentrations from 1 to 100 μM . Log phase promastigotes of *L. infantum* (2×10^6 cells. ml^{-1}) were incubated at 26°C for 24 h in HEPES (25 mM)-buffered RPMI 1640 medium enriched with 10% inactivated FBS in the presence of increasing concentrations of the derivatives or different concentration of DMSO (maximum of 1% as controls).

The effect on the cells viability was tested by MTT (3-[4, 5-methylthiazol-2-yl]-2, 5-diphenyltetrazolium bromide) colorimetric method, based on the reduction of the tetrazolium dye to insoluble formazan by the mitochondrial enzymes [32]. Briefly, 25 μl of MTT (5 mg. ml^{-1}) was added to each well, incubated for 2 h at 37°C and centrifuged at 3000 rpm for 5 min. The supernatant was removed, the cells were washed in PBS, and the precipitated formazan was dissolved in DMSO (250 μl). Cell viability was measured by absorbance at 530 nm on an ELISA plate reader (Synergy HT, Bio-TEK), and calculated using the following formula: $[(L2/L1) \times 100]$, where L1 is the absorbance of




Figure 3. Effects of betulin (A) and betulinic acid (B) derivatives on *Leishmania infantum* promastigotes viability. Cultures of log-phase promastigotes (2×10^6 cells. ml^{-1}) were incubated at 26°C for 24 h at different drug concentrations. Values are expressed as means and SEM.

doi:10.1371/journal.pone.0089939.g003

control cells and L2 is the absorbance of treated cells. Three separate experiments were performed for each drug and the concentration that inhibited viability by 50% (IC_{50}) was determined through dose-response regression analysis, plotted by GraphPad Prism 5.

Isobologram construction and classification of the nature of drug interaction

For the most active betulin and betulinic acid derivatives, combinations were made with them at growing concentrations and




Figure 4. Effects of combinations of betulin and betulinic acid derivatives with miltefosine on *Leishmania infantum* promastigotes viability. (A) Betulin derivative BT06; (B) Betulinic acid derivative AB13. Values are expressed as means and SEM. Cultures of log-phase promastigotes (2×10^6 cells.ml⁻¹) were incubated at 26°C for 24 h at different compound concentrations. doi:10.1371/journal.pone.0089939.g004

miltefosine at fixed concentrations and cells viability was evaluated by MTT as described above.

Classical isobolograms were constructed by plotting drugs concentrations (alone and in combination) that inhibits 50% trophozoites viability, as previously described [33]. The isobologram analysis evaluates the nature of interaction of two drugs, *i.e.*, drug A and drug B. First, the concentrations of drugs A and B required to produce a defined single-agent effect (*e.g.*, IC₅₀), when used as single agents, are placed on the *x* and *y* axes in a two-coordinate plot, corresponding to (CA, 0) and (0, CB), respectively. The line connecting these two points is the line of additivity. Second, the concentrations of the two drugs used in combination to provide the same effect, denoted as (cA, cB), are placed in the same plot. Synergy, additivity, or antagonism are indicated when (cA, cB) is located below, on, or above the line, respectively.

Morphological studies

L. infantum promastigotes were exposed to the most active betulin and betulinic acid derivatives and the morphological alterations were investigated by optical microscopy using direct examination of live microorganism (hanging drop) and after Giemsa stain. Briefly, exponentially grown of *L. infantum* promastigote (2×10^6 cells.ml⁻¹) were treated with BT06 and AB13 at IC₅₀ concentrations for 2 h, 4 h, 6 h and 24 h at 26°C. After incubation, cells were pelleted by centrifugation at 3000 rpm for 5 min and the supernatant was discarded by aspiration. The cell pellet was suspended in fresh medium and approximately 10 μL was placed on a Koch slide and directly observed under the optical microscope phase contrast (Eclipse E400, Nikon coupled with a digital camera 165 DN100 Nikon). In addition, a smear was made which was submitted to Giemsa stain. The smear was fixed with

Table 1. Inhibitory concentrations at 50% (IC₅₀) of the association between betulin and acid betulinic derivatives (BT06, AB13) with miltefosine on *Leishmania infantum* promastigotes.

Drug combination	IC ₅₀ μM (CI)*	R ²
BT06	50.8 (46.9–55.0)	0.85
+miltefosine 2 μM	30.1 (27.2–33.4)	0.94
+miltefosine 4 μM	25.9 (20.1–33.3)	0.81
AB13	25.8 (23.3–28.7)	0.92
+miltefosine 2 μM	7.6 (5.8–10.0)	0.87
+miltefosine 4 μM	6.0 (5.3–6.7)	0.93
Miltefosine	7.6 (6.6–8.8)	0.93

* CI, confidence interval at 95%.

doi:10.1371/journal.pone.0089939.t001




Figure 5. Comparison of the effects of combinations between BT06 and miltefosine with the effects of BT06 alone. Cultures of log-phase promastigotes (2×10^6 cells. ml^{-1}) were incubated at 26°C for 24 h. Significance level of * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. doi:10.1371/journal.pone.0089939.g005

methanol for 5 min, stained with aqueous solution of Giemsa (1/10, v/v) for 10 min at room temperature and finally washed with water and air dried. The stained smear was observed under the microscope with a $100\times$ lens (Eclipse E400, coupled with Nikon digital camera, Nikon DN100 165).

Cell cycle analysis

For analysis of DNA content, exponentially grown of *L. infantum* (2×10^6 cells ml^{-1}) were treated with BT06 and AB13 at IC_{50} concentrations for 2 h, 4 h, 6 h and 24 h at 26°C . At each time point, cells were fixed in 200 μl of 70% ethanol for 30 min. at 4°C . After washing cells with 2 mL of PBS, enriched with 2% of bovine serum albumin (BSA), the pellets were suspended in 0.5 mL of PI solution (PI/Rnase, Immunostep) and incubated for 15 minutes at 37°C [34]. Cells were then analyzed by flow cytometry (FacsCalibur-Beckton-Dickinson). Results were treated using ModFit LT V 2.0 programme.

Phosphatidylserine externalization

Double staining for annexin V-FITC and propidium iodide (PI) was performed as described previously [35]. Briefly, *L. infantum* promastigotes (2×10^6 cells. ml^{-1}) were exposed to BT06 and AB13 at IC_{50} concentrations for 2 h, 4 h, 6 h, and 24 h at 26°C . Cells were then washed with PBS and resuspended in binding buffer (10 mM HEPES-NaOH, pH 7.4, 140 NaCl, 2.5 mM CaCl_2). To 100 μl of this suspension were added 5 μl of Annexin V-FITC and 5 μl of PI (AnnexinV-FITC Apoptosis detection Kit, Immunostep). After 15 min incubation in the dark at room temperature, it was added 400 μl binding buffer and cells were analyzed by flow cytometry (FacsCalibur-Beckton-Dickinson). Data analysis was carried out using the program Paint-a-gate, and values are expressed as a percentage of positive cells for a given marker, relatively to the number of cells analyzed.




Figure 6. Comparison of the effects of combinations between AB13 and miltefosine with the effects of AB13 alone. Cultures of log-phase promastigotes (2×10^6 cells. ml^{-1}) were incubated at 26°C for 24 h. Significance level of * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$. doi:10.1371/journal.pone.0089939.g006




Figure 7. Isobologram analysis for the combinations between BT06 and miltefosine (A) and AB13 and miltefosine (B). The line indicates synergy, additivity or antagonism when the points are located below, on or above the line, respectively. (a) derivative and 2 μM miltefosine; (b) derivative and 4 μM miltefosine; (c) derivative and 6 μM miltefosine. doi:10.1371/journal.pone.0089939.g007

DNA fragmentation assay

Promastigotes of *L. infantum* (2×10^6 cells. ml^{-1}) were exposed to IC_{50} concentration of the most active betulin and betulinic acid

derivatives or to dissolution vehicle (DMSO), and incubated at 26°C for 24 hours. The *Leishmania* DNA extraction was carried out according to the procedure in DNeasy Blood & Tissue (Qiagen).




Figure 8. Optical microscopy observation of *Leishmania infantum* promastigotes in the absence and presence of the betulinic acid derivative AB13. Hanging drop phase contrast (A, B, magnification 200×) and Giemsa staining (C, D, magnification 1000×). Control cells incubated with DMSO at 26°C for 24 h (A, C). Cells exposed to the AB13 derivative at 26°C for 24 h (B, D). doi:10.1371/journal.pone.0089939.g008

Table 2. Flow cytometry analysis of *Leishmania infantum* promastigotes treated with betulin and betulinic acid derivatives (BT06, AB13) showing the percentage of propidium iodide (PI) and annexin-V positive cells.

<i>Leishmania infantum</i> intracellular entities (% of cells)												
	Annexin-V						PI					
	2 h	4 h	6 h	24 h	2 h	4 h	2 h	4 h	6 h	24 h	2 h	4 h
Control (BT06)	3.8	2	1.6	1.5	0.6	1.2	0.6	1.2	0.7	0.9	0.7	0.4
BT06	1.2	2.5	2.6	1.5	0.5	1.4	0.5	0.9	0.9	0.5	0.5	0.6
Control (AB13)	18.8	6.3	3.3	2.4	0.4	0.3	0.4	0.3	0	0.4	3.7	1.5
AB13	17.7	3.4	4.5	2.6	0.1	0	0.1	0	0.2	0	1.9	0.1

doi:10.1371/journal.pone.0089939.t002

DNA integrity analysis was done by electrophoresis, running DNA through an EtBr-treated agarose gel and visualizing it with UV light.

Mammalian cell cytotoxicity

For cytotoxicity assays on mammalian cells, log phase of macrophages (RAW 264.7) were trypsinized and incubated at 37°C in 24-well tissue culture plates in Dulbecco's Modified Eagle Medium (DMEM), enriched with Glutamax and supplemented with 10% fetal bovine serum (FBS), under microaerophilic condition. As soon as the monolayers reached confluence, the medium was removed and the cells were incubated at 37°C for 24 h with fresh medium plus the betulin or betulinic acid derivatives at IC₅₀ concentrations. After incubation, control and treated cells were washed with PBS, pH 7.2., and 450 µl of PBS and 50 µl MTT solution (5 mg.ml⁻¹) were added to each well and incubated at 37°C for 1 h. The cells were then washed with PBS, 500 µl DMSO was added to the wells and absorbance was measured at 530 nm on an ELISA plate reader (Synergy HT, Bio-TEK). The percentage of viable cells was determined as described on viability assay.

Statistical analysis

All experiments were performed in triplicate. The mean and standard deviation (SEM) of three independent assays were determined and statistical analysis between mean values was done by ANOVA test, with a Dunnett's post-test. The significance level was *p<0.05, **p<0.01 and ***p<0.001.

Results and Discussion

This study considers the biological potential of the betulin and betulinic acid derivatives, and is aimed at the evaluation of the anti-*Leishmania* activity of 16 synthetic derivatives of betulin (BT01 to BT09) and betulinic acid (AB10 to AB16). This is followed by a study of the association of the most active derivatives with miltefosine, a drug used in the treatment of leishmaniasis. The activity of the derivatives on *L. infantum* was assessed by cell viability studies. At the highest concentrations, all nine betulin derivatives inhibit promastigote viability (Figure 3A). Generally, the results showed that modifications at C-20 (compounds BT06 and BT07, Figure 1) and at C-30 (compound BT05, Figure 1) have a positive impact on the anti-leishmanial activity of betulin derivatives. The introduction of a carbamate moiety at C-28 in the C-3 substituted derivative BT02 (Figure 1) afforded compound BT03 with lower anti-leishmanial activity (Figure 3A). The betulin derivative BT06 (3β-Hydroxy-(20R)-lupan-29-oxo-28-yl-1H-imidazole-1-carboxylate) was found to be the most effective, with an IC₅₀ of 50.8 µM. The other derivatives did not inhibit the growth of *L. infantum* by 50% at the highest concentration tested (100 µM).

Of the seven betulinic acid derivatives (Figure 3B), the most active compound was AB13 (28-(1H-imidazole-1-yl)-3,28-dioxo-lup-1,20(29)-dien-2-yl-1H-imidazole-1-carboxylate), with an IC₅₀ of 25.8 µM. The remaining betulinic acid derivatives were not able to inhibit viability of *L. infantum* by 50% at the highest concentrations tested (100 µM), however, compounds AB14 with an α,β-unsaturated keto group at ring A (Figure 2), AB15 and AB11 with a keto group and carbamate moiety at C-3 (Figure 2), respectively, showed moderate anti-leishmanial activity. The introduction of the imidazole and methyl ester at C-28 in compounds AB12 and AB16 (Figure 2) respectively, reduce the anti-leishmanial activity (Figure 3B), when compared with derivative AB11 bearing a free carboxylic acid at C-28 position (Figure 2).




Figure 9. Representative cell cycle histograms of *Leishmania infantum*. Promastigotes were incubated at 26°C for 24 h in the absence (A and C) or presence of betulin derivative BT06 (B) and betulinic acid derivative AB13 (D) at IC_{50} concentrations. Propidium iodide staining was performed and samples were analyzed by flow cytometry.
doi:10.1371/journal.pone.0089939.g009

The best activity of AB13 compound, is consistent with published results that report a higher biological activity of betulinic acid derivatives compared with betulin ones [19,31]. The superior anti-*Leishmania* activity of the AB13 derivative could be associated with the higher capacity as a Michael acceptor of this derivative. Considering the Michael addition reaction and according to Sporn and collaborators [36], one possible biological mechanism of action of AB13 may be related to the nucleophilic attack from thiol groups of reduced glutathione or other physiological nucleophiles, including amine nucleophiles, to the α - β -unsaturated-keto group (C1 in ring A) (Figure 2).

The interpretation of results and comparisons between drug activity studies need to take into account the *Leishmania* species and the parasite model cells, i.e., promastigotes or amastigotes. There are only a few reports on the activity of betulin and betulinic acid

derivatives in *Leishmania donovani* amastigotes (IC_{50} values of 8.9 to 30 μ M) [27] and on promastigotes of *L. amazonensis* (IC_{50} of 44.9 μ M to 69.9 μ M) [28]. Comparing these with the present data, we observe that BT06 and AB13 derivatives show antileishmanial activity that is similar or higher than that described for the other triterpenoids. In addition, as well as being active against promastigote cells, BT06 and AB13 are expected to exhibit stronger activity on amastigote forms, as tends to be found among natural extracts and synthetic drugs [37–41].

It is important in the discovery of new drugs that a molecule with anti-*Leishmania* activity does not impart significant toxicity to the host cells. In this study, we have evaluated the toxicity of the most active derivatives in a macrophage cell line (RAW 264.7). We found that BT06 and AB13 derivatives, at IC_{50} concentrations, were not cytotoxic toward the macrophage cell line, suggesting

Table 3. Effects of betulin and betulinic acid derivatives on cellular cycle of *Leishmania infantum* promastigotes.

<i>Leishmania infantum</i> intracellular entities (% of cells)												
	Phase G0/G1				Phase S				Phase G2/M			
	2 h	4 h	6 h	24 h	2 h	4 h	6 h	24 h	2 h	4 h	6 h	24 h
Control (BT06)	83.6	90.7	72	44.2	16.5	9.9	28	33.7	0	0	0	22.1
BT06	92.9	91.8	72.5	67.7	7.1	6.8	28	25.7	0	0	0	6.7
Control (AB13)	75.9	75.3	75.5	48	24.1	24.5	24.5	30	0	0	0	22
AB13	70.6	75.8	60.9	81.2	29.2	24.2	39.1	18.8	0	0	0	0

doi:10.1371/journal.pone.0089939.t003

that they are safe for mammalian cells. Therefore, the two triterpenoids are promising compounds for the discovery of new drugs against *Leishmania* infections.

For each of the two most active derivatives (BT06 and AB13), three different combinations at fixed concentrations of miltefosine (2, 4 and 8 μM) were tested. Dose-response curves showed that the combinations of BT06/miltefosine and AB13/miltefosine were more effective at reducing the viability of promastigotes relative to the derivatives alone (Figure 4). The IC_{50} values obtained for all the tested combinations were less than the values obtained for the pure derivatives (Table 1). The combination of the betulin derivative BT06 with miltefosine at 2 μM and 4 μM induced decreases of the IC_{50} value from 50,8 μM to 30,1 μM and 25,9 μM , respectively. The combination of the betulinic acid derivative AB13 with miltefosine at 2 μM and 4 μM induced decreases of the IC_{50} value from 25,8 μM to 7,6 μM and 6,0 μM , respectively.

The statistical differences between the effects of the combinations of the derivatives BT06 and AB13 with miltefosine and the effects of the derivatives alone were analyzed. The combinations between BT06 and miltefosine (Figure 5) induced a significant reduction ($p < 0.05$, $p < 0.01$ and $p < 0.001$) on promastigote viability relative to the BT06 activity, except for the combination BT06 10 μM with miltefosine 2 μM . The association AB13 and miltefosine is most effective, with high statistical significance ($p < 0.01$ and $p < 0.001$) in all combinations when compared with the effects of the derivative alone (Figure 6). Isobolographic analysis showed that all the interactions of AB13 and BT06 with miltefosine were synergistic, with experimental values (a, b, c) below the theoretical IC_{50} values (line of additivity) (Figure 7). The advantages of combination therapy include increased effectiveness of the drug, reduced dosage, decreased toxicity and a delay or prevention of the onset of drug resistance [3]. The derivatives show a synergistic association with miltefosine, therefore they could be promising molecules for the development of new combination therapies for leishmaniasis.

In this work, we have also looked at the mechanisms of action responsible for the biological activity in *Leishmania* of the most active betulin and betulinic acid derivatives (BT06 and AB13). The effects on morphology, apoptosis/necrosis events, DNA integrity and cell cycle were studied. The morphological modifications were evaluated by optical microscopy using direct examination of live microorganism (hanging drop) and after Giemsa stain. Control cells were very mobile with the characteristic fusiform shape (Figure 8A), while after Giemsa staining it was possible to observe the characteristic shape, long flagellum emerging from the anterior region of the parasite, nucleus and kinetoplast, posterior to the nucleus (Figure 8C). With the BT06 derivative, no significant morphologic alterations were observed (not shown). However, the promastigotes exposed to AB13 suffered significant morphological changes (Figure 8 B,D) compared with the control. These cells showed a marked decrease in mobility, and it was possible to observe changes in the size and shape of the promastigotes, which become smaller and rounder (Figure 8B). The Giemsa stained promastigotes also show alterations in flagellum size (Figure 8D). These effects suggest that the leishmanicidal activity of the AB13 derivative may be associated with changes in the cytoskeletal organization and function, and modification of the activity of mitochondrial bioenergetics.

Cell death by apoptosis/necrosis was assessed by flow cytometry using Annexin V/PI labeling and the study of the DNA integrity. During early apoptosis, phosphatidylserine (PS) is translocated from the cytoplasmic face of the plasma membrane to the external face,

which can be detected using Annexin V. To distinguish apoptotic cell death from necrotic cell death, cells were counterstained with PI, a non-permeable stain with an affinity for nucleic acids, as it selectively enters necrotic cells. Therefore, co-staining of annexin V and PI can differentiate between cells undergoing early apoptosis (annexin V⁺/PI⁻), necrosis or late apoptosis (PI⁺/annexin V⁻) and live cells (PI⁻/annexin V⁻). The *Leishmania* promastigote cells treated during 24 hours with BT06 and AB13 did not show significant differences in annexin V and PI staining compared with the respective controls (Table 2).

The cell cycle analysis was performed by flow cytometry after PI staining of the parasites incubated with betulin derivative BT06 and betulinic acid derivative AB13 for 24 h at IC₅₀ concentrations (Table 3). Figure 9 shows the distribution of cell DNA through cell cycle of parasites in the absence and presence of the BT06 and AB13. After 24 h of incubation, the majority of treated parasite cells were on G0/G1 phase of cell cycle (BT06, 67.7%; AB13, 81.2%), which is the opposite of what occurs in non-treated cells (44.2% and 48%, respectively). Both derivatives promoted retention of *L. infantum* promastigotes in the G0/G1 phase of the cell cycle, suggesting an arrest at this stage of the cycle. We observed a marked reduction in DNA replication and mitosis, with a decrease in the number of cells in G2/M phase and S phase mainly with the derivative AB13. Inhibition of *L. infantum* promastigote proliferation seems to represent a major mechanism of activity of derivatives BT06 and AB13.

It was reported that dihydrobetulinic acid (DHBA), a derivative of betulinic acid, is an excellent inhibitor of *Leishmania* DNA

topoisomerase I and II and induces apoptosis in *L. donovani* [42]. More recently, it was shown that three betulin derivatives (disuccinyl betulin, diglutaryl dihydrobetulin, and disuccinyl dihydrobetulin) inhibit both growth of *Leishmania donovani*, and relaxation activity of the enzyme type IB topoisomerase of the parasite (topoisomerase I; LdTOP1LS) [43]. Considering these potential therapeutic targets in *Leishmania* and based on our results, we suggest that *Leishmania* DNA topoisomerases could be a potential target for BT06 and AB13 derivatives. This hypothesis is also supported in the previous antiproliferative activity of AB13 and BT06 derivatives on cancer cell lines and on their inhibitory properties on DNA topoisomerase [29–31].

In conclusion, the betulin derivative BT06 and betulinic acid derivative AB13 are promising lead compounds which can be used in the discovery of new therapies for *Leishmania* infections, including multidrug treatment schedules with miltefosine.

Acknowledgments

Authors are grateful to José Correia da Costa from Centro de Imunologia e Biologia Parasitária, Instituto Nacional Dr. Ricardo Jorge, Porto for supplying the culture of *Leishmania infantum* Nicolle (zymodeme MON-1).

Author Contributions

Conceived and designed the experiments: MCS JARS. Performed the experiments: MCS RCS RV VA. Analyzed the data: MCS VA JARS. Contributed reagents/materials/analysis tools: MCS MSR JARS. Wrote the paper: MCS RV.

References

- WHO (2010) Control of the leishmaniasis. World Health Organization technical report series (949): xii–xiii, 1–186.
- Croft SL, Barrett MP, Urbina JA (2005) Chemotherapy of trypanosomiasis and leishmaniasis. *Trends Parasitol* 21(11): 508–512.
- Croft SL, Sundar S, Fairlamb AH (2006) Drug Resistance in Leishmaniasis. *Clin Microbiol Rev* 19(1): 111–126.
- Kappagoda S, Singh U, Blackburn BG (2011) Antiparasitic therapy. *Mayo Clin Proc* 86(6): 561–583.
- Guerin PJ, Olliaro P, Sundar S, Boelaert M, Croft SL, et al. (2002) Visceral leishmaniasis: current status of control, diagnosis, and treatment, and a proposed research and development agenda. *Lancet Infect Dis* 2(8): 494–501.
- Gazanion E, Vergnes B, Seveno M, Garcia D, Oury B, et al. (2011) In vitro activity of nicotinamide/antileishmanial drug combinations. *Parasitol Int* 60(1): 19–24.
- Sundar S, Sinha PK, Rai M, Verma DK, Nawin K, et al. (2011) Comparison of short-course multidrug treatment with standard therapy for visceral leishmaniasis in India: an open-label, noninferiority, randomised controlled trial. *Lancet* 377(9764): 477–486.
- Fuertes MA, Nguewa PA, Castilla J, Alonso C, Pérez JM (2008) Anticancer compounds as leishmanicidal drugs: challenges in chemotherapy and future perspectives. *Curr Med Chem* 15:433–439.
- Lee N, Bertholet S, Debrabant A, Muller J, Duncan R, et al. (2002) Programmed cell death in the unicellular protozoan parasite *Leishmania*. *Cell Death Differ* 9(1): 53–64.
- van Zandbergen G, Lüder CGK, Heussler V, Duszenko M (2010) Programmed cell death in unicellular parasites: a prerequisite for sustained infection? *Trends in Parasitology* 26(10): 477–483.
- Moreira W, Leprohon P, Ouellette M (2011) Tolerance to drug-induced cell death favours the acquisition of multidrug resistance in *Leishmania*. *Cell Death Dis* 1;2:e201. doi: 10.1038/cddis.2011.83.
- Sereno D, Holzmüller P, Mangot I, Cuny G, Ouassii A, et al. (2001) Antimonial-mediated DNA fragmentation in *Leishmania infantum* amastigotes. *Antimicrob Agents Chemother* 45(7): 2064–2069.
- Carvalho L, Luque-Ortega JR, López-Martín C, Castans S, Rivas L, et al. (2011) The 8-aminoquinoline analogue sitamaquine causes oxidative stress in *Leishmania donovani* promastigotes by targeting succinate dehydrogenase. *Antimicrob Agents Chemother* 55(9): 4204–4210.
- Sen N, Das BB, Ganguly A, Mukherjee T, Tripathi G, et al. (2004) Camptothecin induced mitochondrial dysfunction leading to programmed cell death in unicellular hemoflagellate *Leishmania donovani*. *Cell Death Differ* 11:924–936.
- Prada CF, Alvarez-Velilla R, Balaña-Fouce R, Prieto C, Calvo-Álvarez E, et al. (2013) Gimnatecan and other camptothecin derivatives poison *Leishmania* DNA-topoisomerase IB leading to a strong leishmanicidal effect. *Biochem Pharmacol* 85(10): 1433–40.
- Croft SL, Coombs GH (2003) Leishmaniasis— current chemotherapy and recent advances in the search for novel drugs. *Trends Parasitol* 19(11): 502–508.
- Paris C, Loiseau PM, Bories C, Bréard J (2004) Miltefosine induces apoptosis-like death in *Leishmania donovani* promastigotes. *Antimicrob Agents Chemother* 48(3): 852–959.
- Baglin I, Mitaine-Offer AC, Nour M, Tan K, Cavé C, et al. (2003) A review of natural and modified betulinic, ursolic and echinocystic acid derivatives as potential antitumor and anti-HIV agents. *Mini Rev Med Chem* 3(6): 525–539.
- Alakurtti S, Mäkelä T, Koskimies S, Yli-Kauhaluoma J (2006) Pharmacological properties of the ubiquitous natural product betulin. *Eur J Pharm Sci* 29(1): 1–13.
- Mukherjee R, Kumar V, Srivastava SK, Agarwal SK, Burman AC (2006) Betulinic acid derivatives as anticancer agents: structure activity relationship. *Anticancer Agents Med Chem* 6(3): 271–9.
- Ryu SY, Oak MH, Yoon SK, Cho DI, Yoo GS, et al. (2000) Anti-allergic and anti-inflammatory triterpenes from the herb of *Prunella vulgaris*. *Planta Med* 66:358–360.
- RajuGautam S, Jachak M (2009) Recent Developments in Inflammatory Natural Products. *Med Res Rev* 29(5): 767–820.
- Cichewicz RH, Kouzi SA (2004) Chemistry, biological activity, and chemotherapeutic potential of betulinic acid for the prevention and treatment of cancer and HIV infection. *Med Res Rev* 24:90–11.
- Xiong J, Kashiwada Y, Chen CH, Qian K, Morris-Natschke SL, et al. (2010) Conjugates of betulin derivatives with AZT as potent anti-HIV agents. *Bioorg Med Chem* 18(17): 6451–69. doi: 10.1016/j.bmc.2010.06.092.
- Yogeeswari P, Sriram D (2005) Betulinic acid and its derivatives: a review on their biological properties. *Curr Med Chem* 12(6): 657–66.
- Filho AAS, Resende DO, Fukui MJ, Santos FF, Pauletti PM, et al. (2009) In vitro antileishmanial, antiplasmodial and cytotoxic activities of phenolics and triterpenoids from *Baccharis dracunculifolia* D. C. (Asteraceae). *Fitoterapia* 80(8): 478–482.
- Alakurtti S, Heiska T, Kiriazis A, Sacerdoti-Sierra N, Jaffe CL, et al. (2010) Synthesis and anti-leishmanial activity of heterocyclic betulin derivatives. *Bioorg Med Chem* 18(4): 1573–1582.
- Gutierrez-Carmona DB, Escalante-Erosa F, Garcia-Sosa K, Ruiz-Pinell G, Dominguez-Yapu D, et al. (2010) Antiprotozoal activity of betulinic acid derivatives. *Phytomedicine* 17(5): 379–82.
- Santos RC, Salvador JAR, Marin S, Cascante M (2009) Novel semisynthetic derivatives of betulin and betulinic acid with cytotoxic activity. *Bioorg Med Chem* 17(17): 6241–6250.
- Santos RC, Salvador JAR, Cortés R, Pachón G, Marin S, et al. (2011) New betulinic acid derivatives induce potent and selective antiproliferative activity

- through cell cycle arrest at the S phase and caspase dependent apoptosis in human cancer cells. *Biochimie* 93(6):1065–1075.
31. Santos RC, Salvador JAR, Marín S, Cascante M, Moreira JN, et al. (2010). Synthesis and structure-activity relationship study of novel cytotoxic carbamate and N-acylheterocyclic bearing derivatives of betulin and betulinic acid. *Bioorg Med Chem* 18(12): 4385–4396.
 32. Denizot F, Lang R (1986) Rapid colorimetric assay for cell growth and survival. Modifications to the tetrazolium dye procedure giving improved sensitivity and reliability. *J Immunol Methods* 89(2): 271–277.
 33. Wagenpfeil S, Treiber U, Lehmer A (2006) Statistical analysis of combined dose effects for experiments with two agents. *Artif Intell Med* 37(1): 65–71.
 34. Darzynkiewicz Z, Juan G, Bedner E (2001) Determining cell cycle stages by flow cytometry. New York: Curr. Protoc. Cell Biol. Chapter 8:Unit 8.4. doi: 10.1002/0471143030.cb0804s01
 35. Vermes I, Haanen C, Steffens-Nakken H, Reutelingsperger CA (1995) Novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *J Immunol Methods* 184:39–51.
 36. Sporn MB, Liby KT, Yore MM, Fu L, Lopchuk JM, et al. (2011) New Synthetic Triterpenoids: Potent Agents for Prevention and Treatment of Tissue Injury Caused by Inflammatory and Oxidative Stress. *J Nat Prod* 74(3): 537–545.
 37. Dutta A, Ghoshal A, Mandal D, Mondal NB, Banerjee S, et al. (2007) Raccemoside A, an anti-leishmanial, water-soluble, natural steroidal saponin, induces programmed cell death in *Leishmania donovani*. *J Med Microbio* 56:1196–1204.
 38. Nakayama H, Desrivot J, Bories C, Franck X, Figadère B, et al. (2007) In vitro and in vivo antileishmanial efficacy of a new nitrilquinoline against *Leishmania donovani*. *Biomed Pharmacother* 61:186–8.
 39. Lakshmi V, Pandey K, Kapil A, Singh N, Samant M, et al. (2007) In vitro and in vivo leishmanicidal activity of *Dysosylum binectariferum* and its fractions against *Leishmania donovani*. *Phytomedicine* 14:36–42.
 40. Santin MR, Dos Santos AO, Nakamura CV, Dias Filho BP, Ferreira IC, et al. (2009) In vitro activity of the essential oil of *Cymbopogon citratus* and its major component (citral) on *Leishmania amazonensis*. *Parasitol Res* 105:1489–96.
 41. Vermeersch M, da Luz RI, Toté K, Timmermans JP, Cos P, et al. (2009) In vitro susceptibilities of *Leishmania donovani* promastigote and amastigote stages to antileishmanial reference drugs: practical relevance of stage-specific differences. *Antimicrob Agents Chemother* 53(9):3855–9.
 42. Chowdhury AR, Mandal S, Goswami A, Ghosh M, Mandal L, et al. (2003) Dihydrobetulinic acid induces apoptosis in *Leishmania donovani* by targeting DNA topoisomerase I and II: implications in antileishmanial therapy. *Mol Med* 9:26–36.
 43. Chowdhury S, Mukherjee T, Sengupta S, Chowdhury SR, Mukhopadhyay S, et al. (2011) Novel Betulin Derivatives as Antileishmanial Agents with Mode of Action Targeting Type IB DNA Topoisomerase. *Mol Pharmacol* 80:694–703.