



Published in final edited form as:

J Hepatol. 2013 August ; 59(2): 387–388. doi:10.1016/j.jhep.2013.01.035.

ALCOHOL AND TOXICITY

Ivan Rusyn¹ and Ramon Bataller^{2,3}

¹Department of Environmental Sciences and Engineering, University of North Carolina at Chapel Hill Chapel Hill, NC, USA

²Department of Medicine, University of North Carolina at Chapel Hill Chapel Hill, NC, USA

³Department of Nutrition, University of North Carolina at Chapel Hill Chapel Hill, NC, USA

Excessive alcohol intake is a major public health challenge worldwide and has been identified as one of the main determinants of a variety of non-communicable diseases [1]. The World Health Organization (WHO) estimates that 4.5% of the global burden of disease and injury, and 4% of all deaths worldwide are attributable to alcohol [2]. Alcohol is the leading risk factor for death among males aged 15-59, particularly in Eastern Europe [2, 3].

Toxic and other adverse effects of alcohol on organs and tissues in humans (Figure 1) are largely a consequence of its metabolism to acetaldehyde, and associated formation of reactive oxygen and nitrogen species, depletion of co-factors (e.g., NAD⁺), and impairment in energy homeostasis [4]. Because of the considerable redundancy in the oxidative enzymatic pathways (alcohol dehydrogenases, CYP2E1 and catalase) that can convert alcohol to acetaldehyde, most tissues are capable of alcohol metabolism, even though liver is the primary site. Likewise, acetaldehyde dehydrogenases are ubiquitous in the mitochondria. A minor, non-oxidative, pathway of alcohol metabolism is to fatty acid ethyl ester (via fatty acid ethyl ester synthase) and phosphatidyl ethanol (via phospholipase D). Alcohol impacts the integrity of the gastrointestinal mucosal barrier. This results in the translocation, via the portal blood flow, of the gut bacteria-derived lipopolysaccharide (endotoxin) and other molecules to the liver and activation of the innate immune response.

The molecular and cellular sequelae of the toxic mediators of alcoholic injury take many forms. Acetaldehyde and oxidants are highly reactive molecules that can damage DNA, proteins and lipids. Changes in hepatic respiration and lipid metabolism lead to tissue hypoxia and impairment in the mitochondrial function. Secondary effects include disruption of signaling pathways and ion channel function, unfolded-protein response and oxidative stress, as well as activation of adaptive immune response largely triggered by acetaldehyde-protein adducts. Cell death triggers additional innate immune response, activation of fibrogenesis, and tissue repair. In addition to pro-inflammatory mediators, other signaling molecules, such as neurotransmitters, are affected by alcohol.

Gross pathological changes associated with alcohol drinking include most or all, depending on the affected tissue, of the following: fat accumulation (steatosis), inflammation, necrosis and fibrosis [5]. Impairment of organ function and carcinogenesis are most terminal pathological effects. As a consequence, alcoholic beverages and ethanol in alcoholic beverages are classified by the WHO International Agency for Research on Cancer as “carcinogenic to humans” (Group 1) [6].

Alcohol is highly diffusible through cell membranes and is metabolized by most tissues. Thus, its toxicity affects most organs. Because liver is the major site of alcohol metabolism, it is one of the major targets for alcohol-induced organ damage. Alcoholic liver diseases include steatosis, different subtypes of steatohepatitis, cirrhosis and hepatocellular carcinoma [5]. Of these, cirrhosis of the liver is a third leading (at 16.6%) cause of alcohol-attributable deaths worldwide [2]. In addition, other anatomical sites in the aero-digestive tract are adversely affected by alcohol, with most important morbidity and mortality due to malignant tumors, all causally related to alcohol consumption, of the oral cavity, pharynx, larynx, esophagus and colorectum [6]. In the pancreas, toxic metabolites of alcohol cause acinar cell injury leading to pancreatitis and subsequent fibrosis [7]. Cardiovascular system is second only to gastrointestinal organs in toxic effects of alcohol [1, 2]. Hypertension, ischemic heart disease, stroke, cardiomyopathy and myocarditis, as well as various arrhythmias, have been associated with alcohol abuse. The linkages between alcohol use and neuropsychiatric disorders have also been established as causal (e.g., for major depression) [8]. In addition, neurobehavioral impairment due to alcohol intoxication is a major contributor to deaths from violence, road traffic accidents and injuries [2]. In the kidney, alcohol has been associated with glomerulonephritis, acute nephropathy and kidney graft failure [9]. Adverse effects of alcohol on human reproduction and development span the wide range of pathological conditions from impaired fertility, to premature and low-weight births, and fetal alcohol syndrome spectrum disorders [2]. The addition of breast cancer to the list of cancers causally related to alcohol consumption suggested that the proportion of malignancies attributable to alcohol consumption is higher than previously estimated [6]. Indeed, recent studies show that up to 5% of breast cancers are attributable to alcohol in northern Europe and North America for a total of approximately 50,000 alcohol-attributable cases of breast cancer worldwide [10].

Factors affecting susceptibility to alcohol toxicity include genetics, gender, lifestyle/nutrition, exposure to environmental chemicals and drugs, and co-morbidities. Variations in genes encoding metabolic pathways for alcohol or triglycerides, as well as many other genes that may be involved in the pathogenesis or protection from alcohol-induced toxicity, modify the individual's response to alcohol abuse and disease outcomes [4-6]. The effect of gender varies by disease outcome with females being more susceptible to alcoholic liver disease, but most alcohol-attributable deaths occur in males [1, 2]. Dietary (e.g., high fat consumption, patterns of alcohol intake), life style (e.g., cigarette smoking, drugs of abuse) and environmental (e.g., nitrosamines, metals and chlorinated solvents) factors also affect both morbidity and mortality related to alcohol abuse. Obesity, viral infections, iron accumulation and other pro-inflammatory conditions, as well as concomitant uptake of certain drugs (e.g., acetaminophen, isoniazid and methotrexate) also compound morbidity and mortality due to alcohol abuse.

Finally, while overwhelming evidence exists to conclude that consumption of alcoholic beverages is harmful to human health, many studies observed some beneficial effects of modest alcohol consumption (usually defined as 1-2 servings/day) [1]. Health conditions that have been associated with "beneficial effects" of alcohol on mortality include cardiovascular diseases (e.g., myocardial infarction), diabetes and chronic kidney disease [1, 9], yet such protection disappears with heavy drinking occasions [2]. No "safe" amount of alcohol consumption has been found for cancer; even consumption of two drinks per day causes a significant increase in risk [6].

References

- [1]. Parry CD, Patra J, Rehm J. Alcohol consumption and non-communicable diseases: epidemiology and policy implications. *Addiction*. 2011; 106:1718–1724. [PubMed: 21819471]

- [2]. World Health Organization. Global status report on alcohol and health. WHO Press; Geneva, Switzerland: 2011.
- [3]. Zaridze D, Brennan P, Boreham J, Boroda A, Karpov R, Lazarev A, et al. Alcohol and cause-specific mortality in Russia: a retrospective case-control study of 48,557 adult deaths. *Lancet*. 2009; 373:2201–2214. [PubMed: 19560602]
- [4]. Zakhari S. Overview: how is alcohol metabolized by the body? *Alcohol Res Health*. 2006; 29:245–254. [PubMed: 17718403]
- [5]. Gao B, Bataller R. Alcoholic liver disease: pathogenesis and new therapeutic targets. *Gastroenterology*. 2011; 141:1572–1585. [PubMed: 21920463]
- [6]. Baan R, Straif K, Grosse Y, Secretan B, el Ghissassi F, Bouvard V, et al. Carcinogenicity of alcoholic beverages. *Lancet Oncol*. 2007; 8:292–293. [PubMed: 17431955]
- [7]. Apte MV, Pirola RC, Wilson JS. Mechanisms of alcoholic pancreatitis. *Journal of gastroenterology and hepatology*. 2010; 25:1816–1826. [PubMed: 21091991]
- [8]. Boden JM, Fergusson DM. Alcohol and depression. *Addiction*. 2011; 106:906–914. [PubMed: 21382111]
- [9]. Schaeffner E, Ritz E. Alcohol and kidney damage: a Janus-faced relationship. *Kidney Int*. 2012; 81:816–818. [PubMed: 22377829]
- [10]. Seitz HK, Pelucchi C, Bagnardi V, La Vecchia C. Epidemiology and pathophysiology of alcohol and breast cancer: Update 2012. *Alcohol Alcohol*. 2012; 47:204–212. [PubMed: 22459019]




Figure 1. Layered graph showing the mechanisms of alcohol-induced toxicity and organ damage
 Layer #1: the amount of *alcohol intake* is a main determinant of toxicity. Layer #2: *Alcohol metabolism* is regulated by several enzymes and is a main determinant of alcohol-induced toxicity. Genetic variations and expression of these enzymes regulate systemic and local effects of alcohol intake. Layer #3: Alcohol metabolites and molecules released in organs damaged by alcohol such as acetaldehyde and ROS are key *toxicity mediators* with powerful biological properties. Layer #4. Such mediators activate several *cellular and molecular mechanisms* such as disrupted lipid metabolism, hypoxia, ER stress, dysregulated immunity, changes in intestinal microbiota and DNA damage. Layer #5. The synergistic effect of the activation of these pathways leads to different histological disturbances in target tissues such

as fat accumulation, inflammation (PMN cells), necrosis/apoptosis, fibrosis and cancer, leading to organ dysfunction. Layer #6. The most susceptible organs to the deleterious effects of alcohol, that account for most of clinical complications, include normal development of the fetus, liver, the kidney, the nervous system, the aero-digestive tract the reproductive system, the pancreas and the cardiovascular system. Layer #7. The individual susceptibility to the toxic effects of alcohol in the human body is determined by genetic (gender, SNPs in target genes) dietary and environmental exposures. Finally, patients with co-morbidities such as viral infections of metabolic disorders are more susceptible to the deleterious effects of alcohol abuse.

EtOH: ethanol; ADH: alcohol dehydrogenase; ALDH: acetaldehyde dehydrogenase; ER: endoplasmic reticulum; SNPs: single nucleotide polymorphism.