

## LETTER

# The gain of hydrogen peroxide resistance benefits growth fitness in mycobacteria under stress

### Dear Editor,

Tuberculosis (TB) is a formidable challenge to global health and is caused by an ancient pathogen, *Mycobacterium tuberculosis* (WHO, 2010). In the pre-antibiotic era, TB killed 20% of adults in Europe and North America in the period from the seventeenth to the nineteenth centuries (Wilson, 2005). Since antibiotics have been included in TB treatment regimens, millions of human lives have been saved by this chemotherapy. However, the standard anti-tuberculosis therapy is only effective for drug-susceptible TB but not for multi-drug resistant and extensively drug-resistant TB. Unfortunately, our knowledge of drug resistance in mycobacteria remains unclear and thus hampers the development of new anti-tubercular drugs. The basic mechanisms of drug resistance in mycobacteria are complicated and the outcome of drug resistance stems from evolution in the final analysis. Understanding the evolutionary strategies used by *M. tuberculosis* might give new insights into designing control strategies against *M. tuberculosis* (Little et al., 2012).


As the most successful bacterial pathogen, *M. tuberculosis* has evolved elegant strategies to evade a host's immune defensive system. For example, D'Arcy Hart and colleagues observed that *M. tuberculosis*-containing phagosomes do not fuse with the lysosome inside murine macrophages (Armstrong and Hart, 1971). Several studies indicated that *M. tuberculosis* succeeds as an intracellular pathogen is partially due to its ability to survive and persist inside macrophages, which provide hostile environments with starvation and other stresses (Podinovskaia et al., 2013). When *M. tuberculosis* infects macrophages, the mycobacteria must overcome exogenous reactive oxygen species (ROS), one of the classical innate defense mechanisms against infection. Additionally, during latency, *M. tuberculosis* continues to be under oxidative stress; thus, the accumulation of mutations caused by oxidative DNA damage was predicted to be a potential risk for drug resistance (Ford et al., 2011). Recently, studies have shown a correlation of ROS damage of DNA with antibiotic resistance (Kohanski et al., 2010), although this notion has been disputed by other recent studies that argued that antibiotic exposure did not induce the formation of ROS (Keren et al., 2013). The role of ROS in the evolution of drug resistance is still an open question.

Currently, the view of evolution is mostly derived from genomic sequencing and lacks direct experimental evidence. In this study, to explore the correlation between hydrogen peroxide ( $H_2O_2$ ) resistance and growth benefits in mycobacteria under stress, we developed a model for evolutionary screening of  $H_2O_2$ -resistant mutations in a pathogenic mycobacteria model, *M. smegmatis*, using a clinically important stressor,  $H_2O_2$  (Fig. 1). This strategy of screening for  $H_2O_2$ -resistant mutations under increased  $H_2O_2$  conditions in *M. smegmatis* was chosen to allow genome-wide identification of essential genes involved in oxidative stress because setting up a library of transposon mutants would fail, by default, to identify candidates that are essential genes. To demonstrate that resistance phenotypes resulted from chromosomal mutations, we continuously sub-cultured resistant strains for 10 generations without  $H_2O_2$  treatment. The actual minimum inhibitory concentrations (MICs) of  $H_2O_2$  in different mutants selected under different  $H_2O_2$  concentrations were different, indicating that gradually increasing  $H_2O_2$  resistance was due to the accumulation of mutated genes rather than a key mutated gene that caused the resistance to  $H_2O_2$  (Fig. S1). After 54 days of selection, the MIC of the mutant strains with the maximum resistance to  $H_2O_2$ , selected at an  $H_2O_2$  concentration of 1.5 mmol/L, was 3.125 mmol/L, while that of the wild type strain mc<sup>2</sup>155 under the same conditions (7H9 medium with 10% ADS) was 0.039 mmol/L. We named the mutant strain mc<sup>2</sup>51. As evolutionary cross-stress protection was observed under various stresses, we compared the growth advantage between mc<sup>2</sup>51 and mc<sup>2</sup>155 in corresponding stresses. We first measured the doubling time of both wild type mc<sup>2</sup>155 and mutant strain mc<sup>2</sup>51. A significant difference in doubling time was detected: the doubling time of wild type mc<sup>2</sup>155 was  $2.734 \pm 0.051$  ( $n = 30$ ), while the doubling time of mc<sup>2</sup>51 was  $3.448 \pm 0.087$  ( $n = 30$ ) (Fig. 2B), and the generation time increased up to 126%, compared to wild type mc<sup>2</sup>155. A growth advantage was observed for mutant strain mc<sup>2</sup>51 in M9 minimal medium compared to wild type mc<sup>2</sup>155, but this did not occur in M9 minimal medium with 10% ADS, suggesting that the mutant strain has a growth advantage under starvation conditions. Additionally, to investigate whether mc<sup>2</sup>51 has a survival advantage in


host cells, cell line THP1 was infected with either mutant strain mc<sup>2</sup>51 or wild type strain mc<sup>2</sup>155 at an MOI of 1:10. The mutant strain mc<sup>2</sup>51 had a statistically significant growth advantage in macrophages compared to wild-type mc<sup>2</sup>155 (Fig. 2D). As a correlation of ROS damage and antibiotic resistance was indicated, we also detected the MICs of various antibiotics in both mc<sup>2</sup>155 and mc<sup>2</sup>51. Previous studies have suggested that H<sub>2</sub>O<sub>2</sub> affected *KatG* mutation frequency and caused changes in resistance to isoniazid (INH) (Timmins and Deretic, 2006). A 100-fold difference in the MIC of INH was detected between mc<sup>2</sup>155 and mc<sup>2</sup>51 (Table S1). To provide insight into the mechanisms involved, we sequenced the whole genomes of both mutant strain mc<sup>2</sup>51 and wild type mc<sup>2</sup>155. Compared with mc<sup>2</sup>155, there are 22 single nucleotide polymorphisms (SNPs) in the mc<sup>2</sup>51 genome, of which 19 are located in coding regions, and 12 of these 19 SNPs are nonsynonymous mutations (Table S2). The data showed that G:C > T:A transversion and G:C > A:T transition were the major mutation types (Fig. 2E). The ratio of G:C > T:A transversion to G:C > A:T transition is an index to measure the contribution of oxidative DNA damage to the development of antibiotic resistance; it was 0.7 in mc<sup>2</sup>51 but is 0.25 under normal growth conditions. This increased ratio indicated that oxidative DNA damage caused the DNA mutations of mc<sup>2</sup>51. It should be noted that the ratio for mutations of *M. tuberculosis* isolated from macaques with latent infection increased drastically to 0.67 (Ford et al., 2011), indicating that oxidative DNA damage is a dominant source of mutation during both active and latent phases of *M. tuberculosis*. To further determine the possible roles of those genes in H<sub>2</sub>O<sub>2</sub> resistance, the sequences of the corresponding genes amplified from wild type mc<sup>2</sup>155 genomic DNA were cloned into the

integrating vector pMV361 (Table S4) and introduced into mc<sup>2</sup>51 via electroporation. Surprisingly, there are no changes in the MIC of H<sub>2</sub>O<sub>2</sub> for most detected transformed strains, except for one strain of mc<sup>2</sup>51 with pMV361-*fur* (*msmeg\_3460*) (Table S3 and data not shown). Furthermore, because *fur* is located upstream of *msmeg\_3461*, which encodes *KatG*, the mutated *Fur* (Ferric uptake regulation protein) with substituted A28 V would influence *KatG* expression. Increased peroxidase and catalase activities were found in mc<sup>2</sup>51 (Fig. 2F), and *KatG* was confirmed by mass spectrometry analysis (Fig. 2F).

As an intracellular pathogen, the success of *M. tuberculosis* is partially due to its ability to survive and persist in macrophages. For example, ROS produced by the host is one of the most important classical innate defense mechanisms against infection. To defend against ROS stress, mycobacteria have evolved many detoxification strategies to scavenge H<sub>2</sub>O<sub>2</sub> (Kumar et al., 2011). Several studies have suggested that accumulation of mutations might link to the emerging drug resistance. In this study, we first selected an H<sub>2</sub>O<sub>2</sub>-resistant strain, mc<sup>2</sup>51, which displayed an 80-fold higher MIC of H<sub>2</sub>O<sub>2</sub> than that in wild type mc<sup>2</sup>155. The mutant strain mc<sup>2</sup>51 has a growth advantage under starvation conditions and within host macrophage cells, but no resistance or susceptibility to the many investigated antibiotics except INH. To characterize the correlation of SNPs in mc<sup>2</sup>51 with the adaptive evolution of *M. smegmatis* resistance to H<sub>2</sub>O<sub>2</sub>, we sequenced the genome of mc<sup>2</sup>51 and showed that the SNP in *fur* contributed the observed resistance to H<sub>2</sub>O<sub>2</sub> (Table S3). Furthermore, the mass spectrometry results indicated that the mutation of *fur* changed the expression of *katG*, causing high resistance to H<sub>2</sub>O<sub>2</sub>. Our



**Figure 1. Overview of the experimental set-up.** *M. smegmatis* mc<sup>2</sup>155 evolved for 25 generations under increasing H<sub>2</sub>O<sub>2</sub> in 7H9 medium supplemented with 10% ADS (50 g/L albumin, 20 g/L dextrose and 8.1 g/L saline). The growth fitness of the final evolved strain was determined by doubling time, growth curve, starvation growth, survival under H<sub>2</sub>O<sub>2</sub> stress, antibiotic resistance, and within macrophage-like cell line THP-1. The genomic DNA of both wild type mc<sup>2</sup>155 and mutant strains mc<sup>2</sup>51 was sequenced to identify single nucleotide polymorphisms (SNPs) in the mutant strain.



**Figure 2. Characterization of H<sub>2</sub>O<sub>2</sub> mutant strain mc<sup>251</sup>.** (A) The concentrations of H<sub>2</sub>O<sub>2</sub> corresponding to the selected date throughout the whole selection process. (B) The H<sub>2</sub>O<sub>2</sub>-resistant strain mc<sup>251</sup> showed reduced growth compared to wild type mc<sup>2155</sup>. The growth rate is indicated by doubling time, which was measured using a fluctuation assay. (C) Growth advantage was determined between mc<sup>2155</sup> and mc<sup>251</sup>. The three panels represent serial dilutions (1:10) of mc<sup>2155</sup> and mc<sup>251</sup> on M9 minimal medium or M9 minimal medium supplemented with 10% ADS, respectively; cultures were spotted (5  $\mu$ L) onto the indicated medium. Photographs were taken after 3-day incubation. (D) The H<sub>2</sub>O<sub>2</sub>-resistant mutant strain mc<sup>251</sup> has a survival advantage over wild type mc<sup>2155</sup> in the macrophage-like cell line THP-1. (E) The proportion of polymorphisms of nucleotide pair mutations from single nucleotide polymorphisms (SNPs). A total of 22 mutations were identified in mc<sup>251</sup> compared to wild type mc<sup>2155</sup>. G:C > T:A transversion and G:C > A:T transition are the major mutation types. (F) The activities of peroxidase and catalase in mc<sup>251</sup> and mc<sup>2155</sup> and the identification of KatG using mass spectrometry. Whole lysates including approximately 50  $\mu$ g protein were separated by non-denaturing 7.4% polyacrylamide gels. By searching the *M. smegmatis* protein database, an MS/MS spectrum matched GPDTITSGLEVVTNTPTK. The highlighted peak is specific to the sequence of KatG. The underlined peptides were identified by MS/MS searching of the band covering 68% of the KatG sequence.

results showed that the development of H<sub>2</sub>O<sub>2</sub> resistance in mycobacteria led to the gain of growth benefits in starvation conditions and within macrophages.

The ratio of G:C > T:A transversion to G:C > A:T transition mutation is used as an index to measure the contribution of oxidative DNA damage to the development of antibiotic resistance (Hershberg et al., 2008). If the oxidative DNA damage contributes to DNA mutation, this ratio will increase significantly. In this study, we calculated the ratio as 0.78, which indicated that oxidative DNA damage is a dominant source of mutation for mycobacteria (Fig. 2E). A similar ratio (0.67) was also reported in mutations of *M. tuberculosis* isolated from macaques with latent infection (Ford et al., 2011). Additionally, previous studies have shown that the

ratio of G:C > T:A transversion to G:C > A:T transition that confers resistance to anti-TB drugs is 0.71 (Wang et al., 2013), which is similar to our results (0.78). However, in our study, the MICs of various antibiotics did not differ between wild type mc<sup>2155</sup> and mc<sup>251</sup> except that of INH, a prodrug that needs bacterial ROS activation. The results suggested that the mutation ratio is an effect rather than a cause for the acquisition of antibiotic resistance.

As ROS is correlated with *M. tuberculosis* pathogenesis, we explored whether mycobacteria obtain any benefits when evolving to adjust their resistance to H<sub>2</sub>O<sub>2</sub>. Our results showed that endogenous H<sub>2</sub>O<sub>2</sub> resistance confers bacterial growth advantages. Newly identified genes, such as *carD*, *moaD1*, and *pp54*, involved in oxidative stress, have been shown to be

important for bacterial survival inside host cells (Mestre et al., 2013). A correlation has also been shown between resistance to oxidative stress and survival in macrophages. In line with this hypothesis, our results showed that H<sub>2</sub>O<sub>2</sub>-resistant strain mc<sup>2</sup>51 had a statistically significant growth advantage in macrophages compared with wild-type mc<sup>2</sup>155 (Fig. 2D). Consistent with a previous study on the relationship of Fur to INH and KatG, the resistance of H<sub>2</sub>O<sub>2</sub> was partially abrogated when wild type Fur was expressed in mc<sup>2</sup>155 (Table S3). In addition, MS data showed that mc<sup>2</sup>51 had increased enzyme activity of KatG (Fig. 2F). Moreover, the SNPs in Fur were also identified in INH-resistant isolates (Ramaswamy et al., 2003). This finding implies that mutant Fur could be produced in humans. Thus, these findings support our hypothesis that the evolutionary adaptation of mycobacteria to H<sub>2</sub>O<sub>2</sub>, accompanied by slow growth, confers growth benefits within macrophages on the mycobacteria.

## FOOTNOTES

This work was supported by the National Basic Research Program (973 Program) (Nos. 2014CB744402 and 2012CB518700), the National Natural Science Foundation of China (Grant Nos. 31270178 and 31070118), and the Key Program of the Chinese Academy of Sciences (KJZD-EW-L02).

Xiaojing Li, Jun Tao, Jiao Han, Xinling Hu, Yuling Chen, Haiteng Deng, Guangchun Zhang, Xiaoxiang Hu, Kaixia Mi declare that they have no conflicts of interest. This article does not contain any studies with human or animal subjects performed by any of the authors.

Xiaojing Li<sup>1,2</sup>, Jun Tao<sup>1</sup>, Jiao Han<sup>1</sup>, Xinling Hu<sup>1</sup>, Yuling Chen<sup>3</sup>, Haiteng Deng<sup>3</sup>, Guangchun Zhang<sup>4</sup>, Xiaoxiang Hu<sup>4</sup>, Kaixia Mi<sup>1,5</sup>✉

<sup>1</sup> CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of Microbiology, CAS, Beijing 100101, China

<sup>2</sup> University of Chinese Academy of Sciences, Beijing 100082, China

<sup>3</sup> School of Life Sciences, Tsinghua University, Beijing 100084, China

<sup>4</sup> School of Life Sciences, China Agriculture University, Beijing 100094, China

<sup>5</sup> Beijing Key Laboratory of Drug Resistance and Resistome, Beijing 100101, China

✉ Correspondence: mik@im.ac.cn (K. Mi)

## OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and

**Electronic supplementary material** The online version of this article (doi:10.1007/s13238-014-0024-5) contains supplementary material, which is available to authorized users.

Xiaojing Li and Jun Tao have contributed to this work equally.

reproduction in any medium, provided the original author(s) and the source are credited.

## REFERENCES

- Armstrong JA, Hart PD (1971) Response of cultured macrophages to *Mycobacterium tuberculosis*, with observations on fusion of lysosomes with phagosomes. *J Exp Med* 134:713–740
- Balbi KJ, Rocha EP, Feil EJ (2009) The temporal dynamics of slightly deleterious mutations in *Escherichia coli* and *Shigella* spp. *Mol Biol Evol* 26:345–355
- Ford CB, Lin PL, Chase MR, Shah RR, Iartchouk O, Galagan J, Mohaideen N, Ioerger TR, Sacchettini JC, Lipsitch M et al (2011) Use of whole genome sequencing to estimate the mutation rate of *Mycobacterium tuberculosis* during latent infection. *Nat Genet* 43:482–486
- Hershberg R, Lipatov M, Small PM, Sheffer H, Niemann S, Homolka S, Roach JC, Kremer K, Petrov DA, Feldman MW et al (2008) High functional diversity in *Mycobacterium tuberculosis* driven by genetic drift and human demography. *PLoS Biol* 6:e311
- Keren I, Wu Y, Inocencio J, Mulcahy LR, Lewis K (2013) Killing by bactericidal antibiotics does not depend on reactive oxygen species. *Science* 339:1213–1216
- Kohanski MA, DePristo MA, Collins JJ (2010) Sublethal antibiotic treatment leads to multidrug resistance via radical-induced mutagenesis. *Mol Cell* 37:311–320
- Kumar A, Farhana A, Guidry L, Saini V, Hondalus M, Steyn AJ (2011) Redox homeostasis in mycobacteria: the key to tuberculosis control? *Expert Rev Mol Med* 13:e39
- Little TJ, Allen JE, Babayan SA, Matthews KR, Colegrave N (2012) Harnessing evolutionary biology to combat infectious disease. *Nat Med* 18:217–220
- Mestre O, Hurtado-Ortiz R, Dos Vultos T, Namouchi A, Cimino M, Pimentel M, Neyrolles O, Gicquel B (2013) High throughput phenotypic selection of *Mycobacterium tuberculosis* mutants with impaired resistance to reactive oxygen species identifies genes important for intracellular growth. *PLoS One* 8:e53486
- Podinovskaia M, Lee W, Caldwell S, Russell DG (2013) Infection of macrophages with *Mycobacterium tuberculosis* induces global modifications to phagosomal function. *Cell Microbiol* 15:843–859
- Ramaswamy SV, Reich R, Dou S-J, Jasperse L, Pan X, Wanger A, Quitugua T, Graviss EA (2003) Single nucleotide polymorphisms in genes associated with isoniazid resistance in *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 47:1241–1250
- Timmins GS, Deretic V (2006) Mechanisms of action of isoniazid. *Mol Microbiol* 62:1220–1227
- Wang ZY, Xiong M, Fu LY, Zhang HY (2013) Oxidative DNA damage is important to the evolution of antibiotic resistance: evidence of mutation bias and its medicinal implications. *J Biomol Struct Dyn* 31:729–733
- WHO (2010). Multidrug and extensively drug-resistant TB (M/XDR-TB): 2010 global report on surveillance and response. WHO/HTM/TB/20103
- Wilson LG (2005) Commentary: medicine, population, and tuberculosis. *Int J Epidemiol* 34:521–524