

The Sites of Photoconversion of Protochlorophyllide to Chlorophyllide in Barley Seedlings¹

Received for publication December 28, 1970

ŞEFİK SÜZER AND KENNETH SAUER

Department of Chemistry and Laboratory of Chemical Biodynamics, University of California, Berkeley, California 94720

ABSTRACT

The photoreduction of protochlorophyllide *a* to chlorophyllide *a* in intact 6-day-old seedlings of etiolated barley (*Hordeum vulgare*) exhibits a small initial phase, followed by an induction period of about 1 hour before a rapid phase of additional chlorophyll formation begins. Cycloheximide, an inhibitor of protein synthesis, has no effect on the initial phase of conversion of preformed protochlorophyllide, but it either abolishes or severely inhibits the subsequent phase of rapid chlorophyll synthesis within 45 minutes of its application to the seedlings. An analysis of the biphasic inhibition process suggests that the lifetime of the enzyme controlling protochlorophyllide synthesis (probably δ -amino-levulinic acid synthetase) is not longer than 10 minutes.

The rapid phase of chlorophyll formation can be effected by a series of brief (15 second) pulses of light spaced at least 5 minutes apart. When longer dark intervals are used, no increase is observed in the yield of chlorophyll per pulse. We interpret the findings to indicate that the photoconversion takes place at distinct enzymatic sites whose concentration does not increase during a period of 4 hours following the initial illumination. The sites can be used repeatedly with a turnover time determined by the removal of the product chlorophyllide and the synthesis and placement of a new protochlorophyllide molecule.

brief flash of activating light (16, 17) or, albeit slowly, by illumination at -80°C (10). In this paper we present evidence that the number of photoconversion sites in a greening seedling does not change significantly during the first 4 hr following the initial illumination. Within this interval the photoconversion sites can be used more than 20 times in succession.

Following an initial conversion of active protochlorophyllide present in dark-grown seedlings, there commonly follows an induction period of 1 hr or more preceding a rapid phase of further chlorophyll synthesis (1, 33). During the induction period the synthesis of protochlorophyllide is limiting, and this limitation can be overcome by feeding the plants with δ -aminolevulinic acid, a precursor of protochlorophyllide (25). Studies using chloramphenicol (18, 19) or cycloheximide (13), which are known to be inhibitors of protein synthesis, indicate that active protein synthesis is normally required in order for protochlorophyllide to be formed and sited on the enzyme where photoconversion to chlorophyllide takes place (13, 20, 26, 28). The action of these and other inhibitors of chlorophyll synthesis has recently been reviewed (14, 27).

The studies of the action of cycloheximide on etiolated barley seedlings described in this paper confirm that the inhibitor does not prevent the initial conversion of preformed protochlorophyllide (20), even when the seedling is exposed to the inhibitor for as long as 48 hr prior to the first illumination. On the other hand, it is highly effective in abolishing chlorophyll synthesis during the subsequent postinduction phase. During this period it acts as soon as 45 min following its application to seedling tips.

MATERIALS AND METHODS

Barley (*Hordeum vulgare*, var. Atlas) seeds were germinated and grown in complete darkness in distilled water on cotton and filter paper. Water was added every 2nd day. The leaves were harvested under a dim green safelight when they were 6 days old, and the upper 3 cm was used in the studies to be described. For spectrometric measurements the tightly curled leaves were flattened between two microscope slides.

Absorption spectra of the mounted leaf segments were measured by placing them directly in front of the photomultiplier of a Unicam SP800 UV-Visible spectrophotometer. The large photosensitive surface of the end-window photomultiplier enabled reasonably sharp spectra to be obtained for the leaves, despite their pronounced light scattering. The amounts of protochlorophyllide and chlorophyll(ide) were estimated from the magnitudes of the absorbance peaks at 650 and 683 (678) nm, respectively.

Solutions of cycloheximide (0.5 mg/ml; Upjohn, Kalamazoo, Mich.) and chloroamphenicol (0.5 mg/ml; as sodium succinate salt, Lepetit) in water were prepared before the start of each experiment and stored at 0 C until used. The inhibitors were ap-

The penultimate stage in the synthesis of chlorophyll *a* in most higher plants is a photochemical step in which two protons are added stereospecifically to the porphin ring system of protochlorophyllide to form chlorophyllide *a*. The initial photoconversion can be monitored spectrophotometrically in intact etiolated leaves (22). There follows a series of dark spectral shifts, during which the chlorophyllide is esterified by phytol to form chlorophyll *a* (24, 34). These reactions have the characteristics of an enzymatic process even apart from the stereospecific nature of the product: they are abolished by mild heating (52 C for 10 min) (23, 29), grinding the leaf tissues with sand and buffer (5), application of a freeze-thaw cycle (9, 29), and extraction of the pigments by organic solvents. Nevertheless, the initial photochemical stage can be completed within a few milliseconds by a

¹The work described in this report was carried out in the Department of Chemistry, Middle East Technical University, Ankara, Turkey.

plied by floating the leaf segments on 10 ml of the solution in covered Petri dishes.

Continuous illumination was provided by a bank of five fluorescent lamps (General Electric, 20 w, 820 lumens, daylight) at a distance of 30 cm from the samples. Brief pulses of illumination were provided by a hand torch held 15 cm from the samples and providing an incident intensity of about 25 ft-c. A duration of 15 sec was sufficient to saturate short term protochlorophyllide conversion. To provide reproducibility for the spectral measurements, the leaves were illuminated in the sample compartment of the spectrophotometer and were left *in situ* during the dark intervals between illumination pulses.

RESULTS

Inhibition of Chlorophyll Formation by Cycloheximide. In preliminary experiments using equal concentrations (0.5 mg/ml) of cycloheximide or of chloramphenicol, the cycloheximide was found to be the more potent inhibitor of chlorophyll formation during extended illumination of etiolated barley seedlings. For this reason cycloheximide was chosen for the experiments which follow.

The normal course of protochlorophyllide to chlorophyll conversion, in the absence of any inhibitor, produced a small initial absorption peak at about 678 nm upon illumination with white light. Associated with a 15-sec illumination with either weak (25 ft-c) or strong (2000 ft-c) light were the following absorbance changes: $\Delta A_{678} = +0.02 \pm 0.005$ and $\Delta A_{650} = -0.01 \pm 0.005$ at the chlorophyll and protochlorophyllide maxima, respectively. Following a lag period of about 1 hr, a rapid phase of chlorophyll synthesis began, and after 6 hr of strong illumination $\Delta A_{678} = 0.72$ was obtained.

The effect of the time of cycloheximide application was studied using a regimen in which leaves were exposed to cycloheximide at different times (0 to 48 hr) before the first illumination. Following a 6-hr period of strong illumination, the absorption spectrum of each leaf was recorded. The results are summarized in Table I. Although some effect of the time of application of inhibitor relative to the time of first illumination was observed, two important findings can be seen: (a) Even when the inhibitor was applied at the same time as the first illumination, there was over 90% inhibition of chlorophyll formation during 6 hr illumination. (b) Continuous application of the inhibitor during 48 hr prior to (and during) the illumination did not abolish completely the formation of chlorophyll in the light.

In order to estimate the time required for cycloheximide to act, we illuminated leaves continuously for 5 hours and then transferred them in the light to a solution containing 0.5 mg/ml of cycloheximide. Leaves for a control were left under the original conditions (distilled water). Samples were removed at intervals (3 leaves per sample), and their absorption spectra were recorded. The results of this experiment are shown in Figure 1. Within the reproducibility of the data, the cycloheximide appears to act very quickly (within 1 hr) to block completely the further formation of chlorophyll. It is significant that neither an increase nor a decrease in chlorophyll absorption was observed during the next 19 hr, whereas the chlorophyll in the control leaves increased steadily to a point beyond which the absorbance at 678 nm could not be measured accurately. The time of action of cycloheximide is measured somewhat more accurately in an experiment described below.

Pulsed Illumination Studies. Because of our concern over possible adverse effects resulting from the high illumination intensity of the experiments described above, the studies to be described now were carried out using relatively low intensity (25 ft-c) white light. A pulse duration of 15 sec was found to saturate the photoconversions and was adopted as a standard period of illumination.

Etiolated leaves were given a preillumination pulse and then

Table I. The Effect of the Application of Cycloheximide before and during Illumination of Etiolated Barley Leaves

Cycloheximide concentration is 0.5 mg/ml.

Time of Application of Cycloheximide	Absorbance of Maximum at 678 nm after 6 Hr of Illumination	Inhibition of Chlorophyll Formation (Relative to Control Minus Cycloheximide)
hr		%
0	0.05	93
-2	0.04	94.5
-4	0.03	96
-6	0.02	97
-8	0.02	97
-12	0.01	98.5
-24	0.01	98.5
-48	0.01	98.5
Control	0.72	




FIG. 1. The effect of the application of cycloheximide on the production of chlorophyll during the rapid phase of synthesis in 6-day-old barley seedlings. The dark-grown seedlings were illuminated beginning at zero time and after 5 hr some of the seedlings were transferred to a solution containing 0.5 mg/ml of cycloheximide. Leaves were removed at intervals from the treated and the control leaves for spectrophotometric measurements. Each point represents the average of measurements on three leaves.

allowed to stand in the dark for periods of 1 to 5 hr before application of a second identical pulse. Absorption spectra were recorded before and after each pulse. In each case an absorbance increase, $\Delta A = +0.02 \pm 0.005$ at 678 nm, and an absorbance decrease, $\Delta A = -0.01 \pm 0.005$ at 650 nm, were observed for each of the two pulses. Within the precision of the measurements, the yields were found to be the same for the first and the second pulse, regardless of whether the dark interval between pulses was 1, 2, 3, 4, or 5 hr. There was no evidence of any additional convertible protochlorophyllide formed after the first hour of darkness.

In order to determine the turnover time of the photoconversion process during the rapid phase of chlorophyll synthesis following the induction period, a series of spaced pulses was applied to a single leaf beginning 1 hr (in darkness) after a preillumination pulse. Spectra of the leaf were measured between each pair of pulses, and the specimens remained in position in the spectrophotometer compartment during the entire course of each experiment. The results of these experiments are summarized in Table II. The yields per pulse are presented as average values over

Table II. Yields of Chlorophyll per Pulse of Illumination as a Function of the Dark Interval between Pulses

Pulses numbered 1 and 2 were given before and after 1 hr of darkness. Dark intervals between subsequent pulses are given in column 1. Yields are given as average values calculated for the pulse numbers indicated at the tops of the following columns. The last column gives the total elapsed time from the second through the last pulse.

Dark Intervals <i>min</i>	Yield of Chlorophyll per Pulse (ΔA_{678}) _{avg}					Total Elapsed Time following Induction <i>min</i>
	Pulse number					
	1 and 2	3-6	7-10	11-15	16-20	
20	0.020	0.014	0.015	160
10	0.020	0.016	0.014	80
5	0.018	0.015	0.013	40
2.5	0.020	0.010	0.011	20
2.0	0.020	0.005	0.006	0.005	0.005	36
1.0	0.020	0.005	0.005	0.004	0.004	18




FIG. 2. The time course of the inhibition of chlorophyll synthesis by cycloheximide during the rapid phase of synthesis in 6-day-old barley seedlings. The dark-grown seedling was illuminated using continuous light for 3 hr, its spectrum was measured, and then it was transferred to a solution containing 0.5 mg/ml of cycloheximide. Three 15-sec pulses of light were given at 5-min intervals. Next, the leaf was put back into the spectrophotometer and spectra were measured between successive light pulses at 5-min intervals.

several pulses (in order to increase the precision). Apart from the first two pulses, each of which occurred following a long dark interval, no significant trends were noted within a single sequence. The results show that for dark intervals between 5 and 20 min the yield of chlorophyll per pulse is independent of the duration of the interval. Shorter dark intervals produce a regular decrease in the yield per pulse, reaching half the maximal value at an interval of 2.0 to 2.5 min of darkness. For the shortest intervals studied, the 15 sec period of illumination is not brief enough to be ignored. Nevertheless, the results point to the facts that the half-time for turnover of the photoconversion is about 2.5 min and that dark intervals longer than 5 min do not produce any increase in the amount of photoconvertible protochlorophyllide. Furthermore, the rate of chlorophyll formation using 15-sec pulses spaced at 5 min intervals, $\Delta A_{678} = 0.015$ per pulse, is comparable with the steady state rate observed under continuous, higher intensity illumination (Fig. 1) of $\Delta A_{678} = 0.016$ per 5 min.

The first two pulses in each experiment described in Table II appear to produce somewhat larger yields per pulse than do those of the subsequent sequence. The effect appears to be real, but is barely outside the experimental uncertainty. Thus, there may be a small increase in the photoconvertible protochlorophyllide produced during prolonged dark periods, but further experiments are required to document this difference.

Time of Action of Cycloheximide using Pulsed Illumination. The effect of single pulses of light on one leaf could be more easily observed than could the increase in chlorophyll content in different leaves under continuous illumination. Therefore, we used the pulse scheme to study the time of action of cycloheximide during the phase of rapid chlorophyll synthesis. For this experiment, a leaf was illuminated continuously (strong light) for 3 hr, its spectrum was measured, and then it was transferred to cycloheximide (0.5 mg/ml) in the dark. Three 15-sec pulses of weak light were given at 5-min intervals. The leaf was then removed and mounted in the spectrophotometer. The total exposure to cycloheximide was 15 min, but no attempt was made to wash off the adhering solution prior to the transfer to the spectrophotometer. The pulses were then continued at 5-min intervals, and the spectrum was recorded in each interval between pulses. The results are presented in Figure 2. During the first 45 min (9 pulses) following the application of the cycloheximide, the yield per pulse was not significantly different from that in the absence of cycloheximide. After the ninth pulse the production of chlorophyll ceased abruptly. The sharp transition observed suggests that the inhibition process occurs in two stages.

DISCUSSION

Inhibitors of protein synthesis, such as chloramphenicol or cycloheximide, have been shown to block the formation of chlorophyll in algal systems and in higher plants (13, 14, 18, 19). Two mechanisms have been proposed to account for the action of these inhibitors. In one of these the inhibitor is thought to block the synthesis of enzymes needed to form the precursors of protochlorophyllide (6). Feeding experiments indicate that the blockage occurs early in the biosynthetic pathway, prior to the formation of δ -aminolevulinic acid (7, 8, 20). The second mechanism, favored by Kirk (12, 13), proposes that the synthesis of new protein is required for the incorporation of chlorophyll into the growing lamellar structures of developing plastids. Because of conflicting reports of the ability of δ -aminolevulinic acid to overcome the inhibition in different organisms, it is difficult at the present time to formulate a single hypothesis encompassing all of the findings.

The results summarized in Table I show that the protochlorophyllide formed in the dark in etiolated barley seedlings is capable of being photoconverted into chlorophyll in the presence of 0.5 mg/ml of cycloheximide, even when the inhibitor was applied as much as 48 hr prior to the first illumination. As the time of prior application is shortened, the initial yield of chlorophyll increases somewhat, owing to the increased content of protochlorophyllide accumulating in the seedlings between 4 and 6 days old. We conclude from these observations that the active protochlorophyllide formed in the dark-grown plastids is in a very stable structure and does not require ongoing synthesis of protochlorophyllide in order to retain its activity over a period of two days.

When cycloheximide is added 3 hr after the start of illumination, there is no significant decrease in chlorophyll formed during the next 45 min. Then, during the next 10 min, the activity falls to zero and no further chlorophyll is formed (Fig. 2). Whether the inhibitor stops protein synthesis quickly and leaves a substantial excess of the essential enzyme(s) or, alternatively, a period of 45 min is required for the inhibitor to act, it is evident that we are dealing with a short-lived (~ 10 min) enzyme that requires

constant and active resynthesis in order to maintain chlorophyll formation in the light. The results of Nadler and Granick (20) previously provided an upper limit of $1\frac{1}{2}$ hr for the lifetime of this enzyme.

The postulate, referred to in the introduction, that protochlorophyllide is converted at specialized sites of an enzymatic nature was supported by the isolation of soluble protochlorophyllide-protein complexes which retain the ability to carry out the photoconversion (2, 15, 21, 30). Because of the low concentration of these complexes relative to the amount of chlorophyll synthesized during the first 2 days of illumination of etiolated seedlings, it is reasonable to suppose that the same photoconversion sites are used repeatedly in building up the chlorophyll content. Both Boardman (3) and Bogorad *et al.* (4) have presented evidence that the chlorophyll, once formed, is translocated to a separate macromolecular structure which sediments in the ultracentrifuge differently from the protochlorophyllide protein. Bogorad *et al.* (4) discuss several possible models for the photoconversion and translocation process. Further support for a limiting number of photoconversion sites was provided by Sundqvist (31, 32) and by Granick and Gassman (11) who found that the presence of an excess of protochlorophyllide, resulting from feeding the leaves with the precursor δ -aminolevulinic acid, does not increase the yield of chlorophyllide over that obtained during a brief illumination of untreated leaves.

In the studies that we have carried out, we find that the yield of chlorophyll formed using brief saturating pulses of illumination given directly to a dark-grown barley seedling is nearly the same as the yield per pulse during the phase of rapid chlorophyll synthesis following the induction period. During the rapid synthesis phase a dark interval of 5 min between light pulses is sufficient to saturate the amount of protochlorophyllide produced and longer dark intervals do not increase the yield of chlorophyll per pulse. The half-time for steps required to replace a converted protochlorophyllide molecule on the active site is about 2.5 min. A similar conclusion can be drawn from the rate of active protochlorophyllide synthesis in the dark following brief illumination of wheat seedlings (32). We find that this process can be repeated at least 20 times without any increase or decrease in the yield per pulse. The turnover time of 2.5 min is significantly longer than the 20 sec required for the regeneration of active protochlorophyllide in bean leaves which had been fed δ -aminolevulinic acid and had built up a pool of inactive protochlorophyllide in the dark (11). Thus, it appears that about 2 min is required for the biosynthesis of protochlorophyllide from its precursors and an additional 20 sec for the placement of the molecule on the photoconversion site.

Our findings are entirely consistent with the postulate that the number of protochlorophyllide photoconversion sites remains constant during the first few hours of illumination of etiolated barley seedlings, and that the same sites can be used repeatedly for the photoreduction.

Acknowledgments—The authors wish to thank Prof. Marcel Bastin of the Middle East Technical University for many fruitful discussions of the results and for graciously making available his laboratory facilities to us. Cycloheximide was made available by courtesy of a gift from the Upjohn Co., Kalamazoo, Michigan. This work was carried out while one of us (KS) was on leave in Ankara and was supported, in part, by the United States Agency for International Development.

LITERATURE CITED

- BLAAUW-JANSEN, G., J. G. KOMEN, AND J. B. THOMAS. 1950. On the relation between the formation of assimilatory pigments and the rate of photosynthesis in etiolated oat seedlings. *Biochim. Biophys. Acta* 5: 179-185.
- BOARDMAN, N. K. 1962. Studies on a protochlorophyll-protein complex. I. Purification and molecular weight determination. *Biochim. Biophys. Acta* 62: 63-79.
- BOARDMAN, N. K. 1967. Chloroplast structure and development. In: A. San Pietro *et al.*, eds., *Harvesting the Sun*. Academic Press, New York. pp. 211-230.
- BOGORAD, L., L. LABER, AND M. GASSMAN. 1968. Aspects of chloroplast development: Transitory pigment-protein complexes and protochlorophyllide regeneration. In: K. Shibata, A. Takamiya, A. Jagendorf, and R. C. Fuller, eds., *Comparative Biochemistry and Biophysics of Photosynthesis*. University Park Press, State College, Pa. pp. 299-312.
- BUTLER, W. L. AND W. R. BRIGGS. 1966. The relation between structure and pigments during the first stages of proplastid greening. *Biochim. Biophys. Acta* 112: 45-53.
- GASSMAN, M. AND L. BOGORAD. 1965. Photocontrol of protochlorophyll synthesis in bean leaves. *Plant Physiol.* 40: lii.
- GASSMAN, M. AND L. BOGORAD. 1967. Control of chlorophyll production in rapidly greening bean leaves. *Plant Physiol.* 42: 774-780.
- GASSMAN, M. AND L. BOGORAD. 1967. Studies on the regeneration of protochlorophyllide after brief illumination of etiolated bean leaves. *Plant Physiol.* 42: 781-784.
- GOEDHEER, J. C. 1961. Effect of changes in chlorophyll concentration on photosynthetic properties. I. Fluorescence and absorption of greening bean leaves. *Biochim. Biophys. Acta* 51: 494-504.
- GOEDHEER, J. C. AND C. A. H. VERHULSDONK. 1970. Fluorescence and photo-transformation of protochlorophyll with etiolated bean leaves from -196 to +20 C. *Biochem. Biophys. Res. Commun.* 39: 260-266.
- GRANICK, S. AND M. GASSMAN. 1970. Rapid regeneration of protochlorophyllide₆₈₀. *Plant Physiol.* 45: 201-205.
- KIRK, J. T. O. 1968. Studies on the dependence of chlorophyll synthesis on protein synthesis in *Euglena gracilis*, together with a nomogram for determination of chlorophyll concentration. *Planta* 78: 200-207.
- KIRK, J. T. O. AND R. L. ALLEN. 1965. Dependence of chloroplast pigment synthesis on protein synthesis: effect of actidione. *Biochem. Biophys. Res. Commun.* 21: 523-530.
- KIRK, J. T. O. AND R. A. E. TILNEY-BASSETT. 1967. *The Plastids*. W. H. Freeman and Co., San Francisco.
- KRASNOVSKY, A. A. AND L. M. KOSOBUTSKAYA. 1952. Spectral investigations of chlorophyll during its formation in plants and in colloidal solutions of material from etiolated leaves. *Dokl. Akad. Nauk SSSR*. 85: 177-180.
- MADSEN, A. 1961. Photochemical conversion of protochlorophyll to chlorophyll-a. In: B. Christensen and B. Buchmann, eds., *Progress in Photobiology*. Elsevier, Amsterdam. pp. 567-569.
- MADSEN, A. 1963. Is the photoconversion of protochlorophyll immediate? *Carnegie Inst. Wash. Year B.* 62: 371-373.
- MARGULIES, M. M. 1962. Effect of chloramphenicol on light dependent development of seedlings of *Phaseolus vulgaris* var. Black Valentine, with particular reference to development of photosynthetic activity. *Plant Physiol.* 37: 473-480.
- MARGULIES, M. M. 1964. Effect of chloramphenicol on light-dependent synthesis of proteins and enzymes of leaves and chloroplasts of *Phaseolus vulgaris*. *Plant Physiol.* 39: 579-585.
- NADLER, K. AND S. GRANICK. 1970. Controls on chlorophyll synthesis in barley. *Plant Physiol.* 46: 240-246.
- SCHOPFER, P. AND H. W. SIEGELMAN. 1968. Purification of protochlorophyllide holochrome. *Plant Physiol.* 43: 990-996.
- SHIBATA, K. 1957. Spectroscopic studies on chlorophyll formation in intact leaves. *J. Biochem.* 44: 147-173.
- SIRONVAL, C. AND M. BROUERS. 1970. The reduction of protochlorophyllide into chlorophyllide. II. The temperature dependence of the $P_{687-687} \rightarrow P_{688-676}$ photo-transformation. *Photosynthetica* 4: 38-47.
- SIRONVAL, C., M. R. MICHEL-WOLWERTZ, AND A. MADSEN. 1965. On the nature and possible functions of the 673- and 684-m μ forms *in vivo* of chlorophyll. *Biochim. Biophys. Acta* 94: 344-354.
- SISLER, E. C. AND W. H. KLEIN. 1963. The effect of age and various chemicals on the lag phase of chlorophyll synthesis in dark grown bean seedlings. *Physiol. Plant.* 16: 315-322.
- SMILLIE, R. M., W. R. EVANS, AND H. LYMAN. 1963. Metabolic events during the formation of a photosynthetic from a nonphotosynthetic cell. *Brookhaven Symp. Biol.* 16: 89-107.
- SMILLIE, R. M. AND N. S. SCOTT. 1969. Organelle biosynthesis: the chloroplast. *Progress in Molecular and Subcellular Biology* 1: 136-202.
- SMILLIE, R. M., N. S. SCOTT, AND D. GRAHAM. 1968. Biogenesis of chloroplasts: roles of chloroplast DNA and chloroplast ribosomes. In: K. Shibata, A. Takamiya, J. Jagendorf, and R. C. Fuller, eds., *Comparative Biochemistry and Biophysics of Photosynthesis*. University Park Press, State College, Pa. pp. 332-53.
- SMITH, J. H. C. AND A. BENITEZ. 1954. The effect of temperature on the conversion of protochlorophyll to chlorophyll a in etiolated barley leaves. *Plant Physiol.* 29: 135-143.
- SMITH, J. H. C. AND D. W. KUPKE. 1956. Some properties of extracted protochlorophyll holochrome. *Nature* 187: 751-752.
- SUNDQVIST, C. 1969. Transformation of protochlorophyllide, formed from exogenous δ -aminolevulinic acid, in continuous light and in flash light. *Physiol. Plant.* 22: 147-156.
- SUNDQVIST, C. 1970. The conversion of protochlorophyllide₆₈₀ to protochlorophyllide₆₈₀ in leaves treated with δ -ALA. *Physiol. Plant.* 23: 412-424.
- VIRGIN, H. I. 1956. Light-induced stomatal transpiration of etiolated wheat leaves as related to chlorophyll a content. *Physiol. Plant.* 9: 482-493.
- WOLFF, J. B. AND L. PRICE. 1957. Terminal steps of chlorophyll a biosynthesis in higher plants. *Arch. Biochem. Biophys.* 72: 293-301.