

Direct observation of Kelvin waves excited by quantized vortex reconnection

Enrico Fonda^{a,b,c}, David P. Meichle^{a,d}, Nicholas T. Ouellette^{a,e}, Sahand Hormoz^f, and Daniel P. Lathrop^{a,d,1}

^aInstitute for Research in Electronics and Applied Physics and ^dDepartment of Physics, University of Maryland, College Park, MD 20742; ^bDipartimento di Matematica e Geoscienze, Università degli Studi di Trieste, 34127 Trieste, Italy; ^cDepartment of Physics, New York University, New York, NY 10003; ^eDepartment of Mechanical Engineering and Materials Science, Yale University, New Haven, CT 06520; and ^fKavli Institute for Theoretical Physics, Kohn Hall, University of California, Santa Barbara, CA 93106

Edited by Katepalli R. Sreenivasan, New York University, New York, NY, and approved September 24, 2013 (received for review July 8, 2013)

Quantized vortices are key features of quantum fluids such as superfluid helium and Bose–Einstein condensates. The reconnection of quantized vortices and subsequent emission of Kelvin waves along the vortices are thought to be central to dissipation in such systems. By visualizing the motion of submicron particles dispersed in superfluid ⁴He, we have directly observed the emission of Kelvin waves from quantized vortex reconnection. We characterize one event in detail, using dimensionless similarity coordinates, and compare it with several theories. Finally, we give evidence for other examples of wavelike behavior in our system.

topological defects | turbulence | cascade

Vorticity in superfluids and Bose–Einstein condensates is constrained to line-like topological defects called quantized vortices (1). The evolution of a tangle of such line vortices defines a dynamical state known as quantum turbulence (see, e.g., refs. 2, 3). Quantum turbulence is in some ways similar to classical turbulence; for example, both show a Kolmogorov energy spectrum (4, 5). However, many features of quantum turbulence, such as its velocity statistics (6, 7), are distinct from classical flows.

A fundamental question in quantum turbulence is the nature of dissipation in the zero-temperature limit (8) where the effects of friction vanish. The vortex-tangle decay, observed experimentally even for $T < 0.1$ K (9), requires a different dissipation mechanism from the classical case; a promising candidate is the excitation of waves by reconnecting vortices (Fig. 1). In his pioneering work (11), Kelvin showed that a helical deformation of a line vortex propagates as a wave. Kelvin waves have long been used to understand a wide range of flow problems, such as airplane wakes (12), tornadoes (13), and the dynamics of neutron stars (14), and are also conceptually related to whistler waves in plasmas (15). Theory and simulations indicate that a cascade of Kelvin waves transfers energy from large length scales (the intervortex spacing) to small scales (the vortex-core size) (16–18), where energy is removed from the system via phonon emission (19–21).

Kelvin waves have been visualized in classical fluids only on thin line vortices (22, 23) and on knotted vortex rings (24). Here we present a unique direct observation of Kelvin waves on quantized vortices and give unique experimental evidence of the emission of Kelvin waves after vortex reconnection. Because our fluid is inviscid and the amplitude H of the waves we observe is much larger than the vortex core size a_0 (we have $H/a_0 > 10^3$), our system satisfies most of the assumptions originally made by Kelvin for his eponymous waves.

Kelvin waves on quantized vortices were first detected indirectly, using torsional oscillators (1, 25), beginning with the work of Hall in 1958 (26). However, the interpretation of such experiments has been criticized (27); additional evidence is therefore needed (1). Ashton and Glaberson (28) measured the velocity of ions passing through the superfluid as a function of an applied electric field and found an anomaly that they associated with the resonant generation of vortex waves. A recent experiment (29) inferred the presence of Kelvin modes in a Bose–Einstein

condensate by examining the damping. However, to our knowledge, no direct observation has shown the existence of Kelvin waves on quantized vortices, and no experimental evidence has shown Kelvin waves launched from vortex reconnection.

Vortex reconnection in superfluid helium, theorized in 1955 by Feynman (30) and studied extensively both numerically and analytically (e.g., refs. 31–40), was characterized experimentally in 2008 by Bewley et al. (41) and Paoletti et al. (6, 42), using micrometer-sized hydrogen particles to visualize the quantized vortices. Because the quantized vortices are local minima of the pressure field, they act as traps for small particles; the motion of the particles can then be used to visualize the evolution of the vortices.

Results

Using a unique technique to create submicron tracer particles, at a mean temperature of 1.981 K we slowly cooled the liquid helium at a rate of 0.2 mK/s, and observed a long, clearly decorated vortex. In Fig. 2A, we show several snapshots in a 2.08×0.4 -mm subregion of our measurement area, with the tracer particles used for further analysis indicated. In multiple frames, captured at 54.3 Hz, several sections of the vortex were visible over a length of about 3 mm, appearing initially straight and nearly horizontal. The normal component of the helium was almost stationary, with particles dragged in different directions at a maximum velocity of 6 μ m/s. Shortly before the snapshots in Fig. 2A, the vortex shown reconnected with another vortex to the right of the field of view. Immediately after the reconnection, we observed a wave propagating down the vortex.

To study the wave motion quantitatively, we track the position of the particles on the vortex (Fig. 2B). The wave-like motion of the vortex is clearly visible in the time evolution of the vertical positions of the particles, plotted in Fig. 3A (see also [Movies S1](#) and [S2](#)). The first peak and trough are clearly visible in most of the tracks, with secondary peaks only in some. We fitted the time evolution of the maximum and minimum z values of each particle, indicated respectively by circles and squares in Fig. 3A, and found $(t-t_0)^{1/2}$ scaling for both, as shown in Fig. 3B. This is consistent with previous theoretical and experimental studies of reconnection (6, 32–34, 41), with the assumption that the quantum of circulation $\kappa = h/m_{He} \simeq 9.97 \times 10^{-8} \text{ m}^2/\text{s}$ is the only relevant dimensional parameter and with the dispersion relation $\omega(k) \propto \kappa k^2$ for Kelvin waves of frequency ω and wavenumber k (43).

In theoretical vortex-filament models, the velocity of the superfluid is calculated from the position of the vortices $s(\sigma, t)$,

Author contributions: E.F., D.P.M., and D.P.L. designed research; E.F. and D.P.M. performed research; E.F., D.P.M., N.T.O., and S.H. contributed new reagents/analytic tools; E.F., D.P.M., N.T.O., S.H., and D.P.L. analyzed data; and E.F., D.P.M., N.T.O., S.H., and D.P.L. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: lathrop@umd.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1312536110/-DCSupplemental.




Fig. 1. (A) Schematic diagram of two vortices reconnecting and exchanging tails, where the red and blue arrows represent the direction of vorticity. (B) After reconnection, the relaxation of the vortex excites Kelvin waves thought to propagate in a self-similar manner. Reprinted with permission from Schwarz KW, *Phys Rev B Condens Matter* 31:5782–5804 (1985). Copyright 1985 by the American Physical Society, http://prb.aps.org/abstract/PRB/v31/i9/p5782_1.

where σ is the arc length along the vortex, using the Biot–Savart integral. Schwarz (10) laid the foundation for studying vortices after reconnection, using the so-called local induction approximation (LIA) (44). This approximation truncates the Biot–Savart integral by neglecting nonlocal terms, reducing it to a much more tractable form. We note that this approximation has in fact been rediscovered several times (45) since the original work of Da Rios in 1906 (46). The LIA implies

$$\frac{\partial \mathbf{s}}{\partial t} \simeq \beta \frac{\partial \mathbf{s}}{\partial \sigma} \times \frac{\partial^2 \mathbf{s}}{\partial \sigma^2}, \quad [1]$$

where $\beta \simeq \kappa$. We neglect logarithmic corrections that depend on the radius of curvature of the vortex and the core size, absorbing




Fig. 2. (A) Four frames of our movie sequence (see [Movie S1](#)) along with circled particles used in the tracking analysis. (B) The positions of the particle tracks on the upper branch show oscillatory behavior after the reconnection event. The cross is the estimated location of the reconnection event.




Fig. 3. (A) The vertical position of the particle tracks on the upper branch shows marked oscillatory behavior just after the observed reconnection. (B) The observed maxima and minima of the vertical positions from A. These are consistent with the expected behavior for spatial scales close to a reconnection event $z - z_0 \sim (t - t_0)^{1/2}$.

them into the constant parameter β . This equation has self-similar solutions for the evolution of the vortex shapes after reconnection (33, 47) and predicts emission of Kelvin waves as an inevitable consequence of the relaxation of any angle-like configuration (8). Note that waves excited by the relaxation of a cusp are localized and polychromatic unlike, for example, the periodic waves artificially excited on straight vortices in recent numerical studies of the Kelvin wave cascade (48, 49). First we define the dimensionless similarity coordinate $\eta = \sigma / \sqrt{\beta t}$. Then, substituting a self-similar solution of the form $s(\sigma, t) = \sqrt{\beta t} \mathbf{G}(\eta)$ into the LIA equation gives

$$\frac{1}{2} \mathbf{G} - \frac{1}{2} \eta \mathbf{G}' = \mathbf{G}' \times \mathbf{G}'', \quad [2]$$

where the primes denote differentiation with respect to η . However, solutions to this simple equation cannot exactly describe our system, which is at finite temperature. Coupling between the superfluid and the residual normal component via the mutual friction must be included. Therefore, we consider a modified LIA equation with a phenomenological temperature-dependent damping term $\alpha(T) d^2 s / d\sigma^2$ as done by Schwarz (10) and Lipniacki (33), which after the substitution used to obtain Eq. 2 results in

$$\frac{1}{2} \mathbf{G} - \frac{1}{2} \eta \mathbf{G}' = \mathbf{G}' \times \mathbf{G}'' + \alpha(T) \mathbf{G}'''. \quad [3]$$

Solutions to Eq. 3 are a two-parameter family of curves that can be completely specified by the temperature-dependent damping term $\alpha(T)$ and the initial curvature c_0 , which determines, via integration of Eq. 3, the opening angle between the two tails of the retracting vortex line (illustration in Fig. 1B). Note that $c_0 = A/4$, where A is the dimensionless prefactor of the scaling law of the interfilament separation distance. Based on our temperature and previous measurements of the mutual friction coefficient (50), we estimate $\alpha = 0.27$.

In addition to the LIA model, we considered novel similarity solutions of a Biot–Savart model akin to that described by Hormoz and Brenner (51) but modified to include the damping term, which leads to an equation analogous to Eq. 3:

$$\frac{1}{2} \mathbf{G} - \frac{1}{2} \eta \mathbf{G}' = \mathbf{G}' \times \mathbf{G}'' + \alpha \mathbf{G}''' + F[\mathbf{G}]. \quad [4]$$

The additional term $F[\mathbf{G}]$ approximates nonlocal contributions to the velocity of the filament. We solve Eq. 4 using coupled-delay differential equations to incorporate the nonlocal interactions. As with LIA, the solutions are a two-parameter family of curves characterized by the prefactor A and the temperature-dependent constant α .




Fig. 4. Particle tracks representing other examples that may be Kelvin waves. Each point corresponds to the particle position in one frame and is separated from the next one by 29 ms. The red triangle represents the end of the track, to clarify the direction of motion.

Following the theoretical prediction that the evolution of the vortex shape is self-similar, we define the similarity coordinates $\zeta = (z - z_0) / \sqrt{\kappa(t - t_0)}$ and $\xi = (x - x_0) / \sqrt{\kappa(t - t_0)}$. This self-similar assumption is reasonable given the $(t - t_0)^{1/2}$ scaling of the wave peak, as shown in Fig. 3. We estimate the spatiotemporal coordinates of the reconnection event (x_0, z_0, t_0) ; those values are adjusted to collapse the tracks. As shown in Fig. 5A, the

trajectories of the individual particles on the vortex collapse well when rescaled into these similarity coordinates.

Fig. 5B and C shows a comparison of the Biot–Savart and LIA curves, with our data rescaled in dimensionless similarity coordinates ξ and ζ . These numerically integrated curves were computed in 3D and rotated to optimize the fit of their projection onto the 2D data. For a dense tangle, the results of LIA are distinctively different from the full Biot–Savart (52) calculation. However, for the vortex density in our system and the resolution of our particle tracks, the two models compare equally well to our similarity-collapsed experimental data.

Although we have focused here on one event, we observed many other tracks with wave-like motions. Some tracks of particles moving in wavy and quasi-circular motion are reported in Fig. 4. Note that a particle trapped on a vortex would appear to move in a circle if observed in a plane normal to the axis of propagation for a helical Kelvin wave. In general, however, care must be taken in interpreting the motion of the particles on a vortex, as the particles are not locked to a specific point along the vortex core.

Using submicron ice particles, we have observed the emission and propagation of Kelvin waves on quantized vortices in superfluid helium following vortex reconnection. The resulting waveforms are in general agreement with those of theoretical similarity theories. These waves carry energy away from the reconnection event and enhance dissipation through mutual friction. Future studies could examine a broader collection of events to characterize the distribution of amplitudes and phase velocities present in quantum turbulence.




Fig. 5. The overlap of the curves described by similarity solutions of LIA and Biot–Savart models (colored curves) to the experimental data in similarity coordinates (blue circles). We report just the Biot–Savart (B-S) curves as the LIA curves are almost indistinguishable. We show two different curves that are both in qualitative agreement with the observation: the red curve for the lower-bound value $A = 3.3$, and the blue curve for the upper-bound value $A = 3.75$, obtained from this particular fit to maximize the overlap of the Biot–Savart curves, respectively, with both the lower and upper branch and just the upper branch of the collapsed data. These values of A correspond, respectively, to intervortex angles of 40° and 29° . In both cases the angles that set the 3D orientation are free parameters selected by hand. We also show a LIA curve without any damping (thin gray line), to demonstrate the necessity of including the mutual friction. (A) shows most of the collapsed data, whereas (C) shows in detail the area close to the first peak of the wave marked by the dashed region in (B). The red cross represents the reconnection estimated origin.

Method

We discovered a unique technique to create submicron tracer particles directly into the superfluid state of helium. In the pioneering work of Bewley et al. (53), the particles were created by injecting a dilute mixture of hydrogen and helium gas into liquid helium above the λ transition. The liquid was then cooled into the superfluid regime. During the cooling process, however, the particles tend to aggregate and float to the surface of the liquid; few particles remain in the measurement region for temperatures below 2 K. Similar effects were seen by La Mantia et al. (54). Attempts to introduce particles directly at superfluid temperatures resulted in excessively large particles that were not acceptable tracers (55).

In contrast, we create frozen tracers directly in the superfluid by injecting a highly diluted mixture of atmospheric air and helium gas (56). We were thus able to keep the temperature below 2 K while resupplying tracer particles. We estimate the particles to be submicron sized by measuring the amount of light they scatter, according to the procedure described in ref. 56.

Our experimental setup consists of an Oxford Instruments STX cryostat with five optical windows. The helium sample section has a $1 \times 1 \times 7.5$ -cm

rectangular glass cell with 1-mm-thick walls, immersed in a 5-cm diameter cylindrical bath. The system is illuminated with a 3-mW 532-nm laser, which is focused into a sheet about 1 cm tall with a full-width at half-maximum width of about 150 μm . We collected 90° scattered light off frozen tracer particles from an 8.2-mm square area, using a 105-mm Micro-Nikkor lens. A Princeton Instrument Pro-EM CCD provides single-photon sensitivity imaging with a 512×512 -pixel resolution.

The vortices we observe are longer (up to several millimeters) and straighter than those from previous studies, because the system is in the superfluid state for about 0.5 h before measuring. In contrast, the system studied by Paoletti et al. (6, 42) was driven strongly out of equilibrium by a thermal counterflow and had a dense tangle of vortices in a larger 5-cm diameter volume.

ACKNOWLEDGMENTS. We thank Katepalli R. Sreenivasan, Matthew S. Paoletti, Michael P. Brenner, Cecilia Rorai, Davide Proment, and Michael E. Fisher for fruitful discussions and comments. E.F. also thanks Katepalli R. Sreenivasan for his scientific interest and support. This work was partially supported by the National Science Foundation.

- Donnelly RJ (1991) *Quantized Vortices in Helium II* (Cambridge Univ Press, Cambridge, UK).
- Vinen WF, Niemela JJ (2002) Quantum turbulence. *J Low Temp Phys* 128(5-6):167–231.
- Skrbek L, Sreenivasan KR (2012) Developed quantum turbulence and its decay. *Phys Fluids* 24:011301.
- Maurer J, Tabeling P (1998) Local investigation of superfluid turbulence. *Europhys Lett* 43(1):29–34.
- Salort J, Chabaud B, L ev eque E, Roche PE (2012) Energy cascade and the four-fifths law in superfluid turbulence. *Europhys Lett* 97:34006.
- Paoletti MS, Fisher ME, Sreenivasan KR, Lathrop DP (2008) Velocity statistics distinguish quantum turbulence from classical turbulence. *Phys Rev Lett* 101(15):154501.
- White AC, Barenghi CF, Proukakis NP, Youd AJ, Wacks DH (2010) Nonclassical velocity statistics in a turbulent atomic Bose-Einstein condensate. *Phys Rev Lett* 104(7):075301.
- Svistunov BV (1995) Superfluid turbulence in the low-temperature limit. *Phys Rev B Condens Matter* 52(5):3647–3653.
- Hendry PC, Lawson NS, Lee RAM, McClintock PVE, Williams CDH (1994) Generation of defects in superfluid 4He as an analogue of the formation of cosmic strings. *Nature* 368:315–317.
- Schwarz KW (1985) Three-dimensional vortex dynamics in superfluid ⁴He: Line-line and line-boundary interactions. *Phys Rev B Condens Matter* 31(9):5782–5804.
- Thomson W (1880) Vibrations of a columnar vortex. *Philos Mag* 10(61):155–168.
- Crow SC (1970) Stability theory for a pair of trailing vortices. *AIAA J* 8:2172–2179.
- Maxworthy T (1973) A vorticity source for large-scale dust devils and other comments on naturally occurring columnar vortices. *J Atmos Sci* 30:1717–1722.
- Epstein RI, Baym G (1992) Vortex drag and the spin-up time scale for pulsar glitches. *Astrophys J* 387:276–287.
- Helliwell RA (2006) *Whistlers and Related Ionospheric Phenomena* (Dover, New York).
- Kivotides D, Vassiliou JC, Samuels DC, Barenghi CF (2001) Kelvin waves cascade in superfluid turbulence. *Phys Rev Lett* 86(14):3080–3083.
- Kozik EV, Svistunov BV (2009) Theory of decay of superfluid turbulence in the low-temperature limit. *J Low Temp Phys* 156:215–267.
- L'vov V, Nazarenko S, Rudenko O (2007) Bottleneck crossover between classical and quantum superfluid turbulence. *Phys Rev B* 76(2):024520.
- Nore C, Abid M, Brachet M (1997) Kolmogorov turbulence in low-temperature superflows. *Phys Rev Lett* 78:3896–3899.
- Leadbeater M, Winiecki T, Samuels DC, Barenghi CF, Adams CS (2001) Sound emission due to superfluid vortex reconnections. *Phys Rev Lett* 86(8):1410–1413.
- Vinen WF, Tsubota M, Mitani A (2003) Kelvin-wave cascade on a vortex in superfluid 4He at a very low temperature. *Phys Rev Lett* 91(13):135301.
- Maxworthy T, Hopfinger EJ, Redekopp LG (1985) Wave motions on vortex cores. *J Fluid Mech* 151:141–165.
- Hopfinger EJ, Browand FK (1982) Vortex solitary waves in a rotating, turbulent flow. *Nature* 295:393–395.
- Kleckner D, Irvine WTM (2013) Creation and dynamics of knotted vortices. *Nat Phys* 9:253–258.
- Sonin E (1987) Vortex oscillations and hydrodynamics of rotating superfluids. *Rev Mod Phys* 59(1):87–155.
- Hall HE (1958) An experimental and theoretical study of torsional oscillations in uniformly rotating liquid helium II. *Proc R Soc A* 245:546–561.
- Raja Gopal E (1964) Oscillations of quantized vortices in rotating liquid helium II. *Ann Phys* 29:350–365.
- Ashton R, Glaberson W (1979) Vortex waves in superfluid ⁴He. *Phys Rev Lett* 42:1062–1064.
- Bretin V, Rosenbusch P, Chevy F, Shlyapnikov GV, Dalibard J (2003) Quadrupole oscillation of a single-vortex Bose-Einstein condensate: Evidence for Kelvin modes. *Phys Rev Lett* 90(10):100403.
- Feynman R (1955) *Progress in Low Temperature Physics*, ed Gorter CJ (North-Holland, Amsterdam), pp 17–53.
- Koplik J, Levine H (1993) Vortex reconnection in superfluid helium. *Phys Rev Lett* 71:1375–1378.
- Aarts RG, Aarts R, de Waele AT (1994) Route to vortex reconnection. *Phys Rev Lett* 72(4):482–485.
- Lipniacki T (2000) Evolution of quantum vortices following reconnection. *Eur J Mech B Fluids* 19:361–378.
- Nazarenko S, West R (2003) Analytical solution for nonlinear Schrödinger vortex reconnection. *J Low Temp Phys* 132(1-2):1–10.
- Berry MV, Dennis MR (2007) Topological events on wave dislocation lines: Birth and death of loops, and reconnection. *J Phys A: Math Theor* 40(1):65–74.
- Kerr RM (2011) Vortex stretching as a mechanism for quantum kinetic energy decay. *Phys Rev Lett* 106(22):224501.
- Kursa M, Bajer K, Lipniacki T (2011) Cascade of vortex loops initiated by a single reconnection of quantum vortices. *Phys Rev B* 83:014515.
- Meichle DP, Rorai C, Fisher ME, Lathrop DP (2012) Quantized vortex reconnection: Fixed points and initial conditions. *Phys Rev B* 86(1):014509.
- Zuccher S, Caliarı M, Baggaley AW, Barenghi CF (2012) Quantum vortex reconnections. *Phys Fluids* 24:125108.
- Hänninen R (2013) Dissipation enhancement from a single vortex reconnection in superfluid helium. *Phys Rev B* 88:054511.
- Bewley GP, Paoletti MS, Sreenivasan KR, Lathrop DP (2008) Characterization of reconnecting vortices in superfluid helium. *Proc Natl Acad Sci USA* 105(37):13707–13710.
- Paoletti MS, Fisher ME, Lathrop DP (2009) Reconnection dynamics for quantized vortices. *Physica D* 239:1367–1377.
- Pocklington HC (1895) The complete system of the periods of a hollow vortex ring. *Philos Trans R Soc Lond A* 186:603–619.
- Arms RJ, Hama FR (1965) Localized-induction concept on a curved vortex and motion of an elliptic vortex ring. *Phys Fluids* 8:553–559.
- Ricca RL (1991) Rediscovery of Da Rios equations. *Nature* 352:561–562.
- Da Rios LS (1906) Sul moto d'un liquido indefinito con un filetto vorticoso di forma qualunque [On the motion of an unbounded fluid with a vortex filament of any shape]. *Rend Circ Mat Palermo* 22(1):117–135. Italian.
- Guti rrez S, Rivas J, Vega L (2003) Formation of singularities and self-similar vortex motion under the localized induction approximation. *Commun Partial Differ Equ* 28:927–968.
- Krstulovic G (2012) Kelvin-wave cascade and dissipation in low-temperature superfluid vortices. *Phys Rev E* 86:055301.
- Proment D, Barenghi CF, Onorato M (2013) Interaction and decay of Kelvin waves in the Gross-Pitaevskii model. ArXiv:1308.0852.
- Barenghi CF, Donnelly RJ, Vinen WF (1983) Friction on quantized vortices in helium II. A review. *J Low Temp Phys* 52(3-4):189–247.
- Hormoz S, Brenner MP (2012) Absence of singular stretching of interacting vortex filaments. *J Fluid Mech* 707:191–204.
- Adachi H, Tsubota M (2009) Numerical studies of counterflow turbulence. *J Low Temp Phys* 158:422–427.
- Bewley GP, Lathrop DP, Sreenivasan KR (2006) Superfluid helium: Visualization of quantized vortices. *Nature* 441(7093):588.
- La Mantia M, Chagovets TV, Rotter M, Skrbek L (2012) Testing the performance of a cryogenic visualization system on thermal counterflow by using hydrogen and deuterium solid tracers. *Rev Sci Instrum* 83(5):055109.
- Bewley GP, Sreenivasan KR, Lathrop DP (2007) Particles for tracing turbulent liquid helium. *Exp Fluids* 44:887–896.
- Fonda E, Sreenivasan KR, Lathrop DP (2012) Liquid nitrogen in fluid dynamics: Visualization and velocimetry using frozen particles. *Rev Sci Instrum* 83(8):085101.