

Salmonella Pathogenicity and Host Adaptation in Chicken-Associated Serovars

Steven L. Foley,^a Timothy J. Johnson,^b Steven C. Ricke,^c Rajesh Nayak,^a Jessica Danzeisen^b

Division of Microbiology, National Center for Toxicological Research, U.S. Food and Drug Administration, Jefferson, Arkansas, USA^a; Department of Veterinary and Biomedical Sciences, University of Minnesota, Saint Paul, Minnesota, USA^b; Center for Food Safety and Department of Food Science, University of Arkansas, Fayetteville, Arkansas, USA^c

SUMMARY	582
INTRODUCTION	582
OVERVIEW OF <i>SALMONELLA</i> COLONIZATION AND PATHOGENESIS	583
Initial Colonization of the Host	584
Invasion of Host Cells	584
Survival in Macrophages and Dendritic Cells	584
<i>Salmonella</i> Pathogenicity Islands	585
Virulence Plasmids	585
Toxins	585
Fimbriae	585
Flagella	585
Other Virulence Factors	585
IMPACT OF HORIZONTAL GENE TRANSFER ON GENOME EVOLUTION AND HOST RANGE	586
Role of Plasmids in <i>Salmonella</i> Genome Evolution	586
Role of Bacteriophages in <i>Salmonella</i> Evolution	588
IMPACT OF THE LOSS OF GENES OR THEIR FUNCTION ON GENOME EVOLUTION AND HOST RANGE	589
EXAMPLES OF <i>SALMONELLA</i> HOST RANGE IN CHICKEN-RELATED SEROVARS	590
THE <i>SALMONELLA</i> -CHICKEN HOST INTERACTION	592
<i>SALMONELLA</i> AND STRESS RESPONSES IN THE CHICKEN GIT	593
<i>Salmonella</i> Acid Tolerance Response and Organic Acids	593
<i>Salmonella</i> Virulence Response and Organic Acids	594
<i>Salmonella</i> Virulence Response and Competition with GIT Microflora	595
CONCLUDING REMARKS	596
ACKNOWLEDGMENTS	597
REFERENCES	597
AUTHOR BIOS	606

SUMMARY

Enteric pathogens such as *Salmonella enterica* cause significant morbidity and mortality. *S. enterica* serovars are a diverse group of pathogens that have evolved to survive in a wide range of environments and across multiple hosts. *S. enterica* serovars such as *S. Typhi*, *S. Dublin*, and *S. Gallinarum* have a restricted host range, in which they are typically associated with one or a few host species, while *S. Enteritidis* and *S. Typhimurium* have broad host ranges. This review examines how *S. enterica* has evolved through adaptation to different host environments, especially as related to the chicken host, and continues to be an important human pathogen. Several factors impact host range, and these include the acquisition of genes via horizontal gene transfer with plasmids, transposons, and phages, which can potentially expand host range, and the loss of genes or their function, which would reduce the range of hosts that the organism can infect. *S. Gallinarum*, with a limited host range, has a large number of pseudogenes in its genome compared to broader-host-range serovars. *S. enterica* serovars such as *S. Kentucky* and *S. Heidelberg* also often have plasmids that may help them colonize poultry more efficiently. The ability to colonize different hosts also involves interactions with the host's immune system and commensal organisms that are present. Thus, the factors that impact the ability of *Salmonella* to

colonize a particular host species, such as chickens, are complex and multifactorial, involving the host, the pathogen, and extrinsic pressures. It is the interplay of these factors which leads to the differences in host ranges that we observe today.

INTRODUCTION

Enteric pathogens are a major source of morbidity and mortality throughout the world. It has been estimated that there are more than 3 million deaths associated with Gram-negative enteric pathogens worldwide due to diarrhea and enteric fever each year (1). Bacteria of genera such as *Escherichia*, *Campylobacter*, *Vibrio*, *Brucella*, *Shigella*, *Yersinia*, and *Salmonella* are responsible for causing enteric diseases. Infections by these pathogens manifest as several disease syndromes, including secretory/noninflammatory diarrhea, inflammatory diarrhea, and enteric fever (2). The differences in disease manifestations are related to the different virulence factors present in the bacteria and the altered phenotypes

Address correspondence to Steven L. Foley, steven.foley@fda.hhs.gov.

Copyright © 2013, American Society for Microbiology. All Rights Reserved.

doi:10.1128/MMBR.00015-13

that these virulence factors allow the organisms to employ during disease pathogenesis.

Noninflammatory or secretory diarrhea is characterized by watery stools without the presence of white blood cells in feces. Patients generally do not develop a high fever. The diarrhea is associated with bacterial release of enterotoxin or their superficial adherence to the lumen of the proximal small intestine (3). Bacteria that are associated with secretory diarrhea include *Vibrio cholerae* and enterotoxigenic, enteroaggregative, or enteropathogenic pathotypes of *Escherichia coli* (2, 4–6). Diarrheal disease is typically associated with the interaction of the bacterial toxins with the host epithelium, leading to disruption of ion transport systems and subsequent fluid secretion that is the hallmark of diarrhea (7).

In contrast to secretory diarrhea, inflammatory diarrhea typically involves invasion of the colonic epithelium by bacterial pathogens, which may produce cytotoxins. Stool samples typically contain white blood cells, especially neutrophils, and patients often have fever associated with the illness (3). Bacteria associated with causing inflammatory diarrhea include *Campylobacter jejuni*, *Shigella* spp., nontyphoidal *Salmonella enterica*, and enteroinvasive *E. coli* (2). These organisms generally utilize mechanisms such as type III secretion systems (T3SSs) (described in detail below) to interact with the host cell's cytoskeletal system to facilitate invasion (8, 9). The cytoskeletal arrangements lead to the pathogens being internalized, where they reside predominantly in either a protected vesicle (*Salmonella* and *Campylobacter*) or the host cell cytoplasm (*Shigella* and enteroinvasive *E. coli*) (9, 10). During the attachment and internalization processes, the innate immune system can become activated through the recognition of pathogen-associated molecular patterns (PAMPs) on the surface of the pathogens (11, 12). These PAMPs include molecules such as lipopolysaccharide (LPS) and flagella that are recognized by Toll-like receptors (TLRs) on the surface of the host cells, which leads to an antibacterial response that is characterized by increased vascular permeability and the attraction of neutrophils to the site of infection (2). This response leads to the development of inflammatory diarrhea, in which the watery stool contains elevated levels of neutrophils.

Enteric fever is associated with invasive organisms that typically initially invade the distal small intestine and spread to underlying tissues and other parts of the host. Patients generally develop a high fever, and fecal samples contain white blood cells, which are predominantly monocytes and lymphocytes (3). Bacteria associated with enteric fever include typhoidal *S. enterica* and *Brucella* spp. (2). These organisms generally cause severe disease that is characterized by a disseminated infection in different parts of the body. The immune response to enteric fever is characterized by interstitial inflammation associated with mononuclear cell infiltration rather than a strong neutrophil response (3, 13). In *S. enterica* serovar Typhi, the presence of *Salmonella* pathogenicity island 7 (SPI-7) and factors encoded in its *viaB* locus, including the Vi capsular antigen, prevents recognition of PAMPs on the cell surface, such as LPS and flagella, by TLR4 and TLR5, respectively. Activation of these TLRs is associated with increased vascular permeability and neutrophil attraction associated with inflammatory diarrhea (14). *Brucella* spp. employ a distinct approach from that employed by *S. Typhi* to evade recognition by TLR4 and TLR5, in that *Brucella* spp. have modified LPS and flagellin structures that are not efficiently recognized by TLRs (15, 16). The ability to evade

recognition by the innate immune system allows the organisms to establish systemic infections associated with enteric fever (2).

Of these enteric pathogens, one of the most commonly detected, in terms of both numbers of human infections and severity of disease (numbers of hospitalizations and deaths associated with infections), is *Salmonella enterica* (17). Salmonellosis can be manifested through different syndromes, including gastroenteritis, enteric fever (typhoid fever), and bacteremia, and as asymptomatic carriage in animals and humans (18). There are approximately 1.03 million cases of nontyphoidal *Salmonella* that occur each year in the United States, costing the economy approximately \$3.31 billion due to premature mortality, disability, and medical and productivity costs, with an annual loss of 16,782 quality-adjusted life years (QALYs) (19). Compared to cost-of-illness figures (medical, employment, and death), the QALY losses also incorporate “psychometric scales” to quantify adverse health status due to people's discomfort and their abilities to engage in routine day-to-day activities following infection (19). *Salmonella* has adapted to survive in a wide range of different environments, such that a large number of human infections are associated not only with food animal sources but also with pets, reptiles, fruits, vegetable, legumes, and other humans (20–22). Among these sources, poultry and poultry-associated products are widely recognized as being among the most important vehicles for human *Salmonella* infections (20, 23). Therefore, *S. enterica* associated with chickens is likely a good model to explore bacterial host adaptation and pathogenesis, which is a key focus of this review.

OVERVIEW OF SALMONELLA COLONIZATION AND PATHOGENESIS

The genus *Salmonella* is broken up into three different species, *S. bongori*, *S. enterica*, and *S. subterranea* (24, 25). Previously, the genus was broken up into many more species, with each individual serotype being considered its own species. The genus was subsequently divided into seven subgenera (I, II, IIIa, IIIb, IV, V, and VI) based on biochemical and genetic properties (24). As more advanced genetic techniques such as DNA-DNA hybridization were used to analyze the members of the genus, it was discovered that many of the serotypes shared a high degree of genetic similarity. Consequently, the genus was divided into two species, *S. enterica* and *S. bongori*, with *S. bongori* containing the members of subgenus V and *S. enterica* containing the members of the remaining six subgenera (24). *S. subterranea* was described as a species in 2005 (25). Serotypes in what is now *S. enterica* subspecies *enterica* (subspecies I) are the predominant pathogens associated with birds and mammals (24).

Within subspecies I, there is a diversity of *Salmonella* serotypes that infect different animal hosts. The ability of different *Salmonella* serotypes to survive and thrive in different host environments involves a number of interconnected factors, including differences in host environments (pH, temperature, and sites of attachment, etc.), the host immune system and its response to different serotypes, the commensal organisms present, and the genetics of the pathogen itself (26, 27). Many serotypes, such as those most commonly associated with human infections, including *S. Enteritidis*, *S. Typhimurium*, *S. Newport*, and *S. Heidelberg*, have a broad host range, with the ability to infect multiple host species (28). Conversely, other serotypes, such as *S. Typhi*, *S. Paratyphi*, *S. Gallinarum*, *S. Choleraesuis*, *S. Abortusovis*, and *S. Dublin*, have restricted host ranges and are associated primarily with

one or a few hosts (29). *S. Dublin* and *S. Choleraesuis* are typically associated with cattle and swine, respectively; however, they also cause infections in humans (30–33). Human infections caused by these narrow-host-range serotypes tend to be fewer (with the exception of human-associated serotypes such as *S. Typhi* and *S. Paratyphi*) yet typically more severe in disease outcome. Data from the Centers for Disease Control and Prevention (CDC) FoodNet Program indicate that over 50% of human cases of *S. Typhi*, *S. Dublin*, *S. Paratyphi A*, and *S. Choleraesuis* infections are invasive rather than limited to gastroenteritis (34). Lower percentages of infections by the broader-host-range serotypes, such as *S. Typhimurium* (6%), *S. Enteritidis* (6%), and *S. Heidelberg* (11%), are invasive in nature (34).

Initial Colonization of the Host

The primary route of infection in humans and animals is through fecal-oral transmission of *Salmonella*. *Salmonella* pathogenesis has been studied mostly as it relates to human infections, while there is more limited information about the mechanisms of colonization and pathogenesis in food animals such as chickens. In general, when food contaminated with *Salmonella* is ingested, the bacteria have to pass through the alimentary system and survive the acidic environment of the stomach. *Salmonella* has been found to respond to the acidic environment through a complex adaptive system, called the acid tolerance response, which requires the synthesis of over 50 acid shock proteins, including the RpoS σ -factor, PhoPQ, Ada, and Fur (35, 36). Bearson et al. reported that *S. Typhimurium* RpoS and PhoPQ provided protection against inorganic acids, while RpoS and FurR offered protection against organic acids (35).

Those *Salmonella* organisms that survive the low-pH environment proceed to the lumen of gastrointestinal tract (GIT) organs, including the small intestine, colon, and cecum (in poultry). Epithelial and immune cells lining these GIT organs provide the initial protective barrier against *Salmonella* in the gut. *Salmonella* competes with the gut microflora to make the initial contact with enterocytes or M cells in order to colonize the GIT (37, 38). Adhesion to the GIT epithelium by *Salmonella* is facilitated by flagella and fimbriae present on the bacterial cell surface (39, 40). Studies have shown that *Salmonella* serovars employ both conserved and host-specific factors that facilitate colonization in the host GIT (41). Signature-tagged mutagenesis studies have reported the ability of multiple *S. Typhimurium* transposon mutants to colonize intestinal tracts of mice, calves (2), chickens, and pigs (42, 43). In addition to the oral-GIT route of invasion, *Salmonella* bacteria have been reported to invade and disseminate in swine and cattle following uptake in the tonsils and respiratory system (44, 45).

Invasion of Host Cells

Once attached to the intestinal epithelium, *Salmonella* bacteria typically express a T3SS, which is a multiprotein complex that facilitates endothelial uptake and invasion (28, 46). The T3SS apparatus acts as a “molecular syringe” (a conduit between the bacterial cytoplasm and the host cell membrane) to transport toxins and other effector proteins into intestinal cells and is associated with over 20 structural and regulatory proteins (47–49). Several effector proteins, such as InvJ, SpaO, PrgI/J, SipA/B/C/D, SptP, AvrA, SopA/B/D/E/E2, SlrP, and SspH1, are also secreted through the T3SS (47, 50, 51). The T3SS machinery is associated with SPI-1, which harbors virulence genes involved in *Salmonella* ad-

hesion, invasion, and toxicity (see below for details on SPIs). The structural apparatus of the SPI-1 T3SS has been reported to be vital for invasion of bovine and porcine ileal mucosa by *Salmonella* (41).

Among the effector proteins that are translocated through the T3SS, SopB plays an important role in the activation of secretory pathways, facilitating inflammation (by attracting neutrophils to the sites of infection) and altering ion balances within cells, leading to secretion of fluids in the GIT and consequent diarrhea (52, 53) in humans (54) and cattle (55). Other translocated proteins, such as SopA, SopD, SopE2, and SipA, have been found to play a role in *Salmonella* gastroenteritis (53). Effector proteins such as SipA, SipC, and SopB can interact with the actin cytoskeleton, causing the host cell membrane to extend outwards, a process referred to as membrane ruffling (50). This ruffling process facilitates engulfment of the *Salmonella* cells by the host cell membrane and internalization in a membrane-bound compartment termed a *Salmonella*-containing vacuole (SCV) (56–58). As the SCV matures, it migrates from the luminal border to the basal membrane, thus avoiding destruction by the phagolysosomal processing pathways (59).

The SCVs play key roles in survival and proliferation of *Salmonella* in intestinal cells and macrophages (60). Once internalized into host cells, *Salmonella* cells express a second T3SS encoded on SPI-2 (SPI-2 T3SS) within the SCV that plays a critical role in causing systemic infections and intracellular pathogenesis (61). The SPI-2 T3SSs are responsible for secreting effector proteins in the SCV, and these proteins interact with the cytoskeleton and motor proteins, leading to the formation of *Salmonella*-induced filaments (SIFs), which project out of the SCV (28). Furthermore, SPI-2 T3SSs have been reported to modulate trafficking of the SCV to avoid fusion with the lysosomes (41). Several effector proteins, such as SpiC, PipB, SseF/G/I/J, SifA/B, SspH1/H2, SlrP, SopD2, and phage-located SrfA/B/C/D/E/G/I/J/K/L/M, are secreted through the SPI-2 T3SS (62). The SIF facilitates fusion of SCVs with other vesicles in the cell and may play a role in *Salmonella* replication (58, 61, 63); the precise role of SIFs in *Salmonella* infections remains unclear (28).

Survival in Macrophages and Dendritic Cells

In a small percentage of cases, *Salmonella* cells are able to replicate within host cells, evade immune responses, and develop invasive and systemic infections (2, 37). These severe manifestations of salmonellosis usually occur when *Salmonella* cells invade macrophages or dendritic (migratory phagocyte) cells. *Salmonella* cells have been shown to be able to multiply within macrophages (64) but do not appear to replicate within dendritic cells, even though they remain viable (60). The exact mechanisms for the differences in *Salmonella* responses within different immune cell types are not entirely clear, as several pathogen and host factors may play roles (65). Dendritic cells are widely distributed in the lymphoid and nonlymphoid tissues and can rapidly facilitate the spread of *Salmonella* cells to various organs of the host body (66). Researchers have shown that the SPI-2 T3SS in *Salmonella* harbors genes that can suppress antigenic presentation in dendritic cells, which limits the host immune response to infected cells (62). In general, the ability of *Salmonella* to cause an infection in humans or animals depends on the innate ability of the bacteria to encode and express a set (or combination) of virulence genes that can evade and neutralize host defenses. These factors are associated with pathogenic-

ity islands, virulence plasmids, toxins, fimbriae, and flagella (28, 40, 67–69).

Salmonella Pathogenicity Islands

The ability of *Salmonella* to efficiently colonize the host has been attributed to gene clusters, such as SPIs, encoding virulence factors that are distributed in the *Salmonella* genome (28). Several major pathogenicity islands have been reported for different serovars, with SPI-1 to -5 being present in most serovars and others being less widely distributed (28, 69–71). In general, SPI-1 is required for invasion of host cells and induction of macrophage apoptosis, SPI-2 is required for systemic infection and replication within macrophages, SPI-3 is required for survival in macrophages and the ability of *Salmonella* to grow in low-magnesium environments, SPI-4 is required for intramacrophage survival and harbors genes for toxin secretion and apoptosis, SPI-5 has been found to cluster genes that encode multiple T3SS effector proteins, and SPI-6 has been found to transport proteins into the environment or host cells in response to external stimuli (28, 40, 41, 69, 70, 72, 73). Amavisit et al. reported genetic variations among SPI-1, -3, and -5, while SPI-2 and -4 were well conserved among 13 different *Salmonella* serovars isolated from warm-blooded animals (bovine, porcine, avian, and equine), the environment, and human patients in their study (73). Those authors found that, with the exception of *S. Typhimurium*, all isolates within the same serovar were identical with regard to the five SPIs that were tested. SPI-1 and SPI-2 have been found to play a role in *Salmonella* persistence and enteritis in chickens (43, 74), cattle (75–77), pigs (42), and humans (37). Fibronectin-binding proteins, encoded by SPI-3, facilitate host-specific *Salmonella* colonization. For example, MisL contributes to *Salmonella* colonization in mice, chickens, and pigs but does not play a significant role in calves (42, 43), while ShdA influenced *S. Typhimurium* persistence in mice but not in pigs (78, 79). SPI-4 has been reported to mediate adherence to and invasion of bovine ileal mucosa, possibly in combination with the SPI-1 T3SS, but not in chickens and pigs (41, 43, 80).

Virulence Plasmids

Strains from several *Salmonella* serovars have serotype-specific virulence plasmids (81). These are low-copy-number plasmids (1 to 2 copies per cell) and range from 50 to 100 kb, depending on the serovar (40). Each of the plasmids contains the *Salmonella* plasmid virulence (*spv*) locus, whose expression has been reported to be important for multiplication of *Salmonella* in the reticuloendothelial system, including the liver and spleen (40, 82). Other plasmids, in addition to the serotype-associated virulence plasmids, also likely contribute to the observed resistance among *Salmonella* bacteria. Recent studies from our laboratories have identified several different plasmids that potentially contribute to virulence in serovars such as *S. Heidelberg*, *S. Kentucky*, and *S. Typhimurium* (83, 84). A more in-depth exploration of plasmid-associated virulence is presented below.

Toxins

Salmonella pathogenicity has also been attributed to the production of endo- and exotoxins. Endotoxins have been found to elicit a wide range of biological responses, while exotoxins, comprising cytotoxins and enterotoxins, have the ability to kill mammalian cells (85). Ashkenazi et al. reported that *Salmonella* serovars Chol-

eraesuis, Enteritidis, and Typhi produced heat-labile, trypsin-sensitive cytotoxins with various molecular masses, including 56 kDa (*S. Typhi*), 70 kDa (*S. Typhimurium*), and 78 kDa (*S. Choleraesuis*) (85). A *Shigella dysenteriae* 1-like cytotoxin has been detected in *Salmonella* serovars Enteritidis, Kapemba, and Thompson (86). Two other types of exotoxins, salmolyisin (encoded by the *slyA* gene) and *Salmonella* enterotoxin (Stn; encoded by the *stn* gene), have been identified in *Salmonella* serovars Typhi, Typhimurium, and Enteritidis (87, 88).

Fimbriae

Fimbriae (pili) are filamentous surface structures that contribute to the colonization of the epithelium by *Salmonella* (89). Each of the fimbrial operons contains multiple genes (typically 8 to 11) that encode the structure and assembly of fimbrins (fimbria proteins) (90). Several fimbrial operons, ranging from 7 to 9 kb in size, have been identified in *Salmonella*. The sequenced strain *S. Enteritidis* PT4 has 13 fimbrial operons (91). Some examples of fimbrial operons include the *agf* and *sef* operons, which encode the *S. Enteritidis* fimbria SEF17 (92, 93); the *pil* operon (located in SPI-7) in *S. Typhi* CT18 (94); and the *lpf* (long polar fimbriae) and *pef* (plasmid-encoded fimbriae) operons in *S. Typhimurium* (95, 96). The SEF14 fimbriae have been found to be expressed by *S. Enteritidis*, *S. Dublin*, and poultry-associated *Salmonella* serovars Berta and Gallinarum, where they appear to be important for adhesion of these serovars to tissues of the reproductive tract (97, 98). Type I fimbriae contribute to *Salmonella* colonization of pigs (99), while 13 major fimbrial subunits of *S. Enteritidis* PT4 have been found to play a role in adherence and colonization of the bacteria in chicken gut (100, 101); the loci on which these subunits were detected were conserved in *S. Paratyphi* and *S. Gallinarum*. A detailed distribution of fimbrial operons among *Salmonella* subspecies and serovars was highlighted in a review by van Asten and van Dijk (40).

Flagella

The majority of *Salmonella* serovars possess up to 10 randomly positioned flagella on their cell surface, which confer motility to these bacteria (40). The ability of certain serovars to display flagellin phase variation provides a potential means for the organisms to minimize the host immune response by creating phenotypic heterogeneity of the flagellar antigens (40). The *fliC* gene, encoding the phase 1 flagellin protein, has been found in *Salmonella* serovars Gallinarum and Enteritidis (102). However, the exact role of flagella (motility and direction of rotation) in *Salmonella* pathogenesis and their possible role in adhesion and invasion of mammalian cells remain unclear (40).

Other Virulence Factors

Virulence factors such as surface polysaccharides may also play a role in persistence of *Salmonella* in the intestinal tract. Multiple mutants affecting lipopolysaccharide (LPS) biosynthesis have been identified in *Salmonella* strains isolated from calves and chickens (41–43, 103). For example, the virulence of the LPS *rfbK*, *dksA*, *hupA*, *sipC*, and *ptsC* mutants and *clpB* and *rfaY* transductants was studied in 1-day-old chicks by Turner and coworkers (103). That study showed that all but the *ptsC* and *rfaY* mutants were attenuated for virulence in chickens. Signature-tagged mutagenesis showed that LPS *S. Typhimurium* mutants (*rfaK*, *rfaB*, *rfaG*, *rfbP*, *rfbN*, *rfbU*, *rfbH*, and *rfaA*) were unable to colonize calf

intestines, suggesting a role of surface polysaccharides and cell envelope proteins as virulence factors contributing to *S. Typhimurium* colonization of calves (43). LPS has been found to contribute to the ability of *S. Enteritidis* to survive in egg albumen (104). That study showed that a mutant strain unable to produce LPS ($\Delta rfbH$) was not able to multiply in eggs at room temperature and did not survive in egg whites at 42°C. Those authors concluded that attenuation increased susceptibility of the $\Delta rfbH$ mutant to undefined antibacterial components of egg albumen (104).

IMPACT OF HORIZONTAL GENE TRANSFER ON GENOME EVOLUTION AND HOST RANGE

There are two major, seemingly opposite, evolutionary approaches in bacteria that impact the range of hosts which an organism can infect and in which it can survive; these approaches are gene acquisition through horizontal transfer and gene or gene function loss (105, 106). *Salmonella* serotypes have expanded their host ranges through the acquisition of new genes through horizontal gene transfer of plasmids and pathogenicity islands, which in turn provide genetic factors that increase the ability to infect new hosts. The addition of these pathogenicity islands to the *Salmonella* genome led to the ability to invade host cells and persist in causing disease. Some transfer events happened more recently, and thus, not all *Salmonella* serovars have acquired the clusters, while other events happened more distantly, and thus, the gene clusters are present in multiple lineages of *S. enterica* (73). There are five SPIs (SPI-1 to -5) that have been found to be important for the ability of *Salmonella* to become pathogenic relative to the commensal *E. coli* (107). It is estimated that 5 to 7% of the genes in the *Salmonella* genome play a role in virulence; interestingly, many of these genes are also present in the commensal *E. coli*; however, they alone are not enough to cause disease (108, 109).

It has been hypothesized that *Salmonella* virulence evolved in a multistep fashion, starting with the acquisition of SPI-1 by all *Salmonella* bacteria and acquisition of additional SPIs distinguishing *S. enterica* from *S. bongori*, followed by expansion of the host range to warm-blooded vertebrates (110). The SPIs, with the exception of a few, are highly conserved across members of the species *S. enterica* but are absent from closely related species such as *S. bongori* and *E. coli* (111). While most of the SPIs are not likely to retain their mobility, the ability of some *Salmonella* genomic islands to excise and transfer has been demonstrated. This excision is also inducible for some regions upon exposure to host conditions, such as macrophage survival and oxidative stress (112).

Integrative conjugative elements (ICEs) are also present in *Salmonella*. ICEs appear to be major players in the horizontal transfer of genetic material between bacterial species, as they carry genes required for their own excision and transfer to recipient bacteria (113). This is exemplified in *Salmonella* by SPI-7, an ICE that is present in *Salmonella* but also among a variety of other *Enterobacteriaceae* (114). SPI-7 ICEs have been found in both *S. enterica* and *S. bongori* and in other bacterial species, and they appear to be evolutionarily related, suggesting that both inter- and intraspecies ICE transfer occurs and further enhances ICE diversity (115).

Role of Plasmids in *Salmonella* Genome Evolution

Considerable research has focused on the plasmids that circulate among *Salmonella* populations of humans and animals, and there is no doubt that plasmids are indigenous to *Salmonella*. Evidence for this exists through a straightforward analysis of the historical

ECOR and SAR reference collections, which reveal a great deal of diversity among their plasmids and an abundance of plasmids in strains of both collections (116). Indeed, a variety of plasmid types have been found among *Salmonella* isolates, including those of incompatibility (Inc) types IncA/C, IncF, IncH, IncI, IncL/M, IncN, IncR, and IncX (117–122). What seems to have changed over time is the accessory gene content that these plasmids carry. It can be argued that genes encoding antimicrobial resistance are omnipresent in microbial populations and that they surface under selective pressure. While this is likely true, it is difficult to argue against the notion that multidrug resistance has increased among clinical *Salmonella* isolates. Historical isolates generally lack phenotypic resistance patterns but still contain plasmids. In contrast, contemporary *Salmonella* isolates still harbor plasmids, but many seem to have evolved in the face of selective pressures to acquire resistance-associated gene arrays. The purpose of this section is not to document the hundreds to thousands of articles that have demonstrated the presence of plasmids among *Salmonella* but to instead present insights into the impact of their presence on genome evolution.

Salmonella virulence plasmids have been considered a defining trait of *Salmonella* serotypes involved in systemic disease. These plasmids range in size from approximately 50 to 100 kb and are found in numerous serovars, including *S. Typhimurium*, *S. Enteritidis*, *S. Dublin*, *S. Choleraesuis*, and *S. Gallinarum* (123). The *Salmonella* virulence plasmid harbors several key virulence factors, including the *spvABCD* system and its regulator *spvR*, which are essential for systemic virulence and are induced upon exposure to nutrient-limited conditions and host cells, particularly macrophages (124–126). These genes are also sufficient to restore systemic virulence in plasmid-cured strains (127). These genes appear to be horizontally acquired on an IncFIIA/IncFIB plasmid backbone, since they have a lower G+C content than the core plasmid (71). There are multiple lineages of the *Salmonella* virulence plasmid that are correlated with a respective serovar, and the virulence plasmids among these lineages have uniquely evolved (Fig. 1). The plasmids sequenced and analyzed thus far can be broadly distributed into three major lineages, and it appears that they have evolved through a combination of gene acquisition and deletion events.

Several possible scenarios for the evolution of *Salmonella* virulence plasmids have been proposed (123, 128). It is possible that an ancestral IncFIIA *spv*⁺ plasmid present in serogroup D isolates diverged into other serogroups and acquired additional traits. A more likely scenario based upon recent work is that an ancestral plasmid type possessed an IncFIIA/IncFIB backbone containing *spv*, *prf*, and *pef* and separated following the divergence of serogroups and that the plasmid lineages subsequently underwent recombination events, resulting in the current virulence plasmid profiles. Interestingly, there is little evidence of extensive transfer of virulence plasmids among the *Salmonella* serovars, even though their transfer *in vivo* has been documented (129). The horizontal transfer of these plasmids is likely a neutral trait in *Salmonella*, and thus, transfer between serovars is not strongly selected (130, 131). In fact, not all strains of serotypes that typically contain virulence plasmids have them; this lack of virulence plasmids is likely reflective of differences in the strain's genetic background that impact the ability to utilize the plasmid for establishing systemic disease (132).

The *S. Typhimurium* virulence plasmid has also been shown to




FIG 1 Evolutionary history of *Salmonella* virulence plasmids inferred by using the maximum likelihood method, based on the Hasegawa-Kishino-Yano method. The bootstrap consensus tree was inferred from 100 bootstrap replicates. Analysis was based on 441 SNPs identified from the conserved sequences of 22 plasmids. Analysis was conducted with MEGA5 (403).

be capable of carrying resistance-associated integrons and transposons (133, 134). Cointegrated plasmids have surfaced in *S. Dublin* which include regions of the *Salmonella* virulence plasmid and plasmids of other incompatibility groups, such as IncX, encoding multidrug resistance (135). This could aid in the expansion of the host range of this plasmid to additional *Salmonella* serovars (136). Cointegrated IncFIB-IncI1 plasmids have also been identified among sequenced *Salmonella* genomes, and they have been found to belong to a unique clade of IncI1-type plasmids and harbor a unique subset of putative virulence genes (115). Certainly, cointegrated plasmids offer an increased potential to harbor more diverse arrays of resistance-associated genes and to expand host range and transmissibility.

In addition to virulence plasmids, *Salmonella* bacteria also harbor a variety of other plasmid types. It seems that the spread of such plasmids, in contrast to the *Salmonella* virulence plasmid, occurs primarily via horizontal gene transfer (137). Resistance-associated plasmids are found among a wide range of *Salmonella* serovars, in a variety of shapes and sizes (138–140). Plasmids belonging to groups such as IncI1, IncH, IncN, and IncX are not considered emergent in *Salmonella*, as they have been isolated from this genus for many years (118, 141). However, they appear to be actively evolving to acquire resistance genes with relevance to therapeutic antibiotics used in humans and animals. In contrast, IncA/C plasmids are not found in historical collections of *Salmonella* isolates but have recently emerged among *Salmonella*, *E. coli*, and *Klebsiella pneumoniae* isolates (142). The emergence of IncA/C plasmids among *Salmonella* isolates has often coincided with the emergence of dominant clonal types, such as serotype 4,[5],12:i:- in Spain (143). In some cases, IncA/C plasmids ap-

pear to have integrated into the *Salmonella* genome. A case in point was highlighted by a study by Shahada et al., which identified a unique 125-kb genomic island among isolates of a dominant clone of *S. Typhimurium* from cattle in Japan. This island, named GI-VII-6, displayed striking similarities to portions of IncA/C plasmids containing three copies of *bla*_{CMY-2} (144).

Other resistance plasmids are also emerging, with increased concern to health. IncN plasmids found in *Salmonella* and *E. coli* were recently associated with possession of genes encoding quinolone resistance (122, 145, 146). IncHI1 and IncHI2 plasmids have been linked to quinolone resistance-encoding genes, extended-spectrum beta-lactamase (ESBL) genes, heavy metal resistance genes, and carriage of typical complex class 1 integrons (119, 147–149). IncI1 plasmids are widespread among *Salmonella* isolates and are notorious for their carriage of *bla*_{CMY} and *bla*_{TEM} genes (137–139). As mentioned above, IncA/C plasmids have been identified more recently among *Salmonella* isolates, and they carry large numbers of resistance-associated genes, including *bla*_{CMY-2}, on a conjugative transposon encoding resistance to third-generation cephalosporins; a gene module containing *floR-strAB-tetAR*, encoding resistance to chloramphenicol-florfenicol, streptomycin-spectinomycin, and tetracycline; and a complex class 1 integron harboring a variety of resistance genes (121, 150, 151).

Small plasmids harboring quinolone resistance genes have been identified in both *Salmonella* and *E. coli* (122, 152, 153). More importantly, it exemplifies a mechanism for plasmid gene swapping via Xer- and *oriT*-mediated site-specific recombination. This mechanism of recombination can occur independently of conjugative transposons or integrons and could represent an additional means of genetic recombination involving plasmids. Other mech-

anisms of additional recombination potential exist on the *Salmonella* chromosome. For example, variant regions of *Salmonella* genomic island 1 (SGI-1) have been identified, which include additional resistance genes driven by IS6100-mediated recombination (154). A number of variants of SGI-1 that confer resistance to a variety of classes of antimicrobial agents have been identified (155).

In addition to plasmids being mobile themselves, they harbor a variety of mobile genetic elements within them, including integrons and conjugative transposons. Class 1 integrons are the most common type among *Salmonella* isolates and can be localized to the bacterial chromosome or plasmids. A large number of novel gene cassette arrays have been identified among class 1 integrons harbored by *Salmonella*, and some appear to have been derived recently from other bacterial species (156). These accessory regions are actively evolving on plasmids, as evidenced by recent comparative genomics-based studies of closely related plasmids with a variety of resistance island variations (149).

SGI-1 is a 43-kb accessory region of the *Salmonella* genome that contains a class 1 integron (*In104*) encoding resistance toward ampicillin, chloramphenicol-florfenicol, tetracycline, sulfamethoxazole, and streptomycin-spectinomycin, also known as classical “pentaresistance.” This region is found in many *Salmonella* isolates and, as mentioned above, is mobilizable. This alone enables a multidrug resistance phenotype, but the carriage of additional plasmids by SGI-1-containing strains is common (157). Interestingly, some plasmids have also been shown to enhance excision and variant formation of *Salmonella* genomic islands, such as the enhanced excision of SGI-1 by plasmids belonging to the *IncA/C* type (158, 159). In addition to enhancing the excision of chromosomal islands, these plasmids may also harbor receptor sites for phage recognition. For example, plasmids such as *IncN* and *IncX* have been shown to convert phage types among *Salmonella* strains (160, 161), and *IncX* plasmids seem to harbor phage recognition sites (162). Thus, when considering the role of plasmids in genome evolution, it is important to consider not only the accessory gene repertoire that they carry but also the modulations that they confer to the *Salmonella* host.

Horizontal transfer of individual mobile elements between different plasmids also occurs. There are numerous examples of this, but one of recent concern involves an emergent gene known as *bla*_{NDM-1}, encoding the New Delhi metallo-beta-lactamase (163). The *bla*_{NDM-1} gene was originally identified among *Klebsiella pneumoniae* isolates from infections originating in India. However, it was quickly realized that this gene was plasmid borne, was present on multiple plasmid types (164–167), and was rapidly spreading between bacteria (168). The first report of *bla*_{NDM} in *Salmonella* involved a patient with a *Klebsiella pneumoniae* infection that was resistant to carbapenems (169). Subsequent culturing identified NDM-positive *Klebsiella* and *Salmonella* within the patient, with *bla*_{NDM} apparently residing on different plasmids in each bacterial isolate. There are many additional examples of genetic transfer between *Salmonella* and other bacterial species. For example, heavy metal resistance operon transfer from *Klebsiella* spp. and environmental bacteria to *Salmonella* has also been hypothesized (170). Certainly, plasmid transfer and plasmid gene transposition play a major role in the evolution of *Salmonella* (171).

Role of Bacteriophages in *Salmonella* Evolution

Bacteriophages are among the most diverse and abundant forms of microbial “life” on earth and have played important roles in bacterial evolution. Phages integrate into the host chromosome as a prophage, where they can impact expression of host factors as well as facilitate the transfer of DNA among bacteria through transduction (172, 173). There are at least 170 known phages that infect *Salmonella*, and these generally fall into five categories based on their genetic sequences, P27-like, P2-like, P22-like, lambdoid, and T7-like, along with outliers such as ϵ 15, KS7, and Felix O1 (174). There is a large amount of diversity that is increasingly being revealed among phage families, which reinforces the idea that recombination is occurring frequently between subgroups of phages and between unrelated groups of phages (175).

The host ranges of *Salmonella* bacteriophages are variable, with some being highly host specific and others being generalists. An example of a generalist is Φ X174, which has been demonstrated to lyse >99% of *Salmonella* isolates tested (176). Thus, the susceptibility of an organism to phage lysis (often used to describe the phage type of the organism) does not necessarily correlate with clonality in all *Salmonella* strains. In contrast, the host specificity of other phages to lyse a narrow range of bacteria has been utilized for subtyping of *Salmonella*. Phage typing has historically been a method of choice for characterizing *Salmonella*. The method is still used for typing of bacteria but typically in conjunction with molecular techniques. For example, single-nucleotide polymorphism (SNP) typing has been used in combination with phage typing to explore relationships between host genetics and phage susceptibility in *S. Typhimurium* (177). This work demonstrated that the majority of isolates of particular phage types, such as definitive type 104 (DT104), had a single evolutionary origin, while *S. Typhimurium* isolates belonging to other phage types, such as DT9 and DT135, seemed to have originated from multiple lineages, indicating a broader host range of the typing phages (177). Phage typing has also identified potential geographical differences among isolates; for example, in the United States, isolates of phage type 8 are the most commonly detected *S. Enteritidis* isolates, compared to phage type 4 in Europe (178).

Phages have played significant roles in *Salmonella* virulence through the regulation of expression of virulence factors and the introduction of a number of effector proteins that are implicated in pathogenesis (179). A detailed understanding of the roles that bacteriophages play in *Salmonella* pathogenesis has been complicated by phage diversity and *Salmonella* strain diversity; however, it is becoming increasingly clear that phages do indeed play major roles in terms of delivering functional effector molecules and adding to the *Salmonella* virulence regulatory networks. For example, SopE2 and SspH2 are effector proteins encoded adjacent to phage-like sequences in *Salmonella*. Others, such as SopE, GogB, SseI, SspH1, and SseK3, are found directly on various prophages of *Salmonella* (179). The distribution of these phage-encoded effector molecules is often serovar and even strain specific, suggesting that they are indeed incorporated via transducing phage (180). In some cases, phage-mediated acquisition of traits may confer a fitness benefit in certain niches. For example, a SopE-encoding phage was identified primarily in cattle-associated epidemic isolates, suggesting that SopE may confer a reservoir advantage in cattle (181–183). Lambdoid phages such as Gifsy-1 and Gifsy-2 have been shown to be important for modulating virulence in *S.*

Typhimurium. The Gifsy-1 prophage carries several potential virulence-associated genes, including *gipA* and *gogB*, which may play a role in colonizing the small intestine (184, 185). The Gifsy-2 prophage also carries multiple virulence factors, including *gtgE*, *gtgB* (*sseI*), and *sodC1*, which have been shown to be important for overall *Salmonella* virulence (184, 186).

Plasmids have been shown to alter the susceptibility of *Salmonella* to bacteriophages. In *S. Enteritidis*, IncN plasmid acquisition resulted in conversion of the phage type from type 4 to type 24 (160). Acquisition of IncX plasmids by *S. Enteritidis* has been shown to induce resistance to multiple bacteriophage types, thereby altering the phage type profile and facilitating phage conversion (161). Low-molecular-weight plasmids have been shown to exert similar effects on *S. Enteritidis* as well (187). In contrast, the acquisition of some plasmids can result in increased phage susceptibility (188). The precise mechanisms of phage conversion relative to plasmid acquisition have not been intensively investigated, but there is certainly a relationship between the two. The clustered regularly interspaced short palindromic repeats (CRISPRs) may also play an important role in the defense of *Salmonella* against foreign DNA from incoming plasmids and phages. A comparative analysis of 28 sequenced *Salmonella* genomes demonstrated that CRISPR-mediated immunity among specific strains might control short-term phenotype changes and mediate long-term sublineage evolution (189). Thus, it is evident that bacteriophages reshape the *Salmonella* genome not only through their direct contributions to the host bacterium but also through indirect interactions with other mobile elements and cellular regulatory networks.

IMPACT OF THE LOSS OF GENES OR THEIR FUNCTION ON GENOME EVOLUTION AND HOST RANGE

The loss of genes or gene function can also impact genome evolution and host range due to the inactivation of genes needed to infect multiple hosts (106). Host-adapted strains of *Salmonella* have a relatively large number of pseudogenes and chromosomal rearrangements associated with recombination of the 16S ribosomal gene (*rrn*) operons that impact their host range (190, 191). Host-adapted serovars often demonstrate a high degree of *rrn* rearrangement within their genome, which is likely the result of the adaptation to the specific host niche that the serovar occupies. This view was consistent with the finding that host-adapted and host generalist strains demonstrated similar rates of *rrn* rearrangement, and as such, the differences were due to adaptation rather than having a more highly mutable genotype (192). Additionally, host-adapted serovars such as *S. Typhi*, *S. Paratyphi A*, *S. Dublin*, *S. Choleraesuis*, and *S. Gallinarum* have a relatively high percentage of pseudogenes in their genomes, especially compared to the broad-host-range *Salmonella* serovars *Enteritidis* and *Typhimurium* (193). Pseudogenes are genes that have been inactivated by insertion, deletion, or nucleotide substitution, leading to a failure to express the gene product associated with the original gene. For example, the genomes of *Salmonella* serovars *Typhi* and *Paratyphi*, which are human-adapted serovars, have >4% of the coding sequences in their genomes as pseudogenes, while for the non-host-restricted *Salmonella* serovar *Typhimurium*, only about 0.9% of the coding sequences are pseudogenes (194). A key consideration related to host adaptation is whether the presence of pseudogenes and the corresponding loss of function led to the host restriction or whether the increased number of pseudogenes

was due to the serovars adapting to their new niches. Indeed, the accumulation of pseudogenes is likely due to host adaptation and subsequent genetic drift following adaptation to the new host niche (194). Approximately one-quarter of the genomes of *Salmonella* serovars *Typhi* and *Paratyphi A* have a high degree of nucleotide similarity to one another, while the remaining portions are more divergent, similar to the differences between the other serovars. It is likely that the region of higher similarity is due to recombination between the two human host-adapted serovars (194). Several (22) of the pseudogenes identified between *Salmonella* serovar *Typhi* and *Paratyphi A* isolates were outside the recombination region and thus likely arose independently in each of the two serovars. However, since the pseudogenes were present in both serotypes, they may be important for survival in the human host niche. The pseudogenes identified included *safe* and *sefD*, which are fimbrial genes, and *shdA*, *ratB*, and *sivH*, which are involved in intestinal colonization and persistence, pathological features not typically associated with typhoid and paratyphoid infections and, as such, not essential for *Salmonella* serovars *Typhi* and *Paratyphi A* (194).

Based on genome analyses, it appears that *Salmonella* pseudogenes are typically removed from the genome relatively early following their formation. The removal is potentially associated with the fitness costs associated with the replication of now noncoding DNA in the bacterial genome: the expression of partial, nonfunctional genes or toxicities associated with partially expressed genes (195). This selective pressure to remove the non-fully active genes is evidenced in that the host-restricted serovars have higher K_a (number of nonsynonymous substitutions per nonsynonymous site)/ K_s (number of synonymous substitutions per synonymous site) ratios than broad-host-range serotypes (195, 196). The higher K_a/K_s ratio is indicative of positive evolution selective pressure for genome change (195). Other studies with *S. Typhi* and *S. Choleraesuis* have found that the majority of pseudogenes are not expressed; however, some pseudogenes appear to have undergone recoding, such that there were secondary mutations that allowed expression of a functional protein (193). In contrast to full recoding, partial expression of truncated, non-fully functional proteins can potentially bind to receptors blocking the fully functional proteins from eliciting their roles in cellular physiology, thereby harming the bacterium (195).

Other factors also likely play a role in host range variability. These factors include the genetic adaptation of organisms to particular nutrients that are abundant in the host organism but limited in other environments (106). For example, in *S. Gallinarum*, there are multiple pathways associated with carbon and energy metabolism that are not functional, including the catabolism of D-glucarate and long-chain maltodextrins, which may limit their ability to survive in certain environments outside the avian host (191). Loss of gene function can also be associated with an increased ability to cause disease in some instances. A good example is *Shigella* species, where *OmpT* is absent, which reduces the early immune response to infection (105). *S. Typhi* is also able to modulate the early immune response through the repression of *fliC* expression, which limits the activation of TLR5 by the flagellin *FliC* (197). In both examples, the loss of the immune response facilitates the intracellular spread of the organism rather than being rapidly cleared with a stronger initial immune response.

EXAMPLES OF *SALMONELLA* HOST RANGE IN CHICKEN-RELATED SEROVARS

Because the diversity of the *Salmonella enterica* species is quite expansive, this section focuses primarily on the host range adaptation of four particular serovars (*S. Enteritidis*, *S. Heidelberg*, *S. Kentucky*, and *S. Gallinarum*) that are associated very commonly with chickens and to various extents with other food animal species and human infections. According to the CDC, in the United States, *S. Enteritidis* was the serovar most commonly implicated in human illness, overtaking *S. Typhimurium* as the most common serovar (198). Likewise, when data from the National Veterinary Services Laboratory of the USDA and from other studies examining the prevalence of *Salmonella* serovars were compared, *S. Enteritidis* was associated most commonly with chickens and eggs and to a much lesser extent with other food animal species (198–202). *S. Heidelberg* is found in most of the major food animal species, eggs, and retail meat samples and is among the top five most common serotypes associated with human illnesses (203, 204). Conversely, *Salmonella* serovars *Kentucky* and *Gallinarum* rarely cause human infections in the United States (although *S. Kentucky* is an emerging serovar in Europe and North Africa) (205). *Salmonella* serovar *Gallinarum* is a host-adapted serovar that is presently made up of two biovars, *Gallinarum* and *Pulorum* (which were previously considered two separate serotypes) (206). This serotype was associated with severe losses to the poultry industry in the United States until it was targeted for eradication by the National Poultry Improvement Plan (NPIP) starting in 1935 (207). After implementation of the NPIP, *S. Gallinarum* was eradicated from commercial poultry flocks in the United States by the mid-1960s (26).

S. Kentucky is currently one of the most common serotypes isolated from broiler chickens in the United States and is detected fairly often in dairy cattle as well (198). The increase in the rate of isolation of *S. Kentucky* in broilers is likely related to a number of factors, including management practices, flock immunity, and genetic changes in the organism (26). When *S. Kentucky* was compared to other serovars with respect to virulence, it was found that the *S. Kentucky* isolates grew more rapidly than other serovars under moderately acidic conditions (pH 5.5) but worse under more highly acidic conditions (pH 2.5) (208). The enhanced ability to grow under these moderately acidic conditions may provide an advantage over other serotypes in environments such as the chicken cecum. This serovar exhibited greater invasiveness in *in vitro* assays using chicken embryo hepatocytes than serovars such as *S. Enteritidis* and *S. Typhimurium* (208). Many *S. Kentucky* isolates also have plasmids with factors such as those associated with antimicrobial and disinfectant resistance, iron acquisition, bacteriocin production, and complement resistance, which may enhance their abilities to survive in birds (83).

It is interesting that *S. Kentucky* has not become a larger public health problem for consumers in the United States, as *Salmonella* from poultry is predicted to be the fourth most important pathogen-food combination associated with food-borne illnesses in the United States (20). It is quite probable that *S. Kentucky* strains most common in poultry are not overly virulent to humans. In analyzing the data from the CDC National *Salmonella* Surveillance Program (<http://www.cdc.gov/ncidod/dbmd/phlisdata/salmonella.htm>), the states with the largest numbers of human *S. Kentucky* infections were generally those associated with higher rates of dairy production rather

than poultry production (198, 209). Likewise, several studies have utilized pulsed-field gel electrophoresis (PFGE) for the molecular characterization of *S. Kentucky* isolates from various sources (chickens, dairy cattle, and human infections); the most common profiles associated with human infections were similar/visually indistinguishable from common profiles from cattle but typically distinct from those of poultry (205, 210–212). Likewise, a majority of *S. Kentucky* strains isolated from poultry-associated sources fall into the sequence type 152 (ST152) clonal complex, while the majority of those isolated from human patients were of ST198 (205). Therefore, a plausible explanation for *S. Kentucky*-associated human infections could be due to occupational exposure to cattle or due to consumption of contaminated products such as raw (unpasteurized) milk or raw-milk cheeses. *S. Kentucky* has been one of the most commonly detected serotypes isolated from prepasteurized milk in the United States (213), and consumption of raw milk and raw-milk cheese has been associated with multiple outbreaks of salmonellosis (214–216).

For *S. Heidelberg*, there does not seem to be a strong association of subtypes with a particular food animal host and subsequent human infection. When *S. Heidelberg* isolates from human patients were compared to those of the major food animal species, there was extensive overlap in PFGE profiles, plasmid types, and antimicrobial susceptibility profiles, indicating a lack of host restriction among *S. Heidelberg* genotypes (203, 217, 218). In a study examining the core genomes of the population structure of many of the prominent serovars, it was concluded that the genomes of *S. Heidelberg* isolates were likely shaped by a high degree of horizontal genetic transfer (219). Consequently, the *S. Heidelberg* strains resided in a lineage distinct from that of the avian-associated *Salmonella* serovars *Enteritidis* and *Gallinarum*, based on genomic comparison. Additionally, the members of *S. Heidelberg* fell outside the lineage containing *Salmonella* serovars *Typhimurium* and *Saintpaul*, yet *S. Heidelberg* isolates shared a relatively high proportion of sequence similarity with the lineage (219).

S. Heidelberg isolates also often contain plasmids. While they lack a serotype-specific virulence plasmid, a common feature in serovars such as *S. Typhimurium*, *S. Enteritidis*, and *S. Gallinarum* (81), they often contain plasmids with virulence genes (83, 84). Interestingly, a number *S. Heidelberg* strains isolated from poultry-associated sources were found to harbor IncFIB plasmids similar to those previously recognized as being important for extraintestinal survival in avian-pathogenic *E. coli* (83, 220–222). Similar plasmids have been found very commonly in *S. Kentucky* isolates from poultry as well. Many IncFIB plasmids contain genes for iron acquisition (aerobactin operon and Sit iron transport systems), colicin production, and serum survival, which likely play a role in increased fitness in the avian environment (83). In many cases, these IncFIB plasmids also contain genes that encode resistance to multiple antimicrobials (often including resistance to tetracycline, streptomycin, chloramphenicol, and sulfonamide). This presents the possibility for coselective pressure, with antimicrobial use selecting for enhanced virulence, or conversely, the increased ability of these bacteria to survive iron-limited environments in the host could select for resistance to one or more antimicrobials (83, 84). *S. Heidelberg* plasmids also contain genes encoding disinfectant and heavy metal resistance, which may provide a selective advantage for survival in the avian production environment where pathogen control strategies are employed (84, 218).

Studies have shown that *Salmonella* serovars Gallinarum and Enteritidis are closely related in both their gene content and their antigenic formula (191, 223–225), suggesting that these two serovars originated from a relatively recent common ancestor (226, 227). Because of their close genetic similarity, they are valuable examples to explore host adaptation. Genomic comparisons of members of these two serovars have indicated that there is a high degree of genetic similarity between the two serotypes, with average nucleotide identities among orthologous genes being 99.7% (191). The genomes of sequenced isolates shared 4,179 predicted coding sequences (CDSs). The differences in CDS content between the strains were associated primarily with bacteriophages incorporated into the respective bacterial genomes (191). The lineage of *S. enterica* that contains these serotypes tends to be one of the most conserved among the species, with an estimated 4% of their core genome sequences originating from recombination with genes from other *Salmonella* serovars (219). Both serovars share the O1, O9, and O12 antigens (223). However, even though *S. Enteritidis* and *S. Gallinarum* are relatively closely related on the genomic level, they are diverse in their numbers of pseudogenes. *S. Gallinarum* isolates are unable to carry out mannose-sensitive hemagglutination and do not express flagellar genes, leading to an observed lack of mobility, while the majority of *S. Enteritidis* strains are motile (110). This lack of motility observed for *S. Gallinarum* isolates is in part associated with mutations in genes associated with flagellar biosynthesis and chemotaxis, including *flhA*, *flhB*, *flgI*, *flgK*, or *cheM* (191). Mutations in *fliC* may also contribute to the lack of motility of isolates of this serovar (226).

One study found that *S. Enteritidis* has 21 pseudogenes, compared to 147 for *S. Gallinarum*, but there are only 5 pseudogenes shared between the two serovars (195). Of these five pseudogenes, three are likely ancestral in origin, and the other two were likely independently acquired. Similarly, the genetically related *S. Dublin* had 212 identified pseudogenes, 177 of which are active genes in *S. Enteritidis*. Many of these functional genes encode surface structures or are involved in the central metabolism of *Salmonella*; thus, their inactivation in *S. Dublin* likely contributes to its host restriction in cattle (91). Likewise, certain pseudogenes are common to multiple host-restricted serotypes; for example, *mglA* and *shdA* are transcribed in *S. Enteritidis* but are present as pseudogenes in *S. Choleraesuis*, *S. Dublin*, *S. Gallinarum*, *S. Paratyphi A*, and *S. Typhi* (91). *MglA* is a small GTP-binding protein subunit of a binding-protein-dependent galactose transport system (228, 229) that likely plays a role as a virulence regulator in intracellular pathogens such as *Francisella* spp. (230) and motility in *Myxococcus xanthus* (231). *ShdA* is a protein that is involved in the colonization of Peyer's patches by *S. Typhimurium* and the shedding of the bacteria following infection (232, 233). The presence of a larger number of functional genes in broader-host-range serovars such as *S. Enteritidis* likely contributed to the ability to colonize and infect a greater number of hosts (234).

There are also several differences in the numbers of the fimbrial genes between *S. Gallinarum* and *S. Enteritidis*. Several fimbrial operons, including *lpf*, *bcf*, *stb*, *stc*, *std*, and *sth*, are important for long-term carriage and shedding of *Salmonella* (37). The *S. Enteritidis* genome has 13 fimbrial operons, 12 of which are present in *S. Gallinarum* (191). However, several of the CDSs within these common operons were identified as pseudogenes due to mutations that potentially prevent the expression of the functional gene product, which consequently impacts the overall function of the

operons. These include mutations in *stiC*, *stfF*, *safC*, *stbC*, *cheM*, *flhB*, *flhA*, *flgK*, *flgI*, *pegC*, *lpfC*, *sefC*, *sefD*, *sthE*, *sthB*, and *sthA* (191). Additionally, the virulence plasmids in the two serotypes contain fimbrial operons; however, they are unique, with the *S. Enteritidis* plasmids carrying the plasmid-borne fimbria (*pef*) operon, while the *S. Gallinarum* plasmids have fimbria genes similar to those of *E. coli* K88 (40, 81, 123, 191, 235).

The sequenced *S. Gallinarum* isolate also lacked some of the T3SS effectors that are present in the *S. Enteritidis* genome, and *sopA* had an early stop codon that likely prevents the expression of a functional SopA protein (191). In addition, there was a mutation in *bscG* of *S. Gallinarum*, whose gene product is predicted to be important for cellulose biosynthesis, which likely contributed to the observed deficiency in its ability to form biofilms (191). Taken together, these mutations may negatively impact the ability of *S. Gallinarum* to colonize mammalian hosts, thereby limiting its ability to cause illnesses in a broad range of host species.

In addition to mutations in *sopA*, several studies have demonstrated the influence of other T3SS effectors on the ability of *Salmonella* serovars to colonize different hosts (206, 236, 237). The T3SSs are multiprotein complexes composed of structural and regulatory elements that transfer functional effectors from the bacterium into host cells, thereby facilitating invasion of and survival within host cells (48, 49). The T3SSs play vital roles in the interaction of *Salmonella* with the host (238). Variations in the gene sequences encoding T3SS factors for both the SPI-1 and SPI-2 systems have been associated with differences in the abilities to invade different hosts. Some of the main variability associated with host specificity is with the secreted effectors and the SipD T3SS tip protein rather than other structural components of the respective T3SSs (206, 236). The SPI-1 T3SS tip protein SipD is important for cell invasion; studies showed that antibodies directed at SipD block the ability of *S. Enteritidis* to invade intestinal epithelial cells (239). Effectors secreted by the SPI-1 T3SS may also play a role in host specificity/range; sequence variability in the genes encoding the effector proteins SipA, SopA, and SopE and the chaperones SicP and InvB showed close evolutionary similarity in *S. Enteritidis* and *S. Gallinarum* compared to other serovars not as closely associated with avian sources (206). Thus, these SPI-1-associated factors may play a role in specificity in the initial invasion of the intestinal epithelium or immune cells in birds. The inflammatory responses to infections by *S. Enteritidis* and *S. Gallinarum* are different, with *S. Enteritidis* eliciting a stronger inflammatory response in chickens, which may prevent a more systemic spread, while *S. Gallinarum* elicits a weaker initial immune response and typically leads to systemic fowl typhoid (240).

On the SPI-2 side, at least 13 SPI-2-associated genes in *S. Enteritidis* and *S. Gallinarum* show close evolutionary similarity to each other compared to other serovars (206). It was hypothesized that the sequences may have coevolved for survival in the avian host, since survival in the reticuloendothelial system has been shown to be important for host specificity in chickens (241). The SPI-2 T3SS plays an important role in survival in the SCVs of macrophages (58). Genetic variability in SPI-1 factors, such as SopE and/or SodC, along with SPI-2 factors, such as SseC, SseD, and/or SseF, may lead to differences in the abilities to survive in different host cells (236). The factors SseG and, possibly, SseF appear to impact the migration of the SCVs to the Golgi apparatus in the host cell, which may serve as a potential source of nutrition for *Salmonella* (242). Hence, differences in the effector proteins

may impact the ability to survive in different host cells and to be transported to sites of systemic infection. In addition to changes in the gene sequences themselves, posttranslational modifications of T3SS effectors can impact the targeting of *Salmonella* or the SCVs to different parts of the host cells. Observed posttranslational modifications, such as lipidation, phosphorylation, and ubiquitylation, can occur following translocation into host cells (243).

THE SALMONELLA-CHICKEN HOST INTERACTION

Each of the serovars discussed above are quite proficient at colonizing chickens; this section provides a more in-depth examination of the interaction of chicken-associated serovars and the avian host environment. One issue that continues to complicate attempts to understand host responses to *Salmonella* infection is the potential variability in host cell interactions among the different serovars. Several studies have been undertaken to gain knowledge of these interactions. Methner et al. conducted a series of colonization inhibition studies in young chicks gavaged with *Salmonella* strains belonging to the same or different serovars. The results demonstrated that single strains did not prevent colonization by other serovars, but mixtures of different serovars were effective against heterologous serovars (244). He et al. compared five poultry-associated serovars (*S. Typhimurium*, *S. Enteritidis*, *S. Heidelberg*, *S. Kentucky*, and *S. Senftenberg*) for their impact on an MC29 virus-transformed chicken macrophage cell line, HD-11, to assess *Salmonella* cell invasion, intracellular survival, and modulation of phorbol myristate acetate (PMA)-stimulated oxidative burst and nitric oxide (NO) activity. Neither *S. Enteritidis* nor *S. Typhimurium* generated detectable NO production by the macrophages, while the other 3 serotypes elicited production of over 100 μM NO. All of the serotypes repressed PMA-mediated oxidative burst and were equally able to invade HD-11 chicken macrophages; however, *S. Enteritidis* was more resistant to intracellular killing than the other 4 serovars (245). Those researchers speculated that these findings, particularly the intracellular survival ability of *S. Enteritidis*, may account for its association with systemic invasion in chickens (245). However, when newly hatched chickens were challenged with each of the 5 serovars, all but *S. Senftenberg* exhibited high rates of organ invasion (>85%, versus 14% for *S. Senftenberg*), which were different than the *in vitro* intracellular survival outcomes (245). These findings were consistent with studies on *S. Enteritidis* strains classified as displaying either low- or high-invasive phenotypes in Caco-2 human colon tissue *in vitro* assays (246, 247). Those studies demonstrated that when mice were orally challenged, low-invasive strains were as virulent as high-invasive strains, even though not all *S. Enteritidis* isolates recovered from poultry are equally pathogenic. Some of the infectivity differences between *S. Enteritidis* and other serovars in avian model systems may be related to the seemingly high epidemiological frequency and long-term association of *S. Enteritidis* with egg-related salmonellosis and a potentially unique evolutionary relationship with laying hens (248–256). This relationship may be evidenced by the limited genetic diversity observed among *S. Enteritidis* strains based on random amplified polymorphic DNA plasmid profiling and phage typing (257). Work with laying hens suggested that because *S. Enteritidis* infects reproductive tissues, namely, the oviduct and ovaries, these tissues represent critical avenues for egg contamination; additionally, this serovar may also have evolved the means to survive the hostile environment of internal egg contents (252, 258–267).

More recently, attempts have been made to identify unique genetic traits which distinguish *S. Enteritidis* from other serovars and lead to its propensity for laying hens and eggs. Methner et al. demonstrated that *S. Enteritidis* strains exhibited substantial colonization inhibition of other *S. Enteritidis* strains but not of other serovars in young chicks, suggesting that there may be differences in colonization targets (244). However, infection studies comparing *S. Enteritidis* and *S. Typhimurium* in adult laying hens by Keller et al. indicated only minimal differences among strains from these two serovars in intestinal, hepatic, splenic, and reproductive tract colonization (261). Likewise, it appears that *S. Enteritidis* genes associated with intestinal colonization are similar to those observed for other *Salmonella* serotypes. Bohez et al. constructed a deletion mutant of the SPI-1 regulator gene *hilA* in *S. Enteritidis* and observed decreases in shedding and cecal colonization to nearly nondetectable levels after 4 weeks in young chicks. Those authors concluded that the HilA regulatory protein was required for continued cecal colonization and long-term shedding (268). Subsequent studies by these researchers demonstrated that the initial colonization by the *S. Enteritidis* *hilA* mutant was sufficient to prevent long-term fecal shedding and cecal and internal organ colonization by the respective *S. Enteritidis* wild-type strain (264, 269). This linkage of *hilA* with systemic infection is supported by the general trends of increased liver, spleen, and ovary colonization observed in adult laying hens when *S. Enteritidis* *hilA* expression levels were increased (270).

However, studies based on infection of young chicks by *S. Enteritidis* deletion mutants for individual SPIs (SPI-1 to -5) revealed that colonization of the liver and spleen was dependent on SPI-1 and SPI-2 but that cecal colonization was not (271). Desin et al. observed less invasion by *S. Enteritidis* SPI-1 mutants of polarized Caco-2 human intestinal epithelial cells and chicken cecal and small intestinal explants but no differences in cecal colonization of 1-week-old chickens, with the exception of an infection delay in livers and spleens (272). Mixed results have also been reported for other serovars and other hosts. *S. Typhimurium* *hilA* deletion mutants were recovered from infected mouse intestinal contents, intestinal tissues, and systemic tissues at a lower frequency than the parental wild-type strain, while SPI-1 deletion mutants were recovered at frequencies similar to those of the wild-type strain (273). Morgan et al. reported a similar lack of an impact on chicken cecal colonization by *S. Typhimurium* SPI-1 mutants, but SPI-1 was required for colonization of calf intestines (43). Dieye et al. used a mixed-infection approach by combining isogenic SPI-1 *S. Typhimurium* deletion mutants with their respective wild-type counterparts and administering them to 1-week-old chicks instead of younger birds. For the most part, the wild-type strain outcompeted the SPI-1 deletion mutant over the 14-day infection period. When cecal and spleen samples were recovered, the level of cecal colonization by the wild-type was substantially higher than that by the mutant by day 14 (74). More recently, Clark et al. used green fluorescent reporter gene constructs for comparison of various *S. Typhimurium* strains in tissue culture infection assays and discovered that only subpopulations expressed SPI-1 genes. Those researchers speculated that genetic heterogeneity of invading *S. Typhimurium* populations could lead to differential SPI-1 expression and heterogeneous *Salmonella* population behavior during infection (274).

There appear to be serovar differences in pathogenicity gene expression as well. Silva et al. constructed an *S. Enteritidis* trans-

poson library and used a microarray-based negative-selection screening approach to isolate mutants deficient for liver and spleen colonization of BALB/c mice. Many of the colonization-deficient strains were identified as having mutations in SPI genes already known to be responsible for *Salmonella* systemic infection; however, mutants that were associated with genes and genomic islands not identified in other *Salmonella* serovars, including *S. Typhimurium*, were also isolated (275). Shah et al. also identified genes and genomic islands in *S. Enteritidis* that were not present in *S. Typhimurium* after screening *S. Enteritidis* mutants generated from a random Tn5 transposon library for their ability to invade Caco-2 and chicken liver (LMH) cells and for survival in chicken macrophages (HD-11) (276). Strains within individual serovars also display differences in the ability to express different genes. Shah et al. screened six *S. Enteritidis* isolates and were able to phenotypically differentiate them as stress sensitive (acid and oxidative stress, survival in egg albumen, and impairment in intestinal colonization and systemic infection) versus stress resistant, even though no changes in genomic content were detectable with genomic hybridization microarrays. Sequencing of the stress-regulatory gene *rpoS* in one of the stress-sensitive strains revealed an insertion mutation and truncated protein, leading those authors to conclude that even minor polymorphisms in certain genes would be sufficient to impact stress or virulence gene responses (277).

As more becomes known about the role of host-pathogen interactions in *Salmonella* pathogenesis, some of the seemingly contradictory pathogen responses may become clear. For example, Knodler et al. demonstrated that hyperreplicating *Salmonella* subpopulations in the cytosol of GIT epithelial cells that are distinct from the SCV populations can mediate their release back into the lumen by inducing inflammation accompanied by epithelial cell death and subsequent shedding (58). Since T3SS-based invasion gene systems are involved, it may be that pathogenicity alterations occurring with different mutants and serovars could be attributed the intracellular subpopulations that are impacted, thereby impacting the abilities to sustain long-term fecal excretion of the respective subpopulation (58, 278). This may partially explain the emergence of individual *Salmonella* carrier chickens within poultry flock populations as well as in other animal species.

In addition, the complexity of the indigenous GIT microflora, interaction of the immune system, and host factors such as age, stress level, and overall health status may play important roles in host-pathogen interactions (279–281). Identifying overarching themes for *Salmonella* pathogenesis and designing effective cross-protective control measures that can be applied across different serovars will require application of combinations of high-throughput omics analyses; animal model responses to sort out shared signals, such as common immune signatures across animal species; along with subsequent mathematical modeling (282–286). Particularly, in poultry, the potential roles of host genes that may impact infection and carrier states of *Salmonella* have been studied for a number of years and have been extensively discussed recently in a comprehensive review by Calenge and Beaumont (283) and will not be discussed here. The remainder of the current review focuses on *Salmonella* serovar responses to the poultry host.

SALMONELLA AND STRESS RESPONSES IN THE CHICKEN GIT

Prior to infecting a potential host, *Salmonella* encounters numerous environmental conditions that represent sufficient stress to influence the organism's ability to grow optimally and in some cases even survive. These stressors can originate as a wide range of physical and chemical challenges either in the environmental niches that *Salmonella* occupies or as manufactured barriers, such as those incorporated during food production and processing, such as antimicrobials, disinfectants, biocides, and preservatives. These intervention approaches have been reviewed extensively elsewhere (273, 287–291).

As noted above, when *Salmonella* spp. come into contact with a prospective host and gain access to its GIT, there are a number of potential barriers that the organism encounters as it travels through the GIT. Ahmer and Gunn classified these host barriers into broad categories of physical, chemical, enzymatic, and immune hurdles that must be surmounted by *Salmonella* spp. (292). These include, in addition to specific host-generated immune- and non-immune-based defenses, iron limitation, acidic and pH shifts, oxidative and osmolarity stresses, and limited nutrient availability that leads to starvation (35, 36, 293–301). Although these barriers are manifested either as a function of the host's defense mechanisms or as a consequence of high population densities and a metabolically active indigenous GIT microbial population, in reality, there is a substantial interaction between the two that makes it difficult to separate them experimentally. For example, the field of avian genetics has advanced sufficiently such that a range of host-innate, cell-mediated, and humoral immune responses that resist and clear pathogens such as *Salmonella* have been identified. The extent of the immune involvement is determined by microbial challenge severity as well as the health and nutritional status of the host (280, 283, 302). The immune system also interacts with the indigenous microflora, and the presence of this microflora is now considered essential for proper immunological development (303). Consequently, nonpathogenic bacteria can be considered active players in modulating immune recognition of pathogens (304, 305). Given the diversity of the GIT microflora, an understanding of the symbiotic relationship between the microbial consortia and the immune system is extremely complex and has been reviewed elsewhere (303, 306, 307). Here the focus is on the microbial metabolic interactions that occur in the GIT that appear to influence *Salmonella* and countermeasures employed by this pathogen to overcome these challenges.

Salmonella Acid Tolerance Response and Organic Acids

The compounds collectively referred to as organic acids are comprised of the saturated straight-chain monocarboxylic acids and their respective derivatives (unsaturated, hydroxylic, phenolic, and multicarboxylic organic acids) and have a long history as antifungal and antimicrobial agents in perishable animal feeds and human foods (308–310). Traditionally, organic acids have been added to elicit bacteriostatic and bactericidal impacts on undesirable organisms in potentially contaminated food products. A wide range of antimicrobial mechanisms, including osmotic stress, anion toxicity, and membrane disruption, have been proposed over the years, but inhibition is generally believed to be closely tied to pH because of its influence on the formation of undissociated organic acids, which can easily cross the cell membrane and dis-

sociate into anions and protons in the presence of the cytoplasmic neutral pH (309–313). The acids not only result in the acidification of the cytoplasm and the accompanying difficulties for cellular constituents to perform essential metabolic functions in the cell but also lead to an energy drain due to the consumption of ATP to actively export protons (310).

Salmonella can encounter short-chain fatty organic acids (SCFAs) (primarily acetate, propionate, butyrate, and lactate) in a variety of environments, because they are used as animal feed additives and antimicrobials in food and meat processing and can be generated by certain starter cultures used for food fermentations (308, 309, 314–317). Once in the GIT, particularly in the lower regions of monogastric hosts and the foregut in ruminants, *Salmonella* must be able to survive the presence of variable SCFA concentrations (mM ranging from <1 to hundreds for particular organic acids) depending on the host animal and digestive tract site associated with fermentation by the indigenous microflora in the GIT (309, 311, 314, 318–321).

Traditionally, it was believed that *Salmonella* spp., much like other bacteria, were sensitive to organic acids and that this sensitivity was dependent on pH (322–325). However, Leyer and Johnson demonstrated that HCl acid-adapted *S. Typhimurium* could also resist organic acids typically found in cheeses (326). Baik et al. induced stationary- and exponential-phase acid (pH 4.4) tolerance responses in *S. Typhimurium* after exposure to a series of organic acids and concluded that both physiological states were protected, but the stationary-phase protection occurred with only some of the organic acids (327). The protection against organic acids did not occur if organic acids were used for the initial acid shock. When Kwon and Ricke exposed *S. Typhimurium* to SCFA at neutral pH for 1 h, they observed that acid tolerance was greater for cells exposed to acetate or propionate than for cells exposed to butyrate or valeric acid and was enhanced under anaerobic incubation conditions (328). Calhoun and Kwon observed substantial enhanced acid tolerance in *S. Enteritidis* when exposed to 100 mM propionate at neutral pH over a longer period of time (16 h) (329). The enhancement may not be surprising since *Salmonella* cells cultured under anaerobic growth conditions can produce fermentation acids (330, 331), suggesting that some of the resistance to SCFA may be related to protection from their own end products. It is also known that *Salmonella* can directly catabolize SCFA. For example, *S. Typhimurium* utilizes the 2-methylcitric acid cycle to convert propionate to pyruvate for reentry into the tricarboxylic acid cycle (332–334).

S. Typhimurium exposure at pH 7.0 to high (mM) levels of SCFA typical of large intestine concentrations induces not only resistance to extreme acidic conditions (pH 3.0) but also cross-protective resistance to high osmolarity (2.5 mM NaCl) and reactive oxygen (20 mM H₂O₂) (335). Similar cross protection against salt stress (KCl and NaCl) has been reported for acetic acid-induced acid tolerance in *S. Typhimurium* but not lactic acid-induced acid tolerance, which appears to be related to sensitivity to hydrogen peroxide (336, 337). Initial pH appears to be a factor, as both growth and survivability are limited when *S. Typhimurium* cells are exposed to propionate at pH 5.0 versus pH 7.0 (328). It appears that even more subtle fluctuations in pH can influence *Salmonella* growth metabolism. For example, Dunkley et al., using anaerobic continuous cultures, demonstrated that *S. Typhimurium* glucose- and ATP-based microbial cell yields (calculated as mg of cell protein synthesized per ATP formed or glucose con-

sumed) were decreased when the pH in the growth vessels was shifted from near-neutral pH to levels approaching pH 6.0 (330). Other stress combinations appear to have more of an impact on SCFA inhibition of *Salmonella*. Milillo and coworkers demonstrated not only that the combination of heat (50°C to 60°C) and organic acids was more effective than single intervention applications when applied to *S. Typhimurium* but also that heated acidified solutions caused substantial membrane damage and leakage of intracellular ions (338, 339). There may also be potential serovar differences. Joerger et al. concluded that *S. Kentucky* was more acid sensitive (pH 2.5) than other *Salmonella* serovars after growth in the presence of acetic acid (340). Gene expression microarray comparisons between *S. Kentucky* and *S. Enteritidis* indicated that regulatory genes associated with acid response (*rpoS*, *fur*, and *phoPQ*) were similar between the two serovars, as were most global transcriptional responses, leading Joerger et al. to speculate that phenotypic differences were perhaps more a function of post-translational or small undetectable transcriptional differences (340).

***Salmonella* Virulence Response and Organic Acids**

While acid tolerance is considered a component of pathogenesis, the presence of SCFA may also directly impact *Salmonella* pathogenesis and virulence gene expression. Durant et al. conducted a series of tissue culture studies with an epidermal carcinoma cell line (HEp-2) to determine the influence of *S. Typhimurium* after growth in either a mixture or separate intestinal concentrations of SCFA (acetate, propionate, and butyrate at pH 6.0 or 7.0) on its subsequent ability to adhere to and invade tissue culture cells (341–343). For *S. Typhimurium* incubated for 4 h in the presence of SCFA, acetate at pH 6.0 and 7.0 did not alter attachment, and only slight increases in invasion occurred, while propionate and butyrate dramatically decreased attachment and invasion at pH 6.0 but not at pH 7.0. There were also indications that these *S. Typhimurium* responses were growth dependent, particularly in the presence of propionate and butyrate at pH 6.0, which reduced the growth rate considerably (341). However, when late-logarithmic-phase (12 h) and stationary-phase (18 h) *S. Typhimurium* cells were compared (342, 343), all SCFAs reduced cell attachment at pH 6.0 and 7.0 but only marginally influenced invasion for late-log-phase cultures of *Salmonella*. The stationary-phase *Salmonella* cells exhibited reduced attachment only in the presence of SCFA at pH 6.0, while actual increases in invasion occurred for lower concentrations of acetate (342, 343). Van Immerseel et al., using avian intestinal and cecal epithelial cell lines, also observed increases in invasion of *S. Enteritidis* exposed to acetate for 4 h and decreases with propionate and butyrate (265, 344). Gantois et al. reported that exposure of *S. Enteritidis* and *S. Typhimurium* to butyrate for 4 h resulted in the lowest percentage of invasion of HeLa cells (345).

In follow-up studies to determine the genetic elements associated with the SCFA influence on *Salmonella* pathogenesis, Durant et al. used *S. Typhimurium* strains constructed to carry chromosomal *lacZY* transcriptional fusions within either *hilA* or *invF* to assess the impact of pH and individual SCFAs on their expression (346). The study concluded that growth of *S. Typhimurium* in the presence of acetate, but not propionate and butyrate, stimulated β -galactosidase expression for both genes, even though growth was inhibited only at pH 6.0. When the *S. Typhimurium* *hilA-lacZY* fusion was incorporated into an *S. Enteritidis* isolate from

poultry, increasing lactate concentrations along with lowering pH reduced expression levels of β -galactosidase (347). Lawhon et al. concluded that acetate provided a signal for invasion through the production of cytoplasmic acetyl-phosphate, while propionate and butyrate repressed invasion genes (348). Gantois et al. conducted comparative transcriptomic analyses of *S. Enteritidis* and *S. Typhimurium* and reported that butyrate downregulated SPI-1 genes, including the regulatory genes *hilD* and *infV*, while Huang et al. identified formate as a diffusible signal to induce SPI-1 genes and invasion (345, 349). Virulence gene responses to organic acids may vary in different serovars as well. Using quantitative reverse transcriptase real-time PCR, Gonzalez-Gil et al. reported that expression of the virulence gene regulator *hilA* was both serovar and strain dependent when isolates were exposed to either mild acetic or HCl acid stress. Responses were dependent on the type of acid used and were time dependent, with *hilA* being upregulated after a 2-h exposure to the respective acid (350). When Milillo et al. compared *S. Typhimurium* transcriptome responses to sodium acetate and sodium propionate exposure after 1 min (55°C at pH 4.0), the respective gene expression patterns were similar, with no increases in expression levels of *rpoS*, *phoPQ*, or virulence factors despite their established roles in organic and inorganic acid stress (35, 327, 328, 338). Conversely, genes related to attachment and/or motility functions were induced, which has been observed previously for *S. Typhimurium* attachment genes after exposure to 42°C (351). There was also a repression of several heat shock response genes, which, in conjunction with cell membrane damage, indicates that membrane integrity is an important component for *Salmonella* cells to accurately assess their extracellular environmental conditions and respond accordingly (351). Clearly, more in-depth genetic analysis is needed to sort out the interactions that occur among different stresses in *Salmonella* versus its collective virulence response capabilities as well as how these may shift over time after initial encounters with the cell's surrounding environment. Serovar and strain differences will also need to be examined to determine whether some of the serovar dissemination patterns are due to the ability of some serovars, but not others, to effectively counter external stresses in their environments.

Salmonella Virulence Response and Competition with GIT Microflora

The GIT systems in most hosts, including food animals and humans, harbor highly complex and metabolically interactive microbial consortia. Particularly in the lower GIT of monogastric animals (colon or cecum) and the foregut (rumen) of ruminants, microbial populations consist of large numbers of mostly saccharolytic anaerobic microorganisms along with methanogenic archaea, which together result in a highly reduced ecosystem with minimal oxygen (318, 320, 352–359). Given the complexity of diets in general and the recalcitrant nature of some of the fiber-containing components (particularly those consumed by herbivores), the GIT microbial consortia are characterized not only by their anaerobic physiology but also by their ability to hydrolyze fiber polymers into soluble carbohydrates that are readily fermentable (360–364). Since energy generation for ATP formation is anaerobic and, therefore, much more limited per carbon source, GIT saccharolytic microorganisms must extract maximum ATP through a multitude of limited-substrate-level and anaerobic respiration phosphorylation pathways (318, 365–368). Given the

complexity of the GIT microflora and the limited energy generation capabilities for microbial growth, each species must catabolize similar carbon sources, necessitating a hierarchy for substrate preferences and competition among microorganisms based on maximum growth, energy efficiency, and substrate affinity (369, 370). If these capabilities are combined with the constant GIT turnover that occurs due to the passage rate of digesta, GIT microbial ecosystems are characterized by their exceptional ability to metabolically scavenge and extract maximum nutritional value from incoming dietary sources.

Consequently, pathogens entering the GIT, in addition to combating the hostile environment derived from fermentation activities of the indigenous microflora and the host barriers, must also directly compete with the GIT microflora for scarce nutrients (298, 371). Pathogen metabolism and virulence are often interconnected to the point that it becomes difficult to distinguish which functions are occurring simultaneously. Rohmer et al. suggested that not only do pathogens acquire virulence factors to access new niches, they may also require new metabolic pathways to sequester available nutrient sources, as evidenced by the presence of metabolic functions encoded on pathogenicity islands (371, 372). In the GIT, pathogens entering the lumen during times of nutritional stress could conceivably employ this strategy by either inducing virulence to escape the GIT altogether or manipulating the GIT environment to outcompete indigenous microflora.

Starvation, malnutrition, or abrupt changes in diet can alter the composition of indigenous GIT microflora and potentially impact the host's immune system and the host's general physiology and well being (373–375). Changes in nutritional status and dietary alteration have also been known for some time to alter the relationship between the GIT and pathogens, such as attachment to the epithelial lining (376). Emptying of the GIT content represents one of the more drastic alterations that occurs either naturally or as part of a prescribed treatment in humans or a management option in animals. Increased populations of *Salmonella* have been consistently observed in a variety of animals undergoing emptying of the GIT due to feed removal, including chickens (377–381), mice (382), and ruminants (383–385). Although the indigenous GIT microflora were not characterized in these early studies, it was speculated that feed removal/GIT emptying potentially influenced the microbial populations or activities, and susceptibility to *Salmonella* colonization was presumed to be related to decreases in levels of antimicrobial organic acids generated during fermentation (250).

One of the most extensively studied host-pathogen systems, which represented a nutritional management strategy linked with *Salmonella* dissemination, was the use of dietary restrictions to induce artificial molting in laying hens. In the U.S. egg industry, laying hens can be induced to molt in order to cease egg production and allow the reproductive tract to undergo a rest phase to allow flock rejuvenation, feather renewal, and a second, higher-production egg-laying cycle (386). Historically, this was accomplished by complete removal of diet for anywhere from 5 to 14 days until cessation of egg production was completed (386, 387). However, *S. Enteritidis* was closely linked with egg contamination, and research conducted during the 1990s with laying hens as the infection model suggested that management practices such as feed removal-induced molts might be partially responsible for the contamination (388, 389). In the following decade, more detailed

studies on *S. Enteritidis* infection in susceptible laying hens concluded that complete removal of feed created a GIT environment in laying hens that not only was easily colonized by *S. Enteritidis* but also led to systemic infection of internal organs, including the ovaries (250, 331, 390). Increased expression levels of the *S. Enteritidis lacZY-hilA* fusion gene in cells incubated in crop contents from birds that had undergone feed removal demonstrated that enhanced systemic invasiveness could be linked with increased expression levels of *hilA* (391). These findings were later replicated and confirmed *in vivo* by quantitative PCR (392). This increase in *S. Enteritidis* colonization and invasiveness also coincided with diminished numbers of lactic acid bacteria in the crop and reduced organic acids in the crop (391) and cecum (270, 393) and led to major shifts in fecal and cecal bacteria compared to fully fed birds (270, 394, 395). Ricke hypothesized that the key to limiting *S. Enteritidis* during molt was to develop low-nutrient dietary strategies, such as formulations with high fiber content, that would still induce molt due to low energy density but were sufficiently fermentable to support an active GIT microflora that would serve as a barrier to pathogen establishment (255). Several molt diets with fermentable substrate combinations have been screened *in vitro* and shown to support optimal avian cecal fermentation (396–398) and retain similar cecal microbial molecular profiles (397). *In vivo* application of some of these diets was shown to retain fermentation and GIT populations during molt while limiting the ability of *S. Enteritidis* to colonize the GIT and become invasive (255, 270, 392, 399, 400). The overall results of the studies demonstrate that there is a complex and important relationship between the microflora of the avian GIT and *Salmonella*, which is key to limiting the pathogen.

CONCLUDING REMARKS

Salmonella serovars display a range of abilities to colonize different animal hosts and cause illnesses in humans. Many of the serovars most commonly implicated in food-borne human infections have a broad range of animal hosts that they can colonize, while others, especially some that cause severe human infections, have much narrower host ranges. Some serovars, such as *S. Typhi* and *S. Paratyphi A*, are human-restricted serovars and can cause severe disease manifestations such as enteric fever. The factors that impact *Salmonella* pathogenesis and host range are complex and are associated with differences in variables such as the genetics of the bacterium and expression of these genetic factors, the host immune status and specific pathogen responses, the host environment, and interactions with intestinal microflora.

A great deal of work has been undertaken to understand the genetics of *Salmonella* and explore the factors that contribute to the evolution of pathogenesis. Based on genome sequencing results and pangenome analyses of *Salmonella* serovars, it is evident that two distinct, yet potentially complementary, evolutionary approaches that impact host range and virulence are occurring. These approaches include gene acquisition through horizontal transfer (associated with SPIs, transposable elements, phages, and plasmids) and the loss of genes or a loss of their function, which impacts host range (105, 106). As the species *S. enterica* evolved, its members' genomes acquired the multiple SPIs that allowed the organism to become pathogenic. Several serovars also acquired specific virulence plasmids that harbor the *spv* operon, which increases virulence (126). Other serovars, while not having serovar-specific virulence plasmids, have other plasmids that may encode

factors that are important for survival in a potential host niche. Plasmids have played a significant role in the evolution of host specificity in some *Salmonella* serovars. A key example includes some of the plasmids of *S. Kentucky* and some *S. Heidelberg* isolates that contain genes associated with extraintestinal survival in poultry, which may provide a selective advantage in poultry (83). Many of these plasmids also contain genes for antimicrobial resistance, which is concerning because of the possibility that antimicrobial use could select for enhanced virulence or conversely that the selective advantage to survive in one environment/host may select for resistance to one or more antimicrobials (83, 84). While our knowledge of the genetic composition of these plasmids has greatly increased with the advent of next-generation sequencing, our understanding of the evolution and dissemination of these plasmids and their interactions with one another remains limited. There is not a clear correlation between carriage of the plasmid and the establishment of systemic disease in humans, further suggesting that it may be a requirement of some strains but not others (401).

On the other side of the spectrum of factors that impact virulence and host range is gene loss or the loss of gene function due to mutation, insertion, or deletion in the gene or regulatory element. Many of the host-restricted serovars have a large number of pseudogenes, which inactivate genes that are often involved in aspects of virulence, nutrient utilization, and metabolism (91). The impact of pseudogenes on host restriction seems fairly clear; however, it is less clear whether gene inactivation leads to the host-restricted phenotypes or whether the organisms settled in the niche and those genes not essential for survival are undergoing the process of genome reduction. Interestingly, *Salmonella* serovar Typhi demonstrates aspects of functional gene loss (a high percentage of pseudogenes) (193) and the impact of horizontal gene transfer on pathogenicity (14). *S. Typhi* isolates have mutations in several of the fimbrial genes that are associated with intestinal attachment, which is important for gastroenteritis but not enteric fever (194). Members of the serovar also acquired SPI-7, which harbors genes for the Vi capsule, which prevents recognition of PAMPs on the surface of *S. Typhi*, thereby limiting the initial immune response following infection (2).

Despite the wealth of knowledge associated with *Salmonella* pathogenesis mechanisms in host animals, several key questions remain on the exact roles and extent of involvement of the respective virulence genes and genomic islands of particular serovars and their characteristic pathogenic phenotypes expressed in animal models. Not only do *in vitro* models, such as tissue culture adhesion and invasion assays, sometimes differ from *in vivo* results, there may also be differences in responses depending on the type of cell line used (402). It is also becoming apparent, as more sophisticated genetic analyses are being applied to animal infection studies, that *Salmonella* serovars vary in their virulence gene capacities, and given the tremendous number of serovars identified and the wide range of corresponding strains isolated for some serovars, there is the potential for almost endless nuances of perhaps subtle but different serovar- and even strain-specific genetic strategies for overcoming host barriers.

An in-depth understanding of the mechanisms of *Salmonella* pathogenesis is important for developing intervention strategies to limit their prevalence and spread and aid in the development of newer drugs and vaccines which could result in better treatment and/or prevention of salmonellosis in animals and humans. The

continued use of whole-genome sequencing, gene expression, and bioinformatics analyses of *Salmonella* serovars from host-restricted and host-specific serovars should continue to provide insights into the specific genetic traits that could contribute to symbiotic and/or disease-causing potential in humans and animals and lead to an overall reduction in disease burden associated with *Salmonella*.

ACKNOWLEDGMENTS

We thank Jing Han, Huizhong Chen, and Carl Cerniglia for their insightful reviews of the manuscript.

The views presented in this article do not necessarily reflect those of the U.S. Food and Drug Administration.

REFERENCES

- Donnenberg MS. 2000. Pathogenic strategies of enteric bacteria. *Nature* 406:768–774.
- Tsolis RM, Young GM, Solnick JV, Baumler AJ. 2008. From bench to bedside: stealth of enteroinvasive pathogens. *Nat. Rev. Microbiol.* 6:883–892.
- Guerrant RL, Steiner TS. 2005. Principles and syndromes of enteric infections, p 1215–1230. *In* Mandell GL, Bennett JE, Dolin R (ed), Principles and practice of infectious diseases, 6th ed, vol 1. Elsevier Churchill Livingstone, Philadelphia, PA.
- Herrington DA, Hall RH, Losonsky G, Mekalanos JJ, Taylor RK, Levine MM. 1988. Toxin, toxin-coregulated pili, and the *toxR* regulon are essential for *Vibrio cholerae* pathogenesis in humans. *J. Exp. Med.* 168:1487–1492.
- Nataro JP, Kaper JB. 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* 11:142–201.
- Liu H, Magoun L, Luperchio S, Schauer DB, Leong JM. 1999. The Tir-binding region of enterohaemorrhagic *Escherichia coli* intimin is sufficient to trigger actin condensation after bacterial-induced host cell signalling. *Mol. Microbiol.* 34:67–81.
- Stephen J. 2001. Pathogenesis of infectious diarrhea. *Can. J. Gastroenterol.* 15:669–683.
- Bhavsar AP, Guttman JA, Finlay BB. 2007. Manipulation of host-cell pathways by bacterial pathogens. *Nature* 449:827–834.
- Cossart P, Sansonetti PJ. 2004. Bacterial invasion: the paradigms of enteroinvasive pathogens. *Science* 304:242–248.
- Watson RO, Galan JE. 2008. *Campylobacter jejuni* survives within epithelial cells by avoiding delivery to lysosomes. *PLoS Pathog.* 4:e14. doi:10.1371/journal.ppat.0040014.
- Hoebe K, Janssen E, Beutler B. 2004. The interface between innate and adaptive immunity. *Nat. Immunol.* 5:971–974.
- Tang D, Kang R, Coyne CB, Zeh HJ, Lotze MT. 2012. PAMPs and DAMPs: signal 0s that spur autophagy and immunity. *Immunol. Rev.* 249:158–175.
- Nguyen QC, Everest P, Tran TK, House D, Murch S, Parry C, Connerton P, Phan VB, To SD, Mastroeni P, White NJ, Tran TH, Vo VH, Dougan G, Farrar JJ, Wain J. 2004. A clinical, microbiological, and pathological study of intestinal perforation associated with typhoid fever. *Clin. Infect. Dis.* 39:61–67.
- Sharma A, Qadri A. 2004. Vi polysaccharide of *Salmonella* Typhi targets the prohibitin family of molecules in intestinal epithelial cells and suppresses early inflammatory responses. *Proc. Natl. Acad. Sci. U. S. A.* 101:17492–17497.
- Andersen-Nissen E, Smith KD, Strobe KL, Barrett SL, Cookson BT, Logan SM, Aderem A. 2005. Evasion of Toll-like receptor 5 by flagellated bacteria. *Proc. Natl. Acad. Sci. U. S. A.* 102:9247–9252.
- Lapaque N, Takeuchi O, Corrales F, Akira S, Moriyon I, Howard JC, Gorvel JP. 2006. Differential inductions of TNF-alpha and IGTP, IIGP by structurally diverse classic and non-classic lipopolysaccharides. *Cell. Microbiol.* 8:401–413.
- Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* 17:7–15.
- Andrews-Polymenis HL, Baumler AJ, McCormick BA, Fang FC. 2010. Taming the elephant: *Salmonella* biology, pathogenesis, and prevention. *Infect. Immun.* 78:2356–2369.
- Hoffmann S, Batz MB, Morris JG, Jr. 2012. Annual cost of illness and quality-adjusted life year losses in the United States due to 14 foodborne pathogens. *J. Food Prot.* 75:1292–1302.
- Batz MB, Hoffman S, Morris JG, Jr. 2011. Ranking the risks: the 10 pathogen-food combinations with the greatest burden on public health. University of Florida, Gainesville, FL.
- Batz MB, Doyle MP, Morris G, Jr, Painter J, Singh R, Tauxe RV, Taylor MR, Lo Fo Wong DM, Food Attribution Working Group. 2005. Attributing illness to food. *Emerg. Infect. Dis.* 11:993–999.
- Centers for Disease Control and Prevention. 1997. *Salmonella* serotype Montevideo infections associated with chicks—Idaho, Washington, and Oregon, spring 1995 and 1996. *MMWR Morb. Mortal. Wkly. Rep.* 46:237–239.
- Batz MB, Hoffmann S, Morris JG, Jr. 2012. Ranking the disease burden of 14 pathogens in food sources in the United States using attribution data from outbreak investigations and expert elicitation. *J. Food Prot.* 75:1278–1291.
- Brenner FW. 1998. Modified Kauffmann-White scheme. Centers for Disease Control and Prevention, Atlanta, GA.
- Su LH, Chiu CH. 2007. *Salmonella*: clinical importance and evolution of nomenclature. *Chang Gung Med. J.* 30:210–219.
- Foley SL, Nayak R, Hanning IB, Johnson TJ, Han J, Ricke SC. 2011. Population dynamics of *Salmonella enterica* serotypes in commercial egg and poultry production. *Appl. Environ. Microbiol.* 77:4273–4279.
- Goto Y, Kiyono H. 2011. Epithelial cell microRNAs in gut immunity. *Nat. Immunol.* 12:195–197.
- Foley SL, Lynne AM, Nayak R. 2008. *Salmonella* challenges: prevalence in swine and poultry and potential pathogenicity of such isolates. *J. Anim. Sci.* 86:E149–E162. doi:10.2527/jas.2007-0464.
- Uzzau S, Brown DJ, Wallis T, Rubino S, Leori G, Bernard S, Casadesus J, Platt DJ, Olsen JE. 2000. Host adapted serotypes of *Salmonella enterica*. *Epidemiol. Infect.* 125:229–255.
- McDonough PL, Fogelman D, Shin SJ, Brunner MA, Lein DH. 1999. *Salmonella enterica* serotype Dublin infection: an emerging infectious disease for the northeastern United States. *J. Clin. Microbiol.* 37:2418–2427.
- Chiu CH, Su LH, Chu C. 2004. *Salmonella enterica* serotype Choleraesuis: epidemiology, pathogenesis, clinical disease, and treatment. *Clin. Microbiol. Rev.* 17:311–322.
- Chiu CH, Wu TL, Su LH, Chu C, Chia JH, Kuo AJ, Chien MS, Lin TY. 2002. The emergence in Taiwan of fluoroquinolone resistance in *Salmonella enterica* serotype Choleraesuis. *N. Engl. J. Med.* 346:413–419.
- Lynne AM, Dorsey LL, David DE, Foley SL. 2009. Characterisation of antibiotic resistance in host-adapted *Salmonella enterica*. *Int. J. Antimicrob. Agents* 34:169–172.
- Vugia DJ, Samuel M, Farley MM, Marcus R, Shiferaw B, Shallow S, Smith K, Angulo FJ. 2004. Invasive *Salmonella* infections in the United States, FoodNet, 1996–1999: incidence, serotype distribution, and outcome. *Clin. Infect. Dis.* 38(Suppl 3):S149–S156. doi:10.1086/381581.
- Bearson BL, Wilson L, Foster JW. 1998. A low pH-inducible, PhoPQ-dependent acid tolerance response protects *Salmonella typhimurium* against inorganic acid stress. *J. Bacteriol.* 180:2409–2417.
- Bearson SM, Bearson BL, Rasmussen MA. 2006. Identification of *Salmonella enterica* serovar Typhimurium genes important for survival in the swine gastric environment. *Appl. Environ. Microbiol.* 72:2829–2836.
- Ruby T, McLaughlin L, Gopinath S, Monack D. 2012. *Salmonella*'s long-term relationship with its host. *FEMS Microbiol. Rev.* 36:600–615.
- Velge P, Wiedemann A, Rosselin M, Abed N, Boumart Z, Chausse AM, Grepinet O, Namdari F, Roche SM, Rossignol A, Virlogeux-Payant I. 2012. Multiplicity of *Salmonella* entry mechanisms, a new paradigm for *Salmonella* pathogenesis. *Open Microbiol.* 1:243–258.
- Darwin KH, Miller VL. 1999. Molecular basis of the interaction of *Salmonella* with the intestinal mucosa. *Clin. Microbiol. Rev.* 12:405–428.
- van Asten AJ, van Dijk JE. 2005. Distribution of “classic” virulence factors among *Salmonella* spp. *FEMS Immunol. Med. Microbiol.* 44:251–259.
- Stevens MP, Humphrey TJ, Maskell DJ. 2009. Molecular insights into farm animal and zoonotic *Salmonella* infections. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 364:2709–2723.
- Carnell SC, Bowen A, Morgan E, Maskell DJ, Wallis TS, Stevens MP.

2007. Role in virulence and protective efficacy in pigs of *Salmonella enterica* serovar Typhimurium secreted components identified by signature-tagged mutagenesis. *Microbiology* 153:1940–1952.
43. Morgan E, Campbell JD, Rowe SC, Bispham J, Stevens MP, Bowen AJ, Barrow PA, Maskell DJ, Wallis TS. 2004. Identification of host-specific colonization factors of *Salmonella enterica* serovar Typhimurium. *Mol. Microbiol.* 54:994–1010.
 44. de Jong H, Ekdahl MO. 1965. Salmonellosis in calves—the effect of dose rate and other factors on transmission. *N. Z. Vet. J.* 13:59–64.
 45. Fedorka-Cray PJ, Kelley LC, Stabel TJ, Gray JT, Laufer JA. 1995. Alternate routes of invasion may affect pathogenesis of *Salmonella typhimurium* in swine. *Infect. Immun.* 63:2658–2664.
 46. Winnen B, Schlumberger MC, Sturm A, Schupbach K, Siebenmann S, Jenny P, Hardt WD. 2008. Hierarchical effector protein transport by the *Salmonella* Typhimurium SPI-1 type III secretion system. *PLoS One* 3:e2178. doi:10.1371/journal.pone.0002178.
 47. Schlumberger MC, Hardt WD. 2006. *Salmonella* type III secretion effectors: pulling the host cell's strings. *Curr. Opin. Microbiol.* 9:46–54.
 48. Galan JE, Wolf-Watz H. 2006. Protein delivery into eukaryotic cells by type III secretion machines. *Nature* 444:567–573.
 49. Marlovits TC, Kubori T, Sukhan A, Thomas DR, Galan JE, Unger VM. 2004. Structural insights into the assembly of the type III secretion needle complex. *Science* 306:1040–1042.
 50. Lostroh CP, Lee CA. 2001. The *Salmonella* pathogenicity island-1 type III secretion system. *Microbes Infect.* 3:1281–1291.
 51. Galan JE. 2001. *Salmonella* interactions with host cells: type III secretion at work. *Annu. Rev. Cell Dev. Biol.* 17:53–86.
 52. Norris FA, Wilson MP, Wallis TS, Galyov EE, Majerus PW. 1998. SopB, a protein required for virulence of *Salmonella dublin*, is an inositol phosphate phosphatase. *Proc. Natl. Acad. Sci. U. S. A.* 95:14057–14059.
 53. Wallis TS, Galyov EE. 2000. Molecular basis of *Salmonella*-induced enteritis. *Mol. Microbiol.* 36:997–1005.
 54. Hopkins KL, Threlfall EJ. 2004. Frequency and polymorphism of *sopE* in isolates of *Salmonella enterica* belonging to the ten most prevalent serotypes in England and Wales. *J. Med. Microbiol.* 53:539–543.
 55. Miroid S, Rabsch W, Rohde M, Stender S, Tschape H, Russmann H, Igwe E, Hardt WD. 1999. Isolation of a temperate bacteriophage encoding the type III effector protein SopE from an epidemic *Salmonella typhimurium* strain. *Proc. Natl. Acad. Sci. U. S. A.* 96:9845–9850.
 56. Frost AJ, Bland AP, Wallis TS. 1997. The early dynamic response of the calf ileal epithelium to *Salmonella typhimurium*. *Vet. Pathol.* 34:369–386.
 57. Goosney DL, Knoechel DG, Finlay BB. 1999. Enteropathogenic *E. coli*, *Salmonella*, and *Shigella*: masters of host cell cytoskeletal exploitation. *Emerg. Infect. Dis.* 5:216–223.
 58. Knodler LA, Vallance BA, Celli J, Winfree S, Hansen B, Montero M, Steele-Mortimer O. 2010. Dissemination of invasive *Salmonella* via bacterial-induced extrusion of mucosal epithelia. *Proc. Natl. Acad. Sci. U. S. A.* 107:17733–17738.
 59. Holden DW. 2002. Trafficking of the *Salmonella* vacuole in macrophages. *Traffic* 3:161–169.
 60. Tierrez A, Garcia-del Portillo F. 2005. New concepts in *Salmonella* virulence: the importance of reducing the intracellular growth rate in the host. *Cell. Microbiol.* 7:901–909.
 61. Hensel M. 2000. *Salmonella* pathogenicity island 2. *Mol. Microbiol.* 36:1015–1023.
 62. Waterman SR, Holden DW. 2003. Functions and effectors of the *Salmonella* pathogenicity island 2 type III secretion system. *Cell. Microbiol.* 5:501–511.
 63. Abrahams GL, Hensel M. 2006. Manipulating cellular transport and immune responses: dynamic interactions between intracellular *Salmonella enterica* and its host cells. *Cell. Microbiol.* 8:728–737.
 64. Hensel M, Shea JE, Waterman SR, Mundy R, Nikolaus T, Banks G, Vazquez-Torres A, Gleeson C, Fang FC, Holden DW. 1998. Genes encoding putative effector proteins of the type III secretion system of *Salmonella* pathogenicity island 2 are required for bacterial virulence and proliferation in macrophages. *Mol. Microbiol.* 30:163–174.
 65. Bueno SM, Riquelme S, Riedel CA, Kalergis AM. 2012. Mechanisms used by virulent *Salmonella* to impair dendritic cell function and evade adaptive immunity. *Immunology* 137:28–36.
 66. Sundquist M, Rydstrom A, Wick MJ. 2004. Immunity to *Salmonella* from a dendritic point of view. *Cell. Microbiol.* 6:1–11.
 67. Garai P, Gnanadhas DP, Chakravorty D. 2012. *Salmonella enterica* serovars Typhimurium and Typhi as model organisms: revealing paradigm of host-pathogen interactions. *Virulence* 3:377–388.
 68. Lahiri A, Iyer N, Das P, Chakravorty D. 2010. Visiting the cell biology of *Salmonella* infection. *Microbes Infect.* 12:809–818.
 69. Foley SL, Lynne AM. 2008. Food animal-associated *Salmonella* challenges: pathogenicity and antimicrobial resistance. *J. Anim. Sci.* 86: E173–E187. doi:10.2527/jas.2007-0447.
 70. Leung KY, Siame BA, Snowball H, Mok YK. 2011. Type VI secretion regulation: crosstalk and intracellular communication. *Curr. Opin. Microbiol.* 14:9–15.
 71. Marcus SL, Brummell JH, Pfeifer CG, Finlay BB. 2000. *Salmonella* pathogenicity islands: big virulence in small packages. *Microbes Infect.* 2:145–156.
 72. Bingle LE, Bailey CM, Pallen MJ. 2008. Type VI secretion: a beginner's guide. *Curr. Opin. Microbiol.* 11:3–8.
 73. Amavisit P, Lightfoot D, Browning GF, Markham PF. 2003. Variation between pathogenic serovars within *Salmonella* pathogenicity islands. *J. Bacteriol.* 185:3624–3635.
 74. Dieye Y, Ameiss K, Mellata M, Curtiss R, III. 2009. The *Salmonella* pathogenicity island (SPI) 1 contributes more than SPI2 to the colonization of the chicken by *Salmonella enterica* serovar Typhimurium. *BMC Microbiol.* 9:3. doi:10.1186/1471-2180-9-3.
 75. Zhang S, Kingsley RA, Santos RL, Andrews-Polymenis H, Raffatellu M, Figueiredo J, Nunes J, Tsolis RM, Adams LG, Baumler AJ. 2003. Molecular pathogenesis of *Salmonella enterica* serotype Typhimurium-induced diarrhea. *Infect. Immun.* 71:1–12.
 76. Zhang S, Santos RL, Tsolis RM, Stender S, Hardt WD, Baumler AJ, Adams LG. 2002. The *Salmonella enterica* serotype Typhimurium effector proteins SipA, SopA, SopB, SopD, and SopE2 act in concert to induce diarrhea in calves. *Infect. Immun.* 70:3843–3855.
 77. Coombes BK, Coburn BA, Potter AA, Gomis S, Mirakhor K, Li Y, Finlay BB. 2005. Analysis of the contribution of *Salmonella* pathogenicity islands 1 and 2 to enteric disease progression using a novel bovine ileal loop model and a murine model of infectious enterocolitis. *Infect. Immun.* 73:7161–7169.
 78. Boyen F, Pasmans F, Donne E, Van Immerseel F, Morgan E, Adriaensens C, Hernalsteens JP, Wallis TS, Ducatelle R, Haesebrouck F. 2006. The fibronectin binding protein ShdA is not a prerequisite for long term faecal shedding of *Salmonella* Typhimurium in pigs. *Vet. Microbiol.* 115:284–290.
 79. Kingsley RA, Santos RL, Keestra AM, Adams LG, Baumler AJ. 2002. *Salmonella enterica* serotype Typhimurium ShdA is an outer membrane fibronectin-binding protein that is expressed in the intestine. *Mol. Microbiol.* 43:895–905.
 80. Gerlach RG, Claudio N, Rohde M, Jackel D, Wagner C, Hensel M. 2008. Cooperation of *Salmonella* pathogenicity islands 1 and 4 is required to breach epithelial barriers. *Cell. Microbiol.* 10:2364–2376.
 81. Rotger R, Casadesus J. 1999. The virulence plasmids of *Salmonella*. *Int. Microbiol.* 2:177–184.
 82. Ahmer BM, Tran M, Heffron F. 1999. The virulence plasmid of *Salmonella typhimurium* is self-transmissible. *J. Bacteriol.* 181:1364–1368.
 83. Han J, Lynne AM, David DE, Tang H, Xu J, Nayak R, Kaldhone P, Logue CM, Foley SL. 2012. DNA sequence analysis of plasmids from multidrug resistant *Salmonella enterica* serotype Heidelberg isolates. *PLoS One* 7:e51160. doi:10.1371/journal.pone.0051160.
 84. Johnson TJ, Thorsness JL, Anderson CP, Lynne AM, Foley SL, Han J, Fricke WF, McDermott PF, White DG, Khatri M, Stell AL, Flores C, Singer RS. 2010. Horizontal gene transfer of a ColV plasmid has resulted in a dominant avian clonal type of *Salmonella enterica* serovar Kentucky. *PLoS One* 5:e15524. doi:10.1371/journal.pone.0015524.
 85. Ashkenazi S, Cleary TG, Murray BE, Wanger A, Pickering LK. 1988. Quantitative analysis and partial characterization of cytotoxin production by *Salmonella* strains. *Infect. Immun.* 56:3089–3094.
 86. Ketyi I, Pacsa S, Emody L, Vertenyi A, Kocsis B, Kuch B. 1979. *Shigella dysenteriae* 1-like cytotoxic enterotoxins produced by *Salmonella* strains. *Acta Microbiol. Acad. Sci. Hung.* 26:217–223.
 87. Prager R, Fruth A, Tschape H. 1995. *Salmonella* enterotoxin (*stn*) gene is prevalent among strains of *Salmonella enterica*, but not among *Salmonella bongori* and other Enterobacteriaceae. *FEMS Immunol. Med. Microbiol.* 12:47–50.
 88. Libby SJ, Goebel W, Ludwig A, Buchmeier N, Bowe F, Fang FC, Guiney DG, Songer JG, Heffron F. 1994. A cytolyisin encoded by *Sal-*

- monella* is required for survival within macrophages. Proc. Natl. Acad. Sci. U. S. A. 91:489–493.
89. Collinson SK, Liu SL, Clouthier SC, Banser PA, Doran JL, Sanderson KE, Kay WW. 1996. The location of four fimbriin-encoding genes, agfA, fimA, sefA and sefD, on the *Salmonella enteritidis* and/or *S. typhimurium* XbaI-BlnI genomic restriction maps. Gene 169:75–80.
 90. Clouthier SC, Collinson SK, Kay WW. 1994. Unique fimbriae-like structures encoded by *sefD* of the SEF14 fimbrial gene cluster of *Salmonella enteritidis*. Mol. Microbiol. 12:893–901.
 91. Betancor L, Yim L, Martinez A, Fookes M, Sasia S, Schelotto F, Thomson N, Maskell D, Chabalgoity JA. 2012. Genomic comparison of the closely related *Salmonella enterica* serovars Enteritidis and Dublin. Open Microbiol. J. 6:5–13.
 92. White AP, Gibson DL, Collinson SK, Banser PA, Kay WW. 2003. Extracellular polysaccharides associated with thin aggregative fimbriae of *Salmonella enterica* serovar Enteritidis. J. Bacteriol. 185:5398–5407.
 93. Thorns CJ, Turcotte C, Gemmell CG, Woodward MJ. 1996. Studies into the role of the SEF14 fimbrial antigen in the pathogenesis of *Salmonella enteritidis*. Microb. Pathog. 20:235–246.
 94. Zhang XL, Tsui IS, Yip CM, Fung AW, Wong DK, Dai X, Yang Y, Hackett J, Morris C. 2000. *Salmonella enterica* serovar Typhi uses type IVB pili to enter human intestinal epithelial cells. Infect. Immun. 68:3067–3073.
 95. Baumber AJ, Heffron F. 1995. Identification and sequence analysis of *lpfABCDE*, a putative fimbrial operon of *Salmonella typhimurium*. J. Bacteriol. 177:2087–2097.
 96. Friedrich MJ, Kinsey NE, Vila J, Kadner RJ. 1993. Nucleotide sequence of a 13.9 kb segment of the 90 kb virulence plasmid of *Salmonella typhimurium*: the presence of fimbrial biosynthetic genes. Mol. Microbiol. 8:543–558.
 97. Turcotte C, Woodward MJ. 1993. Cloning, DNA nucleotide sequence and distribution of the gene encoding the SEF14 fimbrial antigen of *Salmonella enteritidis*. J. Gen. Microbiol. 139:1477–1485.
 98. Doran JL, Collinson SK, Clouthier SC, Cebula TA, Koch WH, Burian J, Banser PA, Todd EK, Kay WW. 1996. Diagnostic potential of sefA DNA probes to *Salmonella enteritidis* and certain other O-serogroup D1 *Salmonella* serovars. Mol. Cell. Probes 10:233–246.
 99. Althouse C, Patterson S, Fedorka-Cray P, Isaacson RE. 2003. Type 1 fimbriae of *Salmonella enterica* serovar Typhimurium bind to enterocytes and contribute to colonization of swine in vivo. Infect. Immun. 71:6446–6452.
 100. De Buck J, Van Immerseel F, Haesebrouck F, Ducatelle R. 2005. Protection of laying hens against *Salmonella* Enteritidis by immunization with type 1 fimbriae. Vet. Microbiol. 105:93–101.
 101. Clayton DJ, Bowen AJ, Hulme SD, Buckley AM, Deacon VL, Thomson NR, Barrow PA, Morgan E, Jones MA, Watson M, Stevens MP. 2008. Analysis of the role of 13 major fimbrial subunits in colonisation of the chicken intestines by *Salmonella enterica* serovar Enteritidis reveals a role for a novel locus. BMC Microbiol. 8:228. doi:10.1186/1471-2180-8-228.
 102. Dauga C, Zabrovskaja A, Grimont PA. 1998. Restriction fragment length polymorphism analysis of some flagellin genes of *Salmonella enterica*. J. Clin. Microbiol. 36:2835–2843.
 103. Turner AK, Lovell MA, Hulme SD, Zhang-Barber L, Barrow PA. 1998. Identification of *Salmonella typhimurium* genes required for colonization of the chicken alimentary tract and for virulence in newly hatched chicks. Infect. Immun. 66:2099–2106.
 104. Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Hautefort I, Thompson A, Hinton JC, Van Immerseel F. 2006. Butyrate specifically down-regulates *Salmonella* pathogenicity island 1 gene expression. Appl. Environ. Microbiol. 72:946–949.
 105. Ochman H, Moran NA. 2001. Genes lost and genes found: evolution of bacterial pathogenesis and symbiosis. Science 292:1096–1099.
 106. Maurelli AT. 2007. Black holes, antivirulence genes, and gene inactivation in the evolution of bacterial pathogens. FEMS Microbiol. Lett. 267:1–8.
 107. Groisman EA, Ochman H. 1997. How *Salmonella* became a pathogen. Trends Microbiol. 5:343–349.
 108. Bowe F, Lipps CJ, Tsolis RM, Groisman E, Heffron F, Kusters JG. 1998. At least four percent of the *Salmonella typhimurium* genome is required for fatal infection of mice. Infect. Immun. 66:3372–3377.
 109. Jarvik T, Smillie C, Groisman EA, Ochman H. 2010. Short-term signatures of evolutionary change in the *Salmonella enterica* serovar Typhimurium 14028 genome. J. Bacteriol. 192:560–567.
 110. Baumber AJ, Tsolis RM, Ficht TA, Adams LG. 1998. Evolution of host adaptation in *Salmonella enterica*. Infect. Immun. 66:4579–4587.
 111. Jacobsen A, Hendriksen RS, Aarestrup FM, Ussery DW, Friis C. 2011. The *Salmonella enterica* pan-genome. Microb. Ecol. 62:487–504.
 112. Quiroz TS, Nieto PA, Tobar HE, Salazar-Echegarai FJ, Lizana RJ, Quezada CP, Santiviago CA, Araya DV, Riedel CA, Kalergis AM, Bueno SM. 2011. Excision of an unstable pathogenicity island in *Salmonella enterica* serovar Enteritidis is induced during infection of phagocytic cells. PLoS One 6:e26031. doi:10.1371/journal.pone.0026031.
 113. Wozniak RA, Fouts DE, Spagnoletti M, Colombo MM, Ceccarelli D, Garriss G, Dery C, Burrus V, Waldor MK. 2009. Comparative ICE genomics: insights into the evolution of the SXT/R391 family of ICEs. PLoS Genet. 5:e1000786. doi:10.1371/journal.pgen.1000786.
 114. Seth-Smith HM, Fookes MC, Okoro CK, Baker S, Harris SR, Scott P, Pickard D, Quail MA, Churcher C, Sanders M, Harmse J, Dougan G, Parkhill J, Thomson NR. 2012. Structure, diversity, and mobility of the *Salmonella* pathogenicity island 7 family of integrative and conjugative elements within Enterobacteriaceae. J. Bacteriol. 194:1494–1504.
 115. Moreno Switt AI, den Bakker HC, Cummings CA, Rodriguez-Rivera LD, Govoni G, Raneiri ML, Degoricija L, Brown S, Hoelzer K, Peters JE, Bolchacova E, Furtado MR, Wiedmann M. 2012. Identification and characterization of novel *Salmonella* mobile elements involved in the dissemination of genes linked to virulence and transmission. PLoS One 7:e41247. doi:10.1371/journal.pone.0041247.
 116. Williams LE, Wireman J, Hilliard VC, Summers AO. 2013. Large plasmids of *Escherichia coli* and *Salmonella* encode highly diverse arrays of accessory genes on common replicon families. Plasmid 69:36–48.
 117. Bielak E, Bergenholtz RD, Jorgensen MS, Sorensen SJ, Hansen LH, Hasman H. 2011. Investigation of diversity of plasmids carrying the *bla*TEM-52 gene. J. Antimicrob. Chemother. 66:2465–2474.
 118. Jones C, Stanley J. 1992. *Salmonella* plasmids of the pre-antibiotic era. J. Gen. Microbiol. 138:189–197.
 119. Gilmour MW, Thomson NR, Sanders M, Parkhill J, Taylor DE. 2004. The complete nucleotide sequence of the resistance plasmid R478: defining the backbone components of incompatibility group H conjugative plasmids through comparative genomics. Plasmid 52:182–202.
 120. Tosini F, Visca P, Luzzi I, Dionisi AM, Pezzella C, Petrucca A, Carattoli A. 1998. Class 1 integron-borne multiple-antibiotic resistance carried by IncFI and IncLM plasmids in *Salmonella enterica* serotype Typhimurium. Antimicrob. Agents Chemother. 42:3053–3058.
 121. Welch TJ, Fricke WF, McDermott PF, White DG, Rosso ML, Rasko DA, Mammel MK, Eppinger M, Rosovitz MJ, Wagner D, Rahalison L, Leclerc JE, Hinshaw JM, Lindler LE, Cebula TA, Carniel E, Ravel J. 2007. Multiple antimicrobial resistance in plague: an emerging public health risk. PLoS One 2:e309. doi:10.1371/journal.pone.0000309.
 122. Garcia-Quintanilla M, Ramos-Morales F, Casadesus J. 2008. Conjugal transfer of the *Salmonella enterica* virulence plasmid in the mouse intestine. J. Bacteriol. 190:1922–1927.
 123. Rychlik I, Karasova D, Sebkova A, Volf J, Sisak F, Havlickova H, Kummer V, Imre A, Szmolka A, Nagy B. 2009. Virulence potential of five major pathogenicity islands (SPI-1 to SPI-5) of *Salmonella enterica* serovar Enteritidis for chickens. BMC Microbiol. 9:268. doi:10.1186/1471-2180-9-268.
 124. Guiney DG, Fang FC, Krause M, Libby S. 1994. Plasmid-mediated virulence genes in non-typhoid *Salmonella* serovars. FEMS Microbiol. Lett. 124:1–9.
 125. Wilson JA, Gulig PA. 1998. Regulation of the *spvR* gene of the *Salmonella typhimurium* virulence plasmid during exponential-phase growth in intracellular salts medium and at stationary phase in L broth. Microbiology 144:1823–1833.
 126. Gulig PA, Doyle TJ, Hughes JA, Matsui H. 1998. Analysis of host cells associated with the Spv-mediated increased intracellular growth rate of *Salmonella typhimurium* in mice. Infect. Immun. 66:2471–2485.
 127. Gulig PA, Danbara H, Guiney DG, Lax AJ, Norel F, Rhen M. 1993. Molecular analysis of *spv* virulence genes of the *Salmonella* virulence plasmids. Mol. Microbiol. 7:825–830.
 128. Chu C, Feng Y, Chien AC, Hu S, Chu CH, Chiu CH. 2008. Evolution of genes on the *Salmonella* virulence plasmid phylogeny revealed from sequencing of the virulence plasmids of *S. enterica* serotype Dublin and comparative analysis. Genomics 92:339–343.
 129. Garcia P, Guerra B, Bances M, Mendoza MC, Rodicio MR. 2011. IncA/C plasmids mediate antimicrobial resistance linked to virulence

- genes in the Spanish clone of the emerging *Salmonella enterica* serotype 4,[5],12:i. J. Antimicrob. Chemother. 66:543–549.
130. Garcia Fernandez A, Cloeckaert A, Bertini A, Praud K, Doublet B, Weill FX, Carattoli A. 2007. Comparative analysis of IncHI2 plasmids carrying blaCTX-M-2 or blaCTX-M-9 from *Escherichia coli* and *Salmonella enterica* strains isolated from poultry and humans. Antimicrob. Agents Chemother. 51:4177–4180.
 131. Yu H, Wang J, Ye J, Tang P, Chu C, Hu S, Chiu CH. 2006. Complete nucleotide sequence of pSCV50, the virulence plasmid of *Salmonella enterica* serovar Choleraesuis SC-B67. Plasmid 55:145–151.
 132. Olsen JE, Brown DJ, Thomsen LE, Platt DJ, Chadfield MS. 2004. Differences in the carriage and the ability to utilize the serotype associated virulence plasmid in strains of *Salmonella enterica* serotype Typhimurium investigated by use of a self-transferable virulence plasmid, pOG669. Microb. Pathog. 36:337–347.
 133. Villa L, Carattoli A. 2005. Integrations and transposons on the *Salmonella enterica* serovar Typhimurium virulence plasmid. Antimicrob. Agents Chemother. 49:1194–1197.
 134. Herrero A, Mendoza MC, Rodicio R, Rodicio MR. 2008. Characterization of pUO-StVR2, a virulence-resistance plasmid evolved from the pSLT virulence plasmid of *Salmonella enterica* serovar Typhimurium. Antimicrob. Agents Chemother. 52:4514–4517.
 135. Rodriguez I, Rodicio MR, Guerra B, Hopkins KL. 2012. Potential international spread of multidrug-resistant invasive *Salmonella enterica* serovar Enteritidis. Emerg. Infect. Dis. 18:1173–1176.
 136. Fluit AC. 2005. Towards more virulent and antibiotic-resistant *Salmonella*? FEMS Immunol. Med. Microbiol. 43:1–11.
 137. Folster JP, Pecic G, McCullough A, Rickert R, Whichard JM. 2011. Characterization of bla(CMY)-encoding plasmids among *Salmonella* isolated in the United States in 2007. Foodborne Pathog. Dis. 8:1289–1294.
 138. Folster JP, Pecic G, Singh A, Duval B, Rickert R, Ayers S, Abbott J, McGlinchey B, Bauer-Turpin J, Haro J, Hise K, Zhao S, Fedorka-Cray PJ, Whichard J, McDermott PF. 2012. Characterization of extended-spectrum cephalosporin-resistant *Salmonella enterica* serovar Heidelberg isolated from food animals, retail meat, and humans in the United States 2009. Foodborne Pathog. Dis. 9:638–645.
 139. Johnson TJ, Siek KE, Johnson SJ, Nolan LK. 2005. DNA sequence and comparative genomics of pAPEC-O2-R, an avian pathogenic *Escherichia coli* transmissible R plasmid. Antimicrob. Agents Chemother. 49:4681–4688.
 140. Garcia-Quintanilla M, Casadesus J. 2011. Virulence plasmid interchange between strains ATCC 14028, LT2, and SL1344 of *Salmonella enterica* serovar Typhimurium. Plasmid 65:169–175.
 141. Sampei G, Furuya N, Tachibana K, Saitou Y, Suzuki T, Mizobuchi K, Komano T. 2010. Complete genome sequence of the incompatibility group I1 plasmid R64. Plasmid 64:92–103.
 142. Johnson TJ, Shepard SM, Rivet B, Danzeisen JL, Carattoli A. 2011. Comparative genomics and phylogeny of the IncI1 plasmids: a common plasmid type among porcine enterotoxigenic *Escherichia coli*. Plasmid 66:144–151.
 143. Garcia-Fernandez A, Fortini D, Veldman K, Mevius D, Carattoli A. 2009. Characterization of plasmids harbouring *qnrS1*, *qnrB2* and *qnrB19* genes in *Salmonella*. J. Antimicrob. Chemother. 63:274–281.
 144. Shahada F, Sekizuka T, Kuroda M, Kusumoto M, Ohishi D, Matsumoto A, Okazaki H, Tanaka K, Uchida I, Izumiya H, Watanabe H, Tamamura Y, Iwata T, Akiba M. 2011. Characterization of *Salmonella enterica* serovar Typhimurium isolates harboring a chromosomally encoded CMY-2 beta-lactamase gene located on a multidrug resistance genomic island. Antimicrob. Agents Chemother. 55:4114–4121.
 145. Antunes P, Mourao J, Machado J, Peixe L. 2011. First description of *qnrS1*-IncN plasmid in a ST11 *Salmonella* Enteritidis clinical isolate from Portugal. Diagn. Microbiol. Infect. Dis. 69:463–465.
 146. Szmolka A, Fortini D, Villa L, Carattoli A, Anjum MF, Nagy B. 2011. First report on IncN plasmid-mediated quinolone resistance gene *qnrS1* in porcine *Escherichia coli* in Europe. Microb. Drug Resist. 17:567–573.
 147. Gutierrez B, Herrera-Leon S, Escudero JA, Hidalgo L, Gonzalez-Sanz R, Arroyo M, San Millan A, Echeita MA, Gonzalez-Zorn B. 2009. Novel genetic environment of *qnrB2* associated with TEM-1 and SHV-12 on pB1004, an IncHI2 plasmid, in *Salmonella* Bredeney BB1047 from Spain. J. Antimicrob. Chemother. 64:1334–1336.
 148. Garcia-Fernandez A, Chiaretto G, Bertini A, Villa L, Fortini D, Ricci A, Carattoli A. 2008. Multilocus sequence typing of IncI1 plasmids carrying extended-spectrum beta-lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. J. Antimicrob. Chemother. 61:1229–1233.
 149. Cain AK, Hall RM. 2012. Evolution of a multiple antibiotic resistance region in IncHI1 plasmids: reshaping resistance regions in situ. J. Antimicrob. Chemother. 67:2848–2853.
 150. Fernandez-Alarcon C, Singer RS, Johnson TJ. 2011. Comparative genomics of multidrug resistance-encoding IncA/C plasmids from commensal and pathogenic *Escherichia coli* from multiple animal sources. PLoS One 6:e23415. doi:10.1371/journal.pone.0023415.
 151. Fricke WF, Welch TJ, McDermott PF, Mammel MK, LeClerc JE, White DG, Cebula TA, Ravel J. 2009. Comparative genomics of the IncA/C multidrug resistance plasmid family. J. Bacteriol. 191:4750–4757.
 152. Tran T, Andres P, Petroni A, Soler-Bistue A, Albornoz E, Zorreguieta A, Reyes-Lamotho R, Sherratt DJ, Corso A, Tolmasky ME. 2012. Small plasmids harboring *qnrB19*: a model for plasmid evolution mediated by site-specific recombination at *oriT* and Xer sites. Antimicrob. Agents Chemother. 56:1821–1827.
 153. Hammerl JA, Beutlich J, Hertwig S, Mevius D, Threlfall EJ, Helmuth R, Guerra B. 2010. pSGI15, a small ColE-like *qnrB19* plasmid of a *Salmonella enterica* serovar Typhimurium strain carrying *Salmonella* genomic island 1 (SGI1). J. Antimicrob. Chemother. 65:173–175.
 154. Targant H, Doublet B, Aarestrup FM, Cloeckaert A, Madec JY. 2010. IS6100-mediated genetic rearrangement within the complex class 1 integron In104 of the *Salmonella* genomic island 1. J. Antimicrob. Chemother. 65:1543–1545.
 155. Siebor E, Neuwirth C. 2011. The new variant of *Salmonella* genomic island 1 (SGI1-V) from a *Proteus mirabilis* French clinical isolate harbours *blaVEB-6* and *qnrA1* in the multiple antibiotic resistance region. J. Antimicrob. Chemother. 66:2513–2520.
 156. Krauland M, Harrison L, Paterson D, Marsh J. 2010. Novel integron gene cassette arrays identified in a global collection of multi-drug resistant non-typhoidal *Salmonella enterica*. Curr. Microbiol. 60:217–223.
 157. Beutlich J, Jahn S, Malorny B, Hauser E, Huhn S, Schroeter A, Rodicio MR, Appel B, Threlfall J, Mevius D, Helmuth R, Guerra B. 2011. Antimicrobial resistance and virulence determinants in European *Salmonella* genomic island 1-positive *Salmonella enterica* isolates from different origins. Appl. Environ. Microbiol. 77:5655–5664.
 158. Kiss J, Nagy B, Olasz F. 2012. Stability, entrapment and variant formation of *Salmonella* genomic island 1. PLoS One 7:e32497. doi:10.1371/journal.pone.0032497.
 159. Douard G, Praud K, Cloeckaert A, Doublet B. 2010. The *Salmonella* genomic island 1 is specifically mobilized in trans by the IncA/C multidrug resistance plasmid family. PLoS One 5:e15302. doi:10.1371/journal.pone.0015302.
 160. Frost JA, Ward LR, Rowe B. 1989. Acquisition of a drug resistance plasmid converts *Salmonella enteritidis* phage type 4 to phage type 24. Epidemiol. Infect. 103:243–248.
 161. Brown DJ, Baggesen DL, Platt DJ, Olsen JE. 1999. Phage type conversion in *Salmonella enterica* serotype Enteritidis caused by the introduction of a resistance plasmid of incompatibility group X (IncX). Epidemiol. Infect. 122:19–22.
 162. Coetzee JN, Bradley DE, du Toit L, Hedges RW. 1988. Bacteriophage X-2: a filamentous phage lysing IncX-plasmid-harboring bacterial strains. J. Gen. Microbiol. 134:2535–2541.
 163. Yong D, Toleman MA, Giske CG, Cho HS, Sundman K, Lee K, Walsh TR. 2009. Characterization of a new metallo-beta-lactamase gene, bla(NDM-1), and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. Antimicrob. Agents Chemother. 53:5046–5054.
 164. Ho PL, Lo WU, Yeung MK, Lin CH, Chow KH, Ang I, Tong AH, Bao JY, Lok S, Lo JY. 2011. Complete sequencing of pNDM-HK encoding NDM-1 carbapenemase from a multidrug-resistant *Escherichia coli* strain isolated in Hong Kong. PLoS One 6:e17989. doi:10.1371/journal.pone.0017989.
 165. Poirel L, Bonnin RA, Nordmann P. 2011. Analysis of the resistome of a multidrug-resistant NDM-1-producing *Escherichia coli* strain by high-throughput genome sequencing. Antimicrob. Agents Chemother. 55:4224–4229.
 166. Poirel L, Dortet L, Bernabeu S, Nordmann P. 2011. Genetic features of

- bla*NDM-1-positive *Enterobacteriaceae*. *Antimicrob. Agents Chemother.* 55:5403–5407.
167. Carattoli A, Villa L, Poirel L, Bonnin RA, Nordmann P. 2012. Evolution of IncA/C *bla*CMY-2-carrying plasmids by acquisition of the *bla*NDM-1 carbapenemase gene. *Antimicrob. Agents Chemother.* 56:783–786.
 168. Chen Y, Zhou Z, Jiang Y, Yu Y. 2011. Emergence of NDM-1-producing *Acinetobacter baumannii* in China. *J. Antimicrob. Chemother.* 66:1255–1259.
 169. Savard P, Gopinath R, Zhu W, Kitchel B, Rasheed JK, Tekle T, Roberts A, Ross T, Razeq J, Landrum BM, Wilson LE, Limbago B, Perl TM, Carroll KC. 2011. First NDM-positive *Salmonella* sp. strain identified in the United States. *Antimicrob. Agents Chemother.* 55:5957–5958.
 170. Reva O, Bezuidt O. 2012. Distribution of horizontally transferred heavy metal resistance operons in recent outbreak bacteria. *Mob. Genet. Elements* 2:96–100.
 171. Brilli M, Mengoni A, Fondi M, Bazzicalupo M, Lio P, Fani R. 2008. Analysis of plasmid genes by phylogenetic profiling and visualization of homology relationships using Blast2Network. *BMC Bioinformatics* 9:551. doi:10.1186/1471-2105-9-551.
 172. Figueroa-Bossi N, Bossi L. 1999. Inducible prophages contribute to *Salmonella* virulence in mice. *Mol. Microbiol.* 33:167–176.
 173. Lemire S, Figueroa-Bossi N, Bossi L. 2011. Bacteriophage crosstalk: coordination of prophage induction by trans-acting antirepressors. *PLoS Genet.* 7:e1002149. doi:10.1371/journal.pgen.1002149.
 174. Kropinski AM, Sulakvelidze A, Konczyk P, Poppe C. 2007. *Salmonella* phages and prophages—genomics and practical aspects. *Methods Mol. Biol.* 394:133–175.
 175. Casjens SR, Thuman-Commike PA. 2011. Evolution of mosaic related tailed bacteriophage genomes seen through the lens of phage P22 virion assembly. *Virology* 411:393–415.
 176. Poppe C. 1999. Epidemiology of *Salmonella enterica* serovar Enteritidis, p 3–18. In Saeed AM, Gast RK, Potter ME, Wall PG (ed), *Salmonella enterica* serovar Enteritidis in humans and animals—epidemiology, pathogenesis, and control. Iowa State University Press, Ames, IA.
 177. Pang S, Octavia S, Reeves PR, Wang Q, Gilbert GL, Sintchenko V, Lan R. 2012. Genetic relationships of phage types and single nucleotide polymorphism typing of *Salmonella enterica* serovar Typhimurium. *J. Clin. Microbiol.* 50:727–734.
 178. Altekruse S, Koehler J, Hickman-Brenner F, Tauxe RV, Ferris K. 1993. A comparison of *Salmonella enteritidis* phage types from egg-associated outbreaks and implicated laying flocks. *Epidemiol. Infect.* 110:17–22.
 179. Boyd EF. 2012. Bacteriophage-encoded bacterial virulence factors and phage-pathogenicity island interactions. *Adv. Virus Res.* 82:91–118.
 180. Brown NF, Coombes BK, Bishop JL, Wickham ME, Lowden MJ, Gal-Mor O, Goode DL, Boyle EC, Sanderson KL, Finlay BB. 2011. *Salmonella* phage ST64B encodes a member of the SseK/NleB effector family. *PLoS One* 6:e17824. doi:10.1371/journal.pone.0017824.
 181. Wray C, Beedell YE, McLaren IM. 1991. A survey of antimicrobial resistance in salmonellae isolated from animals in England and Wales during 1984–1987. *Br. Vet. J.* 147:356–369.
 182. Threlfall EJ, Ward LR, Rowe B. 1978. Spread of multiresistant strains of *Salmonella typhimurium* phage types 204 and 193 in Britain. *Br. Med. J.* 2:997.
 183. Kuhn H, Rabsch W, Tschape H, Tietze E. 1982. Characterization and epidemiology of a *Salmonella typhimurium* epidemic strain. *Z. Arztl. Fortbild. (Jena)* 76:607–610.
 184. Slominski B, Calkiewicz J, Golec P, Wegrzyn G, Wrobel B. 2007. Plasmids derived from Gifsy-1/Gifsy-2, lambdoid prophages contributing to the virulence of *Salmonella enterica* serovar Typhimurium: implications for the evolution of replication initiation proteins of lambdoid phages and enterobacteria. *Microbiology* 153:1884–1896.
 185. Figueroa-Bossi N, Uzzau S, Maloriol D, Bossi L. 2001. Variable assortment of prophages provides a transferable repertoire of pathogenic determinants in *Salmonella*. *Mol. Microbiol.* 39:260–271.
 186. Ho TD, Figueroa-Bossi N, Wang M, Uzzau S, Bossi L, Slauch JM. 2002. Identification of GtgE, a novel virulence factor encoded on the Gifsy-2 bacteriophage of *Salmonella enterica* serovar Typhimurium. *J. Bacteriol.* 184:5234–5239.
 187. Rychlik I, Sebkova A, Gregorova D, Karpiskova R. 2001. Low-molecular-weight plasmid of *Salmonella enterica* serovar Enteritidis codes for retron reverse transcriptase and influences phage resistance. *J. Bacteriol.* 183:2852–2858.
 188. Smarda J, Doroszkiewicz W, Lachowicz TM. 1990. Sensitivity of *Shigella flexneri* and *Escherichia coli* bacteria to bacteriophages and to colicins, lost or established by the acquisition of R plasmids. *Acta Microbiol. Pol.* 39:23–35.
 189. Fricke WF, Mammel MK, McDermott PF, Tartera C, White DG, Leclerc JE, Ravel J, Cebula TA. 2011. Comparative genomics of 28 *Salmonella enterica* isolates: evidence for CRISPR-mediated adaptive sublineage evolution. *J. Bacteriol.* 193:3556–3568.
 190. Matthews TD, Edwards R, Maloy S. 2010. Chromosomal rearrangements formed by rrn recombination do not improve replicore balance in host-specific *Salmonella enterica* serovars. *PLoS One* 5:e13503. doi:10.1371/journal.pone.0013503.
 191. Thomson NR, Clayton DJ, Windhorst D, Vernikos G, Davidson S, Churcher C, Quail MA, Stevens M, Jones MA, Watson M, Barron A, Layton A, Pickard D, Kingsley RA, Bignell A, Clark L, Harris B, Ormond D, Abdellah Z, Brooks K, Cherevach I, Chillingworth T, Woodward J, Norberczak H, Lord A, Arrowsmith C, Jagels K, Moule S, Mungall K, Sanders M, Whitehead S, Chabalgoity JA, Maskell D, Humphrey T, Roberts M, Barrow PA, Dougan G, Parkhill J. 2008. Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91 provides insights into evolutionary and host adaptation pathways. *Genome Res.* 18:1624–1637.
 192. Helm RA, Lee AG, Christman HD, Maloy S. 2003. Genomic rearrangements at rrn operons in *Salmonella*. *Genetics* 165:951–959.
 193. Feng Y, Chien KY, Chen HL, Chiu CH. 2012. Pseudogene recoding revealed from proteomic analysis of *Salmonella* serovars. *J. Proteome Res.* 11:1715–1719.
 194. Holt KE, Thomson NR, Wain J, Langridge GC, Hasan R, Bhutta ZA, Quail MA, Norbertczak H, Walker D, Simmonds M, White B, Bason N, Mungall K, Dougan G, Parkhill J. 2009. Pseudogene accumulation in the evolutionary histories of *Salmonella enterica* serovars Paratyphi A and Typhi. *BMC Genomics* 10:36. doi:10.1186/1471-2164-10-36.
 195. Kuo CH, Ochman H. 2010. The extinction dynamics of bacterial pseudogenes. *PLoS Genet.* 6:e1001050. doi:10.1371/journal.pgen.1001050.
 196. Hurst LD. 2002. The Ka/Ks ratio: diagnosing the form of sequence evolution. *Trends Genet.* 18:486.
 197. Winter SE, Raffatellu M, Wilson RP, Russmann H, Baumler AJ. 2008. The *Salmonella enterica* serotype Typhi regulator TviA reduces interleukin-8 production in intestinal epithelial cells by repressing flagellin secretion. *Cell. Microbiol.* 10:247–261.
 198. Centers for Disease Control and Prevention. 2011. *Salmonella* surveillance: annual summary, 2009. Centers for Disease Control and Prevention, Atlanta, GA.
 199. Elson R, Little CL, Mitchell RT. 2005. *Salmonella* and raw shell eggs: results of a cross-sectional study of contamination rates and egg safety practices in the United Kingdom catering sector in 2003. *J. Food Prot.* 68:256–264.
 200. Hennessy TW, Cheng LH, Kassenborg H, Ahuja SD, Mohle-Boetani J, Marcus R, Shiferaw B, Angulo FJ. 2004. Egg consumption is the principal risk factor for sporadic *Salmonella* serotype Heidelberg infections: a case-control study in FoodNet sites. *Clin. Infect. Dis.* 38:S237–S243. doi:10.1086/381593.
 201. Kuehn BM. 2010. *Salmonella* cases traced to egg producers: findings trigger recall of more than 500 million eggs. *JAMA* 304:1316. doi:10.1001/jama.2010.1330.
 202. Roy P, Dhillon AS, Lauerman LH, Schaberg DM, Bandli D, Johnson S. 2002. Results of *Salmonella* isolation from poultry products, poultry, poultry environment, and other characteristics. *Avian Dis.* 46:17–24.
 203. Han J, David DE, Deck J, Lynne AM, Kaldhove P, Nayak R, Stefanova R, Foley SL. 2011. Comparison of *Salmonella enterica* serovar Heidelberg isolated from human patients with those from animal and food sources. *J. Clin. Microbiol.* 49:1130–1133.
 204. Hennessy TW, Hedberg CW, Slutsker L, White KE, Besser-Wiek JM, Moen ME, Feldman J, Coleman JW, Edmonson LM, MacDonald KL, Osterholm MT. 1996. A national outbreak of *Salmonella enteritidis* infections from ice cream. The Investigation Team. *N. Engl. J. Med.* 334:1281–1286.
 205. Le Hello S, Hendriksen RS, Doublet B, Fisher I, Nielsen EM, Whichard JM, Bouchrif B, Fashae K, Granier SA, Jourdan-Da Silva N, Cloeckaert A, Threlfall EJ, Angulo FJ, Aarestrup FM, Wain J, Weill FX. 2011. International spread of an epidemic population of *Salmonella enterica* serotype Kentucky ST198 resistant to ciprofloxacin. *J. Infect. Dis.* 204:675–684.

206. Eswarappa SM, Janice J, Nagarajan AG, Balasundaram SV, Karnam G, Dixit NM, Chakravorty D. 2008. Differentially evolved genes of *Salmonella* pathogenicity islands: insights into the mechanism of host specificity in *Salmonella*. PLoS One 3:e3829. doi:10.1371/journal.pone.0003829.
207. US Department of Agriculture. 2010. Overview of the United States dairy industry. US Department of Agriculture, National Agricultural Statistics Service, Washington, DC.
208. Joerger RD, Sartori CA, Kniel KE. 2009. Comparison of genetic and physiological properties of *Salmonella enterica* isolates from chickens reveals one major difference between serovar Kentucky and other serovars: response to acid. Foodborne Pathog. Dis. 6:503–512.
209. Johnson TJ, Johnson SJ, Nolan LK. 2006. Complete DNA sequence of a ColBM plasmid from avian pathogenic *Escherichia coli* suggests that it evolved from closely related ColV virulence plasmids. J. Bacteriol. 188:5975–5983.
210. US Department of Agriculture. 1997. The national poultry improvement plan and auxiliary provisions. US Department of Agriculture, Animal and Plant Health Inspection Service, Washington, DC.
211. Marrero-Ortiz R, Han J, Lynne AM, David DE, Stemper ME, Farmer D, Burkhardt W, III, Nayak R, Foley SL. 2012. Genetic characterization of antimicrobial resistance in *Salmonella enterica* serovars isolated from dairy cattle in Wisconsin. Food Res. Int. 45:962–967.
212. Melendez SN, Hanning I, Han J, Nayak R, Clement AR, Wooming A, Herrera P, Jones FT, Foley SL, Ricke SC. 2010. *Salmonella enterica* isolates from pasture-raised poultry exhibit antimicrobial resistance and class I integrons. J. Appl. Microbiol. 109:1957–1966.
213. Van Kessel JA, Karns JS, Wolfgang DR, Hovingh E, Schukken YH. 2012. Dynamics of *Salmonella* serotype shifts in an endemically infected dairy herd. Foodborne Pathog. Dis. 9:319–324.
214. Van Kessel JA, Karns JS, Lombard JE, Koprak CA. 2011. Prevalence of *Salmonella enterica*, *Listeria monocytogenes*, and *Escherichia coli* virulence factors in bulk tank milk and in-line filters from U.S. dairies. J. Food Prot. 74:759–768.
215. Dominguez M, Jourdan-Da Silva N, Vaillant V, Pihier N, Kermin C, Weill FX, Delmas G, Kerouanton A, Brisabois A, de Valk H. 2009. Outbreak of *Salmonella enterica* serotype Montevideo infections in France linked to consumption of cheese made from raw milk. Foodborne Pathog. Dis. 6:121–128.
216. Centers for Disease Control and Prevention. 2007. *Salmonella* Typhimurium infection associated with raw milk and cheese consumption—Pennsylvania, 2007. MMWR Morb. Mortal. Wkly. Rep. 56:1161–1164.
217. Mazurek J, Salehi E, Propes D, Holt J, Bannerman T, Nicholson LM, Bundesen M, Duffy R, Moolenaar RL. 2004. A multistate outbreak of *Salmonella enterica* serotype Typhimurium infection linked to raw milk consumption—Ohio, 2003. J. Food Prot. 67:2165–2170.
218. Kaldhove P, Nayak R, Lynne AM, David DE, McDermott PF, Logue CM, Foley SL. 2008. Characterization of *Salmonella enterica* serovar Heidelberg from turkey-associated sources. Appl. Environ. Microbiol. 74:5038–5046.
219. Lynne AM, Kaldhove P, David D, White DG, Foley SL. 2009. Characterization of antimicrobial resistance in *Salmonella enterica* serotype Heidelberg isolated from food animals. Foodborne Pathog. Dis. 6:207–215.
220. Didelot X, Bowden R, Street T, Golubchik T, Spencer C, McVean G, Sangal V, Anjum MF, Achtman M, Falush D, Donnelly P. 2011. Recombination and population structure in *Salmonella enterica*. PLoS Genet. 7:e1002191. doi:10.1371/journal.pgen.1002191.
221. Johnson TJ, Siek KE, Johnson SJ, Nolan LK. 2006. DNA sequence of a ColV plasmid and prevalence of selected plasmid-encoded virulence genes among avian *Escherichia coli* strains. J. Bacteriol. 188:745–758.
222. Johnson TJ, Lang KL. 2012. IncA/C plasmids: an emerging threat to human and animal health? Mob. Genet. Elements 2:55–58.
223. Brenner FW, Villar RG, Angulo FJ, Tauxe R, Swaminathan B. 2000. *Salmonella* nomenclature. J. Clin. Microbiol. 38:2465–2467.
224. Porwollik S, Boyd EF, Choy C, Cheng P, Florea L, Proctor E, McClelland M. 2004. Characterization of *Salmonella enterica* subspecies I genovars by use of microarrays. J. Bacteriol. 186:5883–5898.
225. Porwollik S, Santiviago CA, Cheng P, Florea L, Jackson S, McClelland M. 2005. Differences in gene content between *Salmonella enterica* serovar Enteritidis isolates and comparison to closely related serovars Gallinarum and Dublin. J. Bacteriol. 187:6545–6555.
226. Li J, Smith NH, Nelson K, Crichton PB, Old DC, Whittam TS, Selander RK. 1993. Evolutionary origin and radiation of the avian-adapted non-motile salmonellae. J. Med. Microbiol. 38:129–139.
227. Olsen JE, Skov MN, Christensen JP, Bisgaard M. 1996. Genomic lineage of *Salmonella enterica* serotype Gallinarum. J. Med. Microbiol. 45:413–418.
228. Richarme G, el Yaagoubi A, Kohiyama M. 1993. The MglA component of the binding protein-dependent galactose transport system of *Salmonella typhimurium* is a galactose-stimulated ATPase. J. Biol. Chem. 268:9473–9477.
229. Zhang Y, Franco M, Ducret A, Mignot T. 2010. A bacterial Ras-like small GTP-binding protein and its cognate GAP establish a dynamic spatial polarity axis to control directed motility. PLoS Biol. 8:e1000430. doi:10.1371/journal.pbio.1000430.
230. Nano FE, Schmerk C. 2007. The *Francisella* pathogenicity island. Ann. N. Y. Acad. Sci. 1105:122–137.
231. Mauriello EM, Mouhamar F, Nan B, Ducret A, Dai D, Zusman DR, Mignot T. 2010. Bacterial motility complexes require the actin-like protein, MreB and the Ras homologue, MglA. EMBO J. 29:315–326.
232. Kingsley RA, van Amsterdam K, Kramer N, Baumler AJ. 2000. The *shdA* gene is restricted to serotypes of *Salmonella enterica* subspecies I and contributes to efficient and prolonged fecal shedding. Infect. Immun. 68:2720–2727.
233. Kingsley RA, Abi Ghanem D, Puebla-Osorio N, Keestra AM, Berghman L, Baumler AJ. 2004. Fibronectin binding to the *Salmonella enterica* serotype Typhimurium ShdA autotransporter protein is inhibited by a monoclonal antibody recognizing the A3 repeat. J. Bacteriol. 186:4931–4939.
234. Wain J, House D, Parkhill J, Parry C, Dougan G. 2002. Unlocking the genome of the human typhoid bacillus. Lancet Infect. Dis. 2:163–170.
235. Edwards RA, Olsen GJ, Maloy SR. 2002. Comparative genomics of closely related salmonellae. Trends Microbiol. 10:94–99.
236. Eswarappa SM, Janice J, Balasundaram SV, Dixit NM, Chakravorty D. 2009. Host-specificity of *Salmonella enterica* serovar Gallinarum: insights from comparative genomics. Infect. Genet. Evol. 9:468–473.
237. Araya DV, Quiroz TS, Tobar HE, Lizana RJ, Quezada CP, Santiviago CA, Riedel CA, Kalergis AM, Bueno SM. 2010. Deletion of a prophage-like element causes attenuation of *Salmonella enterica* serovar Enteritidis and promotes protective immunity. Vaccine 28:5458–5466.
238. Schleker S, Sun J, Raghavan B, Srnc M, Muller N, Koepfinger M, Murthy L, Zhao Z, Klein-Seetharaman J. 2012. The current *Salmonella*-host interactome. Proteomics Clin. Appl. 6:117–133.
239. Desin TS, Mickael CS, Lam PK, Potter AA, Koster W. 2010. Protection of epithelial cells from *Salmonella enterica* serovar Enteritidis invasion by antibodies against the SPI-1 type III secretion system. Can. J. Microbiol. 56:522–526.
240. Kaiser P. 2010. Advances in avian immunology—prospects for disease control: a review. Avian Pathol. 39:309–324.
241. Barrow PA, Huggins MB, Lovell MA. 1994. Host specificity of *Salmonella* infection in chickens and mice is expressed in vivo primarily at the level of the reticuloendothelial system. Infect. Immun. 62:4602–4610.
242. Salcedo SP, Holden DW. 2005. Bacterial interactions with the eukaryotic secretory pathway. Curr. Opin. Microbiol. 8:92–98.
243. van der Heijden J, Finlay BB. 2012. Type III effector-mediated processes in *Salmonella* infection. Future Microbiol. 7:685–703.
244. Methner U, Haase A, Berndt A, Martin G, Nagy B, Barrow PA. 2011. Exploitation of intestinal colonization-inhibition between *Salmonella* organisms for live vaccines in poultry: potential and limitations. Zoonoses Public Health 58:540–548.
245. He H, Genovese KJ, Swaggerty CL, Nisbet DJ, Kogut MH. 2012. A comparative study on invasion, survival, modulation of oxidative burst, and nitric oxide responses of macrophages (HD11), and systemic infection in chickens by prevalent poultry *Salmonella* serovars. Foodborne Pathog. Dis. 9:1104–1110.
246. Shah DH, Zhou X, Addwebi T, Davis MA, Orfe L, Call DR, Guard J, Besser TE. 2011. Cell invasion of poultry-associated *Salmonella enterica* serovar Enteritidis isolates is associated with pathogenicity, motility and proteins secreted by the type III secretion system. Microbiology 157:1428–1445.
247. Shah DH, Zhou X, Addwebi T, Davis MA, Call DR. 2011. In vitro and in vivo pathogenicity of *Salmonella* Enteritidis clinical strains isolated from North America. Arch. Microbiol. 193:811–821.
248. St Louis ME, Morse DL, Potter ME, DeMelfi TM, Guzewish JJ, Tauxe RV, Blake PA. 1988. The emergence of grade A eggs as a major source of

- Salmonella enteritidis* infections. New implications for the control of salmonellosis. *JAMA* 259:2103–2107.
249. Guard J, Gast RK, Guraya R. 2010. Colonization of avian reproductive-tract tissues by variant subpopulations of *Salmonella* Enteritidis. *Avian Dis.* 54:857–861.
 250. Ricke SC, Jones DR, Gast RK. Eggs and egg products. In Doores S, Tortorello ML (ed), *Compendium of methods for the microbiological examinations of foods*, 5th ed, in press. American Public Health Association, Washington, DC.
 251. Golden NJ, Marks HH, Coleman ME, Schroeder CM, Bauer NE, Jr, Schlosser WD. 2008. Review of induced molting by feed removal and contamination of eggs with *Salmonella enterica* serovar Enteritidis. *Vet. Microbiol.* 131:215–228.
 252. Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Van Immerseel F. 2009. The *Salmonella* Enteritidis lipopolysaccharide biosynthesis gene *rfbH* is required for survival in egg albumen. *Zoonoses Public Health* 56:145–149.
 253. Howard ZR, O'Bryan CA, Crandall PG, Ricke SC. 2012. *Salmonella* Enteritidis in shell eggs: current issues and prospects for control. *Food Res. Int.* 45:755–764.
 254. Barrow PA, Jones MA, Smith AL, Wigley P. 2013. The long view: *Salmonella*—the last forty years. *Avian Pathol.* 41:413–420.
 255. Ricke SC. 2003. The gastrointestinal tract ecology of *Salmonella* Enteritidis colonization in molting hens. *Poult. Sci.* 82:1003–1007.
 256. Ricke SC, Dunkley CS, Durant JA. 2013. A review on development of novel strategies for controlling *Salmonella* Enteritidis colonization in laying hens: fiber-based molt diets. *Poult. Sci.* 92:502–525.
 257. Botteldoorn N, Van Coillie E, Goris J, Werbrouck H, Piessens V, Godard C, Scheldeman P, Herman L, Heyndrickx M. 2010. Limited genetic diversity and gene expression differences between egg- and non-egg-related *Salmonella* Enteritidis strains. *Zoonoses Public Health* 57: 345–357.
 258. Gast RK, Beard CW. 1990. Isolation of *Salmonella enteritidis* from internal organs of experimentally infected hens. *Avian Dis.* 34:991–993.
 259. De Buck J, Van Immerseel F, Meulemans G, Haesebrouck F, Ducatelle R. 2003. Adhesion of *Salmonella enterica* serotype Enteritidis isolates to chicken isthmal glandular secretions. *Vet. Microbiol.* 93:223–233.
 260. Keller LH, Benson CE, Krotke K, Eckroade RJ. 1995. *Salmonella enteritidis* colonization of the reproductive tract and forming and freshly laid eggs of chickens. *Infect. Immun.* 63:2443–2449.
 261. Keller LH, Schifferli DM, Benson CE, Aslam S, Eckroade RJ. 1997. Invasion of chicken reproductive tissues and forming eggs is not unique to *Salmonella enteritidis*. *Avian Dis.* 41:535–539.
 262. Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Gast R, Humphrey TJ, Van Immerseel F. 2009. Mechanisms of egg contamination by *Salmonella* Enteritidis. *FEMS Microbiol. Rev.* 33:718–738.
 263. Guard-Petter J. 2001. The chicken, the egg and *Salmonella enteritidis*. *Environ. Microbiol.* 3:421–430.
 264. Bohez L, Dewulf J, Ducatelle R, Pasmans F, Haesebrouck F, Van Immerseel F. 2008. The effect of oral administration of a homologous *hilaA* mutant strain on the long-term colonization and transmission of *Salmonella* Enteritidis in broiler chickens. *Vaccine* 26:372–378.
 265. Van Immerseel F, De Buck J, Pasmans F, Velge P, Bottreau E, Fievez V, Haesebrouck F, Ducatelle R. 2003. Invasion of *Salmonella enteritidis* in avian intestinal epithelial cells in vitro is influenced by short-chain fatty acids. *Int. J. Food Microbiol.* 85:237–248.
 266. Raspoet R, Gantois I, Devloo R, Martel A, Haesebrouck F, Pasmans F, Ducatelle R, Van Immerseel F. 2011. *Salmonella* Enteritidis universal stress protein (*usp*) gene expression is stimulated by egg white and supports oviduct colonization and egg contamination in laying hens. *Vet. Microbiol.* 153:186–190.
 267. De Vylder J, Raspoet R, Dewulf J, Haesebrouck F, Ducatelle R, Van Immerseel F. 2013. *Salmonella* Enteritidis is superior in egg white survival compared with other *Salmonella* serotypes. *Poult. Sci.* 92:842–845.
 268. Bohez L, Ducatelle R, Pasmans F, Botteldoorn N, Haesebrouck F, Van Immerseel F. 2006. *Salmonella enterica* serovar Enteritidis colonization of the chicken caecum requires the *HilA* regulatory protein. *Vet. Microbiol.* 116:202–210.
 269. Bohez L, Ducatelle R, Pasmans F, Haesebrouck F, Van Immerseel F. 2007. Long-term colonisation-inhibition studies to protect broilers against colonisation with *Salmonella* Enteritidis, using *Salmonella* pathogenicity island 1 and 2 mutants. *Vaccine* 25:4235–4243.
 270. Dunkley KD, McReynolds JL, Hume ME, Dunkley CS, Callaway TR, Kubena LF, Nisbet DJ, Ricke SC. 2007. Molting in *Salmonella* Enteritidis-challenged laying hens fed alfalfa crumble. II. Fermentation and microbial ecology response. *Poult. Sci.* 86:2101–2109.
 271. Rychlik I, Gregorova D, Hradecka H. 2006. Distribution and function of plasmids in *Salmonella enterica*. *Vet. Microbiol.* 112:1–10.
 272. Desin TS, Lam PK, Koch B, Mickael C, Berberov E, Wisner AL, Townsend HG, Potter AA, Koster W. 2009. *Salmonella enterica* serovar Enteritidis pathogenicity island 1 is not essential for but facilitates rapid systemic spread in chickens. *Infect. Immun.* 77:2866–2875.
 273. Murray RA, Lee CA. 2000. Invasion genes are not required for *Salmonella enterica* serovar Typhimurium to breach the intestinal epithelium: evidence that *Salmonella* pathogenicity island 1 has alternative functions during infection. *Infect. Immun.* 68:5050–5055.
 274. Clark L, Perrett CA, Malt L, Harward C, Humphrey S, Jepson KA, Martinez-Argudo I, Carney LJ, La Ragione RM, Humphrey TJ, Jepson MA. 2011. Differences in *Salmonella enterica* serovar Typhimurium strain invasiveness are associated with heterogeneity in SPI-1 gene expression. *Microbiology* 157:2072–2083.
 275. Silva CA, Blondel CJ, Quezada CP, Porwollik S, Andrews-Polymenis HL, Toro CS, Zaldivar M, Contreras I, McClelland M, Santiviago CA. 2012. Infection of mice by *Salmonella enterica* serovar Enteritidis involves additional genes that are absent in the genome of serovar Typhimurium. *Infect. Immun.* 80:839–849.
 276. Shah DH, Zhou X, Kim HY, Call DR, Guard J. 2012. Transposon mutagenesis of *Salmonella enterica* serovar Enteritidis identifies genes that contribute to invasiveness in human and chicken cells and survival in egg albumen. *Infect. Immun.* 80:4203–4215.
 277. Shah DH, Casavant C, Hawley Q, Addwebi T, Call DR, Guard J. 2012. *Salmonella* Enteritidis strains from poultry exhibit differential responses to acid stress, oxidative stress, and survival in the egg albumen. *Foodborne Pathog. Dis.* 9:258–264.
 278. Knodler LA, Steele-Mortimer O. 2003. Taking possession: biogenesis of the *Salmonella*-containing vacuole. *Traffic* 4:587–599.
 279. Beal RK, Wigley P, Powers C, Hulme SD, Barrow PA, Smith AL. 2004. Age at primary infection with *Salmonella enterica* serovar Typhimurium in the chicken influences persistence of infection and subsequent immunity to re-challenge. *Vet. Immunol. Immunopathol.* 100:151–164.
 280. Kaiser P, Rothwell L, Galyov EE, Barrow PA, Burnside J, Wigley P. 2000. Differential cytokine expression in avian cells in response to invasion by *Salmonella typhimurium*, *Salmonella enteritidis* and *Salmonella gallinarum*. *Microbiology* 146:3217–3226.
 281. Tsolis RM, Xavier MN, Santos RL, Baumler AJ. 2011. How to become a top model: impact of animal experimentation on human *Salmonella* disease research. *Infect. Immun.* 79:1806–1814.
 282. De Keersmaecker SC, Thijs IM, Vanderleyden J, Marchal K. 2006. Integration of omics data: how well does it work for bacteria? *Mol. Microbiol.* 62:1239–1250.
 283. Calenge F, Beaumont C. 2012. Toward integrative genomics study of genetic resistance to *Salmonella* and *Campylobacter* intestinal colonization in fowl. *Front. Genet.* 3:261. doi:10.3389/fgene.2012.00261.
 284. Lee SJ, Liang L, Juarez S, Nanton MR, Gondwe EN, Msefula CL, Kayala MA, Necchi F, Heath JN, Hart P, Tsolis RM, Heyderman RS, MacLennan CA, Felgner PL, Davies DH, McSorley SJ. 2012. Identification of a common immune signature in murine and human systemic salmonellosis. *Proc. Natl. Acad. Sci. U. S. A.* 109:4998–5003.
 285. Thiele I, Hyduke DR, Steeb B, Fankam G, Allen DK, Bazzani S, Charusanti P, Chen FC, Fleming RM, Hsiung CA, De Keersmaecker SC, Liao YC, Marchal K, Mo ML, Ozdemir E, Raghunathan A, Reed JL, Shin SI, Sigurbjornsdottir S, Steinmann J, Sudarsan S, Swainston N, Thijs IM, Zengler K, Palsson BO, Adkins JN, Bumann D. 2011. A community effort towards a knowledge-base and mathematical model of the human pathogen *Salmonella* Typhimurium LT2. *BMC Syst. Biol.* 5:8. doi:10.1186/1752-0509-5-8.
 286. Gast RK, Guraya R, Guard-Bouldin J, Holt PS, Moore RW. 2007. Colonization of specific regions of the reproductive tract and deposition at different locations inside eggs laid by hens infected with *Salmonella enteritidis* or *Salmonella heidelberg*. *Avian Dis.* 51:40–44.
 287. Jones FT. 2011. A review of practical *Salmonella* control measures in animal feed. *J. Appl. Poult. Res.* 20:102–113.
 288. Park SY, Woodward CL, Kubena LF, Nisbet DJ, Birkhold SG, Ricke SC. 2008. Environmental dissemination of foodborne *Salmonella* in pre-harvest poultry production: reservoirs, critical factors and research strategies. *Crit. Rev. Environ. Sci. Technol.* 38:73–111.

289. Sirsat SA, Muthaiyan A, Ricke SC. 2009. Antimicrobials for pathogen reduction in organic and natural poultry production. *J. Appl. Poult. Res.* 18:379–388.
290. Hur J, Jawale C, Lee JW. 2013. Antimicrobial resistance of *Salmonella* isolated from food animals: a review. *Food Res. Int.* 45:819–830.
291. Moretro T, Heir E, Nesse LL, Vestby LK, Langsrud S. 2012. Control of *Salmonella* in food related environments by chemical disinfection. *Food Res. Int.* 45:532–544.
292. Ahmer BM, Gunn JS. 2011. Interaction of *Salmonella* spp. with the intestinal microbiota. *Front. Microbiol.* 2:101. doi:10.3389/fmicb.2011.00101.
293. Farr SB, Kogoma T. 1991. Oxidative stress responses in *Escherichia coli* and *Salmonella typhimurium*. *Microbiol. Rev.* 55:561–585.
294. Evans MR, Fink RC, Vazquez-Torres A, Porwollik S, Jones-Carson J, McClelland M, Hassan HM. 2011. Analysis of the ArcA regulon in anaerobically grown *Salmonella enterica* sv. Typhimurium. *BMC Microbiol.* 11:58. doi:10.1186/1471-2180-11-58.
295. Troxell B, Fink RC, Porwollik S, McClelland M, Hassan HM. 2011. The Fur regulon in anaerobically grown *Salmonella enterica* sv. Typhimurium: identification of new Fur targets. *BMC Microbiol.* 11:236. doi:10.1186/1471-2180-11-236.
296. Foster JW, Spector MP. 1995. How *Salmonella* survive against the odds. *Annu. Rev. Microbiol.* 49:145–174.
297. Wooldridge KG, Williams PH. 1993. Iron uptake mechanisms of pathogenic bacteria. *FEMS Microbiol. Rev.* 12:325–348.
298. Foster JW. 1991. *Salmonella* acid shock proteins are required for the adaptive acid tolerance response. *J. Bacteriol.* 173:6896–6902.
299. Spector MP. 1998. The starvation-stress response (SSR) of *Salmonella*. *Adv. Microb. Physiol.* 40:233–279.
300. Smith JL. 2003. The role of gastric acid in preventing foodborne disease and how bacteria overcome acid conditions. *J. Food Prot.* 66:1292–1303.
301. Bearson SMD, Bearson BL. 2010. Traversing the swine gastrointestinal tract: *Salmonella* survival and pathogenesis, p 35–48. *In* Ricke SC, Jones FT (ed), *Perspectives on food-safety issues of animal-derived foods*. University of Arkansas Press, Fayetteville, AR.
302. Korver DR. 2006. Overview of the immune dynamics of the digestive system. *J. Appl. Poult. Res.* 15:123–135.
303. Kosiewicz MM, Zirnheld AL, Alard P. 2011. Gut microbiota, immunity, and disease: a complex relationship. *Front. Microbiol.* 2:180. doi:10.3389/fmicb.2011.00180.
304. Kelly D, Conway S, Aminov R. 2005. Commensal gut bacteria: mechanisms of immune modulation. *Trends Immunol.* 26:326–333.
305. Kelly D, Mulder IE. 2012. Microbiome and immunological interactions. *Nutr. Rev.* 70:S18–S30. doi:10.1111/j.1753-4887.2012.00498.x.
306. Hallstrom K, McCormick BA. 2011. *Salmonella* interaction with and passage through the intestinal mucosa: through the lens of the organism. *Front. Microbiol.* 2:88. doi:10.3389/fmicb.2011.00088.
307. Reading NC, Kasper DL. 2011. The starting lineup: key microbial players in intestinal immunity and homeostasis. *Front. Microbiol.* 2:148. doi:10.3389/fmicb.2011.00148.
308. Cherrington CA, Hinton M, Mead GC, Chopra I. 1991. Organic acids: chemistry, antibacterial activity and practical applications. *Adv. Microb. Physiol.* 32:87–108.
309. Ricke SC, Koo OK, Keeton JT. 2013. Fermented meat, poultry, and fish products, p 857–880. *In* Doyle MP, Buchanan RL (ed), *Food microbiology: fundamentals and frontiers*, 4th ed. ASM Press, Washington, DC.
310. Davidson PM, Taylor TM, Schmidt SE. 2013. Chemical preservatives and natural antimicrobial compounds, p 765–801. *In* Doyle MP, Buchanan RL (ed), *Food microbiology: fundamentals and frontiers*, 4th ed. ASM Press, Washington, DC.
311. Russell JB, Diez-Gonzalez F. 1998. The effects of fermentation acids on bacterial growth. *Adv. Microb. Physiol.* 39:205–234.
312. Hirshfield IN, Terzulli S, O'Byrne C. 2003. Weak organic acids: a panoply of effects on bacteria. *Sci. Prog.* 86:245–269.
313. Van Immerseel F, Russell JB, Flythe MD, Gantois I, Timbermont L, Pasmans F, Haesebrouck F, Ducatelle R. 2006. The use of organic acids to combat *Salmonella* in poultry: a mechanistic explanation of the efficacy. *Avian Pathol.* 35:182–188.
314. Russell JB. 1992. Another explanation for the toxicity of fermentation acids at low pH—anion accumulation versus uncoupling. *J. Appl. Bacteriol.* 73:363–370.
315. Ricke SC. 2003. Perspectives on the use of organic acids and short chain fatty acids as antimicrobials. *Poult. Sci.* 82:632–639.
316. Maciorowski KG, Jones FT, Pillai SD, Ricke SC. 2004. Incidence and control of food-borne *Salmonella* spp. in poultry feeds—a review. *Worlds Poult. Sci. J.* 60:446–457.
317. Jones FT, Ricke SC. 2003. Observations on the history of the development of antimicrobials and their use in poultry feeds. *Poult. Sci.* 82:613–617.
318. Hungate RE. 1966. *The rumen and its microbes*. Academic Press, Inc, New York, NY.
319. Annison EF, Hill KJ, Kenworthy R. 1968. Volatile fatty acids in the digestive tract of the fowl. *Br. J. Nutr.* 22:207–216.
320. Cummings JH, Macfarlane GT. 1991. The control and consequences of bacterial fermentation in the human colon. *J. Appl. Bacteriol.* 70:443–459.
321. Cummings JH, Pomare EW, Branch WJ, Naylor CP, Macfarlane GT. 1987. Short chain fatty acids in human large intestine, portal, hepatic and venous blood. *Gut* 28:1221–1227.
322. Goepfert JM, Hicks R. 1969. Effect of volatile fatty acids on *Salmonella typhimurium*. *J. Bacteriol.* 97:956–958.
323. Khan M, Katamay M. 1969. Antagonistic effect of fatty acids against *Salmonella* in meat and bone meal. *Appl. Microbiol.* 17:402–404.
324. Eklund T. 1983. The antimicrobial effect of dissociated and undissociated sorbic acid at different pH levels. *J. Appl. Bacteriol.* 54:383–389.
325. McHan F, Shotts EB. 1993. Effect of short-chain fatty acids on the growth of *Salmonella typhimurium* in an in vitro system. *Avian Dis.* 37:396–398.
326. Leyer GJ, Johnson EA. 1992. Acid adaptation promotes survival of *Salmonella* spp. in cheese. *Appl. Environ. Microbiol.* 58:2075–2080.
327. Baik HS, Bearson S, Dunbar S, Foster JW. 1996. The acid tolerance response of *Salmonella typhimurium* provides protection against organic acids. *Microbiology* 142:3195–3200.
328. Kwon YM, Ricke SC. 1998. Induction of acid resistance of *Salmonella typhimurium* by exposure to short-chain fatty acids. *Appl. Environ. Microbiol.* 64:3458–3463.
329. Calhoun LN, Kwon YM. 2010. The effect of long-term propionate adaptation on the stress resistance of *Salmonella* Enteritidis. *J. Appl. Microbiol.* 109:1294–1300.
330. Dunkley KD, Callaway TR, O'Bryan C, Kunding MM, Dunkley CS, Anderson RC, Nisbet DJ, Crandall PG, Ricke SC. 2009. Cell yields and fermentation responses of a *Salmonella* Typhimurium poultry isolate at different dilution rates in an anaerobic steady state continuous culture. *Antonie Van Leeuwenhoek* 96:537–544.
331. Dunkley KD, Callaway TR, Chalova VI, Anderson RC, Kunding MM, Dunkley CS, Nisbet DJ, Ricke SC. 2008. Glucose yields and genetic responses in a poultry isolate of *Salmonella* Typhimurium in an anaerobic continuous culture during shifts in pH. *Anaerobe* 14:35–42.
332. Horswill AR, Escalante-Semerena JC. 1999. *Salmonella typhimurium* LT2 catabolizes propionate via the 2-methylcitric acid cycle. *J. Bacteriol.* 181:5615–5623.
333. Horswill AR, Dudding AR, Escalante-Semerena JC. 2001. Studies of propionate toxicity in *Salmonella enterica* identify 2-methylcitrate as a potent inhibitor of cell growth. *J. Biol. Chem.* 276:19094–19101.
334. Rocco CJ, Escalante-Semerena JC. 2010. *In Salmonella enterica*, 2-methylcitrate blocks gluconeogenesis. *J. Bacteriol.* 192:771–778.
335. Kwon YM, Park SY, Birkhold SG, Ricke SC. 2000. Induction of resistance of *Salmonella typhimurium* to environmental stresses by exposure to short-chain fatty acids. *J. Food Sci.* 65:1037–1040.
336. Greenacre EJ, Brocklehurst TF. 2006. The acetic acid tolerance response induces cross-protection to salt stress in *Salmonella typhimurium*. *Int. J. Food Microbiol.* 112:62–65.
337. Greenacre EJ, Lucchini S, Hinton JC, Brocklehurst TF. 2006. The lactic acid-induced acid tolerance response in *Salmonella enterica* serovar Typhimurium induces sensitivity to hydrogen peroxide. *Appl. Environ. Microbiol.* 72:5623–5625.
338. Milillo SR, Martin E, Muthaiyan A, Ricke SC. 2011. Immediate reduction of *Salmonella enterica* serotype Typhimurium viability via membrane destabilization following exposure to multiple-hurdle treatments with heated, acidified organic acid salt solutions. *Appl. Environ. Microbiol.* 77:3765–3772.
339. Milillo SR, Ricke SC. 2010. Synergistic reduction of *Salmonella* in a model raw chicken media using a combined thermal and acidified organic acid salt intervention treatment. *J. Food Sci.* 75:M121–M125. doi:10.1111/j.1750-3841.2009.01510.x.
340. Joerger RD, Sartori C, Frye JG, Turpin JB, Schmidt C, McClelland M,

- Porwollik S. 2012. Gene expression analysis of *Salmonella enterica* Enteritidis Nal(R) and *Salmonella enterica* Kentucky 3795 exposed to HCl and acetic acid in rich medium. *Foodborne Pathog. Dis.* 9:331–337.
341. Durant JA, Lowry VK, Nisbet DJ, Stanker LH, Corrier DE, Ricke SC. 1999. Short-chain fatty acids affect cell-association and invasion of HEp-2 cells by *Salmonella typhimurium*. *J. Environ. Sci. Health B* 34: 1083–1099.
342. Durant JA, Lowry VK, Nisbet DJ, Stanker LH, Corrier DE, Ricke SC. 2000. Late logarithmic *Salmonella typhimurium* HEp-2 cell-association and invasion response to short chain volatile fatty acid addition. *J. Food Saf.* 20:1–11.
343. Durant JA, Lowry VK, Nisbet DJ, Stanker LH, Corrier DE, Ricke SC. 2000. Short-chain fatty acids alter HEp-2 cell association and invasion by stationary growth phase *Salmonella typhimurium*. *J. Food Sci.* 65:1206–1209.
344. Van Immerseel F, De Buck J, De Smet I, Pasmans F, Haesebrouck F, Ducatelle R. 2004. Interactions of butyric acid- and acetic acid-treated *Salmonella* with chicken primary cecal epithelial cells in vitro. *Avian Dis.* 48:384–391.
345. Gantois I, Ducatelle R, Pasmans F, Haesebrouck F, Van Immerseel F. 2008. *Salmonella enterica* serovar Enteritidis genes induced during oviduct colonization and egg contamination in laying hens. *Appl. Environ. Microbiol.* 74:6616–6622.
346. Durant JA, Corrier DE, Ricke SC. 2000. Short-chain volatile fatty acids modulate the expression of the *hilA* and *invF* genes of *Salmonella typhimurium*. *J. Food Prot.* 63:573–578.
347. Durant JA, Corrier DE, Stanker LH, Ricke SC. 2000. Expression of the *hilA* *Salmonella typhimurium* gene in a poultry *Salm. enteritidis* isolate in response to lactate and nutrients. *J. Appl. Microbiol.* 89:63–69.
348. Lawhon SD, Maurer R, Suyemoto M, Altier C. 2002. Intestinal short-chain fatty acids alter *Salmonella typhimurium* invasion gene expression and virulence through BarA/SirA. *Mol. Microbiol.* 46:1451–1464.
349. Huang Y, Suyemoto M, Garner CD, Cicconi KM, Altier C. 2008. Formate acts as a diffusible signal to induce *Salmonella* invasion. *J. Bacteriol.* 190:4233–4241.
350. Gonzalez-Gil F, Le Bolloch A, Pendleton S, Zhang N, Wallis A, Hanning I. 2012. Expression of *hilA* in response to mild acid stress in *Salmonella enterica* is serovar and strain dependent. *J. Food Sci.* 77: M292–M297. doi:10.1111/j.1750-3841.2012.02684.x.
351. Sirsat SA, Burkholder KM, Muthaiyan A, Dowd SE, Bhunia AK, Ricke SC. 2011. Effect of sublethal heat stress on *Salmonella Typhimurium* virulence. *J. Appl. Microbiol.* 110:813–822.
352. Hungate RE. 1975. The rumen microbial ecosystem. *Annu. Rev. Ecol. Syst.* 6:39–66.
353. Macfarlane GT, Macfarlane S. 1997. Human colonic microbiota: ecology, physiology and metabolic potential of intestinal bacteria. *Scand. J. Gastroenterol. Suppl.* 222:3–9.
354. Jósefiak D, Rutowski A, Martin SA. 2004. Carbohydrate fermentation in the avian ceca: a review. *Anim. Feed Sci. Technol.* 113:1–15.
355. Burr G, Ricke S, Gatlin D. 2005. Microbial ecology of the gastrointestinal tract of fish and the potential application of prebiotics and probiotics in finfish. *J. World Aquacult. Soc.* 36:425–436.
356. Muegge BD, Kuczynski J, Knights D, Clemente JC, Gonzalez A, Fontana L, Henrissat B, Knight R, Gordon JI. 2011. Diet drives convergence in gut microbiome functions across mammalian phylogeny and within humans. *Science* 332:970–974.
357. Saengerdsu S, Ricke SC. Ecology and characteristics of methanogenic archaea in animals and humans. *Crit. Rev. Microbiol.*, in press.
358. Mead GC. 1989. Microbes of the avian cecum: types present and substrates utilized. *J. Exp. Zool. Suppl.* 3:48–54.
359. Cahill MM. 1990. Bacterial flora of fishes: a review. *Microb. Ecol.* 19:21–41.
360. Wolin MJ, Miller TL. 1983. Interactions of microbial populations in cellulose fermentation. *Fed. Proc.* 42:109–113.
361. Ricke SC, Martin SA, Nisbet DJ. 1996. Ecology, metabolism, and genetics of ruminal selenomonads. *Crit. Rev. Microbiol.* 22:27–56.
362. Mackie RI. 2002. Mutualistic fermentative digestion in the gastrointestinal tract: diversity and evolution. *Integr. Comp. Biol.* 42:319–326.
363. Weimer PJ, Russell JB, Muck RE. 2009. Lessons from the cow: what the ruminant animal can teach us about consolidated bioprocessing of cellulosic biomass. *Bioresour. Technol.* 100:5323–5331.
364. Weimer PJ. 2002. Cellulose degradation by ruminal microorganisms. *Crit. Rev. Biotechnol.* 12:189–223.
365. Hespell RB, Bryant MP. 1979. Efficiency of rumen microbial growth: influence of some theoretical and experimental factors of YATP. *J. Anim. Sci.* 49:1640–1659.
366. Hespell RB. 1979. Efficiency of growth by ruminal bacteria. *Fed. Proc.* 38:2707–2712.
367. Russell JB. 1984. Factors influencing competition and composition of the rumen bacterial flora, p 313–345. *In* Gilchrist FMC, Mackie RI (ed), *Herbivore nutrition in the subtropics and tropics*. The Science Press, Craighall, South Africa.
368. Demeyer D, Van Nevel C. 1986. Influence of substrate and microbial interaction on efficiency of rumen microbial growth. *Reprod. Nutr. Dev.* 26:161–179.
369. Russell JB, Hespell RB. 1981. Microbial rumen fermentation. *J. Dairy Sci.* 64:1153–1169.
370. Coleman ME, Dreesen DW, Wiegert RG. 1996. A simulation of microbial competition in the human colonic ecosystem. *Appl. Environ. Microbiol.* 62:3632–3639.
371. Rohmer L, Hocquet D, Miller SI. 2011. Are pathogenic bacteria just looking for food? Metabolism and microbial pathogenesis. *Trends Microbiol.* 19:341–348.
372. Schmidt H, Hensel M. 2004. Pathogenicity islands in bacterial pathogenesis. *Clin. Microbiol. Rev.* 17:14–56.
373. Tannock GW. 1984. Effect of dietary and environmental stress on the gastrointestinal microbiota, p 517–539. *In* Hentges DJ (ed), *Human intestinal microflora and disease*. Academic Press, Inc, New York, NY.
374. Hashimoto T, Perlot T, Rehman A, Trichereau J, Ishiguro H, Paolino M, Sigl V, Hanada T, Hanada R, Lipinski S, Wild B, Camargo SM, Singer D, Richter A, Kuba K, Fukamizu A, Schreiber S, Clevers H, Verrey F, Rosenstiel P, Penninger JM. 2012. ACE2 links amino acid malnutrition to microbial ecology and intestinal inflammation. *Nature* 487:477–481.
375. Smith MI, Yatsunenko T, Manary MJ, Trehan I, Mkakosya R, Cheng J, Kau AL, Rich SS, Concannon P, Mychaleckyj JC, Liu J, Houpt E, Li JV, Holmes E, Nicholson J, Knights D, Ursell LK, Knight R, Gordon JI. 2013. Gut microbiomes of Malawian twin pairs discordant for kwashiorkor. *Science* 339:548–554.
376. Alverdy J, Stern E. 1998. Effect of immunonutrition on virulence strategies in bacteria. *Nutrition* 14:580–584.
377. Moran ET, Jr, Bilgili SF. 1990. Influence of feeding and fasting broilers prior to marketing on cecal access of orally administered *Salmonella*. *J. Food Prot.* 53:205–207.
378. Humphrey TJ, Baskerville A, Whitehead A, Rowe B, Henley A. 1993. Influence of feeding patterns on the artificial infection of laying hens with *Salmonella enteritidis* phage type 4. *Vet. Rec.* 132:407–409.
379. Ramirez GA, Sarlin LL, Caldwell DJ, Yezak CR, Jr, Hume ME, Corrier DE, Deloach JR, Hargis BM. 1997. Effect of feed withdrawal on the incidence of *Salmonella* in the crops and ceca of market age broiler chickens. *Poult. Sci.* 76:654–656.
380. Corrier DE, Byrd JA, Hargis BM, Hume ME, Bailey RH, Stanker LH. 1999. Survival of *Salmonella* in the crop contents of market-age broilers during feed withdrawal. *Avian Dis.* 43:453–460.
381. Corrier DE, Byrd JA, Hargis BM, Hume ME, Bailey RH, Stanker LH. 1999. Presence of *Salmonella* in the crop and ceca of broiler chickens before and after preslaughter feed withdrawal. *Poult. Sci.* 78:45–49.
382. Miller CP, Bohnhoff M. 1962. A study of experimental *Salmonella* infection in the mouse. *J. Infect. Dis.* 111:107–116.
383. Brownlie LE, Grau FH. 1967. Effect of food intake on growth and survival of salmonellas and *Escherichia coli* in the bovine rumen. *J. Gen. Microbiol.* 46:125–134.
384. Grau FH, Brownlie LE, Roberts EA. 1968. Effect of some preslaughter treatments on the *Salmonella* population in the bovine rumen and faeces. *J. Appl. Bacteriol.* 31:157–163.
385. Grau FH, Brownlie LE, Smith MG. 1969. Effects of food intake on numbers of salmonellae and *Escherichia coli* in rumen and faeces of sheep. *J. Appl. Bacteriol.* 32:112–117.
386. Bell DD. 2003. Historical and current molting practices in the U.S. table egg industry. *Poult. Sci.* 82:965–970.
387. Park SY, Kim WK, Birkhold SG, Kubena LF, Nisbet DJ, Ricke SC. 2004. Induced moulting issues and alternative dietary strategies for the egg industry in the United States. *Worlds Poultry Sci. J.* 60:196–209.
388. Holt PS. 1999. Impact of induced molting on immunity and *Salmonella enterica* serovar Enteritidis infection in laying hens, p 367–375. *In* Saeed AM, Gast RK, Potter ME, Wall PG (ed), *Salmonella enterica* serovar En-

- teritidis in humans and animals—epidemiology, pathogenesis, and control. Iowa State University Press, Ames, IA.
389. Poppe C, McFadden KA, Demczuk WH. 1996. Drug resistance, plasmids, biotypes and susceptibility to bacteriophages of *Salmonella* isolated from poultry in Canada. *Int. J. Food Microbiol.* **30**:325–344.
390. Holt PS. 2003. Molting and *Salmonella enterica* serovar Enteritidis infection: the problem and some solutions. *Poult. Sci.* **82**:1008–1010.
391. Durant JA, Corrier DE, Byrd JA, Stanker LH, Ricke SC. 1999. Feed deprivation affects crop environment and modulates *Salmonella enteritidis* colonization and invasion of leghorn hens. *Appl. Environ. Microbiol.* **65**:1919–1923.
392. Dunkley KD, McReynolds JL, Hume ME, Dunkley CS, Callaway TR, Kubena LF, Nisbet DJ, Ricke SC. 2007. Molting in *Salmonella* Enteritidis-challenged laying hens fed alfalfa crumbles. I. *Salmonella* Enteritidis colonization and virulence gene *hilA* response. *Poult. Sci.* **86**:1633–1639.
393. Corrier DE, Nisbet DJ, Hargis BM, Holt PS, DeLoach JR. 1997. Provision of lactose to molting hens enhances resistance to *Salmonella enteritidis* colonization. *J. Food Prot.* **60**:10–15.
394. Hume ME, Kubena LF, Edrington TS, Donskey CJ, Moore RW, Ricke SC, Nisbet DJ. 2003. Poultry digestive microflora biodiversity as indicated by denaturing gradient gel electrophoresis. *Poult. Sci.* **82**:1100–1107.
395. Ricke SC, Hume ME, Park SY, Moore RW, Birkhold SG, Kubena LF, Nisbet DJ. 2004. Denaturing gradient gel electrophoresis (DGGE) as a rapid method for assessing gastrointestinal tract microflora responses in laying hens fed similar zinc molt induction diets. *J. Rapid Methods Autom. Microbiol.* **12**:69–81.
396. Saengkerdsub S, Kim WK, Andersen RC, Woodward CL, Nisbet DJ, Ricke SC. 2006. Effects of nitrocompounds and feedstuffs on in vitro methane production in chicken cecal contents and rumen fluid. *Anaerobe* **12**:85–92.
397. Dunkley KD, Dunkley CS, Njongmeta NL, Callaway TR, Hume ME, Kubena LF, Nisbet DJ, Ricke SC. 2007. Comparison of in vitro fermentation and molecular microbial profiles of high-fiber feed substrates incubated with chicken cecal inocula. *Poult. Sci.* **86**:801–810.
398. Donalson LM, Kim WK, Chalova VI, Herrera P, McReynolds JL, Gotcheva VG, Vidanovic D, Woodward CL, Kubena LF, Nisbet DJ, Ricke SC. 2008. In vitro fermentation response of laying hen cecal bacteria to combinations of fructooligosaccharide prebiotics with alfalfa or a layer ration. *Poult. Sci.* **87**:1263–1275.
399. Woodward CL, Kwon YM, Kubena LF, Byrd JA, Moore RW, Nisbet DJ, Ricke SC. 2005. Reduction of *Salmonella enterica* serovar Enteritidis colonization and invasion by an alfalfa diet during molt in Leghorn hens. *Poult. Sci.* **84**:185–193.
400. Seo KH, Holt PS, Gast RK. 2001. Comparison of *Salmonella* Enteritidis infection in hens molted via long-term feed withdrawal versus full-fed wheat middling. *J. Food Prot.* **64**:1917–1921.
401. Chiu CH, Chu C, Ou JT. 2000. Lack of evidence of an association between the carriage of virulence plasmid and the bacteremia of *Salmonella typhimurium* in humans. *Microbiol. Immunol.* **44**:741–748.
402. Hurley BP, McCormick BA. 2003. Translating tissue culture results into animal models: the case of *Salmonella typhimurium*. *Trends Microbiol.* **11**:562–569.
403. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**:2731–2739.

Steven L. Foley is a Research Microbiologist with the U.S. Food and Drug Administration's National Center for Toxicological Research. Dr. Foley earned his Ph.D. in Cellular and Molecular Biology from North Dakota State University and completed a postdoctoral fellowship with FDA's Center for Veterinary Medicine. His research program focuses on understanding the role of plasmids in *Salmonella* virulence and antimicrobial resistance and factors that contribute to their spread. He also has an interest in the development and evaluation of molecular subtyping methods for enteric pathogens.



Timothy J. Johnson received his Ph.D. in 2004 from North Dakota State University in Fargo, ND, and his postdoctoral work was performed at the College of Veterinary Medicine at Iowa State University. From 2007 to 2013, he worked as an Assistant Professor of Microbiology within the Department of Veterinary and Biomedical Sciences, College of Veterinary Medicine, University of Minnesota, and is currently an Associate Professor in the same department. Dr. Johnson's research interests involve genomics-based approaches towards understanding the evolution of virulence and antimicrobial resistance in enteric bacteria of production animals and humans and the impact of microbial community succession on animal gut health.



Steven C. Ricke received his B.S. from the University of Illinois and his Ph.D. from the University of Wisconsin and was a USDA-ARS postdoctoral fellow at North Carolina State University. He was a professor at Texas A&M until 2005, when he became the Wray Endowed Chair and Director of the Center for Food Safety at the University of Arkansas. He has received the Poultry Science Association Research Award and American Egg Board Award, was named a Texas Agricultural Experiment Station Faculty Fellow, and received the Division of Agriculture, University of Arkansas, John White Outstanding Research Award. He is a member and served as cofounder and former President of the Arkansas Association of Food Protection. Dr. Ricke's research program is focused primarily on virulence and pathogenic characteristics of food-borne *Salmonella* spp., with emphasis on the growth, survival, and pathogenesis of the organism under conditions encountered during food animal production and processing.



Rajesh Nayak is a Research Scientist with the U.S. Food and Drug Administration and an Adjunct Professor with the Universities of Arkansas and West Virginia. Dr. Nayak earned his Ph.D. at West Virginia University and postdoctoral fellowship at the National Center for Toxicological Research. His research interests are in (i) molecular epidemiology, surveillance, and microbial source tracking of food animal-infectious pathogens through the food production environment using genotyping methods; (ii) threat assessment of emerging pathogens and their genetic characteristics using microarray, whole-genome sequencing, and bioinformatics platforms; (iii) investigating the mechanisms of antimicrobial resistance and pathogenesis in bacterial pathogens isolated from food, feed, environment, human, and veterinary sources; and (iv) developing molecular tools for detection of bacterial pathogens and their genetic traits. Dr. Nayak serves on the editorial board and is an *ad hoc* reviewer for several national and international journals.



Jessica Danzeisen received her M.S. in Microbiology in 2008 from North Dakota State University in Fargo, ND. She is currently a Research Scientist in the laboratory of Dr. Timothy J. Johnson within the Department of Veterinary and Biomedical Sciences, College of Veterinary Medicine, University of Minnesota. Ms. Danzeisen's research interests include antimicrobial resistance in production animals and the gut microbiome related to production animal health.