

Short Communication

Cyclovirus in nasopharyngeal aspirates of Chilean children with respiratory infections

Tung Gia Phan,^{1,2} Vivian Luchsinger,³ Luis F. Avendaño,³ Xutao Deng¹ and Eric Delwart^{1,2}

Correspondence

Eric Delwart
delwarte@medicine.ucsf.edu

¹Blood Systems Research Institute, San Francisco, CA 94118, USA

²Department of Laboratory Medicine, University of California at San Francisco, San Francisco, CA 94118, USA

³Programa de Virología, Facultad de Medicina, Universidad de Chile, Santiago, Chile

Some respiratory tract infections remain unexplained despite extensive testing for common pathogens. Nasopharyngeal aspirates (NPAs) from 120 Chilean infants from Santiago with acute lower respiratory tract infections were analysed by viral metagenomics, revealing the presence of nucleic acids from anelloviruses, adenovirus-associated virus and 12 known respiratory viral pathogens. A single sequence read showed translated protein similarity to cycloviruses. We used inverse PCR to amplify the complete circular ssDNA genome of a novel cyclovirus we named CyCV-ChileNPA1. Closely related variants were detected using PCR in the NPAs of three other affected children that also contained anelloviruses. This report increases the current knowledge of the genetic diversity of cycloviruses whose detection in multiple NPAs may reflect a tropism for human respiratory tissues.

Received 28 October 2013
Accepted 10 January 2014

Cycloviruses, members of a proposed genus within the family *Circoviridae*, have a circular ssDNA genome of approximately 2 kb (Li *et al.*, 2010). Genetically highly diverse cycloviruses were initially found in the faeces of Pakistani children with and without acute flaccid paralysis (Victoria *et al.*, 2009), in wild chimpanzees (Li *et al.*, 2010) and in tissues of farm animals including cows, goats, bats and chickens (Ge *et al.*, 2011; Li *et al.*, 2010, 2011). Unexpectedly, other cyclovirus species have also been detected in insects, namely dragonflies and cockroaches (Dayaram *et al.*, 2013; Padilla-Rodriguez *et al.*, 2013; Rosario *et al.*, 2011). In 2013, a cyclovirus species (CyCV-CN) was found initially using viral metagenomics and then by PCR in 4% of cerebrospinal fluid (CSF) specimens from Vietnamese children with unexplained central nervous system disorder, but not in CSF from patients with non-neurological problems, as well as in 4.2% of faeces from healthy Vietnamese children (Tan *et al.*, 2013). CyCV-CN DNA was also detected in a throat swab (Tan *et al.*, 2013). In this study, 58% of faecal specimens from pigs and poultry in Vietnam were also positive for the same cyclovirus, suggesting possible sources of human infection (Tan *et al.*, 2013). A related cyclovirus was also detected in 10% of CSF samples and 15% of serum samples from adult patients with paraplegia (leg paralysis) from Malawi (Smits *et al.*, 2013).

Nasopharyngeal aspirates (NPAs) from Chilean children less than 2 years old with acute lower respiratory infections were tested for respiratory syncytial virus (RSV), adenovirus, parainfluenza virus 1–3 and influenza A and B viruses by indirect immunofluorescence assays and virus isolation (Avendaño *et al.*, 2003). From 1998 to 2000, a mean of 29% of acute lower respiratory infections samples were positive for RSV (Avendaño *et al.*, 2003). To initiate the characterization of the viruses in non-reactive NPA samples, viral particles were enriched by filtration, and unprotected DNA and RNA were digested using a combination of nuclease enzymes (Victoria *et al.*, 2009). The remaining nucleic acids were then extracted using a MagMAX Viral RNA Isolation kit (Life Technologies), which recovers both RNA and DNA. A DNA library was constructed using a ScriptSeq v2 RNA-Seq Library Preparation kit (Epicentre), which amplifies both RNA and DNA, and sequenced using the Illumina MiSeq platform. Viral sequences were identified using translated protein sequence similarity searches to annotated viral proteins available in GenBank (using BLASTX) and results were mapped using the NCBI Virus Taxonomy browser (<http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?name=Viruses>). The study was approved by the University of California at San Francisco committee on human research.

A total of 120 respiratory specimens from Chilean infants were analysed in 12 pools of 10 specimens using one Illumina MiSeq run of 250 base paired-ends. Viral sequence reads were identified with amino acid similarity >95% to

The GenBank accession numbers for CyCV-ChileNPA1–4 are KF726984–KF726987, respectively.

known viruses. Most numerous sequences were from anelloviruses, followed by enterovirus C, betacoronavirus, bocavirus 1, RSV, human adenovirus 3, enterovirus B, human rhinoviruses A and C, human parainfluenza 3, adeno-associated virus, human pneumovirus, human rhinovirus B and human parechovirus sequences (Table 1). Anelloviruses were identified in all but one pool (Table 1). Anelloviruses have been reported previously in human respiratory secretions (Burián *et al.*, 2011; Jartti *et al.*, 2012). Anelloviruses are usually considered commensal viruses (Okamoto, 2009a), although increased prevalence was found in bronchoalveolar lavage of children with acute exacerbation idiopathic pulmonary fibrosis (Wootton *et al.*, 2011) and acute respiratory diseases (Maggi *et al.*, 2003), and in lung tissues of pigs infected with known respiratory pathogens (Rammohan *et al.*, 2012). Anellovirus plasma load is also increased in advanced AIDS (Li *et al.*, 2013) and in immunosuppressed patients following organ transplantation (De Vlaminc *et al.*, 2013). Increased anellovirus loads may reflect increased replication in immune cells stimulated by chronic inflammation, rather than indicating a direct pathogenic role (De Vlaminc *et al.*, 2013). Detection of RSV, parainfluenza 3 and adenovirus in four, two and one pools, respectively, was probably the result of viral loads being too low for detection by immunofluorescence assays and cell culture (Avendaño *et al.*, 2003). Except for anelloviruses and adeno-associated virus, all other viruses found have been associated with respiratory symptoms.

One sequence from one sample pool showed significant similarity to cyclovirus proteins (BLASTX E-score of 2×10^{-7} to dragonfly cyclovirus, GenBank accession no. KC512919). The full circular cyclovirus genome, referred to as CyCV-ChileNPA1, was then amplified using inverse PCR with specific primers designed from the Illumina-derived short sequence and directly Sanger sequenced by primer walking. Putative ORFs in the cyclovirus genome were predicted using the NCBI ORF Finder (<http://www.ncbi.nlm.nih.gov/gorf/>) requiring an initiation Met codon.

The complete circular genome of CyCV-ChileNPA1 was 1790 bases, with a G + C content of 43 mol%. This genome consisted of two major ORFs encoding replication-associated protein (Rep) and capsid protein (Cap). The intergenic region was 247 bases and encoded a putative stem-loop structure with a stem length of 13 bases, predicted by the Mfold program (Zuker, 2003). Similarly to other cycloviruses, the highly conserved nonamer (TAGTATTAC) was found in the loop (Fig. 1a).

The International Committee on the Taxonomy of Viruses has also proposed a threshold of 75 % nucleotide identity over the entire genome and 70 % amino acid identity for the capsid protein. Rep showed the closest match (65 %) to that of dragonfly CyV-8, whilst Cap shared a lower identity of 30 % with that of the same virus. The higher level of nucleotide divergence of *cap* relative to *rep* was also observed with sequence alignments of

CyCV-ChileNPA1 with its closest relatives (Fig. 1b). Such a high level of sequence divergence indicated that CyCV-ChileNPA1 may be considered a new species within the genus *Cyclovirus*.

CyCV-ChileNPA1 shared conserved Rep motifs (Dayaram *et al.*, 2013; Rosario *et al.*, 2012). Analysis of the deduced amino acid Rep sequences of CyCV-ChileNPA1 revealed three rolling-circle replication motifs I–III: FTxNN (FTWHD), YCSKxGX (YCSKSGE) and HLQGxxNL (HLQGFCSL), respectively (Fig. 2). In the N terminus of cyclovirus Rep, two consensus high-affinity DNA-binding specificity determinants (SPDs), TxR for SPD-region 1 and PxR for SPD-region 2, were present (Dayaram *et al.*, 2013; Londoño *et al.*, 2010). CyCV-ChileNPA1 showed a mutated VxR for SPD-region 1 of unknown functional consequence (Fig. 2). The C-terminal region of the CyCV-ChileNPA1 Rep protein possessed ATP-dependent helicase motifs Walker A, B and C, or GxxGTGKS (GPPGTGKS), VIIDDFYGW and ITSN, respectively (Fig. 2).

Sequence alignment was performed using CLUSTAL_X (Saitou & Nei, 1987). A phylogenetic tree with 100 bootstrap re-samples of the alignment datasets was generated using MEGA5 and the neighbour-joining method (Tamura *et al.*, 2011). Bootstrap values (based on 100 replicates) for each node are given for values >70 % (Fig. 1c). Phylogenetic analysis confirmed the presence of a highly diverse cyclovirus species.

To determine the prevalence of this virus, a nested PCR assay was designed and used to test all 120 NPA samples. Primers ChileNPA-F1 (5'-TGGGTCAGGCTATTACTGGGAG-3') and ChileNPA-R1 (5'-ACTGAATGTCCGTCCGTTGTCC-3') were used for the first round of PCR, and primers ChileNPA-F2 (5'-CAGTGCCATAGTACAGAGTGCCCA-3') and ChileNPA-R2 (5'-CTCCCCTACTCAAA-GAACTCGCCT-3') for the second round of PCR, resulting in an expected amplicon of ~310 bp. The PCR conditions were as follows: denaturation at 95 °C for 5 min, 35 cycles of 95 °C for 30 s, 53 or 55 °C (for the first or second round, respectively) for 30 s and 72 °C for 1 min, a final extension at 72 °C for 10 min, and then held at 4 °C. Amplicons were then sequenced directly for identification. Three additional cases (CyCV-ChileNPA2–4) were positive for the new cyclovirus, yielding a prevalence of 3.3 % in the studied population (4/120). The full genomes of these three viruses were then acquired by overlapping PCR. The genomic sequences of CyCV-ChileNPA2–4 shared a high nucleotide identity of >99 %, showing two, five and six nucleotide mutations compared with the CyCV-ChileNPA1 genome, respectively. All of these point mutations were synonymous except R25S and K93E in the ORF of CyCV-ChileNPA4 Rep.

The four CyCV-ChileNPA PCR-positive samples were then reanalysed using the same metagenomics approach but individually tagged to identify other viruses in these four samples. A total of 2697 unique sequence reads were generated. Three unique CyCV-ChileNPA1 reads were

Table 1. Distribution of sequence reads to different viral types/species in 12 NPA pools from Chile

Virus	NPA pool no. and no. of reads												Total reads
	1	2	3	4	5	6	7	8	9	10	11	12	
	1 246 305	1 381 713	1 561 246	1 775 509	1 136 293	1 383 673	1 327 383	1 139 854	664 165	621 987	1 033 709	1 462 032	
Anelloviruses	247	1115	605		43	81	907	950	575	58	123	645	5347
Enterovirus C	1759	772											2531
Beta coronavirus	1198												1198
Bocavirus 1		326					12	116					454
Respiratory syncytial virus		10					13	13		119		4	159
Human adenovirus 3							93						93
Enterovirus B	85												85
Human rhinovirus A					5	10				11		11	37
Human rhinovirus C	8					2					2		12
Human parainfluenza 3	2	8											10
Adeno-associated virus									10				10
Human metapneumovirus									6		2		8
Human rhinovirus B								4					4
Human parechovirus		2								2			4
Cyclovirus							1						1




Fig. 1. Details of the novel cyclovirus CyCV-ChileNPA1. (a) Genome organization and its stem-loop structure. The locations of the putative *rep* and *cap* genes are indicated by arrows. (b) Pairwise sequence alignments of CyCV-ChileNPA1 with its closest relatives. The sequence nucleotide similarity (%) is indicated by the height of each point along the *y*-axis. The *x*-axis shows the nucleotide positions in the complete genome. (c) Phylogenetic trees generated with Rep and Cap proteins (concatenated) of cycloviruses. Bars, amino acid substitutions per position.

generated from the NPA sample in which it was originally detected. No other viral sequences were detected. A total of 6362 unique reads were also generated from the other three samples positive only by PCR for the cyclovirus. All three

samples contained anellovirus sequences (a total of 456 reads) and no other close matches to mammalian viruses. Anelloviruses are highly prevalent viruses present in many anatomical sites of different mammals (Okamoto, 2009b)




Fig. 2. Alignment of Rep proteins of the newly identified CyCV-ChileNPA1 and dragonfly cyclovirus-8. Conserved motifs are shown within boxes. SPD-r, specificity determinant region. White boxes show RCR and shaded boxes show helicase motifs.

and are generally considered commensal infections. Anelloviruses have also been found in a significant minority of cases of idiopathic pulmonary fibrosis and in cases of acute lung injury (Wootton *et al.*, 2011), are at higher prevalence in plasma and nasopharyngeal samples of febrile versus non-febrile cases (McElvania TeKippe *et al.*, 2012) and are generally increased in the plasma of immunosuppressed individuals such as advanced AIDS patients (Li *et al.*, 2013) or transplant recipients (De Vlaminc *et al.*, 2013). A porcine anellovirus (torque teno sus virus species 1) has also been associated with porcine respiratory disease complex where it might exacerbate infections caused by porcine circovirus 2 and the arterivirus porcine reproductive and respiratory disease symptom virus (Rammohan *et al.*, 2012).

The detection of cyclovirus DNA in different human samples, including faeces, blood and CSF, and in the muscle tissues of farm animals suggests that cycloviruses may cause systemic infections in mammals (Li *et al.*, 2010; Smits *et al.*, 2013; Tan *et al.*, 2013). The detection of cyclovirus DNA in NPAs (upper respiratory tract) of children with lower tract respiratory problems raises the possibility of a role for these viruses in respiratory illnesses. Further investigations of the host and tissue tropism, the transmission route(s) and any physiological consequences of human cyclovirus infections and possible interactions with anelloviruses are required.

Acknowledgements

We acknowledge NHLBI grant R01 HL105770 to E.L.D. and the Blood Systems Research Institute for sustained support.

References

Avedaño, L. F., Palomino, M. A. & Larrañaga, C. (2003). Surveillance for respiratory syncytial virus in infants hospitalized for acute lower

respiratory infection in Chile (1989 to 2000). *J Clin Microbiol* **41**, 4879–4882.

Burián, Z., Szabó, H., Székely, G., Gyurkovits, K., Pankovics, P., Farkas, T. & Reuter, G. (2011). Detection and follow-up of torque teno midi virus ('small anelloviruses') in nasopharyngeal aspirates and three other human body fluids in children. *Arch Virol* **156**, 1537–1541.

Dayaram, A., Potter, K. A., Moline, A. B., Rosenstein, D. D., Marinov, M., Thomas, J. E., Breitbart, M., Rosario, K., Argüello-Astorga, G. R. & Varsani, A. (2013). High global diversity of cycloviruses amongst dragonflies. *J Gen Virol* **94**, 1827–1840.

De Vlaminc, I., Khush, K. K., Strehl, C., Kohli, B., Luikart, H., Neff, N. F., Okamoto, J., Snyder, T. M., Cornfield, D. N. & other authors (2013). Temporal response of the human virome to immunosuppression and antiviral therapy. *Cell* **155**, 1178–1187.

Ge, X., Li, J., Peng, C., Wu, L., Yang, X., Wu, Y., Zhang, Y. & Shi, Z. (2011). Genetic diversity of novel circular ssDNA viruses in bats in China. *J Gen Virol* **92**, 2646–2653.

Jartti, T., Jartti, L., Ruuskanen, O. & Söderlund-Venermo, M. (2012). New respiratory viral infections. *Curr Opin Pulm Med* **18**, 271–278.

Li, L., Kapoor, A., Slikas, B., Bamidele, O. S., Wang, C., Shaukat, S., Masroor, M. A., Wilson, M. L., Ndjanga, J. B. & other authors (2010). Multiple diverse circoviruses infect farm animals and are commonly found in human and chimpanzee feces. *J Virol* **84**, 1674–1682.

Li, L., Shan, T., Soji, O. B., Alam, M. M., Kunz, T. H., Zaidi, S. Z. & Delwart, E. (2011). Possible cross-species transmission of circoviruses and cycloviruses among farm animals. *J Gen Virol* **92**, 768–772.

Li, L., Deng, X., Linsuwanon, P., Bangsberg, D., Bwana, M. B., Hunt, P., Martin, J. N., Deeks, S. G. & Delwart, E. (2013). AIDS alters the commensal plasma virome. *J Virol* **87**, 10912–10915.

Londoño, A., Riego-Ruiz, L. & Argüello-Astorga, G. R. (2010). DNA-binding specificity determinants of replication proteins encoded by eukaryotic ssDNA viruses are adjacent to widely separated RCR conserved motifs. *Arch Virol* **155**, 1033–1046.

Maggi, F., Pifferi, M., Tempestini, E., Fornai, C., Lanini, L., Andreoli, E., Vatteroni, M., Presciuttini, S., Pietrobelli, A. & other authors (2003). TT virus loads and lymphocyte subpopulations in children with acute respiratory diseases. *J Virol* **77**, 9081–9083.

McElvania TeKippe, E., Wylie, K. M., Deych, E., Sodergren, E., Weinstock, G. & Storch, G. A. (2012). Increased prevalence of anellovirus in pediatric patients with fever. *PLoS ONE* **7**, e50937.

- Okamoto, H. (2009a).** History of discoveries and pathogenicity of TT viruses. *Curr Top Microbiol Immunol* **331**, 1–20.
- Okamoto, H. (2009b).** TT viruses in animals. *Curr Top Microbiol Immunol* **331**, 35–52.
- Padilla-Rodriguez, M., Rosario, K. & Breitbart, M. (2013).** Novel cyclovirus discovered in the Florida woods cockroach *Eurycotis floridana* (Walker). *Arch Virol* **158**, 1389–1392.
- Rammohan, L., Xue, L., Wang, C., Chittick, W., Ganesan, S. & Ramamoorthy, S. (2012).** Increased prevalence of torque teno viruses in porcine respiratory disease complex affected pigs. *Vet Microbiol* **157**, 61–68.
- Rosario, K., Marinov, M., Stainton, D., Kraberger, S., Wiltshire, E. J., Collings, D. A., Walters, M., Martin, D. P., Breitbart, M. & Varsani, A. (2011).** Dragonfly cyclovirus, a novel single-stranded DNA virus discovered in dragonflies (Odonata: Anisoptera). *J Gen Virol* **92**, 1302–1308.
- Rosario, K., Duffy, S. & Breitbart, M. (2012).** A field guide to eukaryotic circular single-stranded DNA viruses: insights gained from metagenomics. *Arch Virol* **157**, 1851–1871.
- Saitou, N. & Nei, M. (1987).** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.
- Smits, S. L., Zijlstra, E. E., van Hellemond, J. J., Schapendonk, C. M., Bodewes, R., Schürch, A. C., Haagmans, B. L. & Osterhaus, A. D. (2013).** Novel cyclovirus in human cerebrospinal fluid, Malawi, 2010–2011. *Emerg Infect Dis* [Internet] **19**.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011).** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731–2739.
- Tan, L. V., van Doorn, H. R., Nghia, H. D., Chau, T. T., Tu, L. T., de Vries, M., Canuti, M., Deijs, M., Jebbink, M. F. & other authors (2013).** Identification of a new cyclovirus in cerebrospinal fluid of patients with acute central nervous system infections. *mBio* **4**, e00231-13.
- Victoria, J. G., Kapoor, A., Li, L., Blinkova, O., Slikas, B., Wang, C., Naeem, A., Zaidi, S. & Delwart, E. (2009).** Metagenomic analyses of viruses in stool samples from children with acute flaccid paralysis. *J Virol* **83**, 4642–4651.
- Wootton, S. C., Kim, D. S., Kondoh, Y., Chen, E., Lee, J. S., Song, J. W., Huh, J. W., Taniguchi, H., Chiu, C. & other authors (2011).** Viral infection in acute exacerbation of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* **183**, 1698–1702.
- Zuker, M. (2003).** Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* **31**, 3406–3415.