

NANO EXPRESS

Open Access

Development of an aptamer-conjugated fluorescent nanoprobe for MMP2

Myoung-Eun Han^{1,2}, Sungmin Baek^{1,2}, Hyun-Jung Kim¹, Jung Hwan Lee³, Sung-Ho Ryu³ and Sae-Ock Oh^{1,2*}

Abstract

Matrix metalloproteinase 2 (MMP2) plays critical roles in various diseases, such as atherosclerosis and cancer, and has been suggested to contribute to the instability of atherosclerotic plaque. To visualize MMP2 in pathologic tissues, we developed an aptamer targeting MMP2 protein by performing eight rounds of modified DNA systematic evolution of ligands by exponential enrichment (SELEX). The aptamer showed high affinity for MMP2 ($K_d = 5.59 \text{ nM}$), precipitated MMP2, and detected MMP2 protein in pathological tissues such as atherosclerotic plaque and gastric cancer tissues. Furthermore, a MMP2 aptamer-conjugated fluorescent nanoprobe successfully visualized atherosclerotic plaques in apolipoprotein E (ApoE) knockout mice. These results suggest that the devised MMP2 aptamer could be useful for the development of various diagnostic tools.

Keywords: MMP2; Aptamer; Nanoprobe; Atherosclerosis; Cancer

Background

Matrix metalloproteinases (MMPs) are zinc- and calcium-dependent proteolytic enzymes [1,2]. MMPs can digest extracellular matrix proteins, such as collagen and fibronectin, and many other proteins, such as proteinases, growth factors, cytokines, chemokines, and cell receptors and thus regulate their activities. MMP was first identified in 1962 [3], and since then, other MMPs have been identified. Interestingly, whereas many MMPs are secreted by cells, others are anchored on cellular membranes. Members of this family play important roles in various cellular processes, such as migration, differentiation, and proliferation. Furthermore, they have been associated with pathophysiologies of various diseases, such as cancer, atherosclerosis, and arthritis.

During the progression of atherosclerosis, inflammatory cells such as monocytes and lymphocytes [4] play critical roles. Monocytes are recruited into atherosclerotic sites and differentiate into macrophages. After excessive lipid uptake, they become foamy cells. Notably, plaque macrophages secrete critical molecules such as MMPs

and prothrombotic tissue factor. Then, MMPs destabilize atherosclerotic plaque by degrading extracellular matrix [5,6]. In addition to the roles in atherosclerosis, MMPs can aid the metastasis of cancer cells [2,7].

Information about the stability of atherosclerotic plaque is critical for the stratification and management of patients [8], and unfortunately, anatomical imaging modalities, such as CT or MRI, do not provide this type of information. Because MMPs are associated with the stability of atherosclerotic plaque, their visualization will be helpful in the stratification and management of patients.

The above mentioned actions of MMPs encouraged us to develop a molecular imaging modality that specifically targets MMP molecules. As compared with antibodies, aptamers have several beneficial characteristics, such as low immunogenicity, low molecular weight (8 to 15 kDa), high stability, better penetration, high affinity, and ease of production [9]. From these reasons, we decided to develop a MMP2-specific aptamer. By performing modified DNA systematic evolution of ligands by exponential enrichment (SELEX), we successfully developed a MMP2-specific aptamer which had high affinity and specificity and showed the possibility that it can be applied for molecular imaging.

* Correspondence: hedgehog@pusan.ac.kr

¹Department of Anatomy, School of Medicine, Pusan National University, 49 Busandaehak-ro, Mugeum-eup, Yangsan 626-870, Republic of Korea

²Medical Research Center for Ischemic Tissue Regeneration, Pusan National University, 49 Busandaehak-ro, Mugeum-eup, Yangsan 626-870, Republic of Korea

Full list of author information is available at the end of the article




Figure 1 Sequence and secondary structure of the MMP2 aptamer. (a) Sequence of the 40-nucleotide random region (N40, shaded) and of the two constant regions flanking the random region. (b) The hairpin-like secondary structure of the aptamer is presented in the lower panel.

Methods

In vitro selection of MMP2 DNA aptamers

To select MMP2-specific aptamers, a modified DNA SELEX procedure was used, as previously described [10]. Briefly, an ssDNA library template consisting of a 40-nucleotide random region (N40) flanked by two constant regions was prepared and immobilized on streptavidin-coated beads (Pierce, Rockland, MA, USA) via its 5'-OH-end biotin. A primer extension was then performed using the dATP, dCTP, dGTP, and benzyl-dUTP nucleotides. The modified DNA library was detached from the template under high pH conditions

and then incubated with biotin-tagged target, partitioned using Dynabeads MyOne (Invitrogen, Carlsbad, CA, USA) and amplified by conventional PCR using a 5'-OH terminal biotinylated reverse primer. A primer extension was then performed, and an enriched pool was prepared for the next round. After eight rounds of SELEX, the enriched DNA pool was cloned and sequenced using standard procedures. After each round of SELEX, binding assays were performed to measure the dissociation constant (K_d) value of the aptamer pool to ensure that its K_d value exhibited a decreasing trend.




Figure 2 Affinity of the MMP2 aptamer. (a) ^{32}P -labeled aptamers and different MMP2 protein concentrations were used to examine the binding affinity of the MMP2 aptamer. (b) Images of radiolabeled aptamer that interacted with proteins in the binding assay.




Figure 3 Precipitation of MMP2 protein by MMP2 aptamer.
MMP2 protein in buffer containing 10% serum was incubated with the aptamer (0.2 μ g/ml) overnight at 4°C. The protein was detected by immunoblotting with anti-MMP2 antibody.

for 30 min at 37°C. In order to capture MMP-2, the solution was incubated with Zorbax silica beads (Agilent, Santa Clara, CA, USA) for 1 min with shaking. The protein bead complex was then partitioned through nitrocellulose filter plates (Millipore, Billerica, MA, USA), which were then washed in buffer and exposed to photographic film. Amounts of radiolabeled aptamer that interacted with proteins were quantified using a Fuji FLA-5000 Image Analyzer (Tokyo, Japan). Dissociation constants were calculated by plotting bound MMP2 aptamer versus protein concentration using the following equation: $Y = B_{\max}X/(K_d + X)$, where B_{\max} is the extrapolated maximal amount of bound aptamer/protein complex.

Precipitation of MMP2 protein by the aptamer and western blotting

A protein solution containing MMP2 protein (R&D Systems, Minneapolis, MN, USA) was precleared with streptavidin-coupled beads for 2 h. The solution was then precipitated with biotin-labeled MMP2 or control aptamer at 4°C overnight. Beads were washed four times with 1 ml of wash buffer (200 mM Tris at pH 8.0, 100 mM NaCl and 0.5% NP-40), once with ice-cold




Figure 4 Comparison of the tissue staining abilities of anti-MMP2 antibody and MMP2 aptamer. Normal aorta, atherosclerotic plaques, and gastric cancer tissues were stained with anti-MMP2 antibody and MMP2 aptamer. Similar staining patterns were observed. Scale bar, 100 μ m.




Figure 5 Specificity of the aptamer by immunohistochemical staining. After incubating the MMP2 aptamer with MMP2 protein in PBS at room temperature for 2 h, the immunohistochemical staining in gastric cancer tissues was significantly reduced. Scale bar, 100 μ m.

phosphate buffered saline (PBS), and boiled in 2 \times loading buffer. Finally, proteins were resolved by SDS-PAGE before being probed with MMP2 antibody (AB37150, Abcam, Cambridge, England, UK).

Immunohistochemistry

Tissue samples were embedded in OCT (Sakura, Japan), and 4- μ m thin sections prepared using a cryostat (CM3050S, Leica, Wetzlar, Germany) were placed on poly-lysine-coated microscope slides. The sections were then treated with 0.3% hydrogen peroxide for 30 min to

quench endogenous peroxidase activity. Blocking was performed using 10% normal donkey serum (NDS) in 1 \times PBS. MMP2 aptamer or anti-MMP2 antibody (AB37150, Abcam, Cambridge, England, UK) binding was performed at a dilution of 1:200 in blocking buffer overnight at 4°C, and secondary antibody (horseradish peroxidase-conjugated anti-rabbit, 1:5,000) binding was performed for 2 h at RT. The signal was detected with HRP (Jackson ImmunoResearch Laboratories, West Grove, PA, USA) using the DAB substrate kit (Vector Laboratories, Burlingame, CA, USA). Sections were then counterstained




Figure 6 Construction of the MMP2 aptamer-conjugated fluorescent nanoprobe. The MMP2 aptamer was conjugated into magnetic fluorescent nanoprobe using EDC.




Figure 7 *Ex vivo* imaging of atherosclerotic plaques using the MMP2 aptamer-conjugated fluorescent nanoprobe. Atherosclerotic plaques were induced by feeding ApoE knockout mice a high cholesterol diet for 4 months and were confirmed by oilred O staining (middle panels). *Ex vivo* imaging was performed 2 h after intravenously injecting mice with the MMP2 aptamer-conjugated fluorescent nanoprobe. The MMP2 aptamer (right panels) showed much stronger signals in atherosclerotic plaques than the control aptamer (left panels).

with hematoxylin, dehydrated, and mounted. The primary antibody was omitted from negative control.

Construction of an aptamer-conjugated nanoprobe

An aptamer-conjugated nanoprobe was produced as previously described [11]. MNP@SiO₂(RITC)-(PEG)/COOH/pro-NH₂ nanoparticles (MF nanoparticles, 2 mg/mL) were purchased from Biterials (Seoul, Korea). The carboxyl moieties (1.1×10^4 /nanoparticle) of MF nanoparticles (size, approximately 50 nm; hydrodynamic diameter, 58.1 nm) were covalently linked to a 5'-NH₂-modified MMP2 aptamer using N-(3-dimethylaminopropyl)-N-ethylcarbodiimide (EDC) (Sigma, St. Louis, MO, USA). After 1 h of incubation, the aptamer-conjugated nanoprobe was washed twice with Tris buffer (pH 7.4) and briefly sonicated.

Animal experiments and *ex vivo* imaging

To induce atherosclerosis in mice, apolipoprotein E (ApoE) knockout mice (Jackson Lab, Bar Harbor, ME, USA) were fed with a high cholesterol diet for 16 weeks from 8 weeks of age. All mice were housed under specific pathogen-free conditions in box cages at $23^\circ\text{C} \pm 2^\circ\text{C}$ and $60\% \pm 10\%$ humidity under a 12-hlight/12-h dark cycle with free access to food and water. Mice were sacrificed at week 16 of the experimental period. All animal procedures were performed in compliance with the Institute of Laboratory Animal Research Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee of Pusan National University. Atherosclerotic plaques were visualized by oil red O staining (Sigma). Aortas were removed 2 h after intravenously injecting MMP2 aptamer-conjugated fluorescent nanoprobe. Fluorescence from aortas was observed with Optix MX3/Optical Molecular Imaging System (ART, Montreal, Canada).

Results and discussion

To develop a specific aptamer for MMP2 protein, we performed a modified DNA SELEX technique as described in the 'Methods' section. To select a high-affinity aptamer, we used nucleotides chemically modified by benzylaminocarbonyl-dU (Benzyl-dU) at the 5' positions, which mimic amino acid side chains. After eight

rounds of SELEX, the enriched DNA pool was cloned and sequenced according to standard procedures. After each round of SELEX, binding assays were performed to measure the dissociation constant (K_d) value of the aptamer pool using [α -32P] ATP. The sequence and secondary structure of the best aptamer selected in this study were presented in Figure 1. The mean B_{max} and K_d values of the aptamer were $35\% \pm 0.8\%$ and 5.59 ± 0.52 nM, respectively (Figure 2).

To determine whether the MMP2 aptamer could be used to precipitate the target protein, we performed precipitation and then western blotting using anti-MMP2 antibody. To do this, we biotinylated the aptamer and used streptavidin beads for the precipitation. MMP2 in buffer containing 10% serum was incubated with the biotinylated MMP2 aptamer, and the protein-aptamer complex was then precipitated and detected by immunoblotting. The aptamer successfully precipitated MMP2 protein (Figure 3), whereas the biotinylated control aptamer did not (data not shown).

Next, we examined whether the MMP2 aptamer could be applied for immunohistochemical purposes in pathological tissues, that is, atherosclerotic plaques and gastric cancer tissues. In both tissue types, the MMP2 aptamer successfully detected MMP2 (Figure 4), whereas the control aptamer did not (data not shown). To further confirm the specificity of the aptamer for immunohistochemistry, we performed peptide blocking. Immunohistochemistry was performed after incubating the aptamer for 2 h with the bare protein, and the intensities of positive signals were significantly reduced (Figure 5).

Finally, we used the aptamer for *ex vivo* imaging. To do this, the aptamer was conjugated to fluorescent nanoprobe using EDC (Figure 6). To induce atherosclerosis in mice, ApoE knockout mice were fed a high cholesterol diet for 4 months. After injecting the aptamer-conjugated fluorescent nanoprobe into a tail vein, fluorescent signals from atherosclerotic plaques were observed. The presence of atherosclerotic plaques was confirmed by oilred O staining. The MMP2 aptamer-conjugated nanoprobe produced significantly stronger signals in atherosclerotic plaques than the control aptamer-conjugated probe (Figure 7).

Many studies have tried to visualize MMP molecules. Small molecular MMP inhibitors attached to radioisotopes, such as ^{123}I , ^{99m}TC , and ^{18}F have been used for the imaging of atherosclerotic lesions and myocardial infarctions [12-15]. Notably, a peptide substrate, which fluoresces when cleaved by MMPs, was used to visualize MMP activity [16-18]. However, considerable time is required for *in vivo* imaging using this peptide substrate. We considered that aptamers could overcome this problem because aptamers bind directly to target proteins. In addition, due to its small size and easy chemical

modification, it can be easily applied to construct new nanoparticles as presented in this study ([9], Figure 6).

The specificity of the MMP2 aptamer produced during the present study was confirmed *in vitro* and *ex vivo*. Precipitation and immunohistochemistry studies demonstrated specific protein binding by MMP2 aptamer, and in particular, immunohistochemical staining of MMP2 aptamer was blocked by MMP2 protein. Furthermore, *ex vivo* imaging demonstrated that whereas MMP2 aptamer visualized atherosclerotic plaques, control aptamer did not. These results suggest that the devised MMP2 aptamer has clinical merit.

Conclusions

We developed an aptamer targeting MMP2 protein using a modified DNA SELEX technique. The devised MMP2 aptamer precipitated and detected MMP2 protein in pathological tissues, that is, atherosclerotic plaques and gastric cancer tissues. Furthermore, the MMP2 aptamer-conjugated fluorescent nanoprobe allowed the visualization of atherosclerotic plaques in ApoE knock-out mice. These results indicate that the developed MMP2 aptamer provides a suitable basis for the development of diagnostic tools.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

ME carried out conjugation of the aptamer into the fluorescent nanoprobe and all animal experiments and drafted the manuscript. SM carried out immunohistochemistry. HJ carried out western blotting and immunohistochemistry. JH and SH carried out SELEX. SO conceived of the study, participated in its design and coordination, and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by the Medical Research Center Program (NRF-2010-0005930) and a grant from the National R&D Program for Cancer Control, Ministry for Health, Welfare and Family Affairs, Republic of Korea (0920050) and Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology (2012R1A1A3010521). Dr. Han ME was financially supported by the 2011 Post-Doc Development Program of Pusan National University.

Author details

¹Department of Anatomy, School of Medicine, Pusan National University, 49 Busandaehak-ro, Mugeum-eup, Yangsan 626-870, Republic of Korea.

²Medical Research Center for Ischemic Tissue Regeneration, Pusan National University, 49 Busandaehak-ro, Mugeum-eup, Yangsan 626-870, Republic of Korea.

³Medical POSTECH Aptamer Initiative Program, Division of Integrative Bioscience and Biotechnology, Pohang University of Science and Technology, Pohang 790-784, Republic of Korea.

Received: 20 November 2013 Accepted: 24 February 2014

Published: 3 March 2014

References

- Iyer RP, Patterson NL, Fields GB, Lindsey ML: The history of matrix metalloproteinases: milestones, myths, and misperceptions. *Am J Physiol* 2012, **303**:H919–H930.
- Hadler-Olsen E, Winberg JO, Uhlin-Hansen L: Matrix metalloproteinases in cancer: their value as diagnostic and prognostic markers and therapeutic targets. *Tumour Biol* 2013, **34**:2041–2051.

3. Woessner JF Jr: Catabolism of collagen and non-collagen protein in the rat uterus during post-partum involution. *Biochem J* 1962, **83**:304–314.
4. Cybulsky MI, Gimbrone MA Jr: Endothelial expression of a mononuclear leukocyte adhesion molecule during atherogenesis. *Sci (New York, NY)* 1991, **251**:788–791.
5. Jones CB, Sane DC, Herrington DM: Matrix metalloproteinases: a review of their structure and role in acute coronary syndrome. *Cardiovasc Res* 2003, **59**:812–823.
6. Garcia-Touchard A, Henry TD, Sangiorgi G, Spagnoli LG, Mauriello A, Conover C, Schwartz RS: Extracellular proteases in atherosclerosis and restenosis. *Arterioscler Thromb Vasc Biol* 2005, **25**:1119–1127.
7. Itoh T, Tanioka M, Matsuda H, Nishimoto H, Yoshioka T, Suzuki R, Uehira M: Experimental metastasis is suppressed in MMP-9-deficient mice. *Clin Exp Metastasis* 1999, **17**:177–181.
8. Shaw SY: Molecular imaging in cardiovascular disease: targets and opportunities. *Nat Rev* 2009, **6**:569–579.
9. Praseetha PK, Thampy AS, Venugopalan P, Chavali MS: Molecular nanobiotechnological approaches for the detection and therapy of prion related diseases. *Nano Biomed Eng* 2012, **4**:50–57.
10. Lee YJ, Kim IS, Park SA, Kim Y, Lee JE, Noh DY, Kim KT, Ryu SH, Suh PG: Periostin-binding DNA aptamer inhibits breast cancer growth and metastasis. *Mol Ther* 2013, **21**:1004–1013.
11. Do Hwang W, Ko HY, Lee JH, Kang H, Ryu SH, Song IC, Lee DS, Kim S: A nucleolin-targeted multimodal nanoparticle imaging probe for tracking cancer cells using an aptamer. *J Nucl Med* 2010, **51**:98–105.
12. Schafers M, Riemann B, Kopka K, Breyholz HJ, Wagner S, Schafers KP, Law MP, Schober O, Levkau B: Scintigraphic imaging of matrix metalloproteinase activity in the arterial wall in vivo. *Circulation* 2004, **109**:2554–2559.
13. Su H, Spinale FG, Dobrucki LW, Song J, Hua J, Sweterlitsch S, Dione DP, Cavaliere P, Chow C, Bourke BN, Hu XY, Azure M, Yalamanchili P, Liu R, Cheesman EH, Robinson S, Edwards DS, Sinusas AJ: Noninvasive targeted imaging of matrix metalloproteinase activation in a murine model of postinfarction remodeling. *Circulation* 2005, **112**:3157–3167.
14. Breyholz HJ, Wagner S, Levkau B, Schober O, Schafers M, Kopka K: A 18 F-radiolabeled analogue of CGS 27023A as a potential agent for assessment of matrix-metalloproteinase activity in vivo. *Q J Nucl Med Mol Imaging* 2007, **51**:24–32.
15. Lancelot E, Amirbekian V, Brigger I, Raynaud JS, Ballet S, David C, Rousseaux O, Le Greneur S, Port M, Lijnen HR, Bruneval P, Michel JB, Ouimet T, Roques B, Amirbekian S, Hyafil F, Vucic E, Aguinaldo JG, Corot C, Fayad ZA: Evaluation of matrix metalloproteinases in atherosclerosis using a novel noninvasive imaging approach. *Arterioscler Thromb Vasc Biol* 2008, **28**:425–432.
16. Chen J, Tung CH, Allport JR, Chen S, Weissleder R, Huang PL: Near-infrared fluorescent imaging of matrix metalloproteinase activity after myocardial infarction. *Circulation* 2005, **111**:1800–1805.
17. Nahrendorf M, Swirski FK, Aikawa E, Stangenberg L, Wurdinger T, Figueiredo JL, Libby P, Weissleder R, Pittet MJ: The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. *J Exp Med* 2007, **204**:3037–3047.
18. Deguchi JO, Aikawa M, Tung CH, Aikawa E, Kim DE, Ntziachristos V, Weissleder R, Libby P: Inflammation in atherosclerosis: visualizing matrix metalloproteinase action in macrophages in vivo. *Circulation* 2006, **114**:55–62.

doi:10.1186/1556-276X-9-104

Cite this article as: Han et al.: Development of an aptamer-conjugated fluorescent nanoprobe for MMP2. *Nanoscale Research Letters* 2014 **9**:104.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com