

REVIEW

Lipid analogues as potential drugs for the regulation of mitochondrial cell death

Michael Murray¹, Herryawan Ryadi Eziwar Dyari¹, Sarah E Allison¹ and Tristan Rawling²

¹Pharmacogenomics and Drug Development Group, Discipline of Pharmacology, University of Sydney, Sydney, NSW, Australia, and ²School of Pharmacy, Graduate School of Health, University of Technology, Sydney, NSW, Australia

Correspondence

Dr Michael Murray,
Pharmacogenomics and Drug
Development Group, Discipline
of Pharmacology, Medical
Foundation Building, Room 105,
University of Sydney, Sydney,
NSW 2006, Australia. E-mail:
michael.murray@sydney.edu.au

Keywords

N-acyl ethanolamines; cancer cell; ether phospholipids; fatty acid biotransformation; free fatty acids; intrinsic pathway of apoptosis; mitochondrial ATP production; polyunsaturated fatty acid epoxides; reactive oxygen species

Received

29 June 2013

Revised

22 August 2013

Accepted

26 August 2013

The mitochondrion plays an important role in the production of energy as ATP, the regulation of cell viability and apoptosis, and the biosynthesis of major structural and regulatory molecules, such as lipids. During ATP production, reactive oxygen species are generated that alter the intracellular redox state and activate apoptosis. Mitochondrial dysfunction is a well-recognized component of the pathogenesis of diseases such as cancer. Understanding mitochondrial function, and how this is dysregulated in disease, offers the opportunity for the development of drug molecules to specifically target such defects. Altered energy metabolism in cancer, in which ATP production occurs largely by glycolysis, rather than by oxidative phosphorylation, is attributable in part to the up-regulation of cell survival signalling cascades. These pathways also regulate the balance between pro- and anti-apoptotic factors that may determine the rate of cell death and proliferation. A number of anti-cancer drugs have been developed that target these factors and one of the most promising groups of agents in this regard are the lipid-based molecules that act directly or indirectly at the mitochondrion. These molecules have emerged in part from an understanding of the mitochondrial actions of naturally occurring fatty acids. Some of these agents have already entered clinical trials because they specifically target known mitochondrial defects in the cancer cell.

LINKED ARTICLES

This article is part of a themed issue on Mitochondrial Pharmacology: Energy, Injury & Beyond. To view the other articles in this issue visit <http://dx.doi.org/10.1111/bph.2014.171.issue-8>

Abbreviations

Caspase, cysteine-aspartic protease; CLA, conjugated linoleic acid; CYP, cytochrome P450; EET, epoxyeicosatrienoic acid; HETE, hydroxyeicosatetraenoic acid; LOX, lipoxygenase; PARP, poly (ADP ribose) polymerase; PUFA, polyunsaturated fatty acid; ROS, reactive oxygen species; TTA, tetradecylthioacetic acid

Mitochondrial function: introduction

The mitochondrion has a number of critical homeostatic functions, including the production of energy as ATP, the control of apoptotic cell death and viability, and the biosynthesis of molecules such as steroids and lipids that perform regulatory and structural roles in cells (Kroemer *et al.*, 2007). Mitochondria are also the major source of reactive oxygen

species (ROS) that determine the intracellular redox state and modulate cell proliferation and apoptosis. It is increasingly recognized that disease processes, such as tumourigenesis and metabolic syndrome, are associated with mitochondrial dysfunction (Kroemer *et al.*, 2007). A detailed understanding of how the regulation of mitochondrial function is altered during cancer progression may provide opportunities for drug design strategies that target underlying defects in disease.

Mitochondrial energy production and altered ATP generation in cancer cells

The role of the mitochondrion in energy metabolism is well established and glucose is the primary fuel molecule utilized by the cell in ATP production (Kroemer *et al.*, 2007). The initial glycolytic step occurs in the cytoplasm of the cell and generates pyruvate that enters the mitochondrion where it is converted to citrate; together, these two reactions produce four molecules of ATP from each molecule of glucose. In addition, four large multi-protein respiratory complexes in the inner mitochondrial membrane work in concert to generate a much larger number of ATPs by oxidative phosphorylation (~28–32 per glucose molecule). Simultaneously, this builds up the proton gradient across the inner and outer mitochondrial membranes, which drives many mitochondrial processes.

Fats and proteins may also be utilized by the mitochondrion to produce ATP (Kroemer *et al.*, 2007). Triglyceride esters in adipose tissue are hydrolysed to free fatty acids that undergo mitochondrial β -oxidation to acetyl-CoA units that are then able to enter the citric acid cycle and generate ATP. The oxidative deamination of amino acids produces up to 15% of total metabolic energy in animals. In the initial phase, amino acids are converted to a series of intermediary molecules – pyruvate, α -ketoglutarate, succinyl-CoA, fumarate, oxaloacetate, acetyl-CoA and acetoacetate – that may then enter the citric acid cycle and generate ATP.

Whereas normal cells generate much of their ATP by oxidative phosphorylation, aggressive cancer cells exhibit pronounced bioenergetic differences and overproduce lactate even under normoxic conditions by 'aerobic glycolysis' (Rossignol *et al.*, 2004). This occurs because most of the pyruvate formed by glycolysis is unable to enter the mitochondrion, and is instead converted to lactate by cytosolic lactate dehydrogenase.

Mitochondria and ROS production: activation of the intrinsic pathway of apoptosis

The ROS H_2O_2 , superoxide (O_2^-) and hydroxyl radical (OH^-) are generated by mitochondrial respiratory complexes during uncoupled substrate turnover in the process of ATP formation (Cadenas and Davies, 2000; Hanahan and Weinberg, 2011). Premature leakage of electrons from respiratory complexes, rather than coupled transfer during ATP synthesis, leads to superoxide formation (Skulachev, 1998; Di Paola and Lorusso, 2006; Pike *et al.*, 2011). The mitochondrion is not only the main site of ROS generation; it is also their primary target. The generation of ROS has several consequences in cells, including direct peroxidative damage to membranes and modification of DNA bases that may initiate mutagenesis (Larsen *et al.*, 2005; Nathan and Cunningham-Bussel, 2013). ROS also modulate cellular redox homeostasis (Higdon *et al.*, 2012) and decrease the concentration of thiol-containing species, such as glutathione, which are cytoprotective (Mari *et al.*, 2009). Indeed, glutathione depletion decreases the

integrity and activities of mitochondrial respiratory complexes, which compromises ATP production and cell viability (Merad-Boudia *et al.*, 1998). More recently, it has been recognized that ROS are also able to activate the JNK (Trachootham *et al.*, 2006) and p38 MAPK (Ito *et al.*, 2006) signalling pathways that may trigger apoptotic cell death.

Apoptosis is a coordinated cell death programme that is important in normal tissues (Fulda and Debatin, 2006). There are two major pathways of apoptosis: the extrinsic and intrinsic pathways. In the extrinsic pathway, tumour necrosis factor family receptors at the plasma membrane are activated by FasL and related ligands and modulate intracellular signalling pathways leading to cell deletion. The intrinsic, or mitochondrial, pathway of apoptosis is activated intracellularly by ROS, the inhibition of pro-survival signalling cascades or by major cellular stresses, including DNA damage from exposure to cytotoxic anti-cancer drugs.

The relative ratio of pro- and anti-apoptotic Bcl-2 family proteins is a determinant of the response of cells to apoptotic stimuli (Fulda and Debatin, 2006). Thus, pro-apoptotic Bcl-2 proteins, such as Bax, Bak and Bid, form dimers that destabilize the outer mitochondrial membrane to apoptotic stimuli by forming channels that allow mitochondrial factors to exit, whereas anti-apoptotic members such as Bcl-2, Bcl-XL and Bcl-w stabilize the membrane. Thus, cytochrome *c*, Smac/Diablo or HtrA2/Omi are released to the cytosol, which triggers the commitment step of the mitochondrial apoptotic cascade (Kluck *et al.*, 1997; Yang *et al.*, 2003). The intrinsic and extrinsic apoptotic pathways converge on the executioner cysteine-aspartic acid proteases (caspases)-3 and -7, which lyse a number of cellular protein targets, including the DNA-repair protein poly (ADP ribose) polymerase (PARP); inactivation of the latter promotes apoptosis (Fulda and Debatin, 2006). However, release of Smac/Diablo or HtrA2/Omi from the mitochondrion does not directly activate caspase; rather these factors block the action of the protein inhibitor of apoptosis that normally acts to suppress caspases. Finally, there is also a caspase-independent mitochondrial pathway in which apoptosis-inducing factor and endonuclease G are released to the cytosol (Susin *et al.*, 1999). The translocation of these factors leads to DNA fragmentation, which is the hallmark of apoptosis.

Mitochondrial permeability is controlled by the permeability transition pore complex, which regulates the flow of small molecules across the mitochondrial membrane. Pore opening occurs in response to elevated Ca^{2+} concentrations, increased ROS and depleted adenine nucleotides (Kroemer *et al.*, 2007). Permeability transition pore opening in turn dissipates the proton gradient, uncouples oxidative phosphorylation and decreases ATP formation. In consequence, antioxidant molecules such as glutathione exit the mitochondrion, which further decreases the capacity to detoxify locally generated ROS (Mari *et al.*, 2009). The release of cytochrome *c* from the mitochondrion sets in train the activation of apoptotic death.

The permeability transition pore is a complex arrangement of proteins including the adenine nucleotide translocator, which exchanges ATP and ADP (Marzo *et al.*, 1998a; Brenner *et al.*, 2000), the voltage-dependent anion channel (also termed porin), the soluble mitochondrial matrix protein cyclophilin D (Woodfield *et al.*, 1998), a number of Bcl-2

family proteins (Marzo *et al.*, 1998b; Shimizu *et al.*, 1999), the peripheral benzodiazepine receptor and several proteins that regulate energy metabolism (e.g. hexokinase II and creatine kinase; Marzo *et al.*, 1998a). Several genes encoding components of the mitochondrial permeability transition pore including the peripheral benzodiazepine receptor, the associated protein Prax-1 and the energy-metabolizing enzyme creatine kinase, are overexpressed in some tumours (Kanazawa *et al.*, 1998; Venturini *et al.*, 1998; Galiegue *et al.*, 1999).

While the pro-apoptotic Bax may be down-regulated in cancer cells (Brimmell *et al.*, 1998), Bcl-2 or its anti-apoptotic homologues are frequently overexpressed (Kroemer *et al.*, 2007). This shifts the balance in the cancer cell towards the prevention of apoptosis. Thus, an important adaptation is that the PI3K/Akt signalling cascade inhibits pro-apoptotic Bcl-2 factors and positively regulates the anti-apoptotic factor Bcl-2 (Skorski *et al.*, 1997; Pugazhenthii *et al.*, 2000). As described above, this shift in Bcl-2 factor composition stabilizes the outer mitochondrial membrane. PI3K/Akt has additional survival actions in tumour cells in that it is able to impair signalling by the pro-apoptotic JNK MAPK, enhance glucose uptake by transporters and activate glycolytic enzymes, such as hexokinase (Skorski *et al.*, 1997; Pugazhenthii *et al.*, 2000; Kroemer *et al.*, 2007). These adaptations shift the capacity for energy production towards the glycolytic pathway that characterizes the cancer cell.

Biosynthetic roles of the mitochondrion in normal and cancer cells

Some important steps in lipid metabolism occur in the mitochondrion. While fatty acid synthesis occurs primarily in the cytoplasm, β -oxidation to produce ATP occurs in mitochondria. However, the mitochondrion also has a central role in

the biosynthesis of phospholipids and triglycerides. The first step in the phospholipid synthesis pathway is the esterification of α -glycerol phosphate by acyl-CoA to produce lysophosphatidic and phosphatidic acids (Zborowski and Wojtczak, 1969; Bremer *et al.*, 1976). Some phospholipids, such as phosphatidylcholine, phosphatidylserine and phosphatidylinositol, are synthesized in other organelles and transported to the mitochondrion where further biotransformation occurs. Cardiolipin, a characteristic phospholipid of the inner mitochondrial membrane, is synthesized within this organelle (Hostetler and van den Bosch, 1972).

In mammals, fatty acids are activated to acyl-CoAs on the outer mitochondrial membrane before entering the glycerolipid biosynthetic pathway via glycerol-3-phosphate acyltransferase. Highly proliferative cancer cells have an increased requirement for lipids for the assembly of cell membranes (Samudio *et al.*, 2009; Zaugg *et al.*, 2011). Increased expression of fatty acid synthase in tumour cells promotes formation of long-chain fatty acids and confers a growth advantage (Sabine *et al.*, 1967; Ookhtens *et al.*, 1984). Indeed, fatty acid synthase inhibition decreases the rate of cancer cell proliferation and promotes apoptosis, as reflected by increased caspase-3 activation, down-regulation of anti-apoptotic proteins and the release of cytochrome *c* (Pizer *et al.*, 1998; De Schrijver *et al.*, 2003).

Lipid synthesis and biotransformation in cells

Fatty acids perform structural roles in cells by modulating membrane fluidity and the arrangement of receptors and other proteins in plasma membrane lipid rafts (Spector and Yorek, 1985). Naturally occurring long-chain fatty acids of 18–22 carbons in length are stored in cell membranes as triglycerides or phospholipids. Triglycerides are composed of three fatty acid residues connected to the hydroxyl groups of glycerol via ester linkages (Figure 1A). Phosphoglycerides are




Figure 1

Chemical structures of fatty acids, triglycerides and phospholipids. (A) general structure of triglycerides; (B) general structure of phospholipids; (C) phosphatidylcholine; (D) important dietary fatty acids with carbon numbering.

the major phospholipid class, and are comprised of five components: two fatty acids, glycerol, a phosphate and an alcohol (Figure 1B). Subclasses are designated according to the alcohol group; thus phosphatidylcholines contain the choline group (Figure 1C). Other common alcohols include serine, ethanolamine and inositol. Phosphatidates share this same basic structure but lack the alcohol group, and are intermediates in the biosynthesis of many phosphoglycerides. The glycerol group may also be substituted with sphingosine (2-amino-4-octadecene-1,3-diol), giving rise to sphingomyelins.

Constituent fatty acids in membranes are either saturated [contain no carbon-carbon double bonds, as exemplified by palmitic acid (C16:0)], monounsaturated [contain only one olefinic bond; such as oleic acid (18:1 n-9)] or polyunsaturated [PUFAs that possess multiple olefinic bonds, typified by ω -6 arachidonic acid (20:4 n-6) and ω -3 eicosapentaenoic acid (20:5 n-3); Figure 1D]. The double bonds in naturally occurring unsaturated fatty acids are primarily in the *cis* configuration, which kinks the carbon chain and imparts greater membrane fluidity.

The carbon atoms of fatty acid chains are numbered starting from the carboxyl group and the carbon at the distal end of the molecule is the ω -carbon. Monounsaturated fatty acids and PUFA are classified by the position of the olefinic bond that is furthest from the carboxylate, and this is indicated in the lipid number notation. Thus, arachidonic acid (20:4 n-6) is a C20 fatty acid that has four olefinic bonds, with the double bond furthest from the carboxylate located six bonds from the ω -carbon (to produce a ω -6 olefinic bond).

Release of fatty acids from their esterified phospholipid forms in cell membranes is mediated by PLA₂. Free PUFAs are substrates for COX, lipoxygenase (LOX) and cytochrome P450 (CYP) enzymes that generate multiple metabolites that have diverse homeostatic functions (Figure 2; Spector and Yorek, 1985; Oates *et al.*, 1988; Oliw, 1994; Chen *et al.*, 1995; Murray, 1999; Marden *et al.*, 2003). Together, these enzymes produce PGs, thromboxanes, fatty acid peroxides and their downstream LTs, hydroxyfatty acids and epoxides.




Figure 2

PUFA biotransformation pathways. PUFA are released from their esterified form in cell membranes to free fatty acids. The enzymatic actions of COX, LOX and CYP generate signalling molecules with diverse homeostatic actions.

In comparison, saturated fatty acids primarily undergo CYP-dependent oxidation to the corresponding ω - and ω -1 hydroxy acids.

Apart from free fatty acids, there is evidence that certain lipids may be present at increased levels in cancer cells. Thus, phosphocholine derivatives are increased in cancer cells and solid tumours (Ackerstaff *et al.*, 2003) and plasma triglycerides are reportedly higher in women with invasive breast cancers (Goodwin *et al.*, 1997). Some important enzymes involved in fatty acid biotransformation are also differentially expressed in cancers. Overexpression of COX and CYP enzymes promotes the formation of certain PG and epoxide metabolites that influence the fate of cells, including the rate of proliferation and the inhibition of apoptosis (Tsuji *et al.*, 1997; Jiang *et al.*, 2005). Moreover PUFA-derived metabolites modulate signalling pathways such as ERK and PI3K/Akt that have been implicated in tumourigenesis. As discussed below, inhibition of the formation of these PUFA metabolites modulates apoptosis and other aspects of tumour growth (Chen *et al.*, 2009). Moreover, inhibition of the enzymes that mediate PUFA biotransformation has proven successful in the prevention of experimental tumour progression *in vitro* and *in vivo* (Koehne and DuBois, 2004; Chen *et al.*, 2009). However, at present, there is a deficiency of inhibitory agents that are well tolerated over the prolonged treatment periods required for anti-cancer chemotherapy.

The potential of lipids as anti-cancer agents that act at the mitochondrion

Fatty acids are thought to uncouple oxidative phosphorylation and decrease ATP production by facilitating the leakage of protons across the lipid mitochondrial membrane. Uncoupling is greatest with C12–C16 saturated and longer *cis*-unsaturated fatty acids (Korshunov *et al.*, 1998; Bernardi *et al.*, 2002). For example, 5 μ M laurate (C12-saturated) effectively inhibited H₂O₂ production by mitochondria (Korshunov *et al.*, 1998). Free fatty acids also impair electron transport and activate apoptosis by releasing cytochrome *c* from the inner mitochondrial membrane. There is increasing evidence that several classes of naturally occurring and synthetic lipids have the potential for development as anti-cancer agents that mediate their effects, at least in part, at the mitochondrion. However, some of these lipid-mediated anti-mitochondrial actions may be indirect, by modulating intracellular signalling pathways.

Naturally occurring free fatty acids and synthetic analogues

Saturated fatty acids. In the non-ionized state, medium- and long-chain fatty acids readily penetrate the mitochondrial membrane (McLaughlin and Dilger, 1980; Gutknecht, 1988; Kamp and Hamilton, 1992). Butyric acid and similar short-chain fatty acids induce cell cycle arrest and apoptosis by dissipating the mitochondrial membrane potential ($\Delta\Psi$), leading to growth arrest and apoptosis in colon carcinoma cells *in vitro*, as demonstrated by activation of caspase-3 (Heerdt *et al.*, 1997). In a small structure-activity study, the C4-butyric acid inhibited the growth of the human colonic




Figure 3

Naturally occurring fatty acids and analogues with anti-cancer activity.

adenocarcinoma cell lines HT-29, Colo-320 and SW-948 (IC_{50} s 0.55–2.28 mM), while C3-propionic acid was less potent and C5-valeric and C2-acetic acids were ineffective (Figure 3, Table 1; Milovic *et al.*, 2000). The apoptotic mechanism of butyric acid in colon cancer cells is mediated at least in part by inducing overexpression of the pro-apoptotic protein Bak, which alters the Bak/Bcl-2 ratio (Hague *et al.*, 1997; Ruemmele *et al.*, 1999), but not by modulating ROS or ATP production (Heerdt *et al.*, 1997). Bromo-analogues of butyric and propionic acids (IC_{50} s 0.13–0.39 mM) were several-fold more potent pro-apoptotic agents than butyrate, but also produced some cytotoxicity, so cautious development of these molecules as potential mitochondrial targeted agents is warranted (Milovic *et al.*, 2000).

Longer chain saturated fatty acids also act at the mitochondrion to activate apoptosis. Palmitic acid (C16:0) decreased $\Delta\Psi$ and effected cytochrome *c* release, which induced the proteolysis of PARP and the fragmentation of DNA (de Pablo *et al.*, 1999). This may also be a process of endogenous importance because long-chain fatty acids can be generated intracellularly by activation of PLA₂. In addition, such free fatty acids may also accumulate in mitochondria following exposure of cells to stimuli such as oxidative stress or increased Ca^{2+} concentrations (Broekemeier and Pfeiffer, 1995).

13-Methyltetradecanoic acid is an iso-C₁₅ branched-chain saturated fatty acid that has been found to disrupt mitochondrial integrity and induce apoptosis in a wide range of cancer cell lines (Figure 3, Table 1; Yang *et al.*, 2000;

Wongtangintharn *et al.*, 2005; Lin *et al.*, 2012). Induction of apoptosis by 13-methyltetradecanoic acid was rapid and was detected after only 2 h of treatment over the concentration range 0.04–0.35 mM (Lin *et al.*, 2012). In some cell types, apoptosis appeared to be a caspase-independent pathway but, in human bladder cancer cells, 13-methyltetradecanoic acid down-regulated Bcl-2, up-regulated Bax, promoted mitochondrial dysfunction and cytochrome *c* release, and activated caspases (Lin *et al.*, 2012). 13-Methyltetradecanoic acid also inhibited the PI3K/Akt survival cascade and activated the pro-apoptotic p38 and JNK MAPK pathways (Lin *et al.*, 2012).

In vivo growth of the prostate carcinoma cell line DU 145 and hepatocarcinoma-derived LCI-D35 cells after orthotopic implantation of tumour xenografts into nude mice was also inhibited by 13-methyltetradecanoic acid (35–105 mg kg⁻¹ day⁻¹). Apoptosis was induced without evidence of major toxicity, which suggests that 13-methyltetradecanoic acid could be a potential candidate for chemotherapy of human cancers (Yang *et al.*, 2000).

Unsaturated fatty acids. Naturally occurring PUFAs also modulate cell proliferation and apoptosis. The ω-3 PUFA eicosapentaenoic acid (20:5 n-3) is incorporated into mitochondrial phospholipids, which has a number of consequences for mitochondrial function, including decreased mitochondrial membrane potential and ATP production, increased ROS generation and increased apoptosis (Colquhoun, 2009). ω-6 PUFA also have the potential to

Table 1

Naturally occurring and synthetic fatty acids and other lipids: mitochondrial actions of potential value in cancer chemotherapy

Fatty acid derivative	Mitochondrial actions	Pathways effected	Function	References
Butyric acid	Dissipation of the mitochondrial membrane potential ($\Delta\Psi$)	Caspase-3 activation	↑Apoptosis Growth arrest	Heerdt <i>et al.</i> (1998) Milovic <i>et al.</i> (2000)
Bromo-analogues of butyric and propionic acids	↑ROS, DNA damage		↑Apoptosis	Milovic <i>et al.</i> (2000)
Palmitic acid	↓Mitochondrial membrane $\Delta\Psi$, cytochrome c release		↑Apoptosis ↑Cell death	Merrill and Jones (1990) Schlame <i>et al.</i> (2000) Ostrander <i>et al.</i> (2001)
13-Methyltetradecanoic acid	Disrupted mitochondrial integrity and caused dysfunction, cytochrome c release	↑Bax, ↓Bcl-2 Caspase-3 activation ↓pAkt ↑p38 and JNK MAPK	↑Apoptosis Growth inhibition	Yang <i>et al.</i> (2000) Wongtangtintharn <i>et al.</i> (2005) Lin <i>et al.</i> (2012)
Arachidonic acid-derived PGE ₂		↑EGFR/ERK MAPK ↑PI3K/Akt	↑Survival ↑Proliferation	Chen <i>et al.</i> (2009)
Eicosapentaenoic acid (EPA)	↑ATP glycolysis, ↑ROS, ↓mitochondrial membrane $\Delta\Psi$	Caspase-3 activation PKC inhibition, ↓Bcl-2	↑Apoptosis Growth inhibition	Denys <i>et al.</i> (2005) Colquhoun (2009)
EPA-derived epoxides		p38 MAPK	↑Apoptosis Growth inhibition	Cui <i>et al.</i> (2011)
Docosahexaenoic acid CLA	↓DNA synthesis	PKC inhibition, ↓Bcl-2 ↓Bcl-2, ↓PI3K/Akt	↑Apoptosis ↑Apoptosis ↓Proliferation	Denys <i>et al.</i> (2005) Ip <i>et al.</i> (1999) Ip <i>et al.</i> (2000) Kim <i>et al.</i> (2002)
Jacaric acid	↑ROS	↓Bcl-2, caspase-3, -8 and -9 activation, PARP cleavage	↑Apoptosis (intrinsic and/or extrinsic pathways)	Shinohara <i>et al.</i> (2012) Gasmi and Sanderson (2013)
α-Eleostearic acid	↓Mitochondrial membrane $\Delta\Psi$, ↑DNA fragmentation ↑lipid peroxidation		↑Apoptosis	Tsuzuki <i>et al.</i> (2004), Grossmann <i>et al.</i> (2009)
Vaccenic acid	↑DNA fragmentation		Growth inhibition ↓Cytosolic glutathione ↑Cell death	Miller <i>et al.</i> (2003)
Punicic acid	Disrupted the mitochondrial membrane $\Delta\Psi$, ↑DNA fragmentation, ↑lipid peroxidation	Caspase-3 and -9 activation ↓Bcl-2 (decreased Bcl-2:Bax ratio) ↓pAkt	↑Apoptosis	Grossmann <i>et al.</i> (2010), Gasmi and Sanderson (2010)
Tetradecylthioacetic acid	↑Mitochondrial proliferation, ↑oxidative stress, ↑ROS	Caspase-3 activation PARP cleavage	↑Apoptosis ↓Proliferation	Tronstad <i>et al.</i> (2001) Lin <i>et al.</i> (2012) Tronstad <i>et al.</i> (2003)
Jasmonates	Cytochrome c release, membrane depolarization, swelling of mitochondria, ↓ATP		↑Cell death	Fingrut and Flescher (2002) Rotem <i>et al.</i> (2005) Fingrut <i>et al.</i> (2005) Heyfets and Flescher (2007) Goldin <i>et al.</i> (2007)
EETs		↑EGFR/ERK ↑PI3K/Akt	↑Survival ↑Proliferation	Chen <i>et al.</i> (2009)
HETEs and downstream LTs	Inhibition of 5-LOX depleted mitochondrial glutathione and ↑lipid peroxidation	↑apoptotic Bcl-2 family proteins	↑Survival Enhanced growth	Ghosh and Myers (1998) Avis <i>et al.</i> (2001) Hoque <i>et al.</i> (2005)

Table 1

Continued

Fatty acid derivative	Mitochondrial actions	Pathways effected	Function	References
Short chain ceramides	Cytochrome c release, ↓mitochondrial membrane $\Delta\Psi$, ↑ROS	Caspase-3 activation, ↓Bcl-2: Bax ratio, ↑p38 and JNK MAPK phosphorylation, PARP	↑Apoptosis	Fillet <i>et al.</i> (2003) Flowers <i>et al.</i> (2012)
Anandamide and N-acylethanolamines	↑Permeability of inner mitochondrial membrane, ↑↓lipid peroxidation*, energetic and permeability transition alterations, Ca ²⁺ overload and PTP modulation		↑↓Apoptosis ↑↓proliferation	Epps <i>et al.</i> (1982) Gulaya <i>et al.</i> (1993) Schwarz <i>et al.</i> (1994) De Petrocellis <i>et al.</i> (1998) Maccarrone <i>et al.</i> (2000) Sarker <i>et al.</i> (2000) Wasilewski <i>et al.</i> (2004)
Cardiolipin	Involved in ROS-promoted cytochrome c release	Membrane targeting of Bid, activation of Bax	↑Apoptosis	Kuwana <i>et al.</i> (2002)
Perifosine	↑Cytochrome c release, mitochondrial oxidative phosphorylation disruption, induction of permeability transition	Akt inhibition Caspase-3, -8 and -9 activation ↑JNK MAPK translocation	↑Apoptosis	Kondapaka <i>et al.</i> (2003) Nieto-Miguel <i>et al.</i> (2006) Chiarini <i>et al.</i> (2008) Burgeiro <i>et al.</i> (2013)
Edelfosine	Mitochondrial dysfunction via multiple mechanisms Altered mitochondrial membrane $\Delta\Psi$, ↑DNA fragmentation, ↑ROS	Caspase-3 activation PARP	↑Apoptosis	Cabaner <i>et al.</i> (1999) Gajate <i>et al.</i> (2000) Mollinedo <i>et al.</i> (2011)
Isopropylaminopropanol derivatives (C16:0 and C18:1)		Inhibition of multiple pro-survival kinases	Growth inhibition	Cao <i>et al.</i> (2013)

*Dependent on fatty acid chain length and saturation.

activate apoptosis in human cancer cell lines by promoting lipid peroxidation (Cao *et al.*, 2000).

Conjugated linoleic acid (CLA) is a mixture of geometric isomers of linoleic acid (Figure 3) that decreases the viability of various cancer cell types, including those of the skin, forestomach, mammary gland and colon. CLA (32 μM), but not linoleic acid, inhibited growth of rat mammary epithelial cell organoids that was mediated both by a decrease in DNA synthesis and increased apoptosis (Ip *et al.*, 1999). *In vivo* activity was also noted and feeding CLA to rats that harboured premalignant lesions induced apoptosis in a mammary tumour cell line, as determined by a decrease in expression of the anti-apoptotic bcl-2 (Table 1; Ip *et al.*, 2000). Cho *et al.* (2003) showed that CLA inhibited the proliferation of HT-29 human colorectal cancer cell line by activating apoptosis, due in part to inhibition of PI3K/Akt signalling. Similarly, the 10-*trans*,12-*cis* CLA isomer (5 μM) inhibited the proliferation of Caco-2 colon carcinoma cells, and enhanced apoptosis in premalignant lesions, but not in normal cells (Kim *et al.*, 2002).

Other types of naturally occurring conjugated fatty acids that have pro-apoptotic actions include CLAs such as α -eleostearic acid (9-*cis*,11-*trans*,13-*trans*-18:3) from bitter gourd oil and calendic acid (8-*trans*,10-*trans*-12-*cis*-18:3) from pot marigold. α -Eleostearic acid was quite potent relative to

CLA isomers against tumour cells *in vitro* (Tsuzuki *et al.*, 2004). α -Eleostearic acid (40 μM) promoted lipid peroxidation and apoptosis in MDA-MB-231-derived cell lines as evidenced by a loss of mitochondrial membrane potential and the release of apoptotic factors from the mitochondrion. When treated with α -eleostearic acid in the presence of α -tocotrienol (20 μM), growth inhibition and apoptosis did not occur, thus providing further support for involvement of ROS-mediated lipid peroxidation in the apoptotic mechanism (Grossmann *et al.*, 2009).

The monounsaturated fatty acid vaccenic acid (11-*trans*-18:1 n-7) decreased cell growth, induced DNA fragmentation and depleted cytosolic glutathione levels; these findings are consistent with activation of the intrinsic pathway of apoptosis by lipid peroxides (Miller *et al.*, 2003). Punicic acid (18:3 n-5) is an ω -5 long-chain PUFA found in pomegranate seed oil. In MDA-MB-231 and MDA-ER α 7 cells, puniceic acid disrupted the mitochondrial membrane potential and induced apoptosis, apparently also by a prooxidant mechanism (Grossmann *et al.*, 2010). DNA fragmentation was observed after 24 h treatment of cells at concentrations in the range 10–100 μM (Gasmi and Sanderson, 2010).

Jacaric acid is a linolenic acid isomer obtained from jacaranda that has a conjugated triene system and elicits potent anti-tumour effects both *in vitro* and *in vivo* in nude mice into

which DLD-1 cells had been xeno-transplanted (Shinohara *et al.*, 2012). When compared with natural conjugated linolenic acids in DLD-1 adenocarcinoma cells, the anti-tumour effects of jacaric acid were most potent and correlated with increased ROS production. Thus, jacaric acid induced concentration- and time-dependent LNCaP cell death in part through activation of intrinsic apoptotic pathways, resulting in cleavage of caspase-3, -8 and -9, modulation of pro- and anti-apoptotic Bcl-2 family of proteins and increased cleavage of PARP-1 (Gasmi and Sanderson, 2013).

Several dietary C18 unsaturated fatty acids have been tested for cytotoxicity and induction of apoptosis in human prostate cancer cells. These included three octadecatrienoic geometric isomers (α - and β -calendric and catalpic acids) and two monounsaturated C18 fatty acids (*trans*- and *cis*-vaccenic acid) in addition to jacaric and puniceic acids (Figure 3, Table 1). Jacaric acid and four of its octadecatrienoic acid geometric isomers selectively induced apoptosis in both hormone-dependent (LNCaP) and hormone-independent (PC-3) human prostate cancer cells when tested at concentrations around 10 μ M, without affecting the viability of normal human prostate epithelial cells (Gasmi and Sanderson, 2013). Together, these findings suggest that some of the pro-apoptotic actions of anti-tumour fatty acids may be cell type specific.

From the foregoing, it is evident that most studies to date have focused on ROS generation by unsaturated fatty acids as the mechanism of their pro-apoptotic action. However, there may be additional, more selective, mechanisms that could be developed in synthetic anti-cancer fatty acids. In a recent study, a series of n-3 monounsaturated fatty acids of chain length C16-C22 was synthesized and evaluated in MDA-MB-468 breast cancer cells that stably overexpressed COX-2 (Cui *et al.*, 2012). This reflects the situation that may operate in many human cancers in which COX-2 is up-regulated. The longer chain C19-C22 analogues were found to inhibit proliferation and activate apoptosis; C16-C18 analogues were less active. PGE₂ formation was decreased by the C19-C22 analogues, consistent with COX-2 inhibition, which was supported by molecular modelling that revealed effective interactions with specific amino acid residues in the COX-2 active site. Strategies of this type, in which potential fatty acid drugs target a biotransformation enzyme present in tumours, may be worthy of further consideration. Such agents may enable approaches based on COX-2 inhibition to be retained, perhaps without the toxicity associated with conventional non-steroidal anti-inflammatory drugs.

Fatty acid analogues. Tetradecylthioacetic acid (TTA) is a saturated fatty acid that has a sulphur atom inserted at the C3 position in the carbon chain. TTA (0.2–0.5 mM) decreased proliferation and induced apoptosis in a diverse range of tumour cell lines *in vitro* and *in vivo* (Tronstad *et al.*, 2003; Iversen *et al.*, 2006). Long-chain 3-thia-fatty acids in general also uncouple oxidative phosphorylation and dissipate the mitochondrial membrane potential ($\Delta\Psi$) by direct interaction with the adenine nucleotide translocator to open the mitochondrial permeability transition pore, which leads to decreased ATP production (Wieckowski and Wojtczak, 1998). In accord with this mechanism, TTA stimulates mitochondrial ROS production (Tronstad *et al.*, 2001), leading to glu-

tathione depletion, which renders mitochondria susceptible to further damage (Tronstad *et al.*, 2003). Activation of apoptosis was indicated by the release of mitochondrial cytochrome *c* that enhanced caspase-3 activation and PARP cleavage.

TTA is resistant to mitochondrial β -oxidation and, compared with naturally occurring saturated fatty acids, is degraded relatively slowly to dicarboxylic acids by microsomal oxidation at the ω -carbon and sulphur atoms and by peroxisomal β -oxidation (Hvattum *et al.*, 1991). This property could be useful in development of TTA and analogues as drugs since these molecules are likely to have superior durations of action *in vivo*. Indeed, a diet containing TTA increased the vascularization of colon cancer xenografts in mice and improved the survival of mice with leukaemia xenografts (Jensen *et al.*, 2007).

Jasmonates are plant hormones that structurally resemble fatty acid esters (Figure 3). These agents interact directly with mitochondria in cancer cells to detach hexokinase-II from its location on the voltage-dependent anion channel in the mitochondrial permeability transition pore (Goldin *et al.*, 2008). Indeed, the susceptibility in cancer cells to these molecules is dependent on the association of hexokinase with the mitochondrion. Pro-apoptotic mitochondrial actions of low millimolar concentrations of jasmonates include membrane depolarization, mitochondrial swelling, cytochrome *c* release and cell death; interestingly, the agents were inactive in normal cells (Table 1; Fingrut and Flescher, 2002; Rotem *et al.*, 2005).

Correlations have been reported between methyl jasmonate cytotoxicity in a range of cell types and the extent of ATP depletion (Goldin *et al.*, 2007). Glucose protected against this loss of ATP, whereas the glycolysis inhibitor, 2-deoxyglucose, synergized with methyl jasmonate (Fingrut *et al.*, 2005; Heyfets and Flescher, 2007). Similar effects have been elicited by other hexokinase-detaching agents, such as hypericin and clotrimazole, that also deplete ATP and decrease cell viability (Miccoli *et al.*, 1998; Machida *et al.*, 2006). These findings are consistent with the functional importance of hexokinase in glycolysis, which is a major pathway of ATP production in cancer cells.

Taken together, a number of studies have found that endogenous fatty acids, including saturated and certain unsaturated analogues, activate apoptosis via the mitochondrial intrinsic pathway and impair ATP production. Frequently, this may be due to increased ROS production, which alters mitochondrial membrane potential, but the example of jasmonate, involving hexokinase detachment, illustrates the potential for further development of mitochondrially targeted fatty acids that selectively disrupt energy metabolism in cancer cells.

Fatty acid metabolites

Several PUFA-derived metabolites have been shown to modulate signalling pathways that are implicated in tumourigenesis. Thus, inhibition of the enzymic formation of these fatty acid metabolites has been found to decrease cancer cell viability. Important metabolites that have emerged in this regard include COX-mediated PGE₂, CYP-dependent epoxyeicosatrienoic acids (EETs) and certain LOX-mediated hydroxyeicosatetraenoic acids (HETEs) that inhibit apoptosis and enhance

proliferation (Avis *et al.*, 2001; Koehne and DuBois, 2004; Chen *et al.*, 2009). An understanding of the involvement of fatty acid metabolites in the pathogenesis of cancer could open new avenues for the production of new and safer therapeutic and chemopreventive agents.

Certain fatty acid biotransformation enzymes have been detected at high level in human cancers. Thus, COX-2 and CYP2J2 are overexpressed in many invasive human cancers (Tsuji *et al.*, 1997; Jiang *et al.*, 2005). COX-2-derived PGE₂, LOX metabolites and CYP2J2-derived EETs are implicated in aggressive tumour behaviour (Figure 4, Table 1; Avis *et al.*, 2001; Koehne and DuBois, 2004; Hoque *et al.*, 2005; Jiang *et al.*, 2005). Mechanistic information is available for some of these metabolites. Thus, PGE₂ and EETs enhance tumour cell proliferation and survival by activating the proliferative EGF receptor (EGFR)/ERK MAPK and anti-apoptotic PI3K/Akt signalling pathways (Chen *et al.*, 2009).

Inhibition of 5-LOX by MK886 induces apoptosis in both hormone-responsive (LNCaP) and hormone-unresponsive (PC3) prostate cancer cells (Ghosh and Myers, 1998). An immediate and sustained rise in cytosolic calcium is followed by mitochondrial uncoupling and cytochrome *c* release (Maccarrone *et al.*, 2001). Soon after treatment, cells underwent the mitochondrial permeability transition, followed by other apoptotic events, including externalization of phosphatidylserine and degradation of DNA (Ghosh and Myers, 1998). Cell death was prevented by direct addition of 5-HETE and analogues such as 5-HETE-lactone and 5-oxoeicosatetraenoic acid (50–500 nM; Ghosh and Myers,

1998). In breast cancer cells, inhibition of 5-LOX, but not COX, reduced growth, increased apoptosis, down-regulated bcl-2, up-regulated bax and caused G₁ cell cycle arrest (Avis *et al.*, 2001). Lipid peroxidation and the depletion of mitochondrial glutathione have been linked to the activation of apoptotic Bcl-2 proteins in these cells.

While the inhibition of biotransformation enzymes that generate lipid metabolites may be beneficial in the prevention of cancer progression, there is also increasing evidence emerging that ω -3 PUFAs undergo biotransformation to metabolites that inhibit carcinogenesis. Inhibition of PKC by the parent ω -3 PUFAs eicosapentaenoic and docosahexaenoic acids activates apoptosis by down-regulating Bcl-2 (Denys *et al.*, 2005); COX-2-derived ω -3 PGE₃ was recently implicated in anti-cancer actions of ω -3 PUFAs (Szymczak *et al.*, 2008). Epoxides of ω -3 PUFAs have also been shown to exert growth suppressing and anti-cancer effects. The 17,18-epoxide of eicosapentaenoic acid (ω -3-epoxy-eicosatetraenoic acid), but not other regio-isomeric epoxides of eicosatetraenoic acid, was shown to inhibit cell proliferation. At physiologically relevant concentrations, ω -3-epoxy-eicosapentaenoic acid induced apoptosis and cell cycle arrest of brain microvascular endothelial bEND.3 cells through activation of growth-suppressing p38 MAPK and subsequent down-regulation of cyclin D1 (Cui *et al.*, 2011). More recently, Zhang *et al.* (2013) showed that epoxygenase-mediated metabolites of the C22 ω -3 fatty acid docosahexaenoic acid exerted anti-cancer effects by suppressing vascular endothelial growth factor-mediated angiogenesis.




Figure 4

Chemical structures of important fatty acid metabolites and phospholipids, and general structures of ceramides and sphingomyelins.

Inhibition of angiogenesis resulted in a decrease in primary tumour growth and metastasis *in vitro*. By co-administration with *t*-AUCB, a soluble epoxide hydrolase inhibitor, the 19,20-epoxide was active *in vivo*, reducing tumour growth in the Lewis lung carcinoma model (Zhang *et al.*, 2013). These actions were opposite to those of ω -6 arachidonic acid-derived EETs. However, at present, whether PUFA epoxides also interact directly with the mitochondrion, in addition to their indirect actions mediated by signalling cascades, to elicit these actions is unclear.

N-acyl lipids

Ceramides. Ceramides are a class of *N*-acylated sphingoid bases (Figure 4) found at high concentration in cell membranes and cytosolic organelle membranes and act as intracellular second messengers in the regulation of growth, differentiation and apoptosis. Ceramide accumulation occurs after treatment of cells with apoptotic agents including chemotherapeutic agents, or after treatment with saturated fatty acids such as palmitic acid (Merrill and Jones, 1990). Direct addition of $\sim 1 \mu\text{M}$ ceramide to mitochondria and ceramide accumulation in cells both produced changes in the mitochondrial transmembrane potential by forming channels or targeting Bcl-2 family members that leads to translocation of cytochrome *c* from mitochondria to the cytoplasm, and caspase-3 activation (Garcia-Ruiz *et al.*, 1997). Because respiratory complex III is inhibited by C2-ceramide, the proximal effects of ceramide in cells are mediated at least in part at the mitochondrion (Gudz *et al.*, 1997). Inhibition of p38 and JNK MAPKs decreased ceramide-induced apoptosis by preventing the loss of mitochondrial transmembrane potential and inhibition of caspase activation (Chen *et al.*, 2008). This suggests that ceramides may operate by both direct and indirect mechanisms to exert apoptotic actions at the mitochondrion.

In tumour cells, the cell-permeable shorter chain exogenous C2- and C6-ceramide analogues, but not the longer chain, naturally occurring C16-ceramide, activated intrinsic apoptotic events at concentrations $\geq 10 \mu\text{M}$, including caspase-3 activation, PARP degradation and mitochondrial cytochrome *c* release (Fillet *et al.*, 2003; Flowers *et al.*, 2012). C6-ceramide increased ROS levels in MDA-MB-231 cells, shifted the Bax:Bcl-2 ratio and depolarized the mitochondrial membrane (Flowers *et al.*, 2012). Thus, analogues containing fatty acids of medium chain length may be particularly suited to development as putative anti-cancer agents that target the mitochondrion.

***N*-acylethanolamines.** Anandamide (Figure 4) and other *N*-acylethanolamines reportedly promote apoptosis and/or inhibit cell proliferation (Table 1; Schwarz *et al.*, 1994; De Petrocellis *et al.*, 1998; Maccarrone *et al.*, 2000; Sarker *et al.*, 2000). These molecules appear to act in part by increasing the inner mitochondrial membrane permeability at concentrations up to $100 \mu\text{M}$ (Epps *et al.*, 1982). Anandamide and *N*-oleoylethanolamine exerted protonophoric effects in mitochondria due to dissipation of the transmembrane potential and opening of the permeability transition pore (Wasilewski *et al.*, 2004). Long-chain *N*-acylethanolamines, including anandamide, accumulate in mammalian tissues under a

variety of pathological conditions (Schmid *et al.*, 2002). Indeed, they have been detected at levels up to 500 nmol g^{-1} of infarcted canine myocardium. They have also been shown to inhibit the growth of various cancer cell lines *in vitro*. Cancer tissues usually contain substantially higher concentrations of these lipids than adjacent benign tissue, when normalized to wet weight since most tumours also contained higher levels of phospholipids.

Despite these apparent pro-apoptotic effects of *N*-acylethanolamines in cancer cells, there is also evidence for protective effects in cells. The long-chain *N*-palmitoyl- and *N*-stearoylethanolamine ($100 \mu\text{M}$) inhibited lipid peroxidation in hepatic mitochondria, which is consistent with membrane-protective properties (Gulaya *et al.*, 1998). The beneficial effect of *N*-acylethanolamines on cell survival when oxygen availability is low depends in part on the inhibition of lipid oxidation. The effect of different *N*-acylethanolamines on lipid peroxidation seems to be dependent on the length of acyl chain as was found for other effects of *N*-acylethanolamines in the cell (Gulaya *et al.*, 1993). These findings highlight the need for clarification of the relationships between chain length in *N*-acylethanolamines and their cellular actions prior to development of analogues as potential anti-cancer drugs.

Very recently, novel fatty acid derivatives of isopropylaminopropanol containing C16:0 or C18:1 substituents were prepared and were found to be effective against the growth of hepatoma cells *in vitro* ($\text{IC}_{50}\text{s} \sim 5\text{--}10 \mu\text{M}$) and *in vivo* xenografts when dosed at 25 mg kg^{-1} (Cao *et al.*, 2013). These agents inhibited the activity of multiple kinases, including the pro-survival Akt, which increased caspase and PARP cleavage. It will now be of interest to explore in greater detail how these agents modulate mitochondrial activity to elicit anti-cancer actions.

Phospholipid derivatives

Cardiolipin. Cardiolipin is a structurally complex diphosphatidylglycerol lipid that is synthesized by the mitochondrion (Figure 4). Mitochondrial respiratory complexes have been shown to require cardiolipin for full function. Palmitic acid decreased the levels of cardiolipin, which is responsible for insertion and retention of cytochrome *c* in the inner membrane of the mitochondrion (Schlame *et al.*, 2000; Ostrander *et al.*, 2001). Decreased cardiolipin and altered mitochondrial function mediate palmitate-induced breast cancer cell death by promoting ROS production and release of cytochrome *c* by permeabilization of the outer membrane (Petrosillo *et al.*, 2003); overexpression of the anti-apoptotic Bcl-2 family members Bcl-xL and Bcl-w blocked apoptosis (Kuwana *et al.*, 2002).

Ether phospholipids. The ether phospholipids are a promising class of anti-tumour lipids that act in part at the mitochondrion. Ether phospholipids have one or more glycerol carbons bonded to an alkyl chain via an ether linkage, as opposed to the usual ester linkage. Ether phospholipids include miltefosine, ilmofosine, perifosine, edelfosine and erucylphosphocholine (Figure 4).

The novel alkylphospholipid analogue perifosine is a PI3K/Akt inhibitor (Table 1; Kondapaka *et al.*, 2003) that

inhibits growth at low micromolar concentrations and activates the intrinsic pathway of apoptosis in cells as evidenced by increased caspase activity and cleavage of PARP. Activated caspase-8 cleaves Bid, which migrates to mitochondria and induces cytochrome *c* release (Chiarini *et al.*, 2008). Interestingly, these agents are effective in rapidly proliferating cancer cells, but not quiescent normal cells.

Clinical evaluation of perifosine has been conducted, or is continuing, in patients with cancers of the endometrium, breast, prostate, bladder and other tissues. In 2010, perifosine was evaluated in phase II trials for metastatic colon cancer and extended the time taken for tumour progression. However, in 2013, it was announced that the drug failed trials in patients with relapsed or relapsed/refractory multiple myeloma (<http://www.aezsinc.com/en/page.php?p=60&q=550>, accessed 28 June 2013). When used in combination with the multikinase inhibitor sorafenib, perifosine induced intrinsic apoptosis in cells and anti-tumour effects in NOD/SCID mice with Hodgkin lymphoma cell line xenografts (Locatelli *et al.*, 2013). In cell lines, the combination treatment inhibited MAPK, activated PI3K/Akt phosphorylation and suppressed growth; synergistic induction of mitochondrial dysfunction and cell death was noted. In *in vivo* xenograft studies, there was a reduction in tumour burden, increased survival time, increased apoptosis and necrosis in perifosine/sorafenib-treated animals compared with single agents. Subsequently, treatment of human leukaemia T-cells with the PI3K/Akt inhibitor perifosine and etoposide also effected synergistic induction of apoptosis by dual activation of intrinsic and extrinsic pathways (Nyäkern *et al.*, 2006). This combination produced a twofold increase in caspase-8 activation, and a marked increase in caspase-9, caspase-3 and PARP cleavage, as well as increased Bim, Bid and Bcl-XL expression. Etoposide and perifosine induced leukaemic cell death in part by inactivation of the PI3K/Akt pathway that increased mitochondrial dysfunction. However, it is of considerable interest that ether phospholipids proved to be highly effective when used in combination with other anti-cancer agents.

Edelfosine induces changes in mitochondrial membrane permeability and inhibits mitochondrial respiration (Burgeiro *et al.*, 2013). Edelfosine, miltefosine, erucylphosphocholine and perifosine all activate the JNK MAPK pathway, which induces apoptosis (Ruiter *et al.*, 1999; Nieto-Miguel *et al.*, 2006). The mechanism involves direct activation of apoptosis by phosphorylation of the anti-apoptotic Bcl-XL (Kharbanda *et al.*, 2000; Aoki *et al.*, 2002). Edelfosine also disrupts the mitochondrial transmembrane potential, apparently by altering mitochondrial membrane phosphocholine content (Vrablic *et al.*, 2001), promotes the cleavage of caspase-3 and PARP, and enhances production of ROS in human leukaemic T-cells (Table 1; Cabaner *et al.*, 1999; Gajate *et al.*, 2000). Edelfosine-induced apoptosis is blocked by Bcl-2 (Mollinedo *et al.*, 1997; Ruiter *et al.*, 2003; Hideshima *et al.*, 2006).

New fluorescent edelfosine analogues retained the pro-apoptotic activity of the parent, and colocalized with mitochondria (Mollinedo *et al.*, 2011). These agents induced the swelling of isolated mitochondria, consistent with an increase in mitochondrial membrane permeability. Free radical scavengers did not affect swelling, suggesting that ROS

do not contribute. It was suggested that edelfosine promoted a redistribution of lipid rafts from plasma membrane to mitochondria (Mollinedo *et al.*, 2011). In summary, ether phospholipids have potential for development as a novel class of anti-cancer agents that act in part by altering mitochondrial function.

Challenges in the development of lipid molecules as drugs

Fatty acids differ from typical drugs in that there is a large degree of molecular flexibility inherent in fatty acid alkyl chains. Although fewer than 10 rotatable bonds is seen as desirable for adequate oral bioavailability and membrane permeation (Veber *et al.*, 2002), fatty acids possess satisfactory pharmacokinetic profiles. For example, medium- and long-chain dietary fatty acids are readily absorbed and distributed throughout all tissues of the body including the brain. Moreover, the ether phospholipid edelfosine is orally active despite possessing 27 rotatable bonds. The favourable pharmacokinetic profiles of fatty acids may be in part due to the facilitation of uptake and transport (Ramirez *et al.*, 2001), including binding to serum albumin and incorporation into triglycerides, that distribute these essential nutrients throughout the body. However, the flexibility of fatty acids does present challenges in drug design, particularly in ligand-based approaches where the structure of the drug target is unknown. Conformational analysis is a critical step in pharmacophore and pseudo-receptor modelling, and the high number of rotatable bonds can make identification of low-energy and bioactive conformations difficult and time consuming. When the structure of the drug target is known, ligand docking approaches are straightforward, as exemplified by our recent modelling of the interactions of a series of novel ω -3 monounsaturated fatty acids in the COX-2 active site (Cui *et al.*, 2012).

As mentioned earlier, fatty acids are subject to numerous metabolic processes mediated by COX, LOX and CYP enzymes that can present challenges for the development of lipid-derived drugs that have acceptable *in vivo* stability. Furthermore, lipid-derived mediators (e.g. PGs, epoxides, resolvins and others) frequently possess labile functional groups that are also readily metabolized. Improvement of the metabolic as well as chemical stability of lipid-based drugs, particularly those derived from PGs (Collins and Djuric, 1993; Das *et al.*, 2007) and lipoxin (Duffy and Guiry, 2010), has received much attention and a number of strategies have been developed and successfully employed in drug discovery settings. These strategies commonly involve addition of functional groups to block particular metabolic processes, or bioisosteric replacement of labile functional groups with more robust equivalents. For example, incorporation of a heteroatom in alkyl chains at the position β to the carboxylate functionality can improve the duration of action *in vivo*, as shown by the PG analogue cicaprost (Hildebrand *et al.*, 1989). These approaches have led to the development of numerous marketed drugs, which clearly demonstrate the potential of lipids as drugs and as lead compounds in drug discovery.

Development of new mitochondrial targeted inhibitors in cancer chemotherapy

Evidence is increasing that fatty acids and other lipids act in part by modulation of mitochondrial function. These actions may be direct, such as the detachment of hexokinase from the cancer cell mitochondrial transition pore complex or by uncoupling of respiratory complexes that produce ATP. Alternatively, some lipid agents may act indirectly by increasing ROS activity in cells or by interfering with cell signalling pathways, to perturb the balance of pro- and anti-apoptotic bcl-2 proteins that regulate mitochondrial membrane stability. It should be noted, however, that mitochondrial actions of certain lipid-based molecules may operate alongside non-mitochondrial mechanisms, including altered membrane raft composition or altered gene regulation.

A point of major interest that has emerged from cellular studies is that fatty acid derivatives often exhibit activity against cancer cells, but not normal cells. This appears to be a property that augers well for new anti-cancer drug development based on lipids. The available information does not provide a full understanding for this selectivity because the targeted pathways may be present in both cell types. It will now be of major importance to define in detail the defects present in mitochondria of cancer cells so that new drugs may be developed that have optimal potency with fewer off-target actions.

Acknowledgements

Support of research in the authors' laboratory by the Australian National Health and Medical Research Council is gratefully acknowledged.

Conflict of interest

None.

References

- Ackerstaff E, Glunde K, Bhujwala ZM (2003). Choline phospholipid metabolism: a target in cancer cells? *J Cell Biochem* 90: 525–533.
- Aoki H, Kang PM, Hampe J, Yoshimura K, Noma T, Matsuzaki M *et al.* (2002). Direct activation of mitochondrial apoptosis machinery by c-Jun N-terminal kinase in adult cardiac myocytes. *J Biol Chem* 277: 10244–10250.
- Avis I, Hong SH, Martinez A, Moody T, Choi YH, Trepel J *et al.* (2001). Five-lipoxygenase inhibitors can mediate apoptosis in human breast cancer cell lines through complex eicosanoid interactions. *FASEB J* 15: 2007–2009.
- Bernardi P, Penzo D, Wojtczak L (2002). Mitochondrial energy dissipation by fatty acids. *Vitam Horm* 65: 97–126.

Bremer J, Bjerve KS, Borrebaek B, Christiansen R (1976). The glycerophosphate acyltransferases and their function in the metabolism of fatty acids. *Mol Cell Biochem* 12: 113–125.

Brenner C, Cardiou H, Vieira HL, Zamzami N, Marzo I, Xie Z *et al.* (2000). Bcl-2 and Bax regulate the channel activity of the mitochondrial adenine nucleotide translocator. *Oncogene* 19: 329–336.

Brimmell M, Mendiola R, Mangion J, Packham G (1998). BAX frameshift mutations in cell lines derived from human hematopoietic malignancies are associated with resistance to apoptosis and microsatellite instability. *Oncogene* 16: 1803–1812.

Broekemeier KM, Pfeiffer DR (1995). Inhibition of the mitochondrial permeability transition by cyclosporin A during long time frame experiments: relationship between pore opening and the activity of mitochondrial phospholipases. *Biochemistry* 34: 16440–16449.

Burgeiro A, Pereira CV, Carvalho FS, Pereira GC, Mollinedo F, Oliveira PJ (2013). Edelfosine and perifosine disrupt hepatic mitochondrial oxidative phosphorylation and induce the permeability transition. *Mitochondrion* 13: 25–35.

Cabaner C, Gajate C, Macho A, Munoz E, Modolell M, Mollinedo F (1999). Induction of apoptosis in human mitogen-activated peripheral blood T-lymphocytes by the ether phospholipid ET-18-OCH3: involvement of the Fas receptor/ligand system. *Br J Pharmacol* 127: 813–825.

Cadenas E, Davies KJ (2000). Mitochondrial free radical generation, oxidative stress, and aging. *Free Radic Biol Med* 29: 222–230.

Cao M, Prima V, Nelson D, Svetlov S (2013). Composite fatty acid ether amides suppress growth of liver cancer cells *in vitro* and in an *in vivo* allograft mouse model. *Cell Oncol* 36: 247–257.

Cao Y, Pearman AT, Zimmerman GA, McIntyre TM, Prescott SM (2000). Intracellular unesterified arachidonic acid signals apoptosis. *Proc Natl Acad Sci USA* 97: 11280–11285.

Chen C, Li G, Liao W, Wu J, Liu L, Ma D *et al.* (2009). Selective inhibitors of CYP2J2 related to terfenadine exhibit strong activity against human cancers *in vitro* and *in vivo*. *J Pharmacol Exp Ther* 329: 908–918.

Chen CL, Lin CF, Chang WT, Huang WC, Teng CF, Lin YS (2008). Ceramide induces p38 MAPK and JNK activation through a mechanism involving a thioredoxin-interacting protein-mediated pathway. *Blood* 111: 4365–4374.

Chen J, Murray M, Liddle C, Jiang XM, Farrell GC (1995). Down-regulation of male-specific cytochrome P450s 2C11 and 3A2 in bile duct-ligated male rats: importance to reduced hepatic content of cytochrome P450 in cholestasis. *Hepatology* 22: 580–587.

Chiarini F, Del Sole M, Mongiorgi S, Gaboardi GC, Cappellini A, Mantovani I *et al.* (2008). The novel Akt inhibitor, perifosine, induces caspase-dependent apoptosis and downregulates P-glycoprotein expression in multidrug-resistant human T-acute leukemia cells by a JNK-dependent mechanism. *Leukemia* 22: 1106–1116.

Cho HJ, Kim WK, Kim EJ, Jung KC, Park S, Lee HS *et al.* (2003). Conjugated linoleic acid inhibits cell proliferation and ErbB3 signaling in HT-29 human colon cell line. *Am J Physiol Gastrointest Liver Physiol* 284: G996–1005.

Collins PW, Djuric SW (1993). Synthesis of therapeutically useful prostaglandin and prostacyclin analogs. *Chem Rev* 93: 1533–1564.

Colquhoun A (2009). Mechanisms of action of eicosapentaenoic acid in bladder cancer cells *in vitro*: alterations in mitochondrial metabolism, reactive oxygen species generation and apoptosis induction. *J Urol* 181: 1885–1893.

- Cui PH, Petrovic N, Murray M (2011). The ω -3 epoxide of eicosapentaenoic acid inhibits endothelial cell proliferation by p38 MAP kinase activation and cyclin D1/CDK4 downregulation. *Br J Pharmacol* 162: 1143–1155.
- Cui PH, Rawling T, Bourget K, Kim T, Duke CC, Doddareddy MR *et al.* (2012). Antiproliferative and antimigratory actions of synthetic long chain n-3 monounsaturated fatty acids in breast cancer cells that overexpress cyclooxygenase-2. *J Med Chem* 55: 7163–7172.
- Das S, Chandrasekhar S, Yadav JS, Gree R (2007). Recent developments in the synthesis of prostaglandins and analogues. *Chem Rev* 107: 3286–3337.
- De Petrocellis L, Melch D, Palmisano A, Bisogno T, Laezza C, Bifulco M *et al.* (1998). The endogenous cannabinoid anandamide inhibits human breast cancer cell proliferation. *Proc Natl Acad Sci USA* 95: 8375–8380.
- De Schrijver E, Brusselmans K, Heyns W, Verhoeven G, Swinnen JV (2003). RNA interference-mediated silencing of the fatty acid synthase gene attenuates growth and induces morphological changes and apoptosis of LNCaP prostate cancer cells. *Cancer Res* 63: 3799–3804.
- Denys A, Hichami A, Khan NA (2005). n-3 PUFAs modulate T-cell activation via protein kinase C- α and - ϵ and the NF- κ B signaling pathway. *J Lipid Res* 46: 752–758.
- Di Paola MD, Lorusso M (2006). Interaction of free fatty acids with mitochondria: coupling, uncoupling and permeability transition. *Biochem Biophys Acta* 1757: 1330–1337.
- Duffy CD, Guiry PJ (2010). Recent advances in the chemistry and biology of stable synthetic Lipoxin analogues. *MedChemComm* 1: 249–265.
- Epps DE, Palmer JW, Schmid HHO, Pfeiffer DR (1982). Inhibition of permeability-dependent Ca²⁺ release from mitochondria by N-acylethanolamines, a class of lipids synthesized in ischemic heart tissue. *J Biol Chem* 257: 1383–1391.
- Fillet M, Bentires-Alj M, Derogowski V, Greimers R, Gielen J, Piette J *et al.* (2003). Mechanisms involved in exogenous C2- and C6-ceramide-induced cancer cell toxicity. *Biochem Pharmacol* 65: 1633–1642.
- Fingrut O, Flescher E (2002). Plant stress hormones suppress the proliferation and induce apoptosis in human cancer cells. *Leukemia* 16: 608–616.
- Fingrut O, Reischer D, Rotem R, Goldin N, Altboum I, Zan-Bar I *et al.* (2005). Jasmonates induce nonapoptotic death in high-resistance mutant p53-expressing B-lymphoma cells. *Br J Pharmacol* 146: 800–808.
- Flowers M, Fabrias G, Delgado A, Casas J, Abad JL, Cabot MC (2012). C6-ceramide and targeted inhibition of acid ceramidase induce synergistic decreases in breast cancer cell growth. *Breast Cancer Res Treat* 133: 447–458.
- Fulda S, Debatin KM (2006). Extrinsic versus intrinsic apoptosis pathways in anticancer chemotherapy. *Oncogene* 25: 4798–4811.
- Gajate C, Santos-Beneit M, Macho A, del Carmen Lazaro M, Hernandez-De Rojas A, Modolell M *et al.* (2000). Involvement of mitochondria and caspase-3 in Et-18-OCH₃-induced apoptosis of human leukemic cells. *Int J Cancer* 86: 208–218.
- Galiegue S, Jbilo O, Combes T, Bribeis E, Carayon P, Le Fur G *et al.* (1999). Cloning and characterization of PRAX-1. A new protein that specifically interacts with the peripheral benzodiazepin receptor. *J Biol Chem* 274: 2938–2952.
- Garcia-Ruiz C, Colell A, Mari M, Morales A, Fernandez-Checa JC (1997). Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. Role of mitochondrial glutathione. *J Biol Chem* 272: 11369–11377.
- Gasmi J, Sanderson JT (2010). Growth inhibitory, antiandrogenic, and pro-apoptotic effects of puniceic acid in LNCaP human prostate cancer cells. *J Agric Food Chem* 58: 12149–12156.
- Gasmi J, Sanderson JT (2013). Jacaric acid and its octadecatrienoic acid geoisomers induce apoptosis selectively in cancerous human prostate cells: a mechanistic and 3-D structure-activity study. *Phytomedicine* 20: 734–742.
- Ghosh J, Myers CE (1998). Inhibition of arachidonate 5-lipoxygenase triggers massive apoptosis in human prostate cancer cells. *Proc Natl Acad Sci USA* 95: 13182–13187.
- Goldin N, Heyfets A, Reischer D, Flescher E (2007). Mitochondria-mediated ATP depletion by anti-cancer agents of the jasmonate family. *J Bioenerg Biomemb* 39: 51–57.
- Goldin N, Arzoin L, Heyfets A, Israelson A, Zaslavsky Z, Bravman T *et al.* (2008). Methyl jasmonate binds to and detaches mitochondria-bound hexokinase. *Oncogene* 27: 4636–4643.
- Goodwin PJ, Boyd NF, Hanna W, Hartwick W, Murray D, Qizilbash A *et al.* (1997). Elevated levels of plasma triglycerides are associated with histologically defined premenopausal breast cancer risk. *Nutr Cancer* 27: 284–292.
- Grossmann ME, Mizuno NK, Dammen ML, Schuster T, Ray A, Cleary MP (2009). Eleostearic acid inhibits breast cancer proliferation by means of an oxidation-dependent mechanism. *Cancer Prev Res* 2: 879–886.
- Grossmann ME, Mizuno NK, Schuster T, Cleary MP (2010). Puniceic acid is an omega-5 fatty acid capable of inhibiting breast cancer proliferation. *Int J Oncol* 36: 421–426.
- Gudz TI, Tserng KY, Hoppel CL (1997). Direct inhibition of mitochondrial respiratory chain complex III by cell-permeable ceramide. *J Biol Chem* 272: 24154–24158.
- Gulaya NM, Melnik AA, Balkov DI, Volkov GL, Vysotskiy MV, Vaskovsky VE (1993). The effect of long-chain N-acylethanolamines on some membrane-associated functions of neuroblastoma C1300 N18 cells. *Biochim Biophys Acta* 1152: 280–288.
- Gulaya NM, Kuzmenko AI, Margitich VM, Govseeva NM, Melnichuk SD, Goridko TM *et al.* (1998). Long-chain N-acylethanolamines inhibit lipid peroxidation in rat liver mitochondria under acute hypoxic hypoxia. *Chem Phys Lipids* 97: 49–54.
- Gutknecht J (1988). Proton conductance caused by long-chain fatty acids in phospholipid bilayer membranes. *J Membrane Biol* 106: 83–93.
- Hague A, Diaz GD, Hicks DJ, Krajewski S, Reed JC, Paraskeva C (1997). Bcl-2 and bak may play a pivotal role in sodium butyrate-induced apoptosis in colonic epithelial cells; however overexpression of bcl-2 does not protect against bak-mediated apoptosis. *Int J Cancer* 72: 898–905.
- Hanahan D, Weinberg RA (2011). Hallmarks of cancer: the next generation. *Cell* 144: 646–674.
- Heerdts BG, Houston MA, Augenlicht LH (1997). Short-chain fatty acid initiated cell cycle arrest and apoptosis of colonic epithelial cells is linked to mitochondrial function. *Cell Growth Differ* 8: 523–532.
- Heerdts BG, Houston MA, Anthony GM, Augenlicht LH (1998). Mitochondrial membrane potential ($\Delta\Psi$) in the coordination of

- p53-independent proliferation and apoptosis pathways in human colonic carcinoma cells. *Cancer Res* 58: 2869–2875.
- Heyfets A, Flescher E (2007). Cooperative cytotoxicity of methyl jasmonate with anti-cancer drugs and 2-deoxy-D-glucose. *Cancer Lett* 50: 300–310.
- Hideshima T, Catley L, Yasui H, Ishitsuka K, Raje N, Mitsiades C *et al.* (2006). Perifosine, an oral bioactive novel alkylphospholipid, inhibits akt and induces *in vitro* and *in vivo* cytotoxicity in human multiple myeloma cells. *Blood* 107: 4053–4062.
- Higdon A, Diers AR, Oh JY, Landar A, Darley-Usmar VM (2012). Cell signalling by reactive lipid species: new concepts and molecular mechanisms. *Biochem J* 442: 453–464.
- Hildebrand M, Staks T, Schuett A, Matthes H (1989). Pharmacokinetics of ³H-cicaprost in healthy volunteers. *Prostaglandins* 37: 259–273.
- Hoque A, Lippman SM, Wu TT, Xu Y, Liang ZD, Swisher S *et al.* (2005). Increased 5-lipoxygenase expression and induction of apoptosis by its inhibitors in esophageal cancer: a potential target for prevention. *Carcinogenesis* 26: 785–791.
- Hostetler KY, van den Bosch H (1972). Subcellular and submitochondrial localization of the biosynthesis of cardiolipin and related phospholipids in rat liver. *Biochim Biophys Acta* 260: 380–386.
- Hvattum E, Bergseth S, Pedersen CN, Bremer J, Aarsland A, Berge RK (1991). Microsomal oxidation of dodecylthioacetic acid (a 3-thia fatty acid) in rat liver. *Biochem Pharmacol* 41: 945–953.
- Ip C, Ip MM, Loftus T, Shoemaker S, Shea-Eaton W (2000). Induction of apoptosis by conjugated linoleic acid in cultured mammary tumor cells and premalignant lesions of the rat mammary gland. *Cancer Epidemiol Biomarkers Prev* 9: 689–696.
- Ip MM, Masso-Welch PA, Shoemaker SF, Shea-Eaton WK, Ip C (1999). Conjugated linoleic acid inhibits proliferation and induces apoptosis of normal rat mammary epithelial cells in primary culture. *Exp Cell Res* 250: 22–34.
- Ito K, Hirao A, Arai F, Takubo K, Matsuoka S, Miyamoto K *et al.* (2006). Reactive oxygen species act through p38 MAPK to limit the lifespan of hematopoietic stem cells. *Nat Med* 12: 446–451.
- Iversen PO, Sørensen DR, Tronstad KJ, Gudbrandsen OA, Rustan AC, Berge RK *et al.* (2006). A bioactively modified fatty acid improves survival and impairs metastasis in preclinical models of acute leukemia. *Clin Cancer Res* 12: 3525–3531.
- Jensen LR, Berge K, Bathen TF, Wergedahl H, Schonberg SA, Bofin A *et al.* (2007). Effect of dietary tetradecylthioacetic acid on colon cancer growth studied by dynamic contrast enhanced MRI. *Cancer Biol Ther* 6: 1810–1816.
- Jiang JG, Chen CL, Card JW, Yang S, Chen JX, Fu XN *et al.* (2005). Cytochrome P450 2J2 promotes the neoplastic phenotype of carcinoma cells and is up-regulated in human tumors. *Cancer Res* 65: 4707–4715.
- Kamp F, Hamilton JA (1992). pH gradients across phospholipid membranes caused by fast flip-flop of unionized fatty acids. *Proc Natl Acad Sci USA* 89: 11367–11370.
- Kanazawa A, Tanaka A, Iwata S, Satoh S, Hatano E, Shinohara H *et al.* (1998). The beneficial effect of phosphocreatine accumulation in the creatine kinase transgenic mouse liver in endotoxin-induced hepatic cell death. *J Surg Res* 80: 229–235.
- Kharbanda S, Saxena S, Yoshida K, Pandey P, Kaneki M, Wang Q *et al.* (2000). Translocation of SAPK/JNK to mitochondria and interaction with Bcl-x(L) in response to DNA damage. *J Biol Chem* 275: 322–327.
- Kim EJ, Holthuizen PE, Park HS, Ha YL, Jung KC, Park JH (2002). *Trans-10,cis-12*-conjugated linoleic acid inhibits Caco-2 colon cancer cell growth. *Am J Physiol Gastrointest Liver Physiol* 283: G357–G367.
- Kluck RM, Bossy-Wetzel E, Green DR, Newmeyer DD (1997). The release of cytochrome *c* from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science* 275: 1132–1136.
- Koehne CH, Dubois RN (2004). COX-2 inhibition and colorectal cancer. *Semin Oncol* 31 (2 Suppl 7): 12–21.
- Kondapaka SB, Singh SS, Dasmahapatra GP, Sausville EA, Roy KK (2003). Perifosine, a novel alkylphospholipid, inhibits protein kinase B activation. *Mol Cancer Ther* 2: 1093–1103.
- Korshunov SS, Korkina OV, Ruuge EK, Skulachev VP, Starkov AA (1998). Fatty acids as natural uncouplers preventing generation of O₂⁻ and H₂O₂ by mitochondria in the resting state. *FEBS Lett* 435: 215–218.
- Kroemer G, Galluzzi L, Brenner C (2007). Mitochondrial membrane permeabilization in cell death. *Physiol Rev* 87: 99–163.
- Kuwana T, Mackey MR, Perkins G, Ellisman MH, Latterich M, Schneider R *et al.* (2002). Bid, Bax, and lipids cooperate to form supramolecular openings in the outer mitochondrial membrane. *Cell* 111: 331–342.
- Larsen NB, Rasmussen M, Rasmussen LJ (2005). Nuclear and mitochondrial DNA repair: similar pathways? *Mitochondrion* 5: 89–108.
- Lin T, Yin X, Cai Q, Fan X, Xu K, Huang L *et al.* (2012). 13-Methyltetradecanoic acid induces mitochondrial-mediated apoptosis in human bladder cancer cells. *Urol Oncol* 30: 339–345.
- Locatelli SL, Giacomini A, Guidetti A, Cleris L, Mortarini R, Anichini A *et al.* (2013). Perifosine and sorafenib combination induces mitochondrial cell death and antitumor effects in NOD/SCID mice with Hodgkin lymphoma cell line xenografts. *Leukemia* 27: 1677–1687.
- Maccarrone M, Lorenzon T, Bari M, Melino G, Finazzi-Agro A (2000). Anandamide induces apoptosis in human cells via vanilloid receptors. Evidence for a protective role of cannabinoid receptors. *J Biol Chem* 275: 31938–31945.
- Maccarrone M, Melino G, Finazzi-Agro A (2001). Lipoxygenases and their involvement in programmed cell death. *Cell Death Diff* 8: 776–784.
- Machida K, Ohta Y, Osada H (2006). Suppression of apoptosis by cyclophilin D via stabilization of hexokinase II mitochondrial binding in cancer cells. *J Biol Chem* 281: 14314–14320.
- McLaughlin SGA, Dilger JP (1980). Transport of protons across membranes by weak acids. *Physiol Rev* 60: 825–863.
- Marden NY, Fiala-Beer E, Xiang SH, Murray M (2003). Role of activator protein-1 in the downregulation of the human *CYP2J2* gene in hypoxia. *Biochem J* 373: 669–680.
- Mari M, Morales A, Colell A, Garcia-Ruiz C, Fernandez-Checa JC (2009). Mitochondrial glutathione, a key survival antioxidant. *Antioxid Redox Signal* 11: 2685–2700.
- Marzo I, Brenner C, Zamzami N, Susin SA, Beutner G, Brdiczka D *et al.* (1998a). The permeability transition pore complex: a target for apoptosis regulation by caspases and bcl-2-related proteins. *J Exp Med* 187: 1261–1271.
- Marzo I, Brenner C, Zamzami N, Jurgensmeier JM, Susin SA, Vieira HL *et al.* (1998b). Bax and adenine nucleotide translocator cooperate in the mitochondrial control of apoptosis. *Science* 281: 2027–2031.

- Merad-Boudia M, Nicole A, Santiard-Baron D, Saillé C, Ceballos-Picot I (1998). Mitochondrial impairment as an early event in the process of apoptosis induced by glutathione depletion in neuronal cells: relevance to Parkinson's disease. *Biochem Pharmacol* 56: 645–655.
- Merrill AH Jr, Jones DD (1990). An update of the enzymology and regulation of sphingomyelin metabolism. *Biochim Biophys Acta* 1044: 1–12.
- Miccoli L, Beurdeley-Thomas A, De Pinieux G, Sureau F, Oudard S, Dutrillaux B *et al.* (1998). Light-induced photoactivation of hypericin affects the energy metabolism of human glioma cells by inhibiting hexokinase bound to mitochondria. *Cancer Res* 58: 5777–5786.
- Miller A, McGrath E, Stanton C, Devery R (2003). Vaccenic acid (t11-18:1) is converted to c9,t11-CLA in MCF-7 and SW480 cancer cells. *Lipids* 38: 623–632.
- Milovic V, Teller IC, Turchanowa L, Caspary WF, Stein J (2000). Effect of structural analogues of propionate and butyrate on colon cancer cell growth. *Int J Colorectal Dis* 15: 264–270.
- Mollinedo F, Fernandez-Luna JL, Gajate C, Martin-Martin B, Benito A, Martinez-Dalmau R *et al.* (1997). Selective induction of apoptosis in cancer cells by the ether lipid ET-18-OCH3 (Edelfosine): molecular structure requirements, cellular uptake, and protection by Bcl-2 and Bcl-X(L). *Cancer Res* 57: 1320–1328.
- Mollinedo F, Fernandez M, Hornillos V, Delgado J, Amat-Guerri F, Acuna AU *et al.* (2011). Involvement of lipid rafts in the localization and dysfunction effect of the antitumor ether phospholipid edelfosine in mitochondria. *Cell Death Dis* 2: e158.
- Murray M (1999). Mechanisms and significance of inhibitory drug interactions involving cytochrome P450 enzymes. *Int J Mol Med* 3: 227–238.
- Nathan C, Cunningham-Bussell A (2013). Beyond oxidative stress: an immunologist's guide to reactive oxygen species. *Nat Rev Immunol* 13: 349–361.
- Nieto-Miguel T, Gajate C, Mollinedo F (2006). Differential targets and subcellular localization of antitumor alkyl-lysophospholipid in leukemic versus solid tumor cells. *J Biol Chem* 281: 14833–14840.
- Nyäkern M, Cappellini A, Mantovani I, Martelli AM (2006). Synergistic induction of apoptosis in human leukemia T cells by the Akt inhibitor perifosine and etoposide through activation of intrinsic and Fas-mediated extrinsic cell death pathways. *Mol Cancer Ther* 5: 1559–1570.
- Oates JA, FitzGerald GA, Branch RA, Jackson EK, Knapp HR, Roberts LJ II (1988). Clinical implications of prostaglandin and thromboxane A₂ formation. *N Engl J Med* 319: 689–698.
- Oliw EH (1994). Oxygenation of polyunsaturated fatty acids by cytochrome P450 monooxygenases. *Prog Lipid Res* 33: 329–354.
- Ookhtens M, Kannan R, Lyon I, Baker N (1984). Liver and adipose tissue contributions to newly formed fatty acids in an ascites tumor. *Am J Physiol* 247: R146–R153.
- Ostrand DB, Sparagna GC, Amoscato AA, McMillin JB, Dowhan W (2001). Decreased cardiolipin synthesis corresponds with cytochrome c release in palmitate-induced cardiomyocyte apoptosis. *J Biol Chem* 276: 38061–38067.
- de Pablo MA, Susin SA, Jacotot E, Larochette N, Costantini P, Ravagnan L *et al.* (1999). Palmitate induces apoptosis via a direct effect on mitochondria. *Apoptosis* 4: 81–87.
- Petrosillo G, Ruggiero FM, Paradies G (2003). Role of reactive oxygen species and cardiolipin in the release of cytochrome c from mitochondria. *FASEB J* 17: 2202–2208.
- Pike LS, Smift AL, Croteau NJ, Ferrick DA, Wu M (2011). Inhibition of fatty acid oxidation by etomoxir impairs NADPH production and increases reactive oxygen species resulting in ATP depletion and cell death in human glioblastoma cells. *Biochem Biophys Acta* 1807: 726–734.
- Pizer ES, Chrest FJ, DiGiuseppe JA, Han WF (1998). Pharmacological inhibitors of mammalian fatty acid synthase suppress DNA replication and induce apoptosis in tumor cell lines. *Cancer Res* 58: 4611–4615.
- Pugazhenth S, Nesterova A, Sable C, Heidenreich KA, Boxer LM, Heasley LE *et al.* (2000). Akt/protein kinase B up-regulates Bcl-2 expression through cAMP-response element-binding protein. *J Biol Chem* 275: 10761–10766.
- Ramirez M, Amate L, Gil A (2001). Absorption and distribution of dietary fatty acids from different sources. *Early Hum Dev* 65: S95–S101.
- Rosignol R, Gilkerson R, Aggeler R, Yamagata K, Remington SJ, Capaldi RA (2004). Energy substrate modulates mitochondrial structure and oxidative capacity in cancer cells. *Cancer Res* 64: 985–993.
- Rotem R, Heyfets A, Fingrut O, Blickstein D, Shaklai M, Flescher E (2005). Jasmonates: novel anticancer agents acting directly and selectively on human cancer cell mitochondria. *Cancer Res* 65: 1984–1993.
- Ruemmele FM, Dionne S, Qureshi I, Sarma DS, Levy E, Seidman EG (1999). Butyrate mediates Caco-2 cell apoptosis via up-regulation of pro-apoptotic bak and inducing caspase-3 mediated cleavage of poly-(ADP-ribose) polymerase (PARP). *Cell Death Diff* 6: 729–735.
- Ruiter GA, Zerp SF, Bartelink H, van Blitterswijk WJ, Verheij M (1999). Alkyl-lysophospholipids activate the SAPK/JNK pathway and enhance radiation-induced apoptosis. *Cancer Res* 59: 2457–2463.
- Ruiter GA, Zerp SF, Bartelink H, van Blitterswijk WJ, Verheij M (2003). Anti-cancer alkyl-lysophospholipids inhibit the phosphatidylinositol 3-kinase-Akt/PKB survival pathway. *Anticancer Drugs* 14: 167–173.
- Sabine JR, Abraham S, Chaikoff IL (1967). Control of lipid metabolism in hepatomas: insensitivity of rate of fatty acid and cholesterol synthesis by mouse hepatoma BW7756 to fasting and to feedback control. *Cancer Res* 27: 793–799.
- Samudio I, Fiegl M, Andreeff M (2009). Mitochondrial uncoupling and the Warburg effect: molecular basis for the reprogramming of cancer cell metabolism. *Cancer Res* 69: 2163–2166.
- Sarker KP, Obara S, Nakata M, Kitajima I, Maruyama I (2000). Anandamide induces apoptosis of PC-12 cells: involvement of superoxide and caspase-3. *FEBS Lett* 472: 39–44.
- Schlame M, Rua D, Greenberg ML (2000). The biosynthesis and functional role of cardiolipin. *Prog Lipid Res* 39: 257–288.
- Schmid PC, Wold LE, Krebsbach RJ, Berdyshev EV, Schmid HH (2002). Anandamide and other N-acyl ethanolamines in human tumours. *Lipids* 37: 907–912.
- Schwarz H, Blanco FJ, Lotz M (1994). Anandamide, an endogenous cannabinoid receptor agonist inhibits lymphocyte proliferation and induces apoptosis. *J Neuroimmunol* 55: 107–115.
- Shimizu S, Narita M, Tsujimoto Y (1999). Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. *Nature* 399: 483–487.
- Shinohara N, Tsuduki T, Ito J, Honma T, Kijima R, Sugawara S *et al.* (2012). Jacaric acid, a linolenic acid isomer with a conjugated triene system, has a strong antitumor effect *in vitro* and *in vivo*. *Biochim Biophys Acta* 1821: 980–988.

- Skorski T, Bellacosa A, Nieborowska-Skorska M, Majewski M, Martinez R, Choi JK *et al.* (1997). Transformation of hematopoietic cells by BCR/ABL requires activation of a PI-3k/Akt-dependent pathway. *EMBO J* 16: 6151–6161.
- Skulachev VP (1998). Uncoupling: new approaches to an old problem of bioenergetics. *Biochim Biophys Acta* 1363: 100–124.
- Spector AA, Yorek MA (1985). Membrane lipid composition and cellular function. *J Lipid Res* 26: 1015–1035.
- Susin SA, Lorenzo HK, Zamzami N, Marzo I, Snow BE, Brothers GM *et al.* (1999). Molecular characterization of mitochondrial apoptosis-inducing factor. *Nature* 397: 441–446.
- Szymczak M, Murray M, Petrovic N (2008). Modulation of angiogenesis by ω -3 polyunsaturated fatty acids is mediated by cyclooxygenases. *Blood* 111: 3514–3521.
- Trachootham D, Zhou Y, Zhang H, Demizu Y, Chen Z, Pelicano H *et al.* (2006). Selective killing of oncogenically transformed cells through a ROS-mediated mechanism by β -phenylethyl isothiocyanate. *Cancer Cell* 10: 241–252.
- Tronstad KJ, Berge K, Dyroy E, Madsen L, Berge RK (2001). Growth reduction in glioma cells after treatment with tetradecylthioacetic acid: changes in fatty acid metabolism and oxidative status. *Biochem Pharmacol* 61: 639–649.
- Tronstad KJ, Gjertsen BT, Krakstad C, Berge K, Brustugun OT, Doskeland SO *et al.* (2003). Mitochondrial-targeted fatty acid analog induces apoptosis with selective loss of mitochondrial glutathione in promyelocytic leukemia cells. *Chem Biol* 10: 609–618.
- Tsuji M, Kawano S, DuBois RN (1997). Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc Natl Acad Sci USA* 94: 3336–3340.
- Tsuzuki T, Tokuyama Y, Igarashi M, Miyazawa T (2004). Tumor growth suppression by alpha-eleostearic acid, a linolenic acid isomer with a conjugated triene system, via lipid peroxidation. *Carcinogenesis* 25: 1417–1425.
- Veber DF, Johnson SR, Cheng HY, Smith BR, Ward KW, Kopple KD (2002). Molecular properties that influence the oral bioavailability of drug candidates. *J Med Chem* 45: 2615–2623.
- Venturini I, Zeneroli ML, Corsi L, Avallone R, Farina F, Alho H *et al.* (1998). Up-regulation of peripheral benzodiazepine receptor system in hepatocellular carcinoma. *Life Sci* 63: 1269–1280.
- Vrablic AS, Albright CD, Craciunescu CN, Salganic RI, Zeisel SH (2001). Altered mitochondrial function and overgeneration of reactive oxygen species precede the induction of apoptosis by 1-O-octadecyl-2-O-methyl-*rac*-glycero-3-phosphocholine in p53-defective hepatocytes. *FASEB J* 15: 1739–1744.
- Wasilewski M, Wieckowski MR, Dymkowska D, Wojtczak L (2004). Effects of N-acylethanolamines on mitochondrial energetics and permeability transition. *Biochim Biophys Acta* 1657: 151–163.
- Wieckowski MR, Wojtczak L (1998). Fatty acid-induced uncoupling of oxidative phosphorylation is partly due to opening of the mitochondrial permeability transition pore. *FEBS Lett* 423: 339–342.
- Wongtangintharn S, Oku H, Iwasaki H, Inafuku M, Toda T, Yanagita T (2005). Incorporation of branched-chain fatty acid into cellular lipids and caspase-independent apoptosis in human breast cancer cell line, SKBR-3. *Lipids Health Dis* 4: 29.
- Woodfield K, Ruck A, Brdiczka D, Halestrap AP (1998). Direct demonstration of a specific interaction between cyclophilin-D and the adenine nucleotide translocase confirms their role in the mitochondrial permeability transition. *Biochem J* 336: 287–290.
- Yang QH, Church-Hajduk R, Ren J, Newton ML, Du C (2003). Omi/HtrA2 catalytic cleavage of inhibitor of apoptosis (IAP) irreversibly inactivates IAPs and facilitates caspase activity in apoptosis. *Genes Dev* 17: 1487–1496.
- Yang Z, Liu S, Chen X, Chen H, Huang M, Zheng J (2000). Induction of apoptotic cell death and *in vivo* growth inhibition of human cancer cells by a saturated branched-chain fatty acid, 13-methyltetradecanoic acid. *Cancer Res* 60: 505–509.
- Zaugg K, Yao Y, Reilly PT, Kannan K, Kiarash R, Mason J *et al.* (2011). Carnitine palmitoyltransferase 1C promotes cell survival and tumor growth under conditions of metabolic stress. *Genes Dev* 25: 1041–1051.
- Zborowski J, Wojtczak L (1969). Phospholipid synthesis in rat-liver mitochondria. *Biochim Biophys Acta* 187: 73–84.
- Zhang G, Panigrahy D, Mahakian LM, Yang J, Liu JY, Stephen Lee KS *et al.* (2013). Epoxy metabolites of docosahexaenoic acid (DHA) inhibit angiogenesis, tumor growth, and metastasis. *Proc Natl Acad Sci USA* 110: 6530–6535.