

Published in final edited form as:

*Org Lett.* 2014 January 17; 16(2): 366–369. doi:10.1021/ol403186k.

## Absorption of atmospheric CO<sub>2</sub> as carbonate inside the molecular cavity of a new tripodal hexaurea receptor

 Avijit Pramanik<sup>1</sup>, Maryam Emami Khansari<sup>1</sup>, Douglas R. Powell<sup>2</sup>, Frank R. Fronczek<sup>3</sup>, and Md Alamgir Hossain<sup>1</sup>


Alamgir Hossain: alamgir.hossain@jsums.edu

<sup>1</sup>Department of Chemistry and Biochemistry, Jackson State University, Jackson, MS 39217

<sup>2</sup>Department of Chemistry and Biochemistry, University of Oklahoma, Norman, OK 73019

<sup>3</sup>Department of Chemistry, Louisiana State University, Baton Rouge, LA 70803

### Abstract



A new hexaurea receptor has been synthesized, which absorbs atmospheric CO<sub>2</sub> to produce an air-stable solid carbonate complex under normal conditions. Structural analysis of the carbonate complex with this receptor suggests that the carbonate is fully encapsulated within its highly organized intramolecular cavity *via* twelve strong NH···O bonds in the range of 2.703(3) – 2.989(3) Å from six urea units, with each anionic oxygen coordinated *via* four NH···O bonds with two urea groups.

Continuous increase of carbon dioxide (CO<sub>2</sub>) concentration in the atmosphere resulting from various human activities poses a threat to global warming.<sup>1</sup> CO<sub>2</sub> is also a major source of water acidification by dissolving in water to form carbonic acid (H<sub>2</sub>CO<sub>3</sub>).<sup>2</sup> Therefore, there is a current need to develop new chemical devices for the recycling of carbon dioxide to produce chemical commodities.<sup>3</sup> Recently epoxide-based molecules have been used to react with CO<sub>2</sub> in the presence of a catalyst to produce new polycarbonates and cyclic carbonates.<sup>4</sup> Gale *et al.* reported that simple mono-functional urea-based neutral compounds are capable of absorbing CO<sub>2</sub> in the presence of primary aliphatic amines to form carbamates [ $>N(CO_2)-$ ].<sup>5</sup> Increasing the functional groups, tren-based *tris*-ureas/thioureas,<sup>6</sup> such as pentafluorophenyl substituted *tris*-urea,<sup>6a</sup> *m*-nitrophenyl substituted *tris*-urea,<sup>6b</sup> phenyl substituted *tris*-thiourea,<sup>6c</sup> and *p*-cyanophenyl substituted *tris*-thiourea<sup>6d</sup> have recently been shown to bind carbonate anions, forming 2:1 (host:guest) complexes. Although the carbonate is coordinated *via* 12 hydrogen bonds (NH···O < 3.2 Å) in most

Correspondence to: Alamgir Hossain, alamgir.hossain@jsums.edu.

 Supporting Information available: Crystallographic file in CIF format, synthetic procedures and NMR titration spectra in pdf format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

cases, two receptors are required to provide complementary binding sites for the anion. Therefore, the complete coordination for a carbonate anion can ideally be achieved by a receptor possessing 12 complementary binding sites around a single cavity.

It is well-documented that increasing the effective binding sites in a host, leads to the enhancement of its binding ability for a guest due to the chelate effect.<sup>7</sup> From this viewpoint, we have been interested in synthesizing polyurea-functionalized receptors based on the commercially available 'tren' as a core. Herein, we report a highly organized hexaurea receptor possessing 12 H-bond donors, which absorbs atmospheric CO<sub>2</sub> in the form of carbonate encapsulated in a single cavity surrounded by perfectly arranged six urea units. Within this self-generated intramolecular cavity, the unique orientation of 12 binding sites provides an ideal complementarity for the trigonal planar carbonate anion.

The new hexaurea receptor **1** was synthesized by a three-step strategy (Scheme 1). The *tris*(2-aminoethyl) amine (tren) and *o*-nitrophenyl isocyanate were reacted to give a nitro-functionalized *tris*-urea **2** which was reduced with hydrazine hydrate and Pd/C (10%) to produce an amino-functionalized *tris*-urea **3**.<sup>8</sup> The final coupling was achieved by reacting **3** with pentafluorophenyl isocyanate **4** to form an *o*-xylyl bridged hexaurea **1**.

Single crystals of carbonate complex [1(CO<sub>3</sub>)](*n*-Bu<sub>4</sub>N)<sub>2</sub> were obtained from slow evaporation of a DMSO solution of **1** with an excess amount of *n*-tetrabutyl ammonium fluoride (TBAF) in a vial at room temperature in 2 days. The needle-like crystals were isolated by simple decantation technique, and were found to be air stable. Although, the primary goal was to obtain a fluoride complex, crystallographic analysis revealed that, instead of fluoride, one carbonate was encapsulated within the cavity of **1**. The bound carbonate was originated from atmospheric CO<sub>2</sub> absorbed by the basic TBAF-receptor solution which was left open to the atmosphere. The formation of carbonate in the presence of TBAF is well documented in the literature.<sup>6</sup>

Structural analysis from single crystal X-ray diffraction of the carbonate complex reveals that the complex crystallizes in an orthorhombic *Pna*2<sub>1</sub> space group to give a molecular formula, [1(CO<sub>3</sub>)](*n*-Bu<sub>4</sub>N)<sub>2</sub> in which one divalent carbonate is fully encapsulated.<sup>9</sup> Two tetrabutyl ammonium ions remain outside the cavity to balance the anionic charges. In the receptor unit, all protons on NH groups are pointed towards the cavity, and serve as H-bond donors for the bound anion.

As shown in Figure 1, the encapsulated carbonate is held tightly inside the cavity by strong 12 NH...O bonds ( $d_{\text{NH}\cdots\text{O}} = 2.703(3) - 2.989(3) \text{ \AA}$ ) from 12 NH groups of urea units (Table 1). Each oxygen of the carbonate is coordinated to four NH's from two different arms of the hexaurea receptor, with the two inner NH's from one arm and two outer NH's from other arm. Within a single arm of the receptor, the pentafluorophenyl and *o*-xylyl rings linked with an outer urea group, are almost perpendicular to each other, while these two aromatic rings from two different arms are stacked together *via*  $\pi\cdots\pi$  or C—H... $\pi$  interactions ( $\pi_c\cdots\pi_c = 3.602 \text{ \AA}$ , C<sub>8E</sub>—H... $\pi = 3.841 \text{ \AA}$ ; and C<sub>5E</sub>—H... $\pi = 3.855 \text{ \AA}$ ). Such arrangements of the aromatic rings make the receptor preorganized for the complete participation of all six ureas in coordinating the internal anion. The space-filling view of the complex (Figure 1b), illustrates the encapsulated carbonate inside the cavity of **1**, showing the stacking of the aromatic groups. The trigonal planar carbonate is almost perpendicular to the axis of the tertiary nitrogen (N4C) of **1** and the carbon (C1D) of CO<sub>3</sub><sup>2-</sup>, forming a pseudo C<sub>3</sub>-symmetric complex (Figure 1c).<sup>10</sup>

Proton NMR titrations were performed to examine the stability and stoichiometry of the receptor-carbonate complex in DMSO-*d*<sub>6</sub> using Et<sub>4</sub>NHCO<sub>3</sub>. Upon the addition of

Et<sub>4</sub>NHCO<sub>3</sub> (20 mM) to the receptor (2 mM), a new set of NMR spectrum appeared due to slow exchange on the NMR time scale (Figure 2).<sup>11</sup> All four NH signals were shifted significantly to downfield ( $\Delta\delta_{\text{NHa}} = 0.41$  ppm,  $\Delta\delta_{\text{NHb}} = 1.88$  ppm,  $\Delta\delta_{\text{NHc}} = 1.53$  ppm and  $\Delta\delta_{\text{NHd}} = 1.08$  ppm), indicating the interactions of all NH groups with the anion. Presumably, the chelation of the bound anion, as also observed in the solid state structure of the complex of **1**, results into the formation of a kinetically stable complex on the NMR time scale.<sup>8</sup> The relative change in the integration intensity NH resonances of the **1**-bicarbonate complex and the free **1**, allowed us to determine the binding constant (Figure 3).<sup>11</sup> The experimental data provided the best fit to a 1:1 (host: guest) binding model,<sup>12</sup> yielding a binding constant  $K = 1780 \text{ M}^{-1}$ . The 1:1 binding in solution was further supported by a Job's plot analysis (Figure S17). It is noted that, because of the unavailability of a suitable DMSO soluble CO<sub>3</sub><sup>2-</sup> salt, the HCO<sub>3</sub><sup>-</sup> as a tetraethyl ammonium (Et<sub>4</sub>N<sup>+</sup>) salt was used in the NMR titration studies, as previously used by other groups<sup>6</sup> for tren-based ligands. Thus, the determined binding constant ( $K = 1780 \text{ M}^{-1}$ ) is the result of the interactions of **1** with singly charged HCO<sub>3</sub><sup>-</sup> as opposed to CO<sub>3</sub><sup>2-</sup> observed in the crystal. The time dependent NMR spectra of **1** and Et<sub>4</sub>NHCO<sub>3</sub> in DMSO-*d*<sub>6</sub> showed no change in the NMR signals, suggesting that HCO<sub>3</sub><sup>-</sup> was not deprotonated to form CO<sub>3</sub><sup>2-</sup> during the titration process (Figure S18).

The binding affinity of **1** for Et<sub>4</sub>NHCO<sub>3</sub> was compared with its precursor *tris*-urea receptor **2** (Scheme 1) under similar conditions in DMSO-*d*<sub>6</sub>. The results show that this receptor binds HCO<sub>3</sub><sup>-</sup> with  $K = 226 \text{ M}^{-1}$  (Figure S24 in ESI), which is much weaker than  $1780 \text{ M}^{-1}$  observed for **1**. An *n*-butyl-functionalized *tris*-thiourea reported previously was shown to bind bicarbonate with  $K = 564 \text{ M}^{-1}$  in DMSO-*d*<sub>6</sub>.<sup>6c</sup> This observation further demonstrates that the unique orientation of the binding sites in the hexaurea **1**, makes it an excellent binder for the carbonate anions due to the possible chelate effect observed in traditional transition metal complexes.<sup>7b</sup> We also performed <sup>1</sup>H NMR titrations of **1** for other anions, showing strong affinity for HSO<sub>4</sub><sup>-</sup> ( $2250 \text{ M}^{-1}$ ). The association constants of **1** with Cl<sup>-</sup> and CH<sub>3</sub>COO<sup>-</sup> were 256 and 200  $\text{M}^{-1}$ , respectively. However, the receptor showed no significant binding for I<sup>-</sup>, ClO<sub>4</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup>. Determination of binding constants for F<sup>-</sup> and H<sub>2</sub>PO<sub>4</sub><sup>-</sup> was hampered due to the broadening of NMR peaks.

The formation of the complex in DMSO-*d*<sub>6</sub> was further evaluated by a series of <sup>13</sup>C NMR spectra (Figure 4). Partial <sup>13</sup>C NMR of Et<sub>4</sub>NHCO<sub>3</sub> and free **1** are shown in Figure 4a and Figure 4b, respectively. The sharp signal at 157.19 ppm in the free Et<sub>4</sub>NHCO<sub>3</sub> shifted to 168.38 ppm ( $\Delta\delta = 11.19$  ppm) after the addition of one equivalent of the ligand (Figure 4c), indicating the encapsulation of HCO<sub>3</sub><sup>-</sup> in the receptor's cavity.<sup>6a</sup> Figure 4d displays the <sup>13</sup>C NMR of [1(CO<sub>3</sub>)](*n*-Bu<sub>4</sub>N)<sub>2</sub> used for single crystal diffraction, showing a peak on more downfield at 176.7 ppm for CO<sub>3</sub><sup>2-</sup>. This difference could be due to the charges (HCO<sub>3</sub><sup>-</sup> vs CO<sub>3</sub><sup>2-</sup>) and the counter cations (Et<sub>4</sub>N<sup>+</sup> vs *n*-Bu<sub>4</sub>N<sup>+</sup>) present in the two different samples.

The solid state FT-IR analysis was also performed to examine the interactions of the receptor with carbonate. The free receptor shows two peaks at 1650  $\text{cm}^{-1}$  and 1600  $\text{cm}^{-1}$  for two carbonyl groups (-C=O),<sup>13</sup> which shift to higher wave numbers at 1696  $\text{cm}^{-1}$  ( $\Delta\nu_{\text{CO}} = 46 \text{ cm}^{-1}$ ) and 1630  $\text{cm}^{-1}$  ( $\Delta\nu_{\text{CO}} = 30 \text{ cm}^{-1}$ ), respectively (Figure S15). Further, the significant upward shift ( $\Delta\nu_{\text{N-H}} = 74 \text{ cm}^{-1}$ ) of broad NH's stretching frequency from 3330  $\text{cm}^{-1}$  (free **1**) to 3404  $\text{cm}^{-1}$  (complex) was observed, suggesting the strong N—H...O interactions between NH groups and the carbonate.

The ability of **1** to uptake of the atmospheric CO<sub>2</sub> was examined by dissolving the receptor with TBAF in DMSO and keeping the mixture in open air (Figure 5). The solid products deposited at the interface of the liquid and air, were collected to give almost 99% carbonate complex. The complex was characterized by <sup>1</sup>H NMR and also verified from X-ray

diffraction analysis (CCDC 969272), giving an identical structure described in the preceding section. Treating the complex with water and methanol (1:1, v/v) gave over 95% recovery of **1**. In previous reports, a pentafluorophenyl substituted *tris*-urea,<sup>6a</sup> and a *m*-nitrophenyl substituted *tris*-urea<sup>6b</sup> were shown to fix atmospheric CO<sub>2</sub>, forming 98% and 90% complexes, respectively.

In conclusion, we have developed a novel receptor with six urea groups possessing an intramolecular cavity that is suitable to absorb atmospheric CO<sub>2</sub> as a CO<sub>3</sub><sup>2-</sup> under mild basic conditions. The highly functionalized receptor molecule utilizes all twelve NH-binding sites to interact with the internal carbonate anion, forming an air-stable complex. The solid state structure described herein represents an exceptional example of the encapsulated carbonate in a single molecule providing an ideal complementarity of binding sites for the anion. Given the significant role of CO<sub>2</sub> in environmental, chemical and biological systems,<sup>3,14</sup> the present findings hold promise for developing novel chemical devices for the recycling of atmospheric carbon dioxide.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgments

The National Science Foundation is acknowledged for a CAREER award (CHE-1056927) to MAH. NMR core facility at JSU was supported by the National Institutes of Health (G12RR013459).


## References

- (a) Hansen J, Nazarenko L, Ruedy R, Sato M, Willis J, Genio AD, Koch D, Lacis A, Lo K, Menon S, Novakov T, Perlwitz J, Russell G, Schmidt GA, Tausnev N. *Science*. 2005; 308:1431. [PubMed: 15860591] (b) *Climate Change 2007: Synthesis Report*, International Panel on Climate Change. Cambridge University Press; Cambridge, UK: 2007.
- National Research Council. *Ocean Acidification: A National Strategy to Meet the Challenges of a Changing Ocean*. Washington, D.C., USA: The National Academies Press; 2010. p. 1
- Das C, Gomes N, Jacquet O, Villiers C, Thuéry P, Ephritikhine M, Cantat T. *Angew Chem, Int Ed*. 2012; 51:187.
- (a) Darensbourg DJ. *Chem Rev*. 2007; 107:2388. [PubMed: 17447821] (b) Sakakura T, Kohno K. *Chem Commun*. 2009:1312.
- (a) Edwards PR, Hiscock JR, Gale PA. *Tetrahedron Lett*. 2009; 50:4922.(b) Edwards PR, Hiscock JR, Gale PA, Light ME. *Org Biomol Chem*. 2010; 8:100. [PubMed: 20024139]
- (a) Ravikumar I, Ghosh P. *Chem Commun*. 2010; 46:1082.(b) Dey SK, Chutia R, Das G. *Inorg Chem*. 2012; 51:1727. [PubMed: 22233387] (c) Busschaert N, Gale PA, Haynes CJE, Light ME, Moore SJ, Tong CC, Davis JT, Harrell JWA. *Chem Commun*. 2010; 46:6252.(d) Bose P, Dutta R, Santra S, Chowdhury B, Ghosh P. *Eur J Inorg Chem*. 2012:5791.
- (a) Anslyn, EV.; Dougherty, DA. *Modern Physical Organic Chemistry*. University Science Books; California: 2006. (b) Smith, RM.; Martell, AE. *Critical Stability Constants*. Plenum; New York: 1975.
- Jia C, Wu B, Li S, Huang X, Zhao Q, Li QS, Yang XJ. *Angew Chem, Int Ed*. 2011; 50:486.
- Crystal data for [1(CO<sub>3</sub>)]*[n*-Bu<sub>4</sub>N]<sub>2</sub>: *M* = 1720.82, *a* = 41.2339(11) Å, *b* = 13.6303(4) Å, *c* = 15.0005(4) Å, *α* = 90°, *β* = 90°, *γ* = 90°, *V* = 8430.7(4) Å<sup>3</sup>, *T* = 100(2) K, space group *Pna*2<sub>1</sub>, *Z* = 4, *μ*(MoK $\alpha$ ) = 0.112 mm<sup>-1</sup>, 85226 reflections measured, 19877 independent reflections (*R*<sub>int</sub> = 0.0294). The final *R*<sub>I</sub> values were 0.0360 (*I* > 2 $\sigma$ (*I*)). Flack parameter = 0.06(10). CCDC 963688.
- Isiklan M, Saeed MA, Pramanik A, Wong BM, Fronczek FR, Hossain MA. *Cryst Growth Des*. 2011; 11:959.


11. Hossain MA, Kang SO, Llinares JM, Powell DR, Bowman-James K. *Inorg Chem.* 2003; 42:5043. [PubMed: 12924873]
12. (a) Schneider HJ, Kramer R, Simova S, Schneider U. *J Am Chem Soc.* 1988; 110:6442.(b) Mendy JS, Pilate ML, Horne T, Day VW, Hossain MA. *Chem Commun.* 2010; 46:6084.
13. Custelcean R, Remy P, Bonnesen PV, Jiang DE, Moyer BA. *Angew Chem Int Ed.* 2008; 47:1866.
14. Aresta M, Dibenedetto A. *Dalton Trans.* 2007:2975. [PubMed: 17622414]



**Figure 1.** (a) Single crystal X-ray structure of the carbonate complex of **1** (counter cations and hydrogen atoms on carbons are omitted for clarity), (b) space filling model of encapsulated carbonate complex, (c) partial view of pseudo  $C_3$ -symmetric arrangement of tripodal cavity with the encapsulated carbonate.




**Figure 2.** Partial  $^1\text{H}$  NMR spectra of **1** with an increasing amount of  $\text{Et}_4\text{NHCO}_3$  ( $R = [\text{Et}_4\text{NHCO}_3]_0 / [\mathbf{1}]_0$ ) in  $\text{DMSO-}d_6$ .




**Figure 3.** <sup>1</sup>H NMR titration curve of **1** (2 mM) with an increasing amount of Et<sub>4</sub>NHCO<sub>3</sub> ( $R = [Et_4NHCO_3]_0/[1]_0$ ) in DMSO-*d*<sub>6</sub>. The titration plot was obtained from the relative change in the integration intensity  $\Delta\phi$  ( $\Delta\phi = I_{NHc}/[I_{NHf} + I_{NHc}]$ , where  $I_{NHc}$  is the intensity of NHa signal in the carbonate-**1** complex and  $I_{NHf}$  is the intensity of the corresponding NH signal in the free **1**).






**Figure 4.** Partial  $^{13}\text{C}$  NMR spectra (125 MHz,  $\text{DMSO-}d_6$ ) of (a)  $\text{Et}_4\text{NHCO}_3$ , (b) free receptor **1**, (c) **1** in the presence of one equivalent of  $\text{Et}_4\text{NHCO}_3$ , and (d) solid carbonate complex of the receptor ( $[\mathbf{1}(\text{CO}_3)](n\text{-Bu}_4\text{N})_2$ ).



**Figure 5.** The uptake of CO<sub>2</sub> and the recovery of **1**. Magnified **1**.carbonate crystals are shown at the right side.



**Scheme 1.**  
Synthesis of the hexaurea **1**.

**Table 1**H-bonding parameters (Å, °) for  $[\mathbf{1}(\text{CO}_3)]^{2-}$ .

<b>D—H...O</b>	<b>H...O</b>	<b>D...O</b>	<b>∠DHO</b>
N4A — H4A...O1D	2.20(3)	2.966(3)	152(3)
N7A — H7A...O1D	2.01(3)	2.852(2)	165(3)
N14A — H14A ... O2D	2.16(3)	2.935(3)	155(2)
N17A — H17A ... O2D	1.84(3)	2.703(3)	161(3)
N7B — H7B ... O2D	1.99(3)	2.869(3)	162(2)
N4B — H4B ... O2D	2.19(3)	2.949(3)	154(3)
N14B — H14B ... O3D	1.93(3)	2.801(3)	157(2)
N17B — H17B ... O3D	2.00(3)	2.788(3)	154(3)
N4C — H4C ... O3D	2.03(3)	2.822(3)	162(3)
N7C — H7C ... O3D	2.21(3)	2.989(3)	153(3)
N14C — H14C ... O1D	1.97(3)	2.783(2)	160(3)
N17C — H17C ... O1D	1.92(3)	2.746(3)	156(3)