

Published in final edited form as:

*Chem Commun (Camb)*. 2014 March 21; 50(23): 3071–3073. doi:10.1039/c4cc00432a.

## A photochemical method for determining plasma homocysteine with limited sample processing

Lovemore Hakuna, Jorge O. Escobedo, Mark Lowry, Aabha Barve, Naneki McCallum, and Robert M. Strongin\*

Department of Chemistry, Portland State University, Portland, Oregon 97207, USA

### Abstract

The photolytic formation of thiyl radicals allows for the selective detection of total homocysteine (tHcy) in plasma after reduction and filtering. The mechanism is based on the reduction of viologens by the  $\alpha$ -amino carbon centred radical of Hcy generated by intramolecular hydrogen atom transfer (HAT) of its thiyl radical.

---

Several major pathologies including cardiovascular disease (CVD), dementia, osteoporosis and Alzheimer's disease are associated with elevated levels of plasma tHcy.<sup>1,2</sup> Significant research has shown that Hcy is a risk factor at even modestly elevated levels.<sup>3</sup> Published evidence that shows an association between hyperhomocysteinemia ( $> 12 \mu\text{M tHcy}$ )<sup>4</sup> and major diseases renders its determination of clinical significance.<sup>5</sup>

Current commercial Hcy detection methods use separations, relatively fragile and expensive enzymatic or immunogenic materials and complex instrumentation.<sup>3, 6, 7</sup> Thus, there is need to develop selective, yet simple and inexpensive methods that can be used at point of care diagnostics to facilitate the diagnosis and treatment of related diseases. Available kits generally use multi-step washing procedures and/or specialized storage below  $-20^\circ\text{C}$  limiting their use in emerging nations with limited access to refrigeration or electricity. Moreover, even in developed countries point-of-care and kit-based assays are of interest considering rising health care costs and increasing interest in patient-based monitoring.

A wide variety of useful detection probes for biological thiols have been reported.<sup>8, 9</sup> Most have no specificity for Hcy over other related analytes such as cysteine (Cys) and glutathione (GSH). The Cys levels in human plasma from healthy individuals range from 135.8 to 266.5  $\mu\text{M}$ .<sup>10</sup> Consequently, they complicate the determination of plasma tHcy levels. Though some chemosensors or chemodosimeters that selectively respond to Hcy over Cys and other thiols have been reported, they are typically tested at equimolar, rather than more natural ca. 20-fold excess Cys concentrations.<sup>11</sup>

---

This journal is © The Royal Society of Chemistry [year]

strongin@pdx.edu; Fax: + 1 503-725-9525; Tel: + 1 503-725-9724.

†Electronic Supplementary Information (ESI) available: Detailed experimental procedures and additional absorption spectra data. See DOI: 10.1039/b000000x/

In 2004, we developed a selective colorimetric method for the detection of Hcy based on the kinetically-favored formation of  $\alpha$ -amino carbon centred radical for Hcy via a reversible intramolecular hydrogen atom transfer (HAT) with the corresponding thiyl radical.<sup>12</sup> This is attributed to favored formation of a 5-membered ring in the transition state, as opposed to 4- and 9-membered ring configurations for Cys and GSH, respectively (Scheme 1).

The mechanism shown in Scheme 1 was initially proposed and studied by Zhao *et al.*, under basic conditions (pH 10.5).<sup>13</sup> Azide radical was used to oxidize thiols and the formation of reducing radicals was monitored through the UV-Vis absorption spectra via production of the reduced methyl viologen radical cation ( $MV^{\bullet+}$ ). Under the highly basic conditions investigated by Zhao *et al.*, no colorimetric selectivity between GSH, Cys and Hcy was observed. This was due to the presence of significant amounts of thiolate anion promoting the formation of a reducing disulfide radical anion that also reacts with methyl viologen ( $MV^{2+}$ ) independently of the HAT mechanism. Conversely, neutral conditions investigated by us diminish thiolate formation, thereby enabling selective detection of Hcy in human blood plasma via its reducing carbon radical (Scheme 1).<sup>12,14</sup> A protocol for visual detection of Hcy was developed based on this method wherein the Hcy thiyl radical is generated by heat.<sup>15</sup> The colorimetric method was investigated using human serum calibration standards (NIST SRM 1955), and successfully distinguished micromolar concentration differences (3.79, 6.13, 13.4 and 38.73  $\mu\text{M}$ ) of tHcy visually using  $MV^{2+}$ .<sup>8</sup> The assay protocol involved no sample processing. It only required a two-fold dilution, addition of  $MV^{2+}$  and tris (2-carboxyethyl) phosphine (TCEP) and 2 min heating at reflux.

The basis of this current work is the hypothesis that photolytic methods would afford analogous selectivity via the intramolecular HAT mechanism while enabling the assay to be carried out at room temperature. Johnson and co-workers reported the photochemical reduction of viologens in ethanolic solutions.<sup>16</sup> A mechanism based on the abstraction of a methylene hydrogen atom from EtOH to form a free radical that reduced the viologen in direct sunlight was proposed. We envisioned that this approach could be compatible with our HAT mechanism for Hcy via photolytic, rather than thermal generation of the Hcy thiyl radical. Our hypothesis was confirmed by exposing solutions of thiols and  $MV^{2+}$  in Tris buffer at neutral pH to direct sunlight at room temperature. A blue color was observed within 2 minutes in the Hcy sample while other thiols solutions remained unchanged (Fig. 1).

To create a laboratory test, we reasoned that an appropriate light source to generate the thiyl radical should emit around 325 nm based on the reported S-H bond dissociation energy of Cys of 370 kJ/mol.<sup>17</sup> To this end, we selected a very simple and inexpensive compact fluorescent lamp emitting in this region, Reptisun™ consisting of 10% UVB and 30% UVA. The photolysis experiments were performed using this lamp with a light intensity of 6.85  $\text{mW}/\text{cm}^2$  as measured by a Melles Griot Broadband Power/Energy Meter 13PEM001. We were able to detect Hcy selectively in human blood plasma using  $MV^{2+}$  without any interference from Cys and GSH in the range of their physiological concentrations. Upon irradiation of plasma samples spiked with various biothiols, only the Hcy (15  $\mu\text{M}$ ) spiked sample showed significant absorption response whereas Cys and GSH remained unchanged. The absorption spectra are shown in Fig. S1, ESI†.

In previous studies, benzyl viologen ( $BV^{2+}$ ), a significantly less toxic chromogen than  $MV^{2+}$ , was found to be more reactive towards the Hcy  $\alpha$ -amino carbon-centred radical than  $MV^{2+}$  under thermal conditions.  $BV^{2+}$  has a higher reduction potential ( $-370$  mV) compared to  $MV^{2+}$  ( $-446$  mV).<sup>18</sup> Hence we investigated the response of  $BV^{2+}$  to Hcy and other thiols using the photochemical method.

$BV^{2+}$  indeed displayed selectivity towards Hcy under the new photolytic conditions compared to structurally related thiols (Fig. 2). Moreover, changing the chromogen allowed us to lower its concentration from 50 mM ( $MV^{2+}$ ) to 20 mM ( $BV^{2+}$ ). In addition, the Hcy response was greater as compared to photolysis in the presence of  $MV^{2+}$ . An irradiation time of 15 min afforded optimal selectivity and response.

A selective response of  $BV^{2+}$  to Hcy in human plasma was observed after reduction using immobilized tris(2-carboxyethyl) phosphine (TCEP gel), centrifugation and spiking with various thiols (Fig. S2, ESI<sup>†</sup>).

The proposed mechanism is analogous to the thermal reaction for the selective detection of Hcy we have previously reported.<sup>14</sup> It involves the generation of the Hcy thiyl radical by photolysis followed by the HAT reaction to form the  $\alpha$ -amino carbon centred radical that in turn, reduces  $BV^{2+}$  to its corresponding radical cation (Scheme 2).

We optimized the sample processing by replacing the centrifugation step with a simple filtration of the sample through 0.45  $\mu$ m PVDF filter vials. The spectral response and selectivity was comparable in both methods and excellent selectivity towards tHcy was observed (Fig.3). Interestingly, the use of filters improved the overall background interference from the plasma components.

The spectral responses of  $BV^{2+}$  to Hcy concentration changes, in spiked reduced human plasma monitored at 615 nm, increased linearly with increasing Hcy concentration over a physiologically relevant concentration range (Fig. 4). The inset shows the respective spectral data. To test the limits of possible Cys and GSH interference with the assay, further experiments were carried out with added excess amounts of Cys and GSH to reduced human plasma solutions. Cys and GSH were found to generate significant response only when their concentrations reach 400  $\mu$ M (almost double the normal concentration in healthy individuals) and 100  $\mu$ M (almost 20 times the normal plasma levels) respectively. At these concentrations, their absorbance response was equivalent to that of 5  $\mu$ M Hcy (Fig. S3, ESI<sup>†</sup>).

To further evaluate the selectivity of the  $BV^{2+}$  towards Hcy, control experiments using series of other amino acids were performed. Other amino acids produced no significant absorption response as compared to Hcy, further demonstrating that  $BV^{2+}$  is Hcy specific (Fig. 5).

In conclusion, we have developed a new, relatively simple and inexpensive assay for the selective detection of total Hcy directly in human blood plasma. This method has potential practical application in home test kits or point of care diagnostics because it involves the use of a less toxic chromogen, an inexpensive commercial light source, and simple sample

processing, involving only reduction and filtration prior to photolysis and UV-vis monitoring.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgement


This work was supported by the National Institutes of Health (R01 EB002044) and Boulder Diagnostics.

## Notes and references


1. Werder SF. *Neuropsychiatr. Dis. Treat.* 2010; 6:159–195. [PubMed: 20505848]
2. Glushchenko AV, Jacobsen DW. *Antioxid. Redox Signal.* 2007; 9:1883–1898. [PubMed: 17760510]
3. Kaluzna-Czaplinska J, Zurawicz E, Michalska M, Rynkowski J. *Acta Biochim. Pol.* 2013; 60:137–142. [PubMed: 23741716]
4. Marti-Carvajal AJ, Sola I, Lathyris D, Salanti G. *Cochrane Database Syst. Rev.* 2009
5. Smulders YM, Blom HJ. *J. Inherit. Metab. Dis.* 2011; 34:93–99. [PubMed: 20567905]
6. Newton LAA, Sandhu K, Livingstone C, Leslie R, Davis J. *Expert Rev. Mol. Diagn.* 2010; 10:489–500. [PubMed: 20465503]
7. Nekrassova O, Lawrence NS, Compton RG. *Talanta.* 2003; 60:1085–1095. [PubMed: 18969134]
8. Peng H, Chen W, Cheng Y, Hakuna L, Strongin R, Wang B. *Sensors.* 2012; 12:15907–15946. [PubMed: 23202239]
9. Jung HS, Chen XQ, Kim JS, Yoon J. *Chem. Soc. Rev.* 2013; 42:6019–6031. [PubMed: 23689799]
10. Jacobsen DW, Gatautis VJ, Green R, Robinson K, Savon SR, Secic M, Ji J, Otto JM, Taylor LM. *Clin. Chem.* 1994; 40:873–881. [PubMed: 8087981]
11. Yu C, Zeng F, Luo M, Wu S. *Nanotechnology.* 2012; 23
12. Wang WH, Escobedo JO, Lawrence CM, Strongin RM. *J. Am. Chem. Soc.* 2004; 126:3400–3401. [PubMed: 15025448]
13. Zhao R, Lind J, Merenyi G, Eriksen TE. *J. Am. Chem. Soc.* 1994; 116:12010–12015.
14. Wang WH, Rusin O, Xu XY, Kim KK, Escobedo JO, Fakayode SO, Fletcher KA, Lowry M, Schowalter CM, Lawrence CM, Fronczek FR, Warner IM, Strongin RM. *J. Am. Chem. Soc.* 2005; 127:15949–15958. [PubMed: 16277539]
15. Escobedo JO, Wang W, Strongin RM. *Nat. Protocols.* 2006; 1:2759–2762.
16. Johnson C Jr, Gutowsky H. *J. Chem. Phys.* 1963; 39:58.
17. Nauser T, Koppenol WH, Schöneich C. *J. Phys. Chem. B.* 2012; 116:5329–5341. [PubMed: 22483034]
18. Wang D, Crowe WE, Strongin RM, Sibrian-Vazquez M. *Chem. Commun.* 2009:1876–1878.




**Fig. 1.** Response of  $MV^{2+}$  towards various thiols upon exposure to direct sunlight. Solutions of  $MV^{2+}$  (50 mM) were mixed with thiols (20  $\mu$ M) in 0.5 M Tris buffer at pH 7, saturated with argon and exposed to sunlight. Pictures were taken 2 min after exposure to sunlight.




**Fig. 2.** Spectral response of BV<sup>2+</sup> towards various spiked thiols in human blood plasma upon irradiation. Absorption spectra of solutions of BV<sup>2+</sup> (20 mM) in human blood plasma and 0.5 M Tris buffer (pH 7.0) spiked with 1.5 μM Hcy, 25 μM Cys and 0.6 μM GSH. Plasma (10% v/v) was added to an argon-saturated solution of viologen, thiol & buffer and irradiated for 15 min using a Reptisun™ lamp.




**Fig. 3.** Spectral response of BV<sup>2+</sup> towards various spiked thiols in reduced human plasma upon irradiation. Absorption spectra of solutions of BV<sup>2+</sup> (20 mM) in reduced human blood plasma and 0.5 M Tris buffer (pH 7.0) spiked with 1.5  $\mu$ M Hcy, 25  $\mu$ M Cys and 0.6  $\mu$ M GSH. Plasma was incubated for 1 h with TCEP Gel followed by filtration using a *Single StEP*<sup>TM</sup> 0.45  $\mu$ m PVDF filter vial. The reduced, filtered plasma (10% v/v) was added to an argon-saturated solution of viologen, thiol and buffer and irradiated for 15 min using a Reptisun<sup>TM</sup> lamp.




**Fig. 4.** Spectral response of BV<sup>2+</sup> towards increasing levels of spiked Hcy in reduced human plasma upon irradiation. Absorption spectra of solutions of BV<sup>2+</sup> (20 mM) in 25% human blood plasma and 0.5 M Tris buffer (pH 7.0) spiked with 0–5 µM Hcy. Plasma (25% v/v) was added to an argon-saturated solution of viologen, thiol and buffer, and irradiated for 15 min using a Reptisun™ lamp.



**Fig. 5.** Spectral response of BV<sup>2+</sup> towards various spiked amino acids in human plasma upon irradiation. Absorption spectra of solutions of BV<sup>2+</sup> (20 mM) in human blood plasma and 0.5 M Tris buffer (pH 7.0) spiked with 1.5  $\mu$ M Hcy (dashed line) and 500  $\mu$ M amino acids (solid lines = L-Ala, Arg, Gln, Met, Ser, Thr & Phe). Plasma (10% v/v) was added to an argon-saturated solution of viologen, amino acids and buffer and irradiated for 15 min using a Reptisun<sup>TM</sup> lamp.



**Scheme 1.**  
Kinetically favored HAT reaction for Hcy.

**Scheme 2.**

Mechanism for the detection of Hcy involving three steps: (i) photolytic generation of Hcy thiyl radical, (ii) hydrogen atom transfer turning the thiyl radical into a  $\alpha$ -amino carbon-centred radical, (iii) reduction of BV<sup>2+</sup>.