

Review



Cite this article: Leggett HC, Brown SP, Reece SE. 2014 War and peace: social interactions in infections. *Phil. Trans. R. Soc. B* **369**: 20130365.
<http://dx.doi.org/10.1098/rstb.2013.0365>

One contribution of 14 to a Theme Issue 'Inclusive fitness: 50 years on'.

Subject Areas:

behaviour, evolution, health and disease and epidemiology, microbiology, theoretical biology

Keywords:

plasticity, transmission, virulence, relatedness, ecology, inclusive fitness

Author for correspondence:

Helen C. Leggett
e-mail: helen.leggett@zoo.ox.ac.uk

War and peace: social interactions in infections

Helen C. Leggett^{1,2}, Sam P. Brown^{3,4} and Sarah E. Reece^{3,4}

¹Department of Zoology, Oxford University, South Parks Road, Oxford OX1 3PS, UK

²Biosciences, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK

³Institute of Evolutionary Biology, and ⁴Centre for Immunity, Infection and Evolution, School of Biological Sciences, Ashworth Laboratories, University of Edinburgh, Edinburgh EH9 3JT, UK

One of the most striking facts about parasites and microbial pathogens that has emerged in the fields of social evolution and disease ecology in the past few decades is that these simple organisms have complex social lives, indulging in a variety of cooperative, communicative and coordinated behaviours. These organisms have provided elegant experimental tests of the importance of relatedness, kin discrimination, cooperation and competition, in driving the evolution of social strategies. Here, we briefly review the social behaviours of parasites and microbial pathogens, including their contributions to virulence, and outline how inclusive fitness theory has helped to explain their evolution. We then take a mechanistically inspired 'bottom-up' approach, discussing how key aspects of the ways in which parasites and pathogens exploit hosts, namely public goods, mobile elements, phenotypic plasticity, spatial structure and multi-species interactions, contribute to the emergent properties of virulence and transmission. We argue that unravelling the complexities of within-host ecology is interesting in its own right, and also needs to be better incorporated into theoretical evolution studies if social behaviours are to be understood and used to control the spread and severity of infectious diseases.

1. Introduction

Social acts, ranging from minor help to major self-sacrifice, are seen in all walks of life, from humans to microorganisms. It used to be generally assumed that the parasites and microbial pathogens that cause infectious diseases lived relatively independent unicellular lives, without the cooperative behaviours that have provoked interest in mammals, birds and insects [1]. However, a rapidly expanding body of research demonstrates that much of what parasites and microbial pathogens do, they do in groups. Furthermore, parasites and microbial pathogens display some amazing natural history, including behaviours described as mafia strategies, body-snatching, chemical warfare, mass suicide, suicide bombing and weapons of mass destruction (reviewed by [2–15]).

The expanding interest in understanding social evolution in parasites and microbial pathogens has probably occurred for two reasons. First, they are often well-described and tractable experimental systems for studying the ecology and evolution of social traits in real time, under both highly controlled conditions and in a 'real-world context', which for pathogens and parasites involves being exposed to the complex, changeable and hostile environments inside a host or vector. Second, sociality is a driver of the damage pathogens and parasites do to their hosts (virulence) [16,17], it shapes survival of medical interventions (such as antibiotics) [18,19], and underpins between-host transmission (e.g. [20,21]). Thus, examining the behaviours of parasites and microbial pathogens from the perspective of 'a life in society' is one of the most important issues in applied evolutionary biology. However, the mechanisms through which parasites and microbial pathogens interact with each other, the host/vector and the abiotic environment has been largely overlooked within the evolutionary/ecological search for general principles (and their

empirical support) of the often-connected theories for inclusive fitness and virulence evolution.

It is our view that a ‘bottom-up’ approach to studying microbial pathogens and parasites is the next milestone for understanding their social behaviours and for controlling the infectious diseases they cause. In this article, our aims are to showcase recent empirical and theoretical work demonstrating that a biologically informed bottom-up view: (i) illustrates the extremely rich phenotypic landscape of parasites and microbial pathogens at the within-host scale; (ii) enables integration across levels of biological organization, from the molecular mechanisms underpinning social behaviours to population ecology, to capture the biological complexity required to explain social systems; (iii) can provide novel insight into the evolution and ecology of social behaviours in general; and (iv) offers novel approaches to disease control with the potential to be more ‘evolution-proof’ than current therapies.

We also wish to facilitate cross-discipline communication between empirical and theoretical evolutionary ecologists and biologists in more applied disciplines such as microbiology, parasitology and biomedicine. To achieve this, we begin by providing an overview of the basic evolutionary and ecological frameworks for how social behaviours are studied (§2) and why virulence evolves (§3). The aim of these sections is to furnish readers unfamiliar with the social evolution and virulence evolution literatures with the concepts underpinning the recent developments that form the focus of the following sections. Therefore, aficionados in ecology may wish to skip to §4. The figures and tables illustrate the concepts we discuss in the text. Most examples concern malaria (*Plasmodium*) parasites and microbial pathogens (bacteria and bacteriophage) because these groups span the taxonomic diversity of infectious disease causing organisms, and together they offer the opportunity to integrate understanding at multiple levels of biological organization, from genes and molecular pathways, to phenotypes, to epidemiology [22,23]. For brevity when discussing general concepts, we collectively refer to parasites and microbial pathogens as ‘parasites’ owing to their shared lifestyle of exploiting hosts.

2. Social behaviours

All organisms interact with others throughout their lives, including with family members, unrelated conspecifics and hetero-specifics. Social interactions range from extreme conflict (e.g. lethal combat) to extreme cooperation (e.g. altruistic suicide or sterility) but most interactions lie somewhere between these extremes. Social behaviours can be categorized according to their impact on the lifetime reproductive success of the ‘actor’ expressing a particular social phenotype and any ‘recipients’ impacted by the actor’s phenotype (table 1) [24,48–50]. Taking a simple $+/-$ dichotomy for both actor and recipient gives a simple four-part categorization: (i) *mutual benefit* ($+/+$), where the actor and recipient both gain from the actor’s behaviour; (ii) *selfishness* ($+/-$), where the actor gains at the expense of the recipient; (iii) *altruism* ($-/+$), where the behaviour is detrimental to the actor but beneficial for the recipient and (iv) *spite* ($-/-$), where the behaviour is harmful for both actor and recipient. The pioneering work of Bill Hamilton [24,48] provided a foundation to explore how natural selection drives the spread of these four types of social behaviours through a population. The topics outlined below illustrate the key concepts involved in social interactions.

(a) Inclusive fitness: all for one and one for all

Hamilton’s key insight was that genes controlling the social traits of an actor can influence the replication of gene-copies in recipients. In the case of altruistic traits, Hamilton’s logic reveals a simple genetic nepotism—helping neighbours is another way of helping your own genes to reproduce, so long as they carry the helper-genes of interest. Hamilton proposed a critical metric to weight the likelihood that recipients carry the gene of interest, termed relatedness [51]. Common descent or kinship is the most common reason for interacting individuals to share genes with above-average frequency in a population. Consequently, relatedness can be understood as the chance of gene sharing among kin, above and beyond average probability [52]. Inclusive fitness partitions natural selection into direct and indirect effects; direct effects describe the impact of an individual’s own genes on reproductive success, and indirect effects describe the impact of the focal individual’s genes on the fitness of its social partners, weighted by genetic relatedness [24,26,48]. Cooperation may be mutually beneficial if it directly benefits the actor as well as the recipients, for example, by increasing the success of an individual’s own group (table 1). More extreme acts of altruistic cooperation may be selected if the behaviour helps recipients who are very likely to share the altruistic gene (i.e. if relatedness is high such as within families) [24,26,48]; thus indirectly propagating genes for altruism. An important point to note is that many parasite species reproduce asexually (i.e. clonally) during at least one stage of their life cycle [53], and thus each group of clonally related parasites (genotype) within an infection is expected to behave as a multicellular organism [54] because the genotype is the target of selection.

(b) Cheating: playing the system

When relatedness is low, cooperative behaviours are vulnerable to exploitation by cheats that do not contribute to collective action but still benefit from the cooperative behaviours of others [6]. Cheats can proliferate under these conditions because the benefits of cooperation are shared indiscriminately, and consequently genes for cheating will have greater fitness than the genes for cooperation [34,55]. The spread of cheats through a population can in turn lead to a decline in population fitness (an idea encapsulated by Hardin’s ‘Tragedy of the commons’ [56] and by ‘the Prisoners’ dilemma’ [57]). Empirical studies have demonstrated that cheating can indeed occur in numerous cooperative systems of microbial pathogens [7,34,58–60]. Recent years have witnessed a surge in the application of evolutionary theory to explain the ways in which cooperation is maintained (reviewed by [1,3,6,61–63]), which includes mechanisms for kin discrimination and communication.

(c) Kin recognition: deciding who to help

Relatedness is key to understanding the direction and magnitude of selection on social traits, but what shapes relatedness? A commonly cited scenario is that social acts are expressed blindly to neighbours, who tend to be relatives simply because of incomplete mixing of individuals in populations—the population is ‘viscous’ [24,48]. However, altruists in this system may fall victim to ‘cheats’ that lack the gene for altruism. A way to avoid wasting help on cheaters is to display an altruistic or social gene and to recognize the same gene in others, and

Table 1. A classification of social behaviours, after [24–26]. These examples illustrate that the richness of social behaviours observed in multicellular organisms are mirrored in parasites. Moreover, parasite social behaviours often have consequences for the severity and transmission of disease. Note that it is extremely difficult to quantify costs and benefits of many social behaviours, for actors and recipients, so many of these examples are yet to be fully understood.

effect on recipient	
effect on actor	+
<p>mutual benefit</p> <p><i>Multicellular taxa.</i> This ranges from simple cooperation providing safety in numbers, to rewarding helpers or punishing non-helpers. Group activities in the naked mole rat, <i>Heterocephalus glaber</i>, reduce the risks of predation, hypothermia and starvation [27–29]. Unrelated subordinate <i>Polistes</i> wasps cooperate to raise their colony's offspring because there is reasonable chance of inheriting the dominant's breeding position in the near future [30]. In this case, helpers 'pay to stay' because the acquisition of breeding opportunities is a large pay-off. <i>Parasites.</i> The release of secreted public goods molecules into a sufficiently small and/or structured population can generate mutual benefits, so long as the fraction of reward returned to the actor cell exceeds the direct costs of production, and that some reward is felt by cells other than the actor. Positive interactions between species have been documented. An existing infection of <i>Babesia microti</i> in field voles (<i>Microtus agrestis</i>) appears to increase the probability of <i>Anaplasma phagocytophilum</i> infection and vice-versa [31]. Also, if mixed infections are more challenging than single infections for host immune responses, mutual benefit could occur, which may result in greater virulence.</p>	<p>selfishness</p> <p><i>Multicellular taxa.</i> The classic example is male lions killing their predecessor's cubs when they take over a pride [32]. This brings lionesses into season and so hastens the new male's mating opportunities. Selfish acts can also be disguised as cooperation; white-winged choughs cheat by attempting to fool dominants that they are helping at the nest [33]. <i>Parasites.</i> Cheating is rife in bacterial infections; the cost of producing extracellular iron-scavenging siderophores in <i>Pseudomonas aeruginosa</i> selects for non-producing cheats who can outcompete cooperators [34,35]. Because cooperation is required to efficiently acquire the iron, mixed infections can be less virulent [36]. By contrast, competition in co-infections selects for faster replication in <i>Plasmodium</i> and phage, which causes greater virulence to the host [37,38].</p>
<p>altruism</p> <p><i>Multicellular taxa.</i> The classic example is the eusocial insects (including bees, ants, termites) in which a single queen (or few) monopolize the colony's reproduction [39]. All others in the colony are morphologically or behaviourally specialized to altruistically forage, guard or raise offspring. <i>Parasites.</i> Morphological castes have been described in <i>Philopthalmus</i> sp. trematodes, which produce reproductive and non-reproductive morphs. The sterile morph increases the reproductive output of relatives but only when competing strains share the host [40]. Other examples include suicide in <i>Salmonella typhimurium</i> to facilitate the gut invasion of others [41], and in <i>Escherichia coli</i> infected with phage to prevent parasite transmission to others [42]. Forms of suicide have been described in <i>Plasmodium</i> and trypanosomes, which may regulate density, preventing premature death of the host/vector [15,43]. Suicidal release of a public good is not a necessary condition for altruism—the release of a costly public good into a dense and well-mixed population (minimizing the direct return to the actor cell) is likely to be an altruistic act.</p>	<p>spite</p> <p><i>Multicellular taxa.</i> Soldiers of the polyembryonic parasitoid wasp <i>Copidosoma floridanum</i>, are sterile because they never reach sexual maturity. They preferentially kill other, unrelated embryos developing in their host, which frees up resources for their siblings [44]. Workers of the red fire ant <i>Solenopsis invicta</i>, kill related queens who do not have a greenbeard gene, ensuring that the greenbeard gene persists [45]. <i>Parasites.</i> Bacteria secrete anti-competitor toxins that are costly to produce (e.g. toxin release requires cell lysis in many cases). Toxin production and immunity genes are usually linked so that relatives are immune. Virulence is lower in mixed infections in caterpillar hosts compared with single infections when competing strains are susceptible to each other's bacteriocins [46]. However, the impact of spite on virulence becomes more complicated when spiteful behaviours are affected by simultaneous investments in other social traits (e.g. public goods) [47].</p>




Figure 1. Examples of kin discrimination by: (a) direct recognition, e.g. cells of the slime mould *Dictyostelium* determine whether they are interacting with kin or non-relatives during slug and spore formation based on the sequence similarity of their surface adhesion proteins [66,67] (photo credit Owen Gilbert); (b) indirect cues based on familiarity with individuals, e.g. long-tailed tits learn the vocalization patterns of kin during the natal rearing period [68] (photo credit Sarah Reece) or (c) ‘armpits’ which are a mixture of direct and indirect cues, e.g. ground squirrels use olfactory cues which have a genetic component and are also learnt by self-referencing during development [69] (photo credit Alan Vernon). The malaria parasite, *Plasmodium chabaudi* (d), adjusts investment into male and female transmission stages according to how many other conspecific clones share the host, suggesting kin discrimination occurs [21] (photo credit Sarah Reece and Sinclair Stammers). The mechanism is unknown but indirect cues seems unlikely; an obvious candidate would be that parasites can infer the presence of other clones via the host immune response, but sex ratio adjustment is observed in infections before the required strain-specific responses develop.

only direct help to individuals expressing that gene—a notion popularized as a ‘green beard’ [64,65]. Another way to direct cooperative behaviours towards appropriate recipients is through the ability to recognize kin (figure 1). Kin discrimination can occur via: (i) direct recognition, (ii) indirect cues that convey whether a recipient is likely to be a relative; or (iii) a mixture of direct and indirect information. Kin discrimination systems can require additional selective forces to maintain polymorphisms that can be used as accurate identifiers [70]. Host–parasite systems, in which genotype-by-genotype interactions and frequency-dependent selection maintain genetic variation, are candidate motors maintaining the genetic diversity required for kin discrimination (figure 1).

(d) Communication: coordinating collective action

Parasites have evolved sophisticated communication systems to coordinate behaviours across clone-mates and enable collective actions to be efficiently deployed. For example, 6–10% of all genes in the opportunistic microbial pathogen *Pseudomonas aeruginosa* are controlled by cell–cell signalling systems [71]. Coordination is especially important for behaviours that must be expressed by some, but not all individuals. For example, in cases where the suicide of

some individuals benefits survivors, the suicide trait must not be expressed by all individuals otherwise there would be no survivors [15,41]. Equally, undertaking costly cooperative actions may only pay when the numbers of actors exceeds a certain threshold, and so density-sensing mechanisms are often used to ensure behaviours are only switched on at high densities (‘quorum’) [72]. Microbial pathogens are masters of coordinating collective actions; their quorum-sensing system enables density estimation via collectively produced diffusible molecules [72]. The recent discovery that malaria parasites secrete protein and DNA containing microvesicles that influence the sexual differentiation of other parasite cells [73,74] may be a mechanism to organize the density-dependent decisions observed in reproductive effort and sex allocation [21,75,76].

3. Virulence evolution

Parasites engage in clearly selfish acts with the hosts and vectors they exploit; here we give a brief overview of answers to the basic question of why parasites harm the very source of their livelihoods. The development of evolutionary theory to explain virulence (parasite-induced harm to the host) has

a long history. Theories for the evolution of virulence can be categorized into four broad hypotheses [77,78] under which high virulence is attributed variously to: (i) novel host–parasite associations [79,80]; (ii) transmission–virulence trade-offs [81]; (iii) coincidental evolution of virulence factors [82,83]; and (iv) short-term within-host evolution [84]. For many infectious diseases, social interactions among parasites and virulence are coupled, but the nature of this relationship varies according to the type of interactions involved and who the interaction partners are.

The most influential theoretical framework for virulence evolution centres on virulence being maintained as a result of an unavoidable constraint linking the benefits of transmission with the costs of virulence. In this view, virulence (measured as host death) is an unavoidable cost of the host exploitation required for transmission to new hosts [85–88]. If the costs of increasing exploitation accelerate more rapidly than the transmission benefits of increasing exploitation, then natural selection favours an intermediate level of host exploitation (optimal virulence) [81,89]. Following from this premise, the relatedness of co-infecting parasite genotypes can modulate the best or evolutionary stable strategy of virulence, depending on the nature of social interactions among co-infecting parasites (figure 2). When co-infecting genotypes have direct control over their mechanisms of host exploitation, the benefits of increased exploitation are felt by the individuals responsible whereas the costs of virulence are shared by all, favouring greater virulence than that of parasites in single genotype infections [86–89]. By contrast, if co-infecting parasites work collectively to exploit the host (for example, via the secretion of shared extracellular digestive enzymes), then the benefits of exploitation become collectivized and mixed infections can select for ‘non-producer’ cheats that attenuate virulence [90]. In both scenarios, the spread of cheats (either over- or under-exploiters) undermines the productivity of the infection as a whole [56,88] but has opposite consequences for virulence.

4. Interactions in infections

The virulence–transmission trade-off models and their ‘virulence–kin-selection’ offshoots have been influential in the development of a vast body of subsequent theory [81]. Empirical testing has proceeded at a slower pace but only a few systems have provided support for the virulence–transmission trade-off [83,91]. A central theme emerging in the disease evolution literature is that within-host ecological dynamics are critical determinants of parasite sociality and so virulence [83,91,92]. In the following sections, we take a mechanistically inspired bottom-up approach, viewing virulence and transmission as emergent properties of complex within-host processes and highlighting five aspects of infections that can shape parasite social behaviours: (i) public goods, (ii) mobile elements, (iii) phenotypic plasticity, (iv) spatial structure and (v) multi-species interactions. We illustrate that a better understanding of these processes brings new perspectives to the traditional ‘top-down’ frameworks for the evolution and epidemiology of virulence and transmission.

(a) Public goods

A central aspect of interactions between microbial pathogens is the collective engineering of their shared environment via the




Figure 2. Theoretical relationships between virulence and relatedness under conditions of: (a) individual exploitation (virulence maximized at low relatedness) (b) collective exploitation (virulence maximized at high relatedness) (c) spiteful interactions, e.g. when harming competitors trades off against replication that causes virulence. (summarized by [16]).

secretion of costly ‘public goods’ molecules. These molecules generate a range of benefits to any neighbouring cell that are suitably equipped to profit. For example, public goods molecules may scavenge for limiting resources (e.g. siderophores), aid in the construction of biofilms (e.g. adhesive polymers), kill competing lineages (e.g. bacteriocins) or enhance host exploitation (e.g. digestive enzymes, toxins). Because these molecules are individually costly to produce and yet return a collective benefit, they have become a focus in the study of bacterial cooperation. Among the best-studied model system for public goods cooperation is iron scavenging by secreted siderophores in the opportunistic bacterial pathogen *P. aeruginosa* (and related pseudomonads) [34,93,94]. *In vitro* studies in iron-limited environments have demonstrated that the fate of siderophore-producing ‘cooperator’ lineages in competition with non-producer ‘cheats’ is dependent on the degree of strain mixing or relatedness. When relatedness is high (each sub-population founded by a single clone), producers outcompete cheats, because the benefits of cooperation are disproportionately high for other cooperators. By contrast, when relatedness is low (e.g. each sub-population founded by multiple clones), cheats outcompete producers [34] (but see [95] for an exception driven by strong non-social selection).

The general applicability of a public goods framework for microbial interactions mediated by secreted factors has recently been called into question by Zhang & Rainey [96]. Here again the experimental focus was on siderophore-mediated interactions, where the authors illustrated that in certain standard laboratory experimental conditions (KB media), the production of siderophores is redundant and selected against. This result serves as a valuable reminder that the benefits of secreted molecules are undoubtedly environment dependent, and in this particular environment the secreted molecule does not provide benefits to neighbours and therefore does not function as a public good. Kummerli & Ross-Gillespie [97] responded to Zhang & Rainey [96] with an analysis of the iron content of KB, revealing that it is relatively iron-replete, ensuring that siderophore production is unlikely to provide sufficient benefit to merit the costs of production. From this, Kummerli & Ross-Gillespie conclude that there is no difficulty for the public goods framework, so long as the environmental context is adequately accounted for [97].

(b) Mobile elements: infectious cooperation and locus-specific relatedness

The maintenance of cooperation via a single-cell bottleneck for each sub-population (as in [34] discussed earlier) is a

very stringent condition. How is cooperation maintained under more realistic conditions that allow for some strain mixing, and more frequent interactions with cheats due to mutation or migration? The peculiar biology of bacteria points to an intriguing role played by their molecular parasites in maintaining the cooperative phenotypes of their bacterial hosts. Bacteria are prone to infection with a diverse array of molecular parasites that are able to spread infectious via horizontal gene transfer (HGT) through a population, bringing novel genes along for the ride [98]. Key among these molecular parasites are plasmids, vectors of many medically significant alleles including antibiotic resistance and toxins [18]. Initial theoretical work suggested that the invasion of cheats into a population of cooperators could be prevented if the cooperative trait was encoded by an infectious conjugative plasmid [99]. In this scenario, cheats are liable to be re-programmed via infection with the cooperation-inducing plasmid. A key assumption of this model is that all plasmids carry the cooperative trait, so any act of infection will also increase cooperation, by hitch-hiking on the conjugation alleles.

But what if the social dilemma between cooperative and cheating alleles is played out at the level of the mobile element? More recent theory has pointed out that in an unstructured environment, ‘cheat’ plasmids will outcompete ‘cooperative’ plasmids for the same conditions that favour cheating chromosomal alleles over their cooperative rivals [100], because, again, the benefits of cooperation are not preferentially returned to cooperative alleles. However, the picture changes in structured populations, in which bacteria exploit discrete patches (e.g. hosts), linked by migration and/or transmission. Population structure introduces non-zero relatedness, and so the patterns of relatedness are now predicted to vary at different points of the genome depending on the rate of HGT [17,18,101]. Plasmids with high rates of HGT can readily copy themselves into neighbouring cells within a patch, and so if a cooperative plasmid gene generates benefits for neighbouring cells, it is now more likely to aid gene-copies in neighbouring cells to reproduce. In other words, highly conjugative plasmids gain a greater inclusive fitness return from helping neighbouring cells, favouring cooperative investments at these loci. Bio-informatic support for this inclusive fitness hypothesis has been demonstrated across 20 strains of *Escherichia coli*, where genes liable to experience greater HGT were more likely to code for secreted (cooperative) traits [17]. More recently, experiments show that HGT promotes plasmid-specific relatedness and selection for plasmid-encoded cooperation [102].

(c) Phenotypic plasticity: adaptive adjustment of behaviours

An important feature of parasite lifestyles is that their social environments change constantly, and so parasites have evolved mechanisms to regulate their behaviours. HGT is a form of genetic plasticity that enables the loss and gain of locally adapted alleles [18], but parasites also excel at phenotypic plasticity, extracting multiple phenotypes from one genotype. Adaptive phenotypic plasticity—the ability of an organism to change its behaviour or morphology to fit the environment—is a ubiquitous solution to the challenges of life in a changing environment. Plasticity enables organisms to maintain fitness by altering their phenotype, through

mechanisms such as differential gene expression, to best suit their circumstances [103], and here we focus on how plasticity in the behaviours (life-history traits) of parasites are shaped by their social environment within the host. For example, kin discrimination is a plastic response to social circumstances. By ensuring parasites only cooperate under conditions of high relatedness, kin discrimination may maintain cooperation by ensuring that the behaviour is adaptive from an inclusive fitness perspective, by limiting the potential to be exploited by cheats. Moreover, as well as enabling organisms to respond quickly once environmental change has occurred, organisms can also respond to predictors of future environmental change which enables appropriate phenotypes to be adopted in a timely manner [104].

Typically, evolutionary biologists and parasitologists have overlooked the notion that plasticity can produce qualitative and adaptive changes to the genotype-wide social phenotypes of parasites during infections. This is because they assume that parasite responses to environmental perturbation are mostly directed at maintaining homeostasis. As a result, variation in parasite behaviours is often—and potentially incorrectly—attributed to the footprint of host regulation rather than parasites making strategic decisions. For example, when the coordinated cell cycles of the rodent malaria parasite *Plasmodium chabaudi* are perturbed, they become rescheduled during infection and return to matching the host circadian rhythm. Whether parasite cell cycles are passively scheduled by host factors with a circadian basis or by parasites actively and collectively adjusting their timing is unclear [105]. However, evidence suggests that parasites are responsible for collectively coordinating their cell cycle schedules: synchronous and asynchronous malaria parasite species maintain their schedules in the same host environment (i.e. age–sex–strain-matched inbred mice); there are fitness benefits for parasites with cell cycles matched to the host circadian rhythm and matched infections cause greater virulence to the host [106,107]. Clearly, plasticity in parasite social behaviours complicates the understanding of within-host dynamics, but identifying to what extent parasite and/or host genes are responsible is central to interrogating their evolution.

The diversity of phenotypic plasticity in parasite social behaviours is illustrated in table 2. These traits are adjusted in response to social context and have consequences for virulence and transmission. Unfortunately, evolutionary theory has mostly ignored these behaviours, focusing instead on virulence. This is problematic because changes in virulence are achieved by changes in underlying traits (e.g. behaviours) expressed by both the host and parasites. As the social behaviours underpinning virulence and transmission are likely to be linked by genetic correlations (i.e. different traits are shaped by the same genes) and/or resource allocation trade-offs, the nature of these interactions is central to understanding and predicting virulence evolution [23]. Furthermore, when different genotypes respond to the environment in different ways (genotype-by-environment interactions or $G \times E$), environmental change can expose (or hide) genetic variation in plasticity to natural selection [119] (figure 3). Ecological perturbations such as drugs, vaccines and host shifts are all candidate motors for constraining or facilitating evolution, depending on how the perturbation affects the amount of genetic variation underpinning parasite phenotypes. For example, genetic variation for sex ratio adjustment and reproductive effort in response to social context has been documented in malaria parasites [21,76] and these behaviours are determinants of how

Table 2. Examples of phenotypic plasticity in parasite social behaviours. That phenotypes are a product of both genotypes and the environment, and how they interact, is well known, but often the environment is viewed as obscuring the connection between genes and phenotypes. However, how social behaviours are influenced by environmental variation matters because they affect virulence and transmission. Because multiple environmental factors change simultaneously during infections and virulence and transmission phenotypes are products of multiple social behaviours, parasites can produce a wide range of adaptive phenotypes faster by plasticity than when beneficial mutations or recombination are required to generate new phenotypes.

behaviour/ trait	what happens and why?
developmental schedules	In the host blood, cycles of asexual replication in many species of <i>Plasmodium</i> are tightly synchronized; individual parasites transit through each cell cycle stage and ultimately burst out of their red blood cells in unison and at particular times of day. The duration and synchronicity of cell cycles are plastic [105]. An adaptive basis of this plasticity is yet to be established but in-host competition and host immune responses are likely drivers [108]. Disrupted <i>P. chabaudi</i> schedules result in lower virulence (anaemia; [107]) but quiescence can also help <i>Plasmodium falciparum</i> tolerate antimalarial drugs [109].
lysis time	Pi 2 bacteriophage must lyse their bacterial host (<i>Pseudomonas fluorescens</i>) to transmit. They evolve a plastic lysis time in which they kill host cells more rapidly when co-infecting host cells with other phage than when infecting alone [38]. Plasticity in lysis time evolved in phage lines in mixed-infection conditions owing to the frequent variability in whether they encounter co- or single infections in this treatment (the lysis time in single-infection conditions did not change or become plastic in response to selection). This plasticity enhances the competitive ability of phage since non-plastic phage have fewer mature propagules upon cell lysis and suggests virulence and transmission differ according to whether parasites are in single or mixed genotype infections. In addition, lysis inhibition (LIN) is a mechanism of burst-size increase and latent period extension induced by T4 bacteriophage secondary adsorption of T4-infected <i>E. coli</i> cells. This plastic growth strategy is an adaptation to environments containing high densities of T4-infected cells [110]: when T4-infected cell density is high, high densities of free phages are generated, uninfected cells are rapidly infected, secondary adsorption is likely and LIN is induced with high probability [110–113].
public goods	The production of an iron-scavenging molecule (pyoverdine) by <i>P. aeruginosa</i> bacteria is a cooperative trait. Pyoverdine production per bacterium is tightly regulated by the intracellular supply of free iron, leading to decreased <i>per capita</i> production at higher cell densities and increased production in the presence of non-producing cheats. This phenotypic plasticity significantly influences the costs and benefits of cooperation. Specifically, the investment of resources into pyoverdine production is reduced in iron-rich environments and at high cell densities, but increased under iron limitation, and when pyoverdine is exploited by cheats [114,115]. Regulatory control of public goods provisioning can further protect producers from exploitation by cheats by ‘metabolic prudence’, limiting production to environments where the relative costs of production are minimized [60]. More globally, the regulatory control of multiple secreted factors is under the control of quorum-sensing mechanisms in numerous bacteria, including several significant pathogens [116].
reproductive effort	<i>Plasmodium</i> must replicate asexually in the vertebrate host and undergo a round of sexual reproduction in the vector. This means that resources must be divided between growth (the production of asexual stages for in-host survival) and reproduction (the production of sexual stages for transmission). <i>P. chabaudi</i> adopts reproductive restraint when facing in-host competition, which is consistent with investing in asexual replication (a key determinant of competitive ability) to gain future transmission opportunities [20,76]. <i>P. falciparum</i> also adopts reproductive restraint in response to low doses of drugs, suggesting this is a general strategy for coping with stresses encountered in the host [75].
sex allocation	In addition to the growth versus reproduction trade-off described earlier, <i>Plasmodium</i> must also divide resources between male and female transmission stages (sex ratio). Sex ratios in <i>P. chabaudi</i> and <i>P. falciparum</i> are adjusted in response to the inbreeding rate, which is determined by the number of co-infecting genotypes and their relative frequencies [21]. In single infections, female-biased sex ratios maximize zygote production and increasing the proportion of males in mixed infections, especially if a weak competitor, maximizes representation in the zygote population.
suicide	A ‘suicide trait’ cannot be constitutively expressed (if everyone dies before reproducing, genes for the trait cannot be inherited). Thus, the proportion of parasites that die may be precisely adjusted in response to variation in the density and relatedness of co-infecting parasites, or noisy expression of the genes involved may ensure phenotypic variation [41,117]. The release of bacteriocins to kill competitors requires bacterial cells to lyse themselves in many species, including <i>E. coli</i> [14]. The benefits accruing to surviving kin are highest when at low density, but this is when the costs of losing group members are greatest. By contrast, <i>Plasmodium</i> experiences crowding in the vector: high parasite densities reduce per parasite productivity and elevate vector mortality so suicide in the stage infective to the vector is predicted to regulate infection intensity [118].




Figure 3. Phenotypic plasticity and reaction norms. In panel (a), phenotype does not vary with the environment and both genotypes have identical reaction norms. In panel (b) both genotypes are plastic and (c) there is also genetic variation. Panel (d) illustrates a genotype-by-environment interaction ($G \times E$), where both genotypes are plastic but their phenotypic reaction norms vary. Genetic variation and $G \times E$ can complicate how much genetic variation is exposed to selection; in panel (e) the genotypes produce the same phenotype in environment (E) 1 but not in environment 2, so selection can only differentiate between the genotypes in environment 2.

parasites survive drugs and overcome transmission-blocking immunity [75,120,121]. However, how $G \times E$ affects the speed that parasites could respond to selection on these behaviours is not known.

The potential for interactions between plasticity and evolution introduces yet more complications to understanding how social behaviours shape parasite fitness for two additional reasons. First, adaptive plasticity can facilitate parasite evolution simply by providing more time and/or individuals for beneficial mutations to arise because their survival is enhanced [122]. By contrast, when plasticity buffers parasites against the loss of fitness in a novel environment, the strength of selection imposed by environmental change is reduced, and so parasite evolution is constrained. Quantitative theory that makes testable predictions for the opposing effects of plasticity on rates of evolution is urgently needed. For example, the social behaviours of malaria parasites provide tolerance to drugs: plasticity in reproductive restraint helps buffer against the impact of drugs on within-host survival [75]. Thus, selection for other resistance traits (e.g. drug efflux pumps, alternative metabolic or detoxification pathways) is weakened but this clinically beneficial outcome may be undermined because the greater number of surviving parasites offers more opportunities for resistance mutations to occur. Second, while a behaviour may be plastically adjusted in response to social context, the consequences of the action can subsequently feedback to affect social context. For example, bacteriophage plastically speed-up their host lysis time phenotype in response to being in a mixed versus a single infection, i.e. they are responding to the social context encountered within their host [38]. By lysing the host cell before non-plastic phage can transmit, the plastic phage gain a competitive advantage and consequently become increasingly more likely to interact with related phage.

It is our view that incorporating phenotypic plasticity into social evolution theory represents a milestone for bringing theoretical work and empirical observations closer. To some extent, for a few social traits (e.g. sex allocation of malaria parasites [123]) existing theory that predicts what fixed traits should be at equilibrium can also apply to plastic strategies, and so can be used to make quantitative predictions. However, analyses that incorporate phenomena specific to plasticity, such as its costs and limits, are lacking. The costs and limits of plasticity matter because they may maintain genetic variation in natural populations [124] and could offer novel disease intervention targets [23]. The importance of the costs and limits of plasticity are illustrated by parasites for which the host is an infrequent

environment. For example, *P. aeruginosa* is a supreme generalist microbe, able to grow in soil, water and diverse animal and plant hosts, thanks to high investment in regulatory factors [83,125]. While the benefits of extensive and complex regulatory control are easily appreciated in its broad host range, they also raise the potential cost of making ‘bad decisions’, turning on genes inappropriately when faced with a novel environment. During initial human colonization, *P. aeruginosa* turns on an array of virulence factors [126,127] that cause serious damage to the host. However, many of these damaging traits are subsequently lost or turned off during within-host evolution [128], suggesting that the initial plastic responses were maladaptive. It is possible that the loss of these secreted virulence factors is due to social interactions favouring non-producing ‘cheats’ that do not pay the cost of the collectively useful virulence traits [34,129]. However, the continued ability of these ‘cheat’ strains to persist [129] suggests that the virulence factors are redundant in the host lung, and their initial upregulation was a ‘bad decision’ [83].

(d) Spatial structure

A major limitation of both theoretical and experimental work is that, for simplicity, historically most microbial (especially bacterial) studies considered well-mixed groups in liquid where local spatial structure is minimal [34,130]. This view may be a reasonable approximation for taxa like malaria parasites, where social interactions appear to play-out on a host-wide scale. However, hosts are not ‘a well-mixed bag’ of resources and immune defences, and so the reality for many parasites is that infections are far more structured at a local (within-host) scale [131]. For instance, many bacteria stick themselves to host surfaces or attach to each other, in groups called biofilms. Social interactions are most intense when individuals live side-by-side in these structured environments [132]. For example, conflict between cooperating and cheating *P. aeruginosa* is more intense in biofilms than in liquid culture [133]. In a biofilm, the presence of cheats causes a greater reduction in population growth, reduces the structural integrity of biofilms and increases susceptibility to antibiotics [133]. However, the advantages of life in a biofilm may be tempered by a trade-off recently observed in *Vibrio cholerae*, between the benefits of being better competitors within the host and the costs of impaired ability to disperse [134].

Table 3. The potential of ‘Hamiltonian Medicine’: examples and limitations of proposed biomedical applications of parasite sociality.

concept	examples
cheat therapy	A strategy as simple as the introduction of a cheat (non-producer) strain can lead to direct reduction in parasite virulence, as well as a reduced bacterial population size, that may make the infection more susceptible to other intervention strategies. For example, the introduction of cheater mutants with reduced expression of secreted virulence factors into infections of the bacterial pathogen <i>P. aeruginosa</i> reduces mortality in a mouse model [146], at least in the case of simultaneous inoculation of the target wild-type and the cheater ‘treatment’. The ability of cheats to increase in frequency within a wild-type infection while simultaneously decreasing virulence has led to the idea of exploiting cheater invasion to introduce medically beneficial alleles into infections, such as sensitivity to antibiotics or a lethal toxin under the control of an inducible promoter, which when activated would eliminate both cooperators and cheats [19]. This approach resembles phage therapy, where a live and natural enemy is administered to control an infection at a specific site, and shares the benefits of responsive dosing (the treatment can amplify at the target site, unlike chemical therapeutics). However, cheat therapies face many of the obstacles we outline in the main text—they may be vulnerable to ‘reprogramming’ by cooperation-inducing plasmids, they may be unable to exploit established cooperator populations owing to within-host structure, or owing to plastic phenotypic changes in the resident. Finally, rare cheats may be unable to overcome the local-adaptation advantages of established wild-type infections [95,147].
drug resistance	Drug resistance mechanisms are often thought to impose fitness costs in the absence of drugs. Experiments using malaria parasites suggest that these fitness costs include competitive inferiority, and so suppression by wild-type genotypes in mixed infections could constrain the spread of resistance [148]. However, the extent to which suppression impacts on resistance in natural infections and how this could interact with eradication programmes is unclear. This is because as parasite prevalence decreases, infections will increasingly contain highly related parasites, which are more likely to cooperate than compete. Traditional antibiotics act by killing or stopping cell division, and resistant mutants rapidly replace the original susceptible strains. Instead, if a drug attacks a cell’s ability to secrete a public good that contributes to virulence (an ‘anti-virulence’ drug), then resistant mutants that re-evolve secretion will promote the growth of susceptible cells around them, reducing the spread of resistance. Moreover, because the susceptible cells do not pay the cost of secretion (i.e. they cheat), this puts resistant parasites at a competitive disadvantage, further reducing the spread of resistance [148–151].
evolutionary traps	An underexplored avenue concerns manipulating parasite kin recognition and communication systems to ‘trick’ parasites into adopting strategies that are suboptimal for their fitness and of clinical or epidemiological benefit. Evolving resistance to this type of intervention could be difficult because solutions would probably involve losing the benefit of coordinated action in untreated infections. For example, in malaria parasites, investment in asexual stages (which are responsible for disease symptoms) versus sexual stages is plastic. Parasites competing in mixed infections invest relatively less in sexual stages than when in single infections [76]. A drug that mimics being in a single infection (e.g. masks the cues of competition), and so induces parasites to invest more in sexual stages, will result in less virulent infections, and as long as conditions are vector-free there will be no increase in the risk of transmission to other hosts. Furthermore, the additional sexual stages will provide a stronger stimulus to the host immune system and the resulting responses could more effectively block the transmission of future malaria infections [152]. An approach to blocking transmission would be to induce mass suicide in the vector [15].

(e) Multi-species interactions

Most natural parasite communities are characterized by spatial structure, a multitude of co-infecting species and several environments to cope with. For example, the lesson from bacterial metagenomics is that thousands of species are commonly present in any one environment [132,135]. By contrast, the primary focus of parasite social evolution studies has involved examining what happens when multiple genotypes of a single species are mixed (e.g. [21,37,76,136]). Cross-species parasite social interactions are diverse: depending on the species in question, an incoming species can be excluded, facilitated or unaffected by a resident species [137]. For example, an ongoing malaria infection can exclude conspecifics [138,139] but strongly facilitate infection by heterospecific malaria parasites. In the latter case, species preferentially infecting mature red blood cells generate

anaemia to which the host responds by producing young red blood cells, which is predicted to facilitate malaria species that prefer the abundant young age class, resulting in far higher virulence than single-species infection [140]. However, the mechanisms that determine cross-species interactions are highly diverse, ranging from resource competition, interference competition (e.g. the production of antibiotics and bacteriocins), immune-mediated apparent competition and facilitation (e.g. cross-feeding on partner metabolic byproducts, immunomodulation) [141]. Together this menu of interactions contributes to the astounding diversity of communities of commensals, symbionts and parasites found within multicellular organisms.

A major challenge to unravelling the mechanisms underpinning how communities function is the necessity to combine molecular and ecological approaches to study

highly complex assemblies. A measure of the scale of the problem can be seen by the emergent ecological complexity generated by a simple two-species interaction governed by a single mechanism of metabolic exchange—a food for detoxification exchange—where a cross-feeding partner relieves a producer lineage of by-product toxicity. Recent theory has demonstrated that this simple exchange can generate mutualistic, competitive and exploitative functional relationships, and diverse spatial patternings, dependent on the exact parametrization of the molecular exchange [142]. Unravelling the complexity of these interactions—and how they affect evolution—is urgently required because the microbial communities inside vectors are being manipulated to control disease [143].

5. Why the social lives of parasites matter

Parasitism is one of the most successful modes of life, as measured by how often it evolved and how many parasitic species are presently in existence [144]. Consequently, if explaining cooperation is one of the greatest problems for evolutionary biology, then explaining cooperation in parasites is one of the key aspects of this problem. The irreducible mish-mash of proximate causality of social behaviours in traditionally studied animal taxa is far more accessible for parasites, thanks to their relatively simple and manipulatable genotype–phenotype maps. Parasites make excellent model organisms thanks to their short generation times; ability to generate some real-world complexity, even in the laboratory, by studying *in vivo* infections; and well-defined, measurable, social behaviours. Moreover, the applied importance of parasites has resulted in a vast resource of tools and literature on their molecular and cellular biology, so the genetic and molecular mechanisms that underlie social behaviours can be identified and precisely manipulated [8].

Incorporating a ‘bottom-up’ approach provides a novel perspective on the evolution and maintenance of parasite social behaviours and provides new opportunities for theory-led experimental testing. For example, by understanding aspects of interactions in infections such as those highlighted in this article, traditional virulence evolution theory may be better reconciled with data. Research has focused on social interactions between parasites within hosts (probably owing to the greater interest in disease pathology than transmission) and so social interactions inside vectors have been overlooked, but we expect that they are

equally worthy of investigation. Moreover, for parasite species whose life cycles include multiple host species or periods in the abiotic environment, quantifying how social behaviours at these different scales integrate to shape parasite fitness is also a huge challenge, and highlights the need to consider within-host biology in its broader context.

The social behaviours of parasites contribute to virulence, transmission and resistance to anti-parasite drugs, as illustrated throughout the text and tables of this article. The field of ‘Darwinian Medicine’ aims to use ecological and evolutionary principles to inform the treatment of infections to ensure that interventions are as evolution-proof as possible, and prevent the evolution of more harmful parasites in response to anthropogenic pressures. ‘Hamiltonian Medicine’ is emerging as a subset of this endeavour, asking how parasite social systems and interactions might be subverted or manipulated to better control disease [9,145]. By recognizing that parasites rely on social behaviours to infect and transmit, novel strategies for treatment have been revealed (table 3).

6. Conclusion

A key strength of evolutionary biology is that theory is used to motivate experiments. Historically, this has been the case, with many empirical tests stemming from the basic virulence–transmission trade-off models and their ‘virulence–kin-selection’ offshoots. However, for topics such as phenotypic plasticity, empirical work is often ahead of social evolution theory and this disconnect is especially apparent in systems that have applied importance. We recognize that the complexity of within-host parasite ecology may have been off-putting for evolutionary theorists since, on the face of it, generalities seem unlikely and explaining what is going on requires deeper knowledge of the biological details of individual study systems. However, generalities do exist—such as public goods, mobile elements, phenotypic plasticity, within-host spatial structure and multi-species interactions—that will provide rewarding avenues for future theoretical and experimental research.

Acknowledgements. Thanks to A. Buckling, M. Boots and M. Bell for discussion and insight, and S. Alizon and an anonymous reviewer for useful comments on the manuscript.

Funding statement. H.L. is supported by the BBSRC and ERC. S.R. is supported by the Wellcome Trust (WT082234MA), The Royal Society, and the Centre for Infection, Immunity and Evolution (CIIE) at the University of Edinburgh. S.B. is supported by the CIIE.

References

- West SA, Diggle SP, Buckling A, Gardner A, Griffin AS. 2007 The social lives of microbes. *Annu. Rev. Ecol. Evol. Syst.* **38**, 53–77. (doi:10.1146/annurev.ecolsys.38.091206.095740)
- Nadell CD, Bucci V, Drescher K, Levin SA, Bassler BL, Xavier JB. 2013 Cutting through the complexity of cell collectives. *Proc. R. Soc. B* **280**, 20122770. (doi:10.1098/rspb.2012.2770)
- Crespi BJ. 2001 The evolution of social behavior in microorganisms. *Trends Ecol. Evol.* **16**, 178–183. (doi:10.1016/S0169-5347(01)02115-2)
- Webb JS, Givskov M, Kjelleberg S. 2003 Bacterial biofilms: prokaryotic adventures in multicellularity. *Curr. Opin. Microbiol.* **6**, 578–585. (doi:10.1016/j.mib.2003.10.014)
- Kolter R, Greenberg EP. 2006 Microbial sciences: the superficial life of microbes. *Nature* **18**, 300–302. (doi:10.1038/441300a)
- West SA, Griffin AS, Gardner A, Diggle SP. 2006 Social evolution theory for microorganisms. *Nat. Rev. Microbiol.* **4**, 597–607. (doi:10.1038/nrmicro1461)
- Diggle SP, Griffin AS, Campbell GS, West SA. 2007 Cooperation and conflict in quorum-sensing bacterial populations. *Nature* **450**, 411–414. (doi:10.1038/nature0627915)
- Foster KR, Parkinson K, Thompson CRL. 2007 What can microbial genetics teach sociobiology? *Trends Genet.* **23**, 74–80. (doi:10.1016/j.tig.2006.12.003)
- Hense BA, Kuttler C, Muller J, Rothballer M, Hartmann A, Kreft JU. 2007 Does efficiency sensing unify diffusion and quorum sensing? *Nat. Rev. Microbiol.* **5**, 230–239. (doi:10.1038/nrmicro1600)

10. Williams P, Winzer K, Chan WC, Camara M. 2007 Look who's talking: communication and quorum sensing in the bacterial world. *Phil. Trans. R. Soc. B* **362**, 1119–1134. (doi:10.1098/rstb.2007.2039)
11. Brown SP, Buckling A. 2008 A social life for discerning microbes. *Cell* **135**, 600–603. (doi:10.1016/j.cell.2008.10.030)
12. MacLean R. 2008 The tragedy of the commons in microbial populations: insights from theoretical, comparative and experimental studies. *Heredity* **100**, 471–477. (doi:10.1038/sj.hdy.6801073)
13. Popat R, Cruz SA, Diggle SP. 2008 The social behaviours of bacterial pathogens. *Br. Med. Bull.* **87**, 63–75. (doi:10.1093/bmb/ldn030)
14. Brown SP, Inglis R, Taddei F. 2009 SYNTHESIS: evolutionary ecology of microbial wars: within-host competition and (incidental) virulence. *Evol. Appl.* **2**, 32–39. (doi:10.1111/j.1752-4571.2008.00059.x)
15. Reece SE, Pollitt LC, Colegrave N, Gardner A. 2011 The meaning of death: evolution and ecology of apoptosis in protozoan parasites. *PLoS Pathog.* **7**, e1002320. (doi:10.1371/journal.ppat.1002320)
16. Buckling A, Brockhurst MA. 2008 Kin selection and the evolution of virulence. *Heredity* **100**, 484–488. (doi:10.1038/sj.hdy.6801093)
17. Nogueira T, Rankin DJ, Touchon M, Taddei F, Brown SP, Rocha EPC. 2009 Horizontal gene transfer of the secretome drives the evolution of bacterial cooperation and virulence. *Curr. Biol.* **19**, 1683–1691. (doi:10.1016/j.cub.2009.08.056)
18. Rankin DJ, Rocha EPC, Brown SP. 2010 What traits are carried on mobile genetic elements, and why? *Heredity* **106**, 1–10. (doi:10.1038/hdy.2010.24)
19. Brown SP, West SA, Diggle SP, Griffin AS. 2009 Social evolution in microorganisms and a Trojan horse approach to medical intervention strategies. *Phil. Trans. R. Soc. B* **364**, 3157–3168. (doi:10.1098/rstb.2009.0055)
20. Mideo N, Day T. 2008 On the evolution of reproductive restraint in malaria. *Proc. R. Soc. B* **275**, 1217–1224. (doi:10.1098/rspb.2007.1545)
21. Reece SE, Drew DR, Gardner A. 2008 Sex ratio adjustment and kin discrimination in malaria parasites. *Nature* **453**, 609–615. (doi:10.1038/nature06954)
22. Mideo N *et al.* 2013 Life in cells, hosts, and vectors: parasite evolution across scales. *Infect. Genet. Evol.* **13**, 344–347. (doi:10.1016/j.meegid.2012.03.016)
23. Mideo N, Reece SE. 2012 Plasticity in parasite phenotypes: evolutionary and ecological implications for disease. *Future Microbiol.* **7**, 17–24. (doi:10.2217/fmb.11.134)
24. Hamilton WD. 1964 The genetical evolution of social behaviour II. *J. Theor. Biol.* **7**, 17–52. (doi:10.1016/0022-5193(64)90039-6)
25. West SA, Griffin AS, Gardner A. 2007 Evolutionary explanations for cooperation. *Curr. Biol.* **17**, R661–R672. (doi:10.1016/j.cub.2007.06.004)
26. Hamilton WD. 1970 Selfish and spiteful behaviour in an evolutionary model. *Nature* **228**, 1218–1220. (doi:10.1038/2281218a0)
27. Lacey EA, Sherman PW. 1991 Social organisation of naked mole-rat colonies: evidence for divisions of labor. In *The biology of the naked mole-rat* (eds PW Sherman, JUM Jarvis, RD Alexander), pp. 275–336. Princeton, New Jersey: Princeton University Press.
28. Šumbera R, Zelová J, Kunc P, Knížková I, Burda H. 2007 Patterns of surface temperatures in two mole-rats (Bathyergidae) with different social systems as revealed by IR-thermography. *Physiol. Behav.* **92**, 526–532. (doi:10.1016/j.physbeh.2007.04.029)
29. Spinks AC, Plagányi EE. 1999 Reduced starvation risks and habitat constraints promote cooperation in the common mole-rat, *Cryptomys hottentotus hottentotus*: a computer-simulated foraging model. *Oikos* **85**, 435–444. (doi:10.2307/3546693)
30. Queller DC, Zocchi F, Cervo R, Turillazzi S, Henshaw MT, Santorelli LA, Strassmann JE. 2000 Unrelated helpers in a social insect. *Nature* **405**, 784–787. (doi:10.1038/35015552)
31. Telfer S, Lambin W, Birtles R, Beldomenico P, Burthe S, Paterson S, Begon M. 2010 Species interactions in a parasite community drive infection risk in a wildlife population. *Science* **330**, 243–246. (doi:10.1126/science.1190333)
32. Bertram BCR. 1975 Social factors influencing reproduction in wild lions. *J. Zool.* **177**, 463–482. (doi:10.1111/j.1469-7998.1975.tb02246.x)
33. Boland CRJ, Heinsohn R, Cockburn A. 1997 Deception by helpers in cooperatively breeding white-winged choughs and its experimental manipulation. *Behav. Ecol. Sociobiol.* **41**, 251–256. (doi:10.1007/s002650050386)
34. Griffin AS, West SA, Buckling A. 2004 Cooperation and competition in pathogenic bacteria. *Nature* **430**, 1024–1027. (doi:10.1038/nature02744)
35. Harrison F, Browning LE, Vos M, Buckling A. 2006 Cooperation and virulence in acute *Pseudomonas aeruginosa* infections. *BMC Biol.* **4**, 21. (doi:10.1186/1741-7007-4-21)
36. Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. 1996 Pyoverdinin is essential for virulence of *Pseudomonas aeruginosa*. *Infect. Immun.* **64**, 518–523.
37. Bell AS, De Roode JC, Sim D, Read AF. 2006 Within-host competition in genetically diverse malaria infections: parasite virulence and competitive success. *Evolution* **60**, 1358–1371. (doi:10.1554/05-611.1)
38. Leggett HC, Benmayor R, Hodgson DJ, Buckling A. 2013 Experimental evolution of adaptive phenotypic plasticity in a parasite. *Curr. Biol.* **23**, 139–142. (doi:10.1016/j.cub.2012.11.045)
39. Boomsma JJ. 2009 Lifetime monogamy and the evolution of eusociality. *Phil. Trans. R. Soc. B* **364**, 3191–3207. (doi:10.1098/rstb.2009.0101)
40. Kamiya T, Poulin R. 2013 Caste ratios affect the reproductive output of social trematode colonies. *J. Evol. Biol.* **26**, 509–516. (doi:10.1111/jeb.12062)
41. Ackermann M, Stecher B, Freed NE, Songhet P, Hardt WD, Doebeli M. 2008 Self-destructive cooperation mediated by phenotypic noise. *Nature* **454**, 987–990. (doi:10.1038/nature07067)
42. Refardt D, Bergmiller T, Kümmerli R. 2013 Altruism can evolve when relatedness is low: evidence from bacteria committing suicide upon phage infection. *Proc. R. Soc. B* **280**, 20123035. (doi:10.1098/rspb.2012.3035)
43. Pollitt LC, Colegrave N, Khan SM, Sajid M, Reece SE. 2010 Investigating the evolution of apoptosis in malaria parasites: the importance of ecology. *Parasit. Vectors* **16**, 105. (doi:10.1186/1756-3305-3-105)
44. Giron D, Dunn DW, Hardy ICW, Strand MR. 2004 Aggression by polyembryonic wasp soldiers correlates with kinship but not resource competition. *Nature* **430**, 676–679. (doi:10.1038/nature02721)
45. Keller L, Ross KG. 1998 Selfish genes: a green beard in the red fire ant. *Nature* **394**, 573–575. (doi:10.1038/29064)
46. Massey RC, Buckling A, Ffrench-Constant R. 2004 Interference competition and parasite virulence. *Proc. R. Soc. Lond. B* **271**, 785–788. (doi:10.1098/rspb.2004.2676)
47. Inglis RF, Brown SP, Buckling A. 2012 Spite versus cheats: competition among social strategies shapes virulence in *Pseudomonas aeruginosa*. *Evolution* **66**, 3472–3484. (doi:10.1111/j.1558-5646.2012.01706.x)
48. Hamilton WD. 1964 The genetical evolution of social behaviour I. *J. Theor. Biol.* **7**, 1–16. (doi:10.1016/0022-5193(64)90038-4)
49. Trivers RL. 1985 *Social evolution*. Menlo Park, CA: Benjamin/Cummings.
50. West SA, Griffin AS, Gardner A. 2007 Social semantics: altruism, cooperation, mutualism, strong reciprocity and group selection. *J. Evol. Biol.* **20**, 415–432. (doi:10.1111/j.1420-9101.2006.01258.x)
51. Gardner A, West SA, Wild G. 2011 The genetical theory of kin selection. *J. Evol. Biol.* **24**, 1020–1043. (doi:10.1111/j.1420-9101.2011.02236.x)
52. Bourke AFG. 2011 *Principles of social evolution*. Oxford Series in Ecology and Evolution. Oxford, UK: Oxford University Press.
53. Tibayrenc M, Ayala FJ. 2012 Reproductive clonality of pathogens: a perspective on pathogenic viruses, bacteria, fungi, and parasitic protozoa. *Proc. Natl Acad. Sci. USA* **109**, E3305–E3313. (doi:10.1073/pnas.1212452109)
54. Gardner A, Grafen A. 2009 Capturing the superorganism: a formal theory of group adaptation. *J. Evol. Biol.* **22**, 659–671. (doi:10.1111/j.1420-9101.2008.01681.x)
55. Dumas Z, Kummerli R. 2012 Cost of cooperation rules selection for cheats in bacterial metapopulations. *J. Evol. Biol.* **25**, 473–484. (doi:10.1111/j.1420-9101.2011.02437.x)
56. Hardin G. 1988 Tragedy of the commons. *Science* **162**, 1243–1248. (doi:10.1126/science.162.3859.1243)
57. Nowak M. 2011 *Supercooperators: altruism, evolution, and why we need each other to succeed*. New York, NY: The Free Press.
58. Dugatkin LA. 2002 Animal cooperation among unrelated individuals. *Naturwissenschaften* **89**, 533–541. (doi:10.1007/s00114-002-0379-y)
59. Chuang JS, Rivoire O, Leibler S. 2009 Simpson's paradox in a synthetic microbial system. *Science* **323**, 272–275. (doi:10.1126/science.1166739)

60. Xavier JB, Kim W, Foster KR. 2011 A molecular mechanism that stabilizes cooperative secretions in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **79**, 166–179. (doi:10.1111/j.1365-2958.2010.07436.x)
61. Sachs JL, Mueller UG, Wilcox TP, Bull JJ. 2004 The evolution of cooperation. *Q. Rev. Biol.* **79**, 135–160. (doi:10.1086/383541)
62. Lehmann L, Keller L. 2006 The evolution of cooperation and altruism—a general framework and a classification of models. *J. Evol. Biol.* **19**, 1365–1376. (doi:10.1111/j.1420-9101.2006)
63. Nadell CD, Xavier JB, Foster KR. 2009 The sociobiology of biofilms. *FEMS Microbiol. Rev.* **33**, 206–224. (doi:10.1111/j.1574-6976.2008.00150.x)
64. Dawkins R. 1976 *The selfish gene*. Oxford, UK: Oxford University Press.
65. Gardner A, West SA. 2010 Greenbeards. *Evolution* **64**, 25–38. (doi:10.1111/j.1558-5646.2009.00842.x)
66. Benabentos R *et al.* 2009 Polymorphic members of the *lag* gene family mediate kin discrimination in *Dictyostelium*. *Curr. Biol.* **19**, 567–572. (doi:10.1016/j.cub.2009.02.037)
67. Hirose S, Benabentos R, Ho HI, Kuspa A, Shaulsky G. 2011 Self-recognition in social amoebae is mediated by allelic pairs of *tiger* genes. *Science* **333**, 467–470. (doi:10.1126/science.1203903)
68. Sharp SP, McGowan A, Wood MJ, Hatchwell BJ. 2005 Learned kin recognition cues in a social bird. *Nature* **434**, 1127–1130. (doi:10.1038/nature03522)
69. Mateo JM. 2002 Kin-recognition abilities and nepotism as a function of sociality. *Proc. R. Soc. Lond. B* **269**, 721–727. (doi:10.1098/rspb.2001.1947)
70. Rousset F, Roze D. 2007 Constraints on the origin and maintenance of genetic kin recognition. *Evolution* **61**, 2320–2330. (doi:10.1111/j.1558-5646.2007.00191.x)
71. Schuster M, Lostro CP, Ogi T, Greenberg EP. 2003 Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. *J. Bacteriol.* **185**, 2066–2079. (doi:10.1128/JB.185.7.2066-2079.2003)
72. Darch SE, West SA, Winzer K, Diggle SP. 2012 Density-dependent fitness benefits in quorum-sensing bacterial populations. *Proc. Natl Acad. Sci. USA* **109**, 8259–8263. (doi:10.1073/pnas.1118131109)
73. Mantel PY *et al.* 2013 Malaria-infected erythrocyte-derived microvesicles mediate cellular communication within the parasite population and with the host immune system. *Cell Host Microbe* **13**, 521–534. (doi:10.1016/j.chom.2013.04.009)
74. Regev-Rudzki N *et al.* 2013 Cell–cell communication between malaria-infected red blood cells via exosome-like vesicles. *Cell* **153**, 1120–1133. (doi:10.1016/j.cell.2013.04.029)
75. Reece SE, Ali E, Schneider P, Babiker HA. 2010 Stress, drugs and the evolution of reproductive restraint in malaria parasites. *Proc. R. Soc. B* **277**, 3123–3129. (doi:10.1098/rspb.2010.0564)
76. Pollitt LC, Mideo N, Drew D, Schneider P, Colegrave N, Reece SE. 2011 Competition and the evolution of reproductive restraint in malaria parasites. *Am. Nat.* **177**, 358–367. (doi:10.1086/658175)
77. Levin BR. 1996 The evolution and maintenance of virulence in microparasites. *Emerg. Infect. Dis.* **2**, 93–102. (doi:10.3201/eid0202.960203)
78. Margolis E, Levin BR. 2007 Within-host evolution for the invasiveness of commensal bacteria: an experimental study of bacteremias resulting from *Haemophilus influenzae* nasal carriage. *J. Infect. Dis.* **196**, 1068–1075. (doi:10.1086/520934)
79. Ebert D. 1998 The evolution and expression of virulence. In *Evolution in health and disease* (ed. SC Stearns), pp. 153–167. Oxford, UK: Oxford University Press.
80. Bull JJ, Ebert D. 2008 Invasion thresholds and the evolution of nonequilibrium virulence. *Evol. Appl.* **1**, 172–182. (doi:10.1111/j.1752-4571.2007.00003.x)
81. Alizon S, Hurford A, Mideo N, Van Baalen M. 2009 Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future. *J. Evol. Biol.* **22**, 245–259. (doi:10.1111/j.1420-9101.2008.01658.x)
82. Levin BR, Eden CS. 1990 Selection and evolution of virulence in bacteria – an ecumenical excursion and modest suggestion. *Parasitology* **100**, S103–S115. (doi:10.1017/S0031182000073054)
83. Brown SP, Cornforth DM, Mideo N. 2012 Evolution of virulence in opportunistic pathogens: generalism, plasticity, and control. *Trends Microbiol.* **20**, 336–342. (doi:10.1016/j.tim.2012.04.005)
84. Levin BR, Bull JJ. 1994 Short-sighted evolution and the virulence of pathogenic microorganisms. *Trends Microbiol.* **2**, 76–81. (doi:10.1016/0966-842X(94)90538-X)
85. Anderson RM, May RM. 1982 Coevolution of hosts and parasites. *Parasitology* **85**, 411–426. (doi:10.1017/S0031182000055360)
86. Bremermann HJ, Pickering J. 1983 A game-theoretical model of parasite virulence. *J. Theor. Biol.* **100**, 411–426. (doi:10.1016/0022-5193(83)90438-1)
87. Nowak MA, May RM. 1994 Superinfection and the evolution of parasite virulence. *Proc. R. Soc. Lond. B* **255**, 81–89. (doi:10.1098/rspb.1994.0012)
88. Frank SA. 1996 Models of parasite virulence. *Q. Rev. Biol.* **71**, 37–78. (doi:10.1086/419267)
89. de Roode JC, Yates AJ, Altizer S. 2008 Virulence-transmission trade-offs and population divergence in virulence in a naturally occurring butterfly parasite. *Proc. Natl Acad. Sci. USA* **105**, 7489–7494. (doi:10.1073/pnas.0710909105)
90. Brown SP, Hochberg ME, Grenfell BT. 2002 Does multiple infection select for raised virulence? *Trends Microbiol.* **10**, 401–405. (doi:10.1016/S0966-842X(02)02413-7)
91. Leggett HC, Buckling A, Long GH, Boots M. 2013 Generalism and the evolution of parasite virulence. *Trends Ecol. Evol.* **28**, 592–596. (doi:10.1016/j.tree.2013.07.002)
92. Alizon S, de Roode JC, Michalakis Y. 2013 Multiple infections and the evolution of virulence. *Ecol. Lett.* **16**, 556–567. (doi:10.1111/ele.12076)
93. Buckling A, Harrison F, Vos M, Brockhurst MA, Gardner A, West SA, Griffin A. 2007 Siderophore-mediated cooperation and virulence in *Pseudomonas aeruginosa*. *FEMS Microbiol. Ecol.* **62**, 135–141. (doi:10.1111/j.1574-6941.2007.00388.x)
94. Kummerli R, Griffin AS, West SA, Buckling A, Harrison F. 2009 Viscous medium promotes cooperation in the pathogenic bacterium *Pseudomonas aeruginosa*. *Proc. R. Soc. B* **276**, 3531–3538. (doi:10.1098/rspb.2009.0861)
95. Morgan AD, Quigley BJZ, Brown SP, Buckling A. 2012 Selection on non-social traits limits the invasion of social cheats. *Ecol. Lett.* **15**, 841–846. (doi:10.1111/j.1461-0248.2012.01805.x)
96. Zhang XX, Rainey PB. 2013 Exploring the sociobiology of pyoverdinin-producing *Pseudomonas*. *Evolution* **67**, 3161–3174. (doi:10.1111/evo.12183)
97. Kümmerli R, Ross-Gillespie A. In press. Explaining the sociobiology of pyoverdinin-producing *Pseudomonas*: a comment on Zhang and Rainey (2013). *Evolution*. (doi:10.1111/evo.12311)
98. Frost LS, Leplae R, Summers AO, Toussaint A. 2005 Mobile genetic elements: the agents of open source evolution. *Nat. Rev. Microbiol.* **3**, 722–732. (doi:10.1038/nrmicro1235)
99. Smith J. 2001 The social evolution of bacterial pathogenesis. *Proc. R. Soc. Lond. B* **268**, 61–69. (doi:10.1098/rspb.2000.1330)
100. Mc Ginty SE, Rankin DJ, Brown SP. 2011 Horizontal gene transfer and the evolution of bacterial cooperation. *Evolution* **65**, 21–32. (doi:10.1111/j.1558-5646.2010.01121.x)
101. Mc Ginty SE, Lehmann L, Brown SP, Rankin DJ. 2013 The interplay between relatedness and horizontal gene transfer drives the evolution of plasmid-carried public goods. *Proc. R. Soc. B* **280**, 20130400. (doi:10.1098/rspb.2013.0400)
102. Dimitriu T, Lotton C, Bernard-Capelle J, Brown SP, Linder AB, Taddei F. Submitted. Genetic information transfer promotes cooperation by increasing relatedness in bacteria.
103. Pigliucci M. 2001 *Phenotypic plasticity: beyond nature and nurture*. Baltimore, MD: John Hopkins University Press.
104. Mitchell A, Romano GH, Groisman B, Yona A, Dekel E, Kupiec M, Dahan O, Pilpel Y. 2009 Adaptive prediction of environmental changes by microorganisms. *Nature* **460**, 220–224. (doi:10.1038/nature08112)
105. Mideo N, Reece SE, Smith AL, Metcalf CJ. 2013 The Cinderella syndrome: why do malaria-infected cells burst at midnight? *Trends Parasitol.* **29**, 10–16. (doi:10.1016/j.pt.2012.10.006)
106. O'Donnell AJ, Schneider P, McWatters HG, Reece SE. 2011 Fitness costs of disrupting circadian rhythms in malaria parasites. *Proc. R. Soc. B* **278**, 2429–2436. (doi:10.1098/rspb.2010.2457)
107. O'Donnell AJ, Mideo N, Reece SE. 2013 Disrupting rhythms in *Plasmodium chabaudi*: costs accrue quickly and independently of how infections are initiated. *Malaria J.* **12**, 372. (doi:10.1186/1475-2875-12-372)

108. Greischer MA, Read AF, Bjørnstad ON. 2014 Synchrony in malaria infections: how intensifying within-host competition can be adaptive. *Am. Nat.* **183**, E36–E49. (doi:10.1086/674357)
109. Witkowski B, Lelievre J, Barragan MJ, Laurent V, Su XZ, Berry A, Benoit-Vical F. 2010 Increased tolerance to artemisinin in *Plasmodium falciparum* is mediated by a quiescence mechanism. *Antimicrob. Agents Chemother.* **54**, 1872–1877. (doi:10.1128/AAC.01636-09)
110. Abedon ST. 1990 Selection for lysis inhibition in bacteriophage. *J. Theor. Biol.* **146**, 501–511. (doi:10.1016/S0022-5193(05)80375-3)
111. Abedon ST. 1992 Lysis of lysis-inhibited bacteriophage T4-infected cells. *J. Bacteriol.* **174**, 8073–8080.
112. Abedon ST. 1994 Lysis and the interaction between free phages and infected cells. In *Molecular biology of bacteriophage T4* (eds J Karam *et al.*), pp. 397–405. Washington, DC: American Society for Microbiology.
113. Abedon ST. 1999 Bacteriophage T4 resistance to lysis-inhibition collapse. *Genet Res.* **74**, 1–11. (doi:10.1017/S0016672399003833)
114. Kümmerli R, Jiricny N, Clarke LS, West SA, Griffin AS. 2009 Phenotypic plasticity of a cooperative behaviour in bacteria. *J. Evol. Biol.* **22**, 589–598. (doi:10.1111/j.1420-9101.2008.01666.x)
115. Kümmerli R, Brown SP. 2010 Molecular and regulatory properties of a public good shape the evolution of cooperation. *Proc. Natl Acad. Sci. USA* **107**, 18 921–18 926. (doi:10.1073/pnas.1011154107)
116. Rutherford ST, Bassler BL. 2012 Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb. Perspect. Med.* **2**, a012427. (doi:10.1101/cshperspect.a012427)
117. Viney M, Reece SE. 2013 Adaptive noise. *Proc. R. Soc. B* **280**, 20131104. (doi:10.1098/rspb.2013.1104)
118. Pollitt LC, Churcher TS, Dawes EJ, Khan SM, Sajid M, Basáñez MG, Colegrave N, Reece SE. 2013 Costs of crowding for the transmission of malaria parasites. *Evol. Appl.* **6**, 617–629. (doi:10.1111/eva.12048)
119. Schlichting CD, Pigliucci M. 1998 *Phenotypic evolution: a reaction norm perspective*. Sunderland, MA: Sinauer Associates Incorporated.
120. Ramiro RS, Alpedrinha J, Carter L, Gardner A, Reece SE. 2011 Sex and death: the effects of innate immune factors on the sexual reproduction of malaria parasites. *PLoS Pathog.* **7**, e1001309. (doi:10.1371/journal.ppat.1001309)
121. Reece SE, Khan SM, Waters AP, Janse CJ, Kaczanowski S. 2012 Why are male malaria parasites in such a rush? Sex-specific evolution and host-parasite interactions. *Evol. Med. Public Health.* **2013**, 3–13. (doi:10.1093/emph/eos003)
122. West-Eberhard MJ. 2003 *Developmental plasticity and evolution*. Oxford, UK: Oxford University Press.
123. Hamilton WD. 1967 Extraordinary sex ratios: a sex-ratio theory for sex linkage and inbreeding has new implications in cytogenetics and entomology. *Science* **156**, 477–488. (doi:10.1126/science.156.3774.477)
124. DeWitt TJ, Sih A, Wilson DS. 1998 Costs and limits of phenotypic plasticity. *Trends Ecol. Evol.* **13**, 77–81. (doi:10.1016/S0169-5347(97)01274-3)
125. Stover CK *et al.* 2000 Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an opportunistic pathogen. *Nature* **406**, 959–964. (doi:10.1038/35023079)
126. Duan K, Surette MG. 2007 Environmental regulation of *Pseudomonas aeruginosa* PAO1 Las and Rhl quorum-sensing systems. *J. Bacteriol.* **189**, 4827–4836. (doi:10.1128/JB.00043-07)
127. Bielecki P *et al.* 2011 *In vivo* expression profiling of *Pseudomonas aeruginosa* infections reveals niche-specific and strain-independent transcriptional programs. *PLoS ONE* **6**, e24235. (doi:10.1371/journal.pone.0024235)
128. Jiricny N, Diggle SP, West SA, Evans BA, Ballantyne G, Ross-Gillespie A, Griffin AS. 2010 Fitness correlates with the extent of cheating in a bacterium. *J. Evol. Biol.* **23**, 738–747. (doi:10.1111/j.1420-9101.2010.01939.x)
129. Bragonzi A, Paroni M, Nonis A, Cramer N, Montanari S, Rejman J, Di Serio C, Döring G, Tümmler B. 2009 *Pseudomonas aeruginosa* microevolution during cystic fibrosis lung infection establishes clones with adapted virulence. *Am. J. Respir. Crit. Care Med.* **180**, 138–145. (doi:10.1164/rccm.200812-19430C)
130. Harrison F, Paul J, Massey RC, Buckling A. 2008 Interspecific competition and siderophore-mediated cooperation in *Pseudomonas aeruginosa*. *Isme J.* **2**, 49–55. (doi:10.1038/ismej.2007.96)
131. Grant AJ, Restif O, McKinley TJ, Sheppard M, Maskell DJ, Mastroeni P. 2008 Modelling within-host spatiotemporal dynamics of invasive bacterial disease. *PLoS Biol.* **6**, e74. (doi:10.1371/journal.pbio.0060074)
132. Mitrí S, Xavier J, Foster KR. 2011 Social evolution in multispecies biofilms. *Proc. Natl Acad. Sci. USA* **108**(Suppl. 2), 10 839–10 846. (doi:10.1073/pnas.1100292108)
133. Popat R, Cruz SA, Messina M, Williams P, West SA, Diggle SP. 2012 Quorum-sensing and cheating in bacterial biofilms. *Proc. R. Soc. B* **279**, 4765–4771. (doi:10.1098/rspb.2012.1976)
134. Nadell CD, Bassler BL. 2011 A fitness trade-off between local competition and dispersal in *Vibrio cholerae* biofilms. *Proc. Natl Acad. Sci. USA* **108**, 14 181–14 185. (doi:10.1073/pnas.1111147108)
135. The Human Microbiome Project Consortium. 2012 Structure, function and diversity of the healthy human microbiome. *Nature* **486**, 207–214. (doi:10.1038/nature11234)
136. Balmer O, Stearns SC, Schötzau A, Brun R. 2009 Intraspecific competition between co-infecting parasite strains enhances host survival in African trypanosomes. *Ecology* **90**, 3367–3378. (doi:10.1890/08-2291.1)
137. Eswarappa SM, Estrela S, Brown SP. 2012 Within-host dynamics of multi-species infections: facilitation, competition and virulence. *PLoS ONE* **7**, e38730. (doi:10.1371/journal.pone.0038730)
138. Vardo AM, Kaufhold KD, Schall JJ. 2007 Experimental test for preinfection in a lizard malaria parasite (*Plasmodium mexicanum*). *J. Parasitol.* **93**, 280–282. (doi:10.1645/GE-1005R.1)
139. Portugal S *et al.* 2011 Host-mediated regulation of superinfection in malaria. *Nat. Med.* **17**, 732–737. (doi:10.1038/nm.2368)
140. McQueen PG, McKenzie FE. 2006 Competition for red blood cells can enhance *Plasmodium vivax* parasitemia in mixed-species malaria infections. *Am. J. Trop. Med. Hyg.* **75**, 112–125.
141. Pedersen AB, Fenton A. 2007 Emphasizing the ecology in parasite community ecology. *Trends Ecol. Evol.* **22**, 133–139. (doi:10.1016/j.tree.2006.11.005)
142. Estrela S, Brown SP. 2013 Metabolic and demographic feedbacks shape the emergent spatial structure and function of microbial communities. *PLoS Comp. Biol.* **9**, e1003398. (doi:10.1371/journal.pcbi.1003398)
143. Hoffmann AA *et al.* 2011 Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature* **476**, 454–457. (doi:10.1038/nature10356)
144. Poulin R, Morand S. 2000 The diversity of parasites. *Q. Rev. Biol.* **75**, 277–293. (doi:10.1086/393500)
145. Foster KR. 2005 Hamiltonian medicine: why the social lives of pathogens matter. *Science* **308**, 1269–1270. (doi:10.1126/science.1108158)
146. Rumbaugh KP, Diggle SP, Watters CM, Ross-Gillespie A, Griffin AS, West SA. 2009 Quorum sensing and the social evolution of bacterial virulence. *Curr. Biol.* **19**, 341–345. (doi:10.1016/j.cub.2009.01.050)
147. Read AF, Day T, Huijben S. 2011 The evolution of drug resistance and the curious orthodoxy of aggressive chemotherapy. *Proc. Natl Acad. Sci. USA* **108**, 10 871–10 877. (doi:10.1073/pnas.1100299108)
148. André JB, Godelle B. 2005 Multicellular organization in bacteria as a target for drug therapy. *Ecol. Lett.* **8**, 800–810. (doi:10.1111/j.1461-0248.2005.00783.x)
149. Mellbye B, Schuster M. 2011 The sociomicrobiology of antivirulence drug resistance: a proof of concept. *mBio* **2**, e00131. (doi:10.1128/mBio.00131-11)
150. Ross-Gillespie A, Weigert M, Brown SP, Kümmerli R. 2014 Gallium-mediated siderophore quenching as an evolutionarily robust antibacterial treatment. *Evol. Med. Publ. Health* **2014**, 18–29. (doi:10.1093/emph/eou003)
151. Allen RC, Popat R, Diggle SP, Brown SP. In press. Targeting virulence: can we make evolution-proof drugs? *Nat. Rev. Microbiol.* (doi:10.1038/nrmicro3232)
152. Carter LC, Mideo NM, Pollitt LC, Kafack BFC, Llinas M, Reece SE. 2013 Stress and sex in malaria parasites: why does commitment vary? *Evol. Med. Public Health* **2013**, 135–147. (doi:10.1093/emph/eot011)