

Published in final edited form as:

Handb Clin Neurol. 2009 ; 92: 283–305. doi:10.1016/S0072-9752(08)01914-3.

Cerebral vascular dysregulation in the ischemic brain

ALEXANDER KUNZ¹ and COSTANTINO IADECOLA^{2,*}

¹University Hospital Carl Gustav Carus, Dresden, Germany

²Weill Cornell Medical College, New York, NY, USA

The brain has limited fuel reserves and is highly dependent on a constant supply of oxygen and energy substrates delivered through blood flow (Hossmann, 1994). Interruption of cerebral blood flow (CBF) leads to brain dysfunction and, if the interruption is prolonged, brain death. Cerebral blood vessels are endowed with powerful regulatory mechanisms that assure that the brain is perfused at all times. However, during cerebral ischemia these mechanisms become dysfunctional and fail to compensate for the reduction in CBF. Thus cerebral ischemia produces profound alterations of the major control mechanisms governing the cerebral circulation. Such dysregulation, termed vasomotor paralysis (Langfitt et al., 1964; Hoedt-Rasmussen et al., 1967; Paulson, 1971), occurs in the setting of both focal cerebral ischemia, in which CBF is reduced in a restricted brain region, and global cerebral ischemia, in which CBF is globally reduced throughout the brain. The cerebrovascular dysregulation undermines the ability of the brain to maintain CBF, aggravates the intensity of the ischemic insult, and amplifies the tissue damage. Therapies aimed at restoring cerebrovascular regulation after ischemia offer the opportunity to improve cerebral perfusion and limit ischemic injury. In this chapter, we will briefly review the effects of cerebral ischemia on the regulation of CBF, focusing on the potential mechanisms involved in the cerebrovascular dysfunction and on the implications for ischemic brain injury.

14.1. Effects of cerebral ischemia and reperfusion on cerebral blood flow

14.1.1. Cerebral blood flow after arterial occlusion

The introduction of techniques for measuring regional CBF provided new insights into the alterations of CBF distribution following focal or global cerebral ischemia. Occlusion of a major cerebral artery produces a severe reduction in CBF to the area of the brain supplied by that particular artery. Thus, the CBF reduction is greatest in the center of the ischemic territory. Surrounding this ‘core’ area of cerebral ischemia there is an area in which a less severe CBF reduction is found. This peripheral region of the ischemic territory is termed ischemic penumbra. Whereas brain tissue in the ischemic core is irreversibly damaged, the penumbra represents potentially salvageable brain tissue, in which neuronal activity is suppressed but the tissue is potentially viable (Astrup et al., 1981). The fate of the penumbra depends on the residual CBF and the duration of the flow reduction (Astrup et al., 1981; Heiss and Rosner, 1983). Numerous clinical studies have attempted to define the CBF values corresponding to the ischemic penumbra. Using different imaging techniques the range of CBF in the penumbra has been estimated to be between 12 and 22 ml/100 g per minute (Heiss et al., 2001). If CBF drops below this critical threshold for a sufficient period of time the tissue will be damaged.

14.1.2. Cerebral blood flow at reperfusion

In human ischemic stroke, the arterial occlusion can be either permanent or transient. It has been shown that an early re-establishment of the CBF in ischemic areas (reperfusion) can be associated with a favorable functional outcome (Minematsu et al., 1992; Ringelstein et al., 1992). On the other hand, reperfusion can also exacerbate brain injury through the development of space-occupying hemorrhagic transformations or cerebral edema (Nakagawa et al., 1990; Wang and Lo, 2003). Therefore, CBF changes at reperfusion have important pathophysiological implications. Focal cerebral ischemia induced by an acute occlusion of a cerebral artery leads to a pronounced decrease in CBF in the dependent brain area. After reopening the vessel occlusion, reperfusion in the ischemic core typically shows a biphasic pattern: a transient increase in CBF (post-ischemic hyperperfusion) followed by a more sustained reduction in CBF (hypoperfusion; Fig. 14.1). Post-ischemic hyperperfusion, also known as 'luxury perfusion', has long been demonstrated in animal stroke models and human stroke as well (Lassen, 1966; Sundt and Waltz, 1971; Pulsinelli et al., 1982; Traupe et al., 1982; Todd et al., 1986; Heiss et al., 1997; Marchal et al., 1999). Because of a lack of pre-ischemic CBF values, post-ischemic hyperperfusion is usually termed 'post-ischemic hyperemia' (Hoedt-Rasmussen et al., 1967; Olsen et al., 1981; Baron et al., 1989). The length of post-ischemic hyperperfusion is directly proportional to the duration of the ischemic phase (Gourley and Heistad, 1984). The hyperemic phase is not mediated by an increase in oxygen or glucose utilization (Gourley and Heistad, 1984) and can be attributed to abnormal vasodilation in the ischemic territory (Marchal et al., 1996, 1999). Such abnormal vasodilation has multiple causes, including lactic acidosis secondary to ischemia-induced anaerobic glycolysis (Rehncrona et al., 1981) and/or release of vasoactive mediators from the ischemic brain, including ions, metabolites, and reactive oxygen species (Berne et al., 1974; Traystman et al., 1991; Nelson et al., 1992; Silver and Erecinska, 1992; Wei et al., 1996).

The mechanisms leading to post-ischemic hypoperfusion are still unclear. Measurements of intravascular pressure and segmental vascular resistance in cats following global cerebral ischemia have shown that ischemia–reperfusion is followed by an increase in cerebrovascular resistance involving both extra- and intracranial arteries (Schmidt-Kastner et al., 1987). This increase in vascular resistance could be, in part, due to microvascular compression and vasospasm, as suggested by a scanning electron microscopy study of corrosion casts in rats following transient focal ischemia (Ohtake et al., 2004). In this study, microvessels appeared constricted, compressed, and narrowed, effects attributed to edema, hemorrhage, and vasospasm presumably induced by vasoconstrictors released from the ischemic brain (Ohtake et al., 2004). Additional causes of post-ischemic hypoperfusion may include reduced metabolic demand, reduced synthesis of parenchymal and endothelial vasoactive agents, and intravascular plugging by platelets and leukocytes (Hossmann, 1983; del Zoppo and Mabuchi, 2003). A deficit of the potent vasodilator nitric oxide (NO) does not seem to be involved because inhibition of NO synthesis reduces CBF further, suggesting that NO production helps to counteract the post-ischemic increase in vascular resistance (Clavier et al., 1994). Furthermore, administration of prostacyclin does not normalize post-ischemic flow, suggesting that the hypoperfusion is not due to a lack of endothelial-derived prostacyclin (van den Kerckhoff et al., 1983).

14.2. Cerebrovascular autoregulation

Cerebral blood flow is relatively independent of changes in mean arterial pressure with a certain range. The lower and upper limits of CBF autoregulation correspond to mean arterial pressures of approximately 50–60 and 150–160 mmHg respectively (Chillon and Baumbach, 2002; see Lassen, 1959; Heistad and Kontos, 1983; Paulson et al., 1990 for review) (Fig. 14.2). This property of the cerebral circulation, termed auto-regulation, is also a

characteristic of the circulation of other organs (Paulson et al., 1990). The vascular adjustments underlying autoregulation consist of a constriction of cerebral resistance vessels when cerebral perfusion pressure (arterial pressure minus intracranial pressure) increases and a vasodilation of these vessels when perfusion pressure decreases (Kontos et al., 1978). Thus, the changes in cerebral perfusion pressure induced by changes in arterial pressure are counteracted by changes in cerebrovascular resistance, which tend to keep CBF constant. However, if the change in arterial pressure exceeds the capacity of the vessels to compensate, autoregulation is lost. Thus, increases in arterial pressure above the upper limit of autoregulation lead to increases in CBF, while decreases below the lower limit lead to reductions in CBF (Chillon and Baumbach, 2002). Furthermore, rapid variations in arterial pressure are not compensated for by autoregulation and produce changes in CBF (Aaslid et al., 1989; Florence and Seylaz, 1992). This is because the vascular adjustments initiated by arterial pressure are not instantaneous and take several seconds to take effect. The cellular basis of cerebrovascular autoregulation resides in the intrinsic property of vascular smooth muscles to react to changes in transmural pressure (myogenic tone; Bayliss, 1902; Harder et al., 1995; Wellman et al., 2002). Neural activity, hypoxia, hypo- and hypercapnia, or stimulation of cerebrovascular nerves can modulate the intrinsic vascular myogenic response and may influence the rapidity of the vascular adjustments, the slope of the pressure–flow relationship, and the range of pressures over which CBF is autoregulated (Heistad and Kontos, 1983; Busija and Heistad, 1984; Paulson et al., 1990).

14.2.1. Effect of cerebral ischemia on autoregulation: animal studies

It is well established that cerebral ischemia impairs autoregulation. Whereas most experimental studies have investigated the effect of either hypertension or hypotension on CBF, a few studies have tested the full range of pressures over which CBF is autoregulated (Table 14.1). Experimental studies in different species have investigated cerebrovascular autoregulation during reperfusion following transient cerebral ischemia. One such study investigated CBF autoregulation before, during, and after transient bilateral carotid artery ligation in rats (Shiokawa et al., 1986). Resting CBF was markedly reduced 30 min after ligation and autoregulation during hypotension was severely impaired, resulting in almost zero flow. Such autoregulatory dysfunction persisted, even though less pronounced, in the acute phase after restoration of CBF to the ischemic lesion (Shiokawa et al., 1986). In the phase of hyperperfusion following transient global cerebral ischemia in dogs, autoregulation was present but attenuated (Christopherson et al., 1993). In that study, the CBF response to CO₂ was also markedly attenuated. This finding is at variance with other investigations that demonstrated restored autoregulation during the phase of post-ischemic hyperperfusion while the CO₂ response is still reduced (Hossmann et al., 1973; Nemoto et al., 1975). Perhaps, this discrepancy can be attributed to the different methods used for the induction of cerebral ischemia.

The upper limit of autoregulation was impaired within the first hour after permanent middle cerebral artery occlusion in cats (Shima et al., 1983). However, 2 h after ischemia, autoregulation was partially restored (Shima et al., 1983). Measurement of pial artery pressure revealed that autoregulation in non-occluded collateral arteries supplying the ischemic territory is preserved (Shima et al., 1983). Therefore, collateral pathways could partially compensate for the CBF reduction following middle cerebral artery occlusion (Shima et al., 1983). Other studies found that autoregulation remains markedly impaired for several years after cerebral ischemia. For example, Symon and colleagues (1975) investigated autoregulation during hypotension in baboons 3 years after focal cerebral ischemia. They found that autoregulation was abolished in the infarct itself but it was present, albeit significantly impaired, in brain areas surrounding the infarcted tissue.

The degree of impairment of autoregulation is related to the magnitude of the CBF reduction. Dirnagl and Pulsinelli (1990) tested autoregulation during hypertension and hypotension in rats following permanent combined middle cerebral artery and common carotid artery occlusion. Autoregulation was attenuated in areas in which CBF fell to levels between 30% and 60% of preocclusion values and was lost in areas where CBF was below 30%. Similarly, in a study using intra-arterial injection of silicone for the occlusion of intracranial arteries, autoregulation was more or less impaired depending on the proximity of the region to the ischemic core, where the CBF reduction is most pronounced (Lauer et al., 1998). Another important finding that emerged from experimental studies is that in the ischemic brain autoregulation is also impaired in remote non-ischemic areas. For example, moderate hypotension 24 hours after transient focal cerebral ischemia in rats demonstrated that autoregulation is impaired in the normally perfused peri-infarct cortex and subcortical white matter, as well as in the subcortical white matter of the contralateral hemisphere (MacGregor et al., 2000). In another study, the lower limit of autoregulation was reduced in the rat cerebellum after transient bilateral carotid occlusion (Shiokawa et al., 1986). These remote effects are likely to be mechanistically related to the concept of diaschisis (see section 14.4.1).

14.2.2. Human studies

Alterations of cerebrovascular autoregulation following ischemic brain injury have also been investigated in humans (Table 14.2). The use of positron emission tomography (PET) provided valuable insights into the compensatory mechanisms that maintain cerebral perfusion in the initial stages of CBF reduction before brain injury occurs. Powers (1991) categorized these cerebrovascular adjustments into three stages: stage 0, when cerebral perfusion pressure is normal; stage 1, when cerebral perfusion pressure is reduced and autoregulation dilates cerebral vessels to maintain CBF; and stage 2, when the compensatory capacity for cerebral vasodilation is exceeded, CBF begins to decrease, and oxygen extraction fraction increases to maintain the cerebral metabolic rate of O₂ (CMRO₂) (Powers, 1991). In stage 2 cerebrovascular autoregulation is disrupted.

In patients with acute stroke, all studies found an impaired autoregulation in the affected vascular territory, even though different techniques were used for CBF measurement and estimation of autoregulation (Olsen et al., 1981; Heiss et al., 1992; Eames et al., 2002; Immink et al., 2005). Immink et al. demonstrated, using measurement of dynamic cerebral autoregulation, that patients with acute unilateral lacunar strokes have bilaterally impaired autoregulation, whereas, in patients with an acute middle cerebral artery territory stroke, autoregulation is affected only in the ischemic hemisphere (Immink et al., 2005). These results are consistent with the assumption that patients with acute lacunar strokes suffer from bilateral small vessel disease and are therefore at increased risk of further ischemic events. In addition, autoregulation was transiently impaired in patients who had recently suffered a transient ischemic attack, and the autoregulatory impairment outlasted the clinical symptoms (Skinhoj et al., 1970). Moreover, Olsen et al. (1983) found an impairment of autoregulation in collateral arteries supplying the penumbra in patients less than 72 h after middle cerebral artery occlusion. Remote disturbances of autoregulation in the contralateral hemisphere of two patients with subacute (< 6 days) cerebral ischemia have also been described (Paulson et al., 1972). However, these patients had space-occupying lesions and the possibility that the remote effect was due to an increase of intracranial pressure could not be ruled out.

Autoregulation is also impaired by stenosis or occlusion of cerebral arteries. Two investigations using transcranial Doppler sonography found that cerebral autoregulation is impaired in patients with critical internal carotid artery stenoses or occlusions (Gooskens et al., 2003; Reinhard et al., 2003). Using two different paradigms to assess autoregulation, both groups found a correlation between degree of stenosis and loss of autoregulation.

Furthermore, both studies concluded that the impairment in autoregulation is more severe in cases with bilateral internal carotid artery stenoses or occlusions, and that the alteration also depends on the collateral supply through the circle of Willis. Furthermore, Haubrich et al. (2007) demonstrated in a cohort of elderly patients that impaired cerebral auto-regulation may recover after carotid angioplasty. Gong et al. (2006) measured autoregulation in patients with middle cerebral artery stenosis and found impaired dynamic autoregulation, which correlated with the degree of stenosis, insufficient collateralization, and the severity of ischemic stroke. Furthermore, autoregulation improved after middle cerebral artery stenting/angioplasty (Gong et al., 2006). Additional evidence of autoregulatory compromise in patients with internal carotid artery or middle cerebral artery occlusion was provided by Ouchi et al. (2001) using PET to measure CBF. A change in posture from supine to sitting was used as the autoregulatory challenge. Assuming that the sitting position was associated with a reduction in CBF and an increase in oxygen extraction fraction in regions ipsilateral to the stenotic vessels provided evidence for impaired autoregulation in post-stenotic resistance vessels. Similar results were described in a patient with recurrent transient ischemic attacks who had a critical ipsilateral internal carotid artery stenosis and a contralateral internal carotid artery occlusion and in whom CBF was measured during hypotension using xenon-133 inhalation (Tatemichi et al., 1990).

In summary, cerebral ischemia in animals, as in humans, impairs cerebrovascular autoregulation. The degree of impairment depends on the magnitude of the local CBF reduction but autoregulation is also impaired in remote non-ischemic regions. Although the temporal profile of the autoregulatory impairment following ischemic stroke suggests a trend towards improvement over time, severe disturbances in auto-regulation have also been observed in chronic stroke. In addition to acute cerebral ischemia, critical stenosis of cerebral arteries is also capable of compromising cerebrovascular autoregulation. Preliminary evidence suggests that stenting or angioplasty of diseased cerebral vessels may ameliorate the autoregulatory deficit.

14.3. Cerebrovascular reactivity to hypercapnia

Arterial partial pressure of carbon dioxide (P_{CO_2}) is a potent vasoactive stimulus for the cerebral circulation. Hypercapnia produces vasodilation whereas hypocapnia produces vasoconstriction. The cerebrovascular effects of hypercapnia are thought to be mediated by the changes in perivascular pH that accompany increases in P_{CO_2} (Kontos et al., 1977). The mechanisms of the pH effect are multifactorial and involve direct effects on smooth muscle cells via adenosine triphosphate (ATP)-sensitive K^+ channels (Faraci et al., 1994), prostanoids derived from cyclooxygenase-1 (Niwa et al., 2001), and NO (Iadecola, 1992; Faraci et al., 1994; Sandor et al., 1994; Thompson et al., 1996).

14.3.1. Effect of cerebral ischemia on cerebrovascular reactivity: animal studies

The cerebrovascular reactivity to changes in P_{CO_2} is impaired following focal or global cerebral ischemia (Table 14.3). Most studies have tested CBF reactivity to hypercapnia, while a few also examined the effects of hypocapnia (Nemoto et al., 1975; Jones et al., 1989; Christopherson et al., 1993). With permanent ischemia several studies found that CO_2 reactivity is abolished in the ischemic core but is still present but reduced in the brain tissue surrounding the core area (Symon et al., 1975; Jones et al., 1989; Dettmers et al., 1993; Harris et al., 2001). However, in transient ischemia, improvements in CO_2 reactivity have been reported during reperfusion (Koch et al., 1984; Schmidt-Kastner et al., 1986; Ono et al., 1997; Olah et al., 2000). With reperfusion, CO_2 reactivity recovered faster in penumbral regions whereas the ischemic core showed a lack of recovery (Olah et al., 2000). However, in the penumbra CO_2 reactivity, while improved, remained below normal values (Olah et al., 2000). The impairment of CBF reactivity to CO_2 is present despite full recovery of energy

metabolism (Olah et al., 2000). During post-ischemic hypoperfusion, most investigators found an attenuated or even abolished response to CO₂ (Nemoto et al., 1975; van den Kerckhoff et al., 1983; Schmidt-Kastner et al., 1987; Leffler et al., 1989; Christopherson et al., 1993). Most investigations suggest an association between persistently impaired CO₂ reactivity and irreversible brain damage (Seki et al., 1984; Schmidt-Kastner et al., 1986).

In some cases, hypercapnia produced a paradoxical reduction in CBF (Shima et al., 1983; Ono et al., 1997; Olah et al., 2000). Such reversal of hypercapnic vasodilation to vasoconstriction has been attributed to a 'steal' of blood flow into neighboring areas where CO₂ reactivity is retained (Symon et al., 1974). Although this phenomenon has been confirmed in some animal studies (Waltz, 1970; Paulson, 1971; Symon et al., 1971; Ott et al., 1975; Olah et al., 2000), other studies could not find evidence for an intracerebral steal in hypercapnia (Olsen et al., 1983; Jones et al., 1989) or during administration of cerebrovasodilators (Date and Hossmann, 1984; Gogolak et al., 1985). Therefore, the existence of an intracerebral steal phenomenon during hypercapnia or vasodilator administration remains controversial. Like cerebrovascular autoregulation, CBF reactivity to CO₂ is also altered in regions remote from the ischemic lesion. For example, in baboons it has been shown that the CO₂ reactivity 6 h after permanent middle cerebral artery occlusion is also attenuated in non-infarcted, non-penumbra tissue supplied by the affected middle cerebral artery (Dettmers et al., 1993).

14.3.2. Human studies

Consistent with findings obtained in animal studies, investigations in patients with ischemic stroke have demonstrated alterations in the cerebrovascular reactivity to changes in P_{CO_2} (Table 14.4). Olsen et al. (1981) examined cerebrovascular reactivity to hyper- and hypocapnia in hyperemic areas of nine patients with acute ischemic stroke and found an impaired CO₂ response in four of these patients. Other studies have shown a more consistent attenuation in CO₂ reactivity. For example, Novak and colleagues (2003) used transcranial Doppler sonography for the analysis of CO₂ reactivity in 20 patients with minor stroke and found an impaired response in the ipsilateral hemisphere.

Simultaneous measurements of cerebrovascular autoregulation and CO₂ reactivity revealed that CO₂ reactivity is better preserved than autoregulation in hemodynamically compromised brain areas (Paulson et al., 1972). Thus in this study, a state of dissociated vasoparalysis with impaired autoregulation and preserved CO₂ reactivity was described. The authors stated that dissociated vasoparalysis might represent a stage in a gradual change between normal vasomotor function and complete vasoparalysis (Paulson et al., 1972).

As in animal studies, remote alterations in cerebrovascular CO₂ reactivity have also been found in humans. In some cases, these remote alterations could be related to the phenomenon of diaschisis (see section 14.4.1). For example, in patients with capsular infarcts the CBF response to hypocapnia was reduced and the response to hypercapnia was enhanced in a restricted ipsilateral cortical area with reduced basal CBF, presumably because of diaschisis (Takano et al., 1988). More diffuse changes in CBF reactivity to CO₂ have also been reported in patients with cerebrovascular diseases. Maeda and colleagues (1993) compared CO₂ reactivity in patients with cortical or lacunar infarction with those of normal subjects. Patients with cortical infarctions exhibited reduced CO₂ reactivity in the ischemic hemisphere compared to normal control subjects. In contrast, patients with subcortical infarctions exhibited a diffuse attenuation in CO₂ responses in both hemispheres. These findings have been confirmed more recently (de Leeuw et al., 2003). Based on the finding of a reduced CO₂ response in the non-ischemic hemisphere also, it was concluded that this impaired CO₂ reactivity is a marker of generalized small-vessel disease. Another example of generalized impaired CO₂ reactivity was provided by Yamamoto and colleagues

(1980). In this study, the hemispheric response to hypercapnia was significantly reduced in patients with hemispheric infarction as well as in patients with vertebrobasilar ischemia. Similarly, Levine et al. (1988) demonstrated reduced CO₂ reactivity in the ischemic hemisphere of patients with transient ischemic attack or minor ischemic stroke, as well as in the nonischemic hemisphere of stroke patients. The influence of generalized pathological alterations in the cerebral vasculature on the CBF reactivity to CO₂ has also been demonstrated in patients with cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL) in whom a profound attenuation was reported (Pfefferkorn et al., 2001). These findings underline the importance of the functional state of cerebral blood vessels in the mechanisms of the vascular reactivity to CO₂.

As noted for cerebrovascular autoregulation, CBF reactivity to CO₂ is reduced in patients with stenosis or occlusion of one or more large cerebral arteries (Clifton et al., 1988; Keyeux et al., 1988; Tatemichi et al., 1990; Miller et al., 1992; Bakker et al., 2003). Several studies found a correlation between the degree of internal carotid artery stenosis, the number of affected arteries, and the impairment in CO₂ reactivity (Brown et al., 1986; Levine et al., 1991; Gooskens et al., 2003; Reinhard et al., 2003). Furthermore, some studies considered the degree of collateral circulation to be an important factor for the preservation of CO₂ reactivity. Accordingly, it was found that the CO₂ reactivity is more impaired in patients with poor collateral blood supply (Norrving et al., 1982; Bullock et al., 1985; Ringelstein et al., 1988, 1994). Two PET studies correlated CO₂ reactivity with CBF, cerebral blood volume (CBV) and oxygen extraction fraction in patients with occlusive carotid artery disease. In one study, a positive correlation between the CBF/CBV ratio and CO₂ reactivity was found, while oxygen extraction fraction was negatively correlated with CO₂ reactivity in most patients (Herold et al., 1988). In the other study, a negative correlation between oxygen extraction fraction and hypercapnic response was observed, a finding suggesting a relationship between reduced vasodilatory capacity of cerebral arteries and increase in oxygen extraction fraction. Using this correlation, Kanno et al. (1988) defined the oxygen extraction fraction value that corresponds to exhausted vasodilatory capacity in cerebral arteries. Interestingly, this value coincided with the upper limit of the oxygen extraction fraction measured in normal tissue. The robust correlation between CBF response to CO₂ and oxygen extraction fraction indicates that CO₂ reactivity is a suitable method for assessing the cerebrovascular and metabolic reserves of the hemodynamically compromised brain (Herold et al., 1988; Kanno et al., 1988).

An alternative approach to assess cerebrovascular reactivity in humans is the intravenous injection of acetazolamide, a carbonic anhydrase inhibitor. Administration of acetazolamide produces a gradual decline in brain pH (Heuser et al., 1975). Acidosis is a potent stimulus for cerebral arteries to dilate. Considering that the cerebrovascular effects of arterial P_{CO₂} are mediated by the changes in perivascular pH (Kontos et al., 1977), the acetazolamide test simulates the cerebrovascular vasodilation induced by hypercapnia. Unlike CO₂, assessing cerebrovascular reactivity with acetazolamide does not require the assistance of the patient, which is an advantage in acutely ill stroke patients who might be unable to cooperate. In general, CO₂ and acetazolamide responses show similar results in patients with cerebrovascular diseases (Ringelstein et al., 1992). The CBF reactivity to acetazolamide is impaired in patients with transient ischemic attack (Chollet et al., 1989), occlusive internal carotid artery disease (Schreiber et al., 1998), and acute and chronic cerebral ischemic disease (Vorstrup et al., 1986; Nariai et al., 1995). Nemoto et al. (2007) published a PET study in which the relationship between the cerebrovascular acetazolamide reserve and the oxygen extraction fraction reactivity to acetazolamide was analyzed in patients with symptomatic internal carotid artery occlusion. The authors found a close correlation between hemispheric cerebrovascular reserve and oxygen extraction fraction reactivity. They suggest that the oxygen extraction fraction reactivity to acetazolamide may be a valuable measure of

hemodynamic compromise (Nemoto et al., 2007). However, by comparing acetazolamide and hypercapnic challenges, some studies found differences in the cerebrovascular reactivity to acetazolamide and hypercapnia in patients with cerebrovascular occlusive diseases (Kazumata et al., 1996). Therefore, the CBF reactivity to CO₂ and acetazolamide cannot always be considered to be identical.

In summary, cerebral ischemia impairs the reactivity of CBF to CO₂, and the degree of impairment is related to the magnitude of the local CBF reduction. CO₂ reactivity recovers after reperfusion in areas that are not damaged. Following focal cerebral ischemia, CO₂ reactivity is reduced in non-ischemic areas, suggesting a remote effect leading to impairment even in the normally perfused brain. Moreover, CO₂ reactivity is globally impaired in patients with diffuse microvascular disease. Because of its close correlation with indices of brain oxygen metabolism, cerebrovascular CO₂ reactivity may be a valuable tool to assess the vascular and metabolic reserves of the hemodynamically compromised brain.

14.4. Neurovascular coupling

There is substantial evidence that neural activity is a major factor controlling CBF. In the resting brain, there is a general correspondence between the flow of a given brain region and its rate of cerebral glucose utilization, a variable reflecting neural activity (Reivich, 1974; Sokoloff et al., 1977; Kuschinsky, 1987; Iadecola, 1993). Thus, regions with low glucose utilization, such as the white matter, have low flow, whereas regions with high glucose use, such as the auditory cortex, have high flow (Kuschinsky, 1987). In addition, when the activity of the brain is enhanced, either focally or globally, local blood flow increases in proportion to the intensity of the activation (Raichle et al., 1975; Greenberg et al., 1979; Tsubokawa et al., 1980; Iadecola et al., 1983; Fox and Raichle, 1984; Frostig et al., 1990). Conversely, if brain activity is decreased CBF decreases proportionally (Nilsson et al., 1978; Ueki et al., 1988). The close correspondence between CBF and neural activity reflects a homeostatic mechanism by which the delivery of substrates through blood flow, as well as the removal of metabolites produced by brain activity, are coupled to the energy demands of neurons and glia. Neuronal, glial, and vascular factors, acting in concert, mediate the increase in CBF produced by neural activity (Zonta et al., 2003; Cauli et al., 2004; Mulligan and MacVicar, 2004; Takano et al., 2006; Iadecola and Nedergaard, 2007). Neurons and astrocytes release vasoactive agents, such as ions (K⁺, H⁺), neurotransmitters (acetylcholine, vasoactive intestinal polypeptide, etc.), and neuromodulators (adenosine, nitric oxide, prostanoids, p450 metabolites) that act on local vessels to initiate the response (Iadecola, 1993; Iadecola and Nedergaard, 2007). Vascular cells, including endothelial cells, pericytes, and smooth muscle cells, transduce these neuronal and glial signals into changes in vascular resistance that ultimately mediate the increases in CBF (Iadecola and Nedergaard, 2007).

14.4.1. Effect of ischemia on neurovascular coupling: animal studies

Cerebral ischemia attenuates the CBF increase produced by functional activation (Table 14.5). Some studies have focused on the effect of focal or global ischemia on the coupling between CBF and cerebral glucose utilization in the resting state, while other studies have focused on the increase in CBF induced by brain activity. For example, Ginsberg et al. (1989) found reduced resting CBF and cerebral glucose utilization in the left barrel-field cortex and other left cortical regions in rats with a left frontal photothrombotic ischemia. Moreover, following contralateral vibrissa stimulation a marked suppression of the increase in CBF and cerebral glucose utilization produced by activation was observed throughout the barrel-field cortex (Ginsberg et al., 1989). In support of these findings, Kunz et al. (2007b) showed that the increase in CBF produced by whisker stimulation is markedly attenuated 1 h after transient focal cerebral ischemia in mice. Similarly, permanent middle cerebral artery occlusion attenuated the increase in CBF and estimated CMRO₂ produced by somatosensory

activation in rats, in which CBF was measured by magnetic resonance imaging (MRI) with arterial spin labeling (Shen et al., 2005). However, the increases in CBF and CMRO₂ produced by somatosensory activation were not attenuated 30 min after middle cerebral artery occlusion lasting 15 min, an ischemic challenge that did not lead to brain injury (Shen et al., 2005). Therefore, the alterations in neurovascular coupling are graded according to the intensity of the ischemic insult.

Duckrow et al. (1981) examined the effect of transient global ischemia on the activation-induced changes in the redox state of cytochrome a₃ in the rat cerebral cortex. Before ischemia, focal electrical stimulation of the cerebral cortex increased the oxidation of cytochrome a₃, and increased local blood volume, a parameter reflecting stimulus-induced hyperemia. After ischemia, the amplitude of the oxidative response and of the blood volume increase was reduced up to 30 min after ischemia. Also, the length of time required for the oxidative response to return to baseline was increased, despite full recovery of the mitochondrial baseline redox state. These observations provide evidence that increasing the work of the post-ischemic cortex unveils metabolic abnormalities that cannot be appreciated in the resting state. Another study investigated the correlation between hemodynamic parameters and field potentials evoked by somatosensory activation after transient global ischemia in the rat (Ueki et al., 1988). In the normal state, somatosensory stimulation increased CBF, cerebral glucose utilization, and local brain lactate concentration. By 3 h after global ischemia, the increases in CBF were suppressed while somatosensory evoked potentials had already reappeared (Ueki et al., 1988). No local changes in brain pH, lactate, glucose, and ATP were observed. Therefore, after global ischemia, both CBF and metabolic responses are depressed, despite recovery of somatosensory evoked potentials. The data suggest that, after ischemia, the metabolic workload of the cortex during functional activation is reduced and does not require increased delivery of substrates via an increase in CBF. The findings also indicate that ischemia disrupts the coupling between synaptic activity and CBF/cerebral glucose utilization, which may have long-term consequences for brain function.

Dietrich et al. (1986) investigated the time course of the ischemia-induced attenuation of the cerebral glucose utilization increase evoked by stimulation of the facial whiskers in the rat. They found that, 1 day after transient global ischemia, metabolic responses are severely depressed in the neocortex and thalamus but not in the trigeminal complex. Resting cerebral glucose utilization in the cortical barrel field and in the ventrobasal thalamus were also decreased. However, the activation-induced increase in cerebral glucose utilization returned to baseline 5 days after the ischemic insult. At this time, the spatial pattern of activation within the cortical barrel field was different from that of normal rats, suggesting that different local pathways were used. These observations support the idea that ischemia induces reorganization of cortical circuits in an attempt to compensate for the neuronal dysfunction and loss (Nudo, 2003). Alterations in regional blood flow and metabolism have also been described in brain areas remote from the ischemic lesion. These alterations are probably related to the concept of diaschisis (von Monakow, 1914), a term indicating brain dysfunction in areas remote from the ischemic regions, crossed cerebellar diaschisis being a typical example (Feeney and Baron, 1986). Gold and Lauritzen (2002) examined the neurophysiological basis of crossed cerebellar diaschisis in rat. Inactivation of the cerebral cortex by focal ischemia, spreading depression, or topical application of the Na⁺ channel blocker tetrodotoxin reduced contralateral Purkinje cells' spiking activity and cerebellar blood flow (Gold and Lauritzen, 2002). These cerebellar alterations, which represent the neurophysiological basis of crossed cerebellar diaschisis, were not related to reductions in the excitability of Purkinje cells or cerebellar vascular reactivity. Thus, crossed cerebellar diaschisis is mediated by deactivation of the cerebellar cortex resulting from interruption of neocortical excitatory inputs (Gold and Lauritzen, 2002). Therefore, the remote changes in

brain function produced by cerebral ischemia are likely to result from deafferentation rather than intrinsic changes in the neural circuitry or blood vessels of the remote region.

14.4.2. Human studies

Although several investigations have addressed the effects of ischemic stroke on neurovascular coupling, most of these studies have been performed in patients with chronic stroke (Table 14.6). Inao et al. (1998) used PET to examine patients with minor strokes and critical ipsilateral internal carotid–middle cerebral artery stenosis or occlusion. Despite an impaired ipsi-lateral acetazolamide response, the authors found a nearly normal CBF response to neural activation. Based on these results, it can be hypothesized that CBF response to neural activity is more robust than CBF reactivity to CO₂. Unfortunately, the authors only tested the acetazolamide response and not the CO₂ response or cerebrovascular autoregulation. Yamauchi et al. (2005) examined CBF responses in the primary visual cortex during visual stimulation in patients with internal carotid artery stenoses or occlusions using PET. Visual stimulation increased CBF in the primary visual cortex ipsilateral to the internal carotid artery lesion. However, the CBF increase in the surrounding regions was significantly reduced compared to the non-affected hemisphere. This finding raises the possibility that the local hemodynamic response evoked by visual stimulation might occur at the expense of perfusion of the surrounding region in brain areas with an already compromised vascular reactivity (Yamauchi et al., 2005).

Blood-oxygen-level-dependent (BOLD) functional MRI (fMRI) is a powerful tool for studying functional activation in humans. BOLD-fMRI is based on the paramagnetic property of deoxyhemoglobin. During functional activation, a local decrease in the concentration of deoxyhemoglobin is observed (Ogawa et al., 1990). This decrease in deoxyhemoglobin gives rise to the BOLD signal, which reflects the interplay between changes in CBF, CBV, and CMRO₂ during functional activation (Dirnagl et al., 2002). A few studies have examined functional activation using BOLD-fMRI in ischemic stroke patients. Krainik et al. (2005) investigated BOLD signal changes induced by manual tasks in patients who had fully recovered from stroke in the frontal lobe. The authors found that the BOLD signal changes in the sensorimotor cortex and supplementary motor area of the lesioned hemisphere was decreased despite the fact that these regions were anatomically intact. Interestingly, an impaired BOLD response to hypocapnia was a predictor of impaired BOLD response to functional activation as well. Based on these findings, the authors could provide a relationship between a reduced CO₂ reactivity in brain areas surrounding the ischemic infarct and an impaired neurovascular coupling. Similar findings were described in a population of patients with a lacunar stroke (Pineiro et al., 2002). These patients exhibited a lower BOLD signal increase and a lower rate of rise of the BOLD signal during a sequential finger-tapping task performed contralaterally to the ischemic lesion. These changes were also found in the non-lesioned hemisphere for the same task performed with the corresponding hand. Unfortunately, CBF reactivity to CO₂ was not investigated in these patients. Since CO₂ reactivity is globally impaired in patients with lacunar stroke (Maeda et al., 1993; de Leeuw et al., 2003), it can be assumed that cerebral small-vessel disease is a major factor responsible for the impairment of both CO₂ reactivity and neurovascular coupling. Also, Sakatani et al. (2003) described only limited activation areas with BOLD-fMRI in the sensorimotor cortex on the lesion side in ischemic stroke patients performing a motor task. In addition to BOLD-fMRI measurements, the authors performed near-infrared spectroscopy (NIRS) during the motor tasks, and they were able to measure the local changes in deoxyhemoglobin, oxyhemoglobin, and total hemoglobin. During the hand grasping task, increases in the focal concentrations of oxyand total hemoglobin were observed in the sensorimotor cortex, reflecting local increases in CBF. Also, a sustained increase in the concentration of deoxyhemoglobin was observed throughout the task in

stroke patients, whereas a decrease was observed in control subjects. The increased deoxyhemoglobin signal measured by NIRS in stroke patients was accompanied by reduced BOLD signal activation in sensorimotor cortex (Sakatani et al., 2003; Murata et al., 2006). Similar changes in NIRS signals were also found in the left prefrontal cortex of aphasic stroke patients (Sakatani et al., 1998). The mechanism of the increase in deoxyhemoglobin in stroke patients remains unclear, although the possibility of a more pronounced increase in oxygen consumption by neuronal activity in patients with ischemic stroke has been considered (Sakatani et al., 2003).

The relationship between BOLD-fMRI and neural activity after stroke was examined more directly by Rossini et al. (2004). These investigators studied neural activity detected by magnetoencephalography (MEG) and BOLD-fMRI in patients with a history of stroke or transient ischemic attack, using median nerve stimulation to activate the somatosensory cortex. While MEG signals were detected in the affected and the unaffected hemispheres, the corresponding BOLD signal was not observed. Such uncoupling between neurophysiological responses and BOLD signal was related to a reduced vasomotor reactivity as assessed by transcranial Doppler sonography during CO₂ inhalation. Therefore, even though CO₂ reactivity and functional hyperemia are impaired in stroke patients, the neural response evoked by the activation is still preserved (Rossini et al., 2004). Similar findings were described in a patient with bilateral internal carotid artery occlusions and unilateral vertebral artery occlusion (Rother et al., 2002).

In general, the results with BOLD-fMRI in patients with stroke differ from those with PET in that a reduction in the BOLD response to activation was usually found. Changes in regional CBF evoked by neural activation measured by PET are likely to reflect more precisely the hemodynamic changes. This is not surprising considering that the BOLD response is an indirect measure of CBF and also reflects CBV and CMRO₂ (Dirnagl et al., 2002; Ugurbil et al., 2003). This is especially true after stroke, where the normal relationship between CBF, CBV, and CMRO₂ may be altered by the disease process (Sakatani et al., 2003). Therefore, observations with fMRI need to be interpreted with caution because the BOLD signal may not reliably reflect CBF in the injured brain (Sakatani et al., 2003).

A few studies investigated BOLD responses in patients with stenosis or occlusion of major cerebral arteries. Bilecen et al. (2002) performed fMRI of the auditory cortex during auditory stimulation (1000 Hz, sine-tone) in acute stroke patients with unilateral internal carotid artery stenosis or occlusion and found a reduced BOLD response only in patients with ischemic lesions in the watershed areas between the anterior and middle or posterior and middle cerebral arteries. Patients without cerebral lesions or internal carotid artery disease had symmetric responses. The authors concluded that the presence of borderzone lesions in patients with reduced BOLD responses suggests that internal carotid artery stenosis is hemodynamically significant and contributes to the reduced BOLD response (Bilecen et al., 2002). Reduced or negative BOLD effects in patients with occlusive diseases of major cerebral arteries and without ischemic stroke were also described (Carusone et al., 2002; Hamzei et al., 2003). The negative BOLD effect is difficult to interpret and could reflect a more pronounced increase in CMRO₂ compared to CBF, as suggested by the observation that deoxyhemoglobin is increased in patients with stroke during activation (Sakatani et al., 2003).

In summary, acute cerebral ischemia in animals results in an attenuation of the CBF and metabolic responses induced by neural activation. Most studies in humans have been performed in patients with chronic stroke. While in PET and NIRS studies the increases in local CBF and metabolism evoked by activation were found to not be impaired, in studies using BOLD-fMRI an impaired cerebrovascular response to neural activation was found,

often coupled to a reduction in CO₂ reactivity. In addition, alterations in neurovascular coupling are also present in patients with generalized impairment of cerebrovascular reactivity due to cerebral small-vessel disease or occlusive disease of major cerebral arteries. Although BOLD-fMRI seems to be more sensitive to the neurovascular disturbance induced by stroke, the basic mechanisms of the BOLD response in the ischemic brain have not been completely elucidated and BOLD-fMRI findings need to be interpreted with caution.

14.5. Role of reactive oxygen species in the cerebrovascular dysregulation following cerebral ischemia

There is accumulating evidence that reactive oxygen species are a major contributor to the disturbances in the regulation of the cerebral vasculature following ischemic brain injury. Reactive oxygen species are molecules that often have an unpaired electron in the outer orbital ring. Cerebral ischemia results in an overproduction of reactive oxygen species, which overwhelms endogenous antioxidant enzymes (Traystman et al., 1991; Chan, 2001). The production of reactive oxygen species following cerebral ischemia has been demonstrated in several animal studies (Armstead et al., 1988; Cao et al., 1988; Kontos et al., 1992; Zhang and Piantadosi, 1994; Globus et al., 1995; Kil et al., 1996; Piantadosi and Zhang, 1996; Chan et al., 1998; Murakami et al., 1998; Kawase et al., 1999; Kim et al., 2000, 2001, 2002; Kunz et al., 2007a). A peak in reactive oxygen species production occurs in the early reperfusion period after global ischemia in cats (Nelson et al., 1992), raising the possibility that reperfusion provides the oxygen needed for the formation of reactive oxygen species. Similarly, in transient focal cerebral ischemia in mice, reactive oxygen species production exhibits a peak at 2 h after reperfusion and at 72 h as well (Kunz et al., 2007a). The late post-ischemic increase in reactive oxygen species formation may reflect a radical produced by inflammatory cells infiltrating the ischemic brain through the enzyme NADPH oxidase (Kunz et al., 2007a). However, reactive oxygen species production has also been reported to occur before reperfusion (Liu et al., 2003), indicating that reestablishment of CBF is not needed to provide the oxygen for the formation of reactive oxygen species.

14.5.1. Reactive oxygen species and ischemic cerebrovascular dysregulation

Reactive oxygen species, including superoxide, hydrogen peroxide, and peroxynitrite, are potent dilators of cerebral arterioles and may play a role in normal cerebrovascular regulation (Kontos, 2001; Niwa et al., 2001). However, higher concentrations produce vasoconstriction and disrupt cerebrovascular regulation (Traystman et al., 1991; Kontos, 2001; Iadecola, 2004; Faraci, 2005). There is growing evidence that the cerebrovascular dysregulation associated with several brain diseases is mediated by reactive oxygen species (Iadecola, 2004; Faraci, 2005; Girouard and Iadecola, 2006). For example, the cerebrovascular dys-regulation that occurs in Alzheimer's disease, diabetes, hyperhomocystinemia, and angiotensin-II-induced hypertension is mediated by reactive oxygen species (Mayhan and Sharpe, 1998; Zhang et al., 1998; Iadecola et al., 1999; Didion and Faraci, 2003; Kazama et al., 2004; Park et al., 2004, 2005; Girouard et al., 2006).

The impaired cerebrovascular regulation observed after cerebral ischemia is also related to vascular oxidative stress. Cerebral ischemia induces markers of oxidative damage in cerebral blood vessels (Chan et al., 1998; Maneen et al., 2006). The cerebrovascular response to acetylcholine, which is mediated by endothelial NO, is abolished following transient global cerebral ischemia in cats (Nelson et al., 1992) and is prevented by the superoxide scavengers superoxide dismutase and catalase. Additionally, the alterations in CO₂ reactivity are prevented by the administration of deferoxamine, an iron scavenger, suggesting a role also for the hydroxyl radical in cerebrovascular dysregulation after transient ischemia (Nelson et al., 1992). Free radical scavengers prevent the attenuation in

the cerebrovascular response to NMDA induced by global ischemia in piglets (Bari et al., 1996). Furthermore, intravenous administration of polyethylene-glycol-conjugated superoxide dismutase improves post-ischemic hypercapnic CBF after global cerebral ischemia in piglets (Kirsch et al., 1993). In a global ischemia model in dogs, combined administration of superoxide dismutase and deferoxamine reduced post-ischemic hypoperfusion and enhanced recovery of evoked potentials (Cerchiari et al., 1987). In the isolated rat middle cerebral artery, studied after ischemia reperfusion, the attenuation in the smooth muscle relaxation produced by activation of Kir2.x, a barium-sensitive inward rectifier K⁺ channel, is counteracted by a free radical scavenger (Petrault et al., 2004). Interestingly, in this model, the ischemia-induced attenuation in endothelium-dependent vasodilation to acetylcholine was not normalized by the free radical scavenger (Petrault et al., 2004), suggesting that factors other than reactive oxygen species may also be involved. Similar results were obtained in rodents with topical cortical application of acetylcholine *in situ* (Rosenblum and Wormley, 1995; Rosenblum, 1997). Subsequent experiments demonstrated that the endothelium-dependent dysfunction, but not the Kir2.x dysfunction, was ameliorated by inducing neutropenia prior to ischemia, implicating neutrophils in mechanisms of the vascular dysregulation, possibly through metalloprotease-induced disruption of microvessels (Petrault et al., 2004). Supporting a role of neutrophils, activation of intravascular neutrophils induces endothelial cell dysfunction in rabbits (Akopov et al., 1994). Therefore, although reactive oxygen species are important pathogenic effectors in the cerebrovascular dysregulation induced by ischemia, loss of microvascular integrity through metalloprotease activation, disruption of the extracellular matrix, and loss of integrins is also likely to play a role (del Zoppo and Mabuchi, 2003).

14.5.1.1. NADPH oxidase as a potential source of reactive oxygen species mediating cerebrovascular dysfunction—

There are several cellular and enzymatic sources of reactive oxygen species in the ischemic brain (Traystman et al., 1991; Chan, 2001). However, because reactive oxygen species diffuse poorly and are highly reactive, they tend to act at the site of formation (Stamler, 1996; Halliwell and Whiteman, 2004). It is therefore likely that the reactive oxygen species responsible for the cerebrovascular dysfunction are produced in cerebrovascular cells. The enzyme NADPH oxidase has recently emerged as a major source of reactive oxygen species in cerebral blood vessels (Cai et al., 2003). NADPH oxidase is a multi-unit enzyme initially described in phagocytic cells and responsible for reactive oxygen species production in leukocytes and macrophages (Babior, 2004; Lambeth, 2004). NADPH oxidase is composed of membrane-bound subunits (Nox and p22^{phox}) and cytoplasmic subunits (p40^{phox}, p47^{phox}, p67^{phox}, Rac1 and Rac2) (Lambeth, 2004). Activation stimuli lead to phosphorylation of p47^{phox}, which results in the assembly of the enzyme and production of superoxide (Vignais, 2002). NADPH oxidase is involved in production of reactive oxygen species and cerebrovascular dysfunction in several brain conditions associated with vascular oxidative stress, including models of Alzheimer's disease (Park et al., 2005), and with the cerebrovascular complications of angiotensin II hypertension, diabetes, and hyperhomocystinemia (Cooper et al., 2002; Faraci and Lentz, 2004; Kazama et al., 2004; Girouard et al., 2006). Mice lacking the Nox2 subunit of NADPH oxidase have reduced ischemic brain injury (Walder et al., 1997) and reduced reactive oxygen species production 72 h after middle cerebral artery occlusion (Kunz et al., 2007a) but it has not been established whether vascular factors contribute to the protection. Therefore additional studies are needed to clarify the role of NADPH oxidase in the mechanisms of the cerebrovascular dysregulation associated with cerebral ischemia.

14.5.1.2. Mechanisms by which reactive oxygen species mediate

cerebrovascular dysfunction—The mechanisms by which reactive oxygen species disrupt cerebrovascular regulation are diverse (Iadecola, 2004; Faraci, 2006). One

mechanism is based the interaction of superoxide with NO. NO reacts with superoxide highly efficiently, forming peroxynitrite (Beckman et al., 1990; Pacher et al., 2007). Recent studies implicate peroxynitrite in the cerebrovascular dysfunction induced by ischemia. Transient ischemia with reperfusion reduces the ability of the middle cerebral artery to constrict in response to increased intramural pressure (myogenic tone) (Cipolla et al., 1997). Such reperfusion-induced reduction in myogenic tone can account for the profound disruption of autoregulation observed after focal cerebral ischemia (see section 14.2). The effect was more pronounced for longer duration of reperfusion and was related to depolymerization of the vascular smooth muscle protein F-actin (Maneen et al., 2006). The loss of myogenic tone was associated with an increase in the peroxynitrite marker nitrotyrosine and could be reproduced by exogenous peroxynitrite (Maneen et al., 2006). In addition to its actions on myogenic tone, peroxynitrite has other deleterious effects on cerebrovascular regulation. First, peroxynitrite formation reduces the bioavailability of NO, resulting in an impairment of NO-dependent vasodilation (Adachi et al., 2004; Faraci, 2005). Peroxynitrite inhibits the enzyme prostacyclin synthase and reduces the ability of cerebral arteries to dilate through impaired synthesis of the potent vasodilator prostacyclin (Zou et al., 1999). Peroxynitrite also inhibits the mitochondrial isoform of superoxide dismutase, enhancing ischemia-induced oxidative stress (Guo et al., 2003). Furthermore, peroxynitrite induces DNA strand breaks, which activate the DNA repair enzyme poly-ADP-ribose polymerase and lead to endothelial cell dysfunction through energy depletion (Soriano et al., 2001). Reactive oxygen species and peroxynitrite oxidize tetrahydrobiopterin, a cofactor for the enzymatic activity of nitric oxide synthase. Deficiency of tetrahydrobiopterin results in uncoupling of nitric oxide synthase, a process by which this enzyme produces superoxide instead of NO and increases oxidative stress (Katusic, 2001). However, peroxynitrite is not uniformly deleterious and can also be beneficial in certain settings. For example, low levels of peroxynitrite are required for the neuroprotection induced by preconditioning with the proinflammatory mediator lipopolysaccharide (Kunz et al., 2007b). However, relatively large amounts of peroxynitrite produced during ischemia–reperfusion are likely to be damaging to the brain and blood vessels.

In summary, reactive oxygen species can alter post-ischemic cerebrovascular regulation through numerous factors, including depletion of vasodilators (NO, prostacyclin), enhanced oxidative stress and energy depletion. However, reactive oxygen species are not the only factor responsible for the cerebrovascular dysfunction and loss of microvascular integrity through disruption of extracellular matrix and integrins is also likely to play a role.

14.6. Conclusions

Cerebral ischemia results in a profound disruption of cerebrovascular regulatory mechanisms that assure that the brain is adequately perfused. Cerebral ischemia impairs cerebrovascular autoregulation, attenuates functional hyperemia, and alters hypercapnic vasodilation, critical mechanisms by which active brain cells maintain the homeostasis of their microenvironment through delivery of nutrients and removal of waste. These alterations are in large part mediated by a surge of free radicals, which leads to neurovascular oxidative stress and impairs the function of neurons, glia, and vascular cells. These alterations in vascular regulation render the brain more vulnerable to the changes in perfusion pressure and to the metabolic dysfunction induced by ischemia, thereby increasing the susceptibility of the tissue to injury. Therapeutic strategies targeting the cerebrovascular dysfunction induced by ischemia offer the opportunity to preserve post-ischemic vascular regulation and improve the outcome of cerebral ischemia. Such a vasoprotective approach, in combination with cytoprotective therapies, may open new avenues for the treatment of ischemic stroke.

References

- Aaslid R, Lindegaard KF, Sorteberg W, et al. Cerebral autoregulation dynamics in humans. *Stroke*. 1989; 20:45–52. [PubMed: 2492126]
- Adachi T, Weisbrod RM, Pimentel DR, et al. S-glutathiolation by peroxynitrite activates SERCA during arterial relaxation by nitric oxide. *Nat Med*. 2004; 10:1200–1207. [PubMed: 15489859]
- Akopov SE, Sercombe R, Seylaz J. Endothelial dysfunction in cerebral vessels following carotid artery infusion of phorbol ester in rabbits: the role of polymorphonuclear leukocytes. *J Cereb Blood Flow Metab*. 1994; 14:1078–1087. [PubMed: 7929651]
- Armstead WM, Mirro R, Busija DW, et al. Post-ischemic generation of superoxide anion by newborn pig brain. *Am J Physiol*. 1988; 255:H401–H403. [PubMed: 2841877]
- Astrup J, Siesjo BK, Symon L. Thresholds in cerebral ischemia—the ischemic penumbra. *Stroke*. 1981; 12:723–725. [PubMed: 6272455]
- Babior BM. NADPH oxidase. *Curr Opin Immunol*. 2004; 16:42–47. [PubMed: 14734109]
- Bakker FC, Klijn CJ, Jennekens-Schinkel A, et al. Cognitive impairment in patients with carotid artery occlusion and ipsilateral transient ischemic attacks. *J Neurol*. 2003; 250:1340–1347. [PubMed: 14648151]
- Bari F, Errico RA, Louis TM, et al. Differential effects of short-term hypoxia and hypercapnia on N-methyl-D-aspartate-induced cerebral vasodilatation in pigs. *Stroke*. 1996; 27:1634–1639. discussion 1639–1640. [PubMed: 8784141]
- Baron JC, Frackowiak RS, Herholz K, et al. Use of PET methods for measurement of cerebral energy metabolism and hemodynamics in cerebrovascular disease. *J Cereb Blood Flow Metab*. 1989; 9:723–742. [PubMed: 2684991]
- Bayliss WM. On the local reactions of the arterial wall to changes of internal pressure. *J Physiol*. 1902; 28:220–231. [PubMed: 16992618]
- Beckman JS, Beckman TW, Chen J, et al. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci U S A*. 1990; 87:1620–1624. [PubMed: 2154753]
- Berne RM, Rubio R, Curnish RR. Release of adenosine from ischemic brain: effect on cerebral vascular resistance and incorporation into adenine nucleotides. *Circ Res*. 1974; 35:262–271.
- Bilecen D, Radu EW, Schulte AC, et al. fMRI of the auditory cortex in patients with unilateral carotid artery steno-occlusive disease. *J Magn Reson Imaging*. 2002; 15:621–627. [PubMed: 12112511]
- Brown MM, Wade JP, Bishop CC, et al. Reactivity of the cerebral circulation in patients with carotid occlusion. *J Neurol Neurosurg Psychiatry*. 1986; 49:899–904. [PubMed: 3091771]
- Bullock R, Mendelow AD, Bone I, et al. Cerebral blood flow and CO₂ responsiveness as an indicator of collateral reserve capacity in patients with carotid arterial disease. *Br J Surg*. 1985; 72:348–351. [PubMed: 3922463]
- Busija DW, Heistad DD. Factors involved in the physiological regulation of the cerebral circulation. *Rev Physiol Biochem Pharmacol*. 1984; 101:161–211. [PubMed: 6441228]
- Cai H, Griendling KK, Harrison DG. The vascular NAD(P)H oxidases as therapeutic targets in cardiovascular diseases. *Trends Pharmacol Sci*. 2003; 24:471–478. [PubMed: 12967772]
- Cao W, Carney JM, Duchon A, et al. Oxygen free radical involvement in ischemia and reperfusion injury to brain. *Neurosci Lett*. 1988; 88:233–238. [PubMed: 3380359]
- Carusone LM, Srinivasan J, Gitelman DR, et al. Hemodynamic response changes in cerebrovascular disease: implications for functional MR imaging. *AJNR Am J Neuroradiol*. 2002; 23:1222–1228. [PubMed: 12169483]
- Cauli B, Tong XK, Rancillac A, et al. Cortical GABA interneurons in neurovascular coupling: relays for subcortical vasoactive pathways. *J Neurosci*. 2004; 24:8940–8949. [PubMed: 15483113]
- Cerchiari EL, Hoel TM, Safar P, et al. Protective effects of combined superoxide dismutase and deferoxamine on recovery of cerebral blood flow and function after cardiac arrest in dogs. *Stroke*. 1987; 18:869–878. [PubMed: 3629645]
- Chan PH. Reactive oxygen radicals in signaling and damage in the ischemic brain. *J Cereb Blood Flow Metab*. 2001; 21:2–14. [PubMed: 11149664]

- Chan PH, Kawase M, Murakami K, et al. Overexpression of SOD1 in transgenic rats protects vulnerable neurons against ischemic damage after global cerebral ischemia and reperfusion. *J Neurosci*. 1998; 18:8292–8299. [PubMed: 9763473]
- Chillon, JM.; Baumbach, GL. Autoregulation: arterial and intracranial pressure.. In: Edvinsson, L.; Krause, DN., editors. *Cerebral Blood Flow and Metabolism*. 2nd edn.. Lippincott Williams & Wilkins; Philadelphia, PA: 2002.
- Chollet F, Celsis P, Clanet M, et al. SPECT study of cerebral blood flow reactivity after acetazolamide in patients with transient ischemic attacks. *Stroke*. 1989; 20:458–464. [PubMed: 2784599]
- Christopherson TJ, Milde JH, Michenfelder JD. Cerebral vascular autoregulation and CO₂ reactivity following onset of the delayed posts ischemic hypoperfusion state in dogs. *J Cereb Blood Flow Metab*. 1993; 13:260–268. [PubMed: 8436617]
- Cipolla MJ, McCall AL, Lessov N, et al. Reperfusion decreases myogenic reactivity and alters middle cerebral artery function after focal cerebral ischemia in rats. *Stroke*. 1997; 28:176–180. [PubMed: 8996508]
- Clavier N, Kirsch JR, Hurn PD, et al. Effect of post-ischemic hypoperfusion on vasodilatory mechanisms in cats. *Am J Physiol*. 1994; 267:H2012–H2018. [PubMed: 7977832]
- Clifton GL, Haden HT, Taylor JR, et al. Cerebrovascular CO₂ reactivity after carotid artery occlusion. *J Neurosurg*. 1988; 69:24–28. [PubMed: 3132540]
- Cooper D, Stokes KY, Tailor A, et al. Oxidative stress promotes blood cell–endothelial cell interactions in the microcirculation. *Cardiovasc Toxicol*. 2002; 2:165–180. [PubMed: 12665663]
- Date H, Hossmann KA. Effect of vasodilating drugs on intracortical and extracortical vascular resistance following middle cerebral artery occlusion in cats. *Ann Neurol*. 1984; 16:330–336. [PubMed: 6385825]
- De Leeuw FE, Van Huffelen A, Kappelle J. Cerebrovascular reactivity in patients with a recent lacunar infarction. *J Neurol*. 2003; 250:232–233. [PubMed: 12622086]
- Del Zoppo GJ, Mabuchi T. Cerebral microvessel responses to focal ischemia. *J Cereb Blood Flow Metab*. 2003; 23:879–894. [PubMed: 12902832]
- Dettmers C, Young A, Rommel T, et al. CO₂ reactivity in the ischaemic core, penumbra, and normal tissue 6 hours after acute MCA-occlusion in primates. *Acta Neurochir (Wien)*. 1993; 125:150–155. [PubMed: 8122541]
- Didion SP, Faraci FM. Angiotensin II produces superoxide-mediated impairment of endothelial function in cerebral arterioles. *Stroke*. 2003; 34:2038–2042. [PubMed: 12829858]
- Dietrich WD, Ginsberg MD, Busto R. Effect of transient cerebral ischemia on metabolic activation of a somatosensory circuit. *J Cereb Blood Flow Metab*. 1986; 6:405–413. [PubMed: 3733900]
- Dimagl U, Pulsinelli W. Autoregulation of cerebral blood flow in experimental focal brain ischemia. *J Cereb Blood Flow Metab*. 1990; 10:327–336. [PubMed: 2329121]
- Dimagl, U.; Edvinsson, L.; Villringer, A. Measuring cerebral blood flow and metabolism.. In: Edvinsson, LE.; Krause, DN., editors. *Cerebral Blood Flow and Metabolism*. 2nd edn.. Lippincott Williams & Wilkins; Philadelphia, PA: 2002.
- Duckrow RB, Lamanna JS, Rosenthal M. Disparate recovery of resting and stimulated oxidative metabolism following transient ischemia. *Stroke*. 1981; 12:677–686. [PubMed: 6272454]
- Eames PJ, Blake MJ, Dawson SL, et al. Dynamic cerebral autoregulation and beat to beat blood pressure control are impaired in acute ischaemic stroke. *J Neurol Neurosurg Psychiatry*. 2002; 72:467–472. [PubMed: 11909905]
- Faraci FM. Oxidative stress: the curse that underlies cerebral vascular dysfunction? *Stroke*. 2005; 36:186–188. [PubMed: 15653575]
- Faraci FM. Reactive oxygen species: influence on cerebral vascular tone. *J Appl Physiol*. 2006; 100:739–743. [PubMed: 16421281]
- Faraci FM, Lentz SR. Hyperhomocysteinemia, oxidative stress, and cerebral vascular dysfunction. *Stroke*. 2004; 35:345–347. [PubMed: 14757874]
- Faraci FM, Breese KR, Heistad DD. Cerebral vasodilation during hypercapnia. Role of glibenclamide-sensitive potassium channels and nitric oxide. *Stroke*. 1994; 25:1679–1683. [PubMed: 8042220]
- Feeney DM, Baron JC. Diaschisis. *Stroke*. 1986; 17:817–830. [PubMed: 3532434]

- Florence G, Seylaz J. Rapid autoregulation of cerebral blood flow: a laser-Doppler flowmetry study. *J Cereb Blood Flow Metab.* 1992; 12:674–680. [PubMed: 1618945]
- Fox PT, Raichle ME. Stimulus rate dependence of regional cerebral blood flow in human striate cortex, demonstrated by positron emission tomography. *J Neurophysiol.* 1984; 51:1109–1120. [PubMed: 6610024]
- Frostig RD, Lieke EE, Ts'o DY, et al. Cortical functional architecture and local coupling between neuronal activity and the microcirculation revealed by in vivo high-resolution optical imaging of intrinsic signals. *Proc Natl Acad Sci U S A.* 1990; 87:6082–6086. [PubMed: 2117272]
- Ginsberg MD, Castella Y, Dietrich WD, et al. Acute thrombotic infarction suppresses metabolic activation of ipsilateral somatosensory cortex: evidence for functional diaschisis. *J Cereb Blood Flow Metab.* 1989; 9:329–341. [PubMed: 2715205]
- Girouard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease. *J Appl Physiol.* 2006; 100:328–335. [PubMed: 16357086]
- Girouard H, Park L, Anrather J, et al. Angiotensin II attenuates endothelium-dependent responses in the cerebral microcirculation through nox-2-derived radicals. *Arterioscler Thromb Vasc Biol.* 2006; 26:826–832. [PubMed: 16439707]
- Globus MY, Busto R, Lin B, et al. Detection of free radical activity during transient global ischemia and recirculation: effects of intraischemic brain temperature modulation. *J Neurochem.* 1995; 65:1250–1256. [PubMed: 7643104]
- Gogolak I, Gotoh F, Tomita M, et al. No intracerebral steal phenomenon in the ischemic brain following papaverine administration. *Stroke.* 1985; 16:114–117. [PubMed: 3966254]
- Gold L, Lauritzen M. Neuronal deactivation explains decreased cerebellar blood flow in response to focal cerebral ischemia or suppressed neocortical function. *Proc Natl Acad Sci U S A.* 2002; 99:7699–7704. [PubMed: 12032346]
- Gong XP, Li Y, Jiang WJ, et al. Impaired dynamic cerebral autoregulation in middle cerebral artery stenosis. *Neurol Res.* 2006; 28:76–81. [PubMed: 16464367]
- Gooskens I, Schmidt EA, Czosnyka M, et al. Pressure-autoregulation, CO₂ reactivity and asymmetry of haemodynamic parameters in patients with carotid artery stenotic disease. A clinical appraisal. *Acta Neurochir (Wien).* 2003; 145:527–532. [PubMed: 12910394]
- Gourley JK, Heistad DD. Characteristics of reactive hyperemia in the cerebral circulation. *Am J Physiol.* 1984; 246:H52–H58. [PubMed: 6696088]
- Greenberg J, Hand P, Sylvestro A, et al. Localized metabolic-flow couple during functional activity. *Acta Neurol Scand.* 1979; 72:12–13.
- Guo W, Adachi T, Matsui R, et al. Quantitative assessment of tyrosine nitration of manganese superoxide dismutase in angiotensin II-infused rat kidney. *Am J Physiol Heart Circ Physiol.* 2003; 285:H1396–H1403. [PubMed: 12791589]
- Halliwell B, Whiteman M. Measuring reactive species and oxidative damage in vivo and in cell culture: how should you do it and what do the results mean? *Br J Pharmacol.* 2004; 142:231–255. [PubMed: 15155533]
- Hamzei F, Knab R, Weiller C, et al. The influence of extra- and intracranial artery disease on the BOLD signal in fMRI. *Neuroimage.* 2003; 20:1393–1399. [PubMed: 14568508]
- Harder DR, Narayanan J, Gebremedhin D, et al. Transduction of physical forces by the vascular wall: role of phospholipase C and cytochrome P450 metabolites of arachidonic acid. *Trends Cardiovasc Med.* 1995; 5:7–14. [PubMed: 21232232]
- Harris NG, Lythgoe MF, Thomas DL, et al. Cerebrovascular reactivity following focal brain ischemia in the rat: a functional magnetic resonance imaging study. *Neuroimage.* 2001; 13:339–350. [PubMed: 11162274]
- Haubrich C, Kruska W, Diehl RR, et al. Recovery of the blood pressure—cerebral flow relation after carotid stenting in elderly patients. *Acta Neurochir (Wien).* 2007; 149:131–136. discussion 137. [PubMed: 16964557]
- Heiss WD, Rosner G. Functional recovery of cortical neurons as related to degree and duration of ischemia. *Ann Neurol.* 1983; 14:294–301. [PubMed: 6314871]
- Heiss WD, Huber M, Fink GR, et al. Progressive derangement of periinfarct viable tissue in ischemic stroke. *J Cereb Blood Flow Metab.* 1992; 12:193–203. [PubMed: 1548292]

- Heiss WD, Graf R, Lottgen J, et al. Repeat positron emission tomographic studies in transient middle cerebral artery occlusion in cats: residual perfusion and efficacy of postischemic reperfusion. *J Cereb Blood Flow Metab.* 1997; 17:388–400. [PubMed: 9143221]
- Heiss WD, Forsting M, Diener HC. Imaging in cerebrovascular disease. *Curr Opin Neurol.* 2001; 14:67–75. [PubMed: 11176220]
- Heistad, DD.; Kontos, H. Cerebral circulation.. In: Abboud, F.; Shephard, J., editors. *Handbook of Physiology: The Cardiovascular System.* American Physiological Society; Bethesda, MA: 1983.
- Herold S, Brown MM, Frackowiak RS, et al. Assessment of cerebral haemodynamic reserve: correlation between PET parameters and CO₂ reactivity measured by the intravenous 133 xenon injection technique. *J Neurol Neurosurg Psychiatry.* 1988; 51:1045–1050. [PubMed: 3145962]
- Heuser D, Astrup J, Lassen NA, et al. Brain carbonic acid acidosis after acetazolamide. *Acta Physiol Scand.* 1975; 93:385–390. [PubMed: 238362]
- Hoedt-Rasmussen K, Skinhoj E, Paulson O, et al. Regional cerebral blood flow in acute apoplexy. The ‘luxury perfusion syndrome’ of brain tissue. *Arch Neurol.* 1967; 17:271–281. [PubMed: 6053570]
- Hossmann, KA. Hemodynamics of postischemic reperfusion of the brain.. In: Weinstein, PR.; Faden, AL., editors. *Protection of the Brain from Ischemia.* Williams & Wilkins; Baltimore, MD.: 1983.
- Hossmann KA. Viability thresholds and the penumbra of focal ischemia. *Ann Neurol.* 1994; 36:557–565. [PubMed: 7944288]
- Hossmann KA, Lechtape-Gruter H, Hossmann V. The role of cerebral blood flow for the recovery of the brain after prolonged ischemia. *Z Neurol.* 1973; 204:281–299. [PubMed: 4155720]
- Iadecola C. Does nitric oxide mediate the increases in cerebral blood flow elicited by hypercapnia? *Proc Natl Acad Sci U S A.* 1992; 89:3913–3916. [PubMed: 1570313]
- Iadecola C. Regulation of the cerebral microcirculation during neural activity: is nitric oxide the missing link? *Trends Neurosci.* 1993; 16:206–214. [PubMed: 7688160]
- Iadecola C. Neurovascular regulation in the normal brain and in Alzheimer's disease. *Nat Rev Neurosci.* 2004; 5:347–360. [PubMed: 15100718]
- Iadecola C, Nedergaard M. Glial regulation of the cerebral microvasculature. *Nat Neurosci.* 2007; 10:1369–1376. [PubMed: 17965657]
- Iadecola C, Nakai M, Mraovitch S, et al. Global increase in cerebral metabolism and blood flow produced by focal electrical stimulation of dorsal medullary reticular formation in rat. *Brain Res.* 1983; 272:101–114. [PubMed: 6616188]
- Iadecola C, Zhang F, Niwa K, et al. SOD1 rescues cerebral endothelial dysfunction in mice overexpressing amyloid precursor protein. *Nat Neurosci.* 1999; 2:157–161. [PubMed: 10195200]
- Immink RV, Van Montfrans GA, Stam J, et al. Dynamic cerebral autoregulation in acute lacunar and middle cerebral artery territory ischemic stroke. *Stroke.* 2005; 36:2595–2600. [PubMed: 16254228]
- Inao S, Tadokoro M, Nishino M, et al. Neural activation of the brain with hemodynamic insufficiency. *J Cereb Blood Flow Metab.* 1998; 18:960–967. [PubMed: 9740099]
- Jones SC, Bose B, Furlan AJ, et al. CO₂ reactivity and heterogeneity of cerebral blood flow in ischemic, border zone, and normal cortex. *Am J Physiol.* 1989; 257:H473–H482. [PubMed: 2504058]
- Kanno I, Uemura K, Higano S, et al. Oxygen extraction fraction at maximally vasodilated tissue in the ischemic brain estimated from the regional CO₂ responsiveness measured by positron emission tomography. *J Cereb Blood Flow Metab.* 1988; 8:227–235. [PubMed: 3125186]
- Katusic ZS. Vascular endothelial dysfunction: does tetrahydrobiopterin play a role? *Am J Physiol Heart Circ Physiol.* 2001; 281:H981–H986. [PubMed: 11514262]
- Kawase M, Murakami K, Fujimura M, et al. Exacerbation of delayed cell injury after transient global ischemia in mutant mice with CuZn superoxide dismutase deficiency. *Stroke.* 1999; 30:1962–1968. [PubMed: 10471451]
- Kazama K, Anrather J, Zhou P, et al. Angiotensin II impairs neurovascular coupling in neocortex through NADPH oxidase-derived radicals. *Circ Res.* 2004; 95:1019–1026. [PubMed: 15499027]

- Kazumata K, Tanaka N, Ishikawa T, et al. Dissociation of vasoreactivity to acetazolamide and hypercapnia. Comparative study in patients with chronic occlusive major cerebral artery disease. *Stroke*. 1996; 27:2052–2058. [PubMed: 8898815]
- Keyeux A, Laterre C, Beckers C. Resting and hyper-capnic rCBF in patients with unilateral occlusive disease of the internal carotid artery. *J Nucl Med*. 1988; 29:311–319. [PubMed: 3126276]
- Kil HY, Zhang J, Piantadosi CA. Brain temperature alters hydroxyl radical production during cerebral ischemia/reperfusion in rats. *J Cereb Blood Flow Metab*. 1996; 16:100–106. [PubMed: 8530542]
- Kim GW, Sugawara T, Chan PH. Involvement of oxidative stress and caspase-3 in cortical infarction after photothrombotic ischemia in mice. *J Cereb Blood Flow Metab*. 2000; 20:1690–1701. [PubMed: 11129785]
- Kim GW, Lewen A, Copin J, et al. The cytosolic antioxidant, copper/zinc superoxide dismutase, attenuates blood–brain barrier disruption and oxidative cellular injury after photothrombotic cortical ischemia in mice. *Neuroscience*. 2001; 105:1007–1018. [PubMed: 11530238]
- Kim GW, Kondo T, Noshita N, et al. Manganese superoxide dismutase deficiency exacerbates cerebral infarction after focal cerebral ischemia/reperfusion in mice: implications for the production and role of superoxide radicals. *Stroke*. 2002; 33:809–815. [PubMed: 11872908]
- Kirsch JR, Helfaer MA, Haun SE, et al. Polyethylene glycol-conjugated superoxide dismutase improves recovery of postischemic hypercapnic cerebral blood flow in piglets. *Pediatr Res*. 1993; 34:530–537. [PubMed: 8255689]
- Koch KA, Jackson DL, Schmiedl M, et al. Total cerebral ischemia: effect of alterations in arterial PCO₂ on cerebral microcirculation. *J Cereb Blood Flow Metab*. 1984; 4:343–349. [PubMed: 6432807]
- Kontos HA. Oxygen radicals in cerebral ischemia: the 2001 Willis lecture. *Stroke*. 2001; 32:2712–2716. [PubMed: 11692043]
- Kontos CD, Wei EP, Williams JI, et al. Cytochemical detection of superoxide in cerebral inflammation and ischemia in vivo. *Am J Physiol*. 1992; 263:H1234–H1242. [PubMed: 1329563]
- Kontos HA, Raper AJ, Patterson JL. Analysis of vasoactivity of local pH, PCO₂ and bicarbonate on pial vessels. *Stroke*. 1977; 8:358–360. [PubMed: 16363]
- Kontos HA, Wei EP, Navari RM, et al. Responses of cerebral arteries and arterioles to acute hypotension and hypertension. *Am J Physiol*. 1978; 234:H371–H383. [PubMed: 645875]
- Krainik A, Hund-Georgiadis M, Zysset S, et al. Regional impairment of cerebrovascular reactivity and BOLD signal in adults after stroke. *Stroke*. 2005; 36:1146–1152. [PubMed: 15879326]
- Kunz A, Anrather J, Zhou P, et al. Cyclooxygenase-2 does not contribute to postischemic production of reactive oxygen species. *J Cereb Blood Flow Metab*. 2007a; 27:545–551. [PubMed: 16820798]
- Kunz A, Park L, Abe T, et al. Neurovascular protection by ischemic tolerance: role of nitric oxide and reactive oxygen species. *J Neurosci*. 2007b; 27:7083–7093. [PubMed: 17611261]
- Kuschinsky W. Coupling of function metabolism and blood flow in the brain. *News Physiol Sci*. 1987; 2:217–220.
- Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol*. 2004; 4:181–189. [PubMed: 15039755]
- Langfitt TW, Weinstein JD, Kassell NF. Cerebral vasomotor paralysis as a cause of brain swelling. *Trans Am Neurol Assoc*. 1964; 89:214–215. [PubMed: 4953351]
- Lassen NA. Cerebral blood flow and oxygen consumption in man. *Physiol Rev*. 1959; 39:183–238. [PubMed: 13645234]
- Lassen NA. The luxury-perfusion syndrome and its possible relation to acute metabolic acidosis localised within the brain. *Lancet*. 1966; 2:1113–1115. [PubMed: 4162534]
- Lauer, KK.; Shen, H.; Hudetz, AG., et al. Regional cerebral blood flow autoregulation in a silicone embolus model of focal brain ischemia as assessed by multichannel laser Doppler flowmetry.. In: Hudetz, AG.; Bruley, DF., editors. *Oxygen Transport to Tissue XX*. Plenum Press; New York: 1998.
- Leffler CW, Busija DW, Mirro R, et al. Effects of ischemia on brain blood flow and oxygen consumption of newborn pigs. *Am J Physiol*. 1989; 257:H1917–H1926. [PubMed: 2513731]

- Levine RL, Sunderland JJ, Lagreze HL, et al. Cerebral perfusion reserve indexes determined by fluoromethane positron emission scanning. *Stroke*. 1988; 19:19–27. [PubMed: 3122375]
- Levine RL, Dobkin JA, Rozental JM, et al. Blood flow reactivity to hypercapnia in strictly unilateral carotid disease: preliminary results. *J Neurol Neurosurg Psychiatry*. 1991; 54:204–209. [PubMed: 1903147]
- Liu S, Liu M, Peterson S, et al. Hydroxyl radical formation is greater in striatal core than in penumbra in a rat model of ischemic stroke. *J Neurosci Res*. 2003; 71:882–888. [PubMed: 12605415]
- MacGregor DG, Carswell HV, Graham DI, et al. Impaired cerebral autoregulation 24 h after induction of transient unilateral focal ischaemia in the rat. *Eur J Neurosci*. 2000; 12:58–66. [PubMed: 10651860]
- Maeda H, Matsumoto M, Handa N, et al. Reactivity of cerebral blood flow to carbon dioxide in various types of ischemic cerebrovascular disease: evaluation by the transcranial Doppler method. *Stroke*. 1993; 24:670–675. [PubMed: 8488521]
- Maneen MJ, Hannah R, Vitullo L, et al. Peroxynitrite diminishes myogenic activity and is associated with decreased vascular smooth muscle F-actin in rat posterior cerebral arteries. *Stroke*. 2006; 37:894–899. [PubMed: 16456123]
- Marchal G, Furlan M, Beaudouin V, et al. Early spontaneous hyperperfusion after stroke. A marker of favourable tissue outcome? *Brain*. 1996; 119:409–419. [PubMed: 8800936]
- Marchal G, Young AR, Baron JC. Early postischemic hyperperfusion: pathophysiologic insights from positron emission tomography. *J Cereb Blood Flow Metab*. 1999; 19:467–482. [PubMed: 10326714]
- Mayhan WG, Sharpe GM. Superoxide dismutase restores impaired histamine-induced increases in venular macromolecular efflux during diabetes mellitus. *Microcirculation*. 1998; 5:211–218. [PubMed: 9789261]
- Miller JD, Smith RR, Holaday HR. Carbon dioxide reactivity in the evaluation of cerebral ischemia. *Neuro-surgery*. 1992; 30:518–521.
- Minematsu K, Yamaguchi T, Omae T. ‘Spectacular shrinking deficit’: rapid recovery from a major hemispheric syndrome by migration of an embolus. *Neurology*. 1992; 42:157–162. [PubMed: 1734297]
- Mulligan SJ, MacVicar BA. Calcium transients in astrocyte endfeet cause cerebrovascular constrictions. *Nature*. 2004; 431:195–199. [PubMed: 15356633]
- Murakami K, Kondo T, Kawase M, et al. Mitochondrial susceptibility to oxidative stress exacerbates cerebral infarction that follows permanent focal cerebral ischemia in mutant mice with manganese superoxide dismutase deficiency. *J Neurosci*. 1998; 18:205–213. [PubMed: 9412501]
- Murata Y, Sakatani K, Katayama Y, et al. Increase in focal concentration of deoxyhaemoglobin during neuronal activity in cerebral ischaemic patients. *J Neurol Neurosurg Psychiatry*. 2002; 73:182–184. [PubMed: 12122179]
- Murata Y, Sakatani K, Hoshino T, et al. Effects of cerebral ischemia on evoked cerebral blood oxygenation responses and BOLD contrast functional MRI in stroke patients. *Stroke*. 2006; 37:2514–2520. [PubMed: 16946162]
- Nakagawa Y, Fujimoto N, Matsumoto K, et al. Morphological changes in acute cerebral ischemia after occlusion and reperfusion in the rat. *Adv Neurol*. 1990; 52:21–27. [PubMed: 2396514]
- Nariai T, Suzuki R, Hirakawa K, et al. Vascular reserve in chronic cerebral ischemia measured by the acetazolamide challenge test: comparison with positron emission tomography. *AJNR Am J Neuroradiol*. 1995; 16:563–570. [PubMed: 7793382]
- Nelson CW, Wei EP, Povlishock JT, et al. Oxygen radicals in cerebral ischemia. *Am J Physiol*. 1992; 263:H1356–H1362. [PubMed: 1332509]
- Nemoto EM, Snyder JV, Carroll RG, et al. Global ischemia in dogs: cerebrovascular CO₂ reactivity and autoregulation. *Stroke*. 1975; 6:425–431. [PubMed: 1154479]
- Nemoto EM, Yonas H, Pindzola RR, et al. PET OEF reactivity for hemodynamic compromise in occlusive vascular disease. *J Neuroimaging*. 2007; 17:54–60. [PubMed: 17238870]
- Nilsson, B.; Rehncrona, S.; Siesjo, BK. Ciba Foundation Symposium 56 (New series). Elsevier North-Holland; Amsterdam: 1978. Coupling of cerebral metabolism and blood flow in epileptic seizures, hypoxia and hypoglycemia. In: *Cerebral Vascular Smooth Muscle and Its Control*.

- Niwa K, Haensel C, Ross ME, et al. Cyclooxygenase-1 participates in selected vasodilator responses of the cerebral circulation. *Circ Res.* 2001; 88:600–608. [PubMed: 11282894]
- Norrving B, Nilsson B, Risberg J. rCBF in patients with carotid occlusion. Resting and hypercapnic flow related to collateral pattern. *Stroke.* 1982; 13:155–162. [PubMed: 6801822]
- Novak V, Chowdhary A, Farrar B, et al. Altered cerebral vasoregulation in hypertension and stroke. *Neurology.* 2003; 60:1657–1663. [PubMed: 12771258]
- Nudo RJ. Functional and structural plasticity in motor cortex: implications for stroke recovery. *Phys Med Rehabil Clin North Am.* 2003; 14:S57–S76.
- Ogawa S, Lee TM, Kay AR, et al. Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc Natl Acad Sci U S A.* 1990; 87:9868–9872. [PubMed: 2124706]
- Ohtake M, Morino S, Kaidoh T, et al. Three-dimensional structural changes in cerebral microvessels after transient focal cerebral ischemia in rats: scanning electron microscopic study of corrosion casts. *Neuropathology.* 2004; 24:219–227. [PubMed: 15484700]
- Olah L, Franke C, Schwandt W, et al. CO₂ reactivity measured by perfusion MRI during transient focal cerebral ischemia in rats. *Stroke.* 2000; 31:2236–2244. [PubMed: 10978058]
- Olsen TS, Larsen B, Skriver EB, et al. Focal cerebral hyperemia in acute stroke. Incidence, pathophysiology and clinical significance. *Stroke.* 1981; 12:598–607. [PubMed: 6975512]
- Olsen TS, Larsen B, Herning M, et al. Blood flow and vascular reactivity in collaterally perfused brain tissue. Evidence of an ischemic penumbra in patients with acute stroke. *Stroke.* 1983; 14:332–341. [PubMed: 6658900]
- Ono Y, Morikawa S, Inubushi T, et al. T2*-weighted magnetic resonance imaging of cerebrovascular reactivity in rat reversible focal cerebral ischemia. *Brain Res.* 1997; 744:207–215. [PubMed: 9027380]
- Ott EO, Abraham J, Meyer JS, et al. Regional cerebral blood flow measured by the gamma camera after direct injection of ¹³³Xe into the distal stump of the occluded middle cerebral artery. *Stroke.* 1975; 6:376–381. [PubMed: 1154474]
- Ouchi Y, Nobezawa S, Yoshikawa E, et al. Postural effects on brain hemodynamics in unilateral cerebral artery occlusive disease: a positron emission tomography study. *J Cereb Blood Flow Metab.* 2001; 21:1058–1066. [PubMed: 11524610]
- Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007; 87:315–424. [PubMed: 17237348]
- Park L, Anrather J, Forster C, et al. Ab-induced vascular oxidative stress and attenuation of functional hyper-emia in mouse somatosensory cortex. *J Cereb Blood Flow Metab.* 2004; 24:334–342. [PubMed: 15091114]
- Park L, Anrather J, Zhou P, et al. NADPH-oxidase-derived reactive oxygen species mediate the cerebrovascular dysfunction induced by the amyloid beta peptide. *J Neurosci.* 2005; 25:1769–1777. [PubMed: 15716413]
- Paulson OB. Cerebral apoplexy (stroke): pathogenesis, pathophysiology and therapy as illustrated by regional blood flow measurements in the brain. *Stroke.* 1971; 2:327–360. [PubMed: 4398838]
- Paulson OB, Olesen J, Christensen MS. Restoration of autoregulation of cerebral blood flow by hypocapnia. *Neurology.* 1972; 22:286–293. [PubMed: 5062263]
- Paulson OB, Strandgaard S, Edvinsson L. Cerebral autoregulation. *Cerebrovasc Brain Metab Rev.* 1990; 2:161–192. [PubMed: 2201348]
- Petrault O, Bastide M, Cotellet N, et al. The neuro-protective effect of the antioxidant flavonoid derivative ditert-butylhydroxyphenyl is parallel to the preventive effect on post-ischemic Kir2.x impairment but not to post-ischemic endothelial dysfunction. *Naunyn Schmiedebergs Arch Pharmacol.* 2004; 370:395–403. [PubMed: 15502971]
- Pfefferkorn T, Von Stuckrad-Barre S, Herzog J, et al. Reduced cerebrovascular CO₂ reactivity in CADASIL: a transcranial Doppler sonography study. *Stroke.* 2001; 32:17–21. [PubMed: 11136908]
- Piantadosi CA, Zhang J. Mitochondrial generation of reactive oxygen species after brain ischemia in the rat. *Stroke.* 1996; 27:327–331. discussion 332. [PubMed: 8571432]
- Pineiro R, Pendlebury S, Johansen-Berg H, et al. Altered hemodynamic responses in patients after subcortical stroke measured by functional MRI. *Stroke.* 2002; 33:103–109. [PubMed: 11779897]

- Powers WJ. Cerebral hemodynamics in ischemic cerebrovascular disease. *Ann Neurol.* 1991; 29:231–240. [PubMed: 2042939]
- Pulsinelli WA, Levy DE, Duffy TE. Regional cerebral blood flow and glucose metabolism following transient forebrain ischemia. *Ann Neurol.* 1982; 11:499–502. [PubMed: 7103426]
- Raichle ME, Gado MH, Grubb RL Jr, et al. In vivo correlation between regional cerebral blood flow and oxygen utilization in man. *Trans Am Neurol Assoc.* 1975; 100:68–71. [PubMed: 1226643]
- Rehncrona S, Rosen I, Siesjo BK. Brain lactic acidosis and ischemic cell damage: 1. Biochemistry and neuro-physiology. *J Cereb Blood Flow Metab.* 1981; 1:297–311. [PubMed: 7328145]
- Reinhard M, Muller T, Roth M, et al. Bilateral severe carotid artery stenosis or occlusion—cerebral autoregulation dynamics and collateral flow patterns. *Acta Neurochir (Wien).* 2003; 145:1053–1059. [PubMed: 14663562]
- Reivich M. Blood flow metabolism couple in brain. *Res Publ Assoc Res Nerv Ment Dis.* 1974; 53:125–140. [PubMed: 4216058]
- Ringelstein EB, Sievers C, Ecker S, et al. Noninvasive assessment of CO₂-induced cerebral vasomotor response in normal individuals and patients with internal carotid artery occlusions. *Stroke.* 1988; 19:963–969. [PubMed: 3135641]
- Ringelstein EB, Biniek R, Weiller C, et al. Type and extent of hemispheric brain infarctions and clinical outcome in early and delayed middle cerebral artery recanalization. *Neurology.* 1992; 42:289–298. [PubMed: 1736156]
- Ringelstein EB, Weiller C, Weckesser M, et al. Cerebral vasomotor reactivity is significantly reduced in low-flow as compared to thromboembolic infarctions: the key role of the circle of Willis. *J Neurol Sci.* 1994; 121:103–109. [PubMed: 8133304]
- Rosenblum WI. Selective impairment of response to acetylcholine after ischemia/reperfusion in mice. *Stroke.* 1997; 28:448–451. discussion 451–452. [PubMed: 9040704]
- Rosenblum WI, Wormley B. Selective depression of endothelium-dependent dilations during cerebral ischemia. *Stroke.* 1995; 26:1877–1881. discussion 1882. [PubMed: 7570742]
- Rossini PM, Altamura C, Ferretti A, et al. Does cerebrovascular disease affect the coupling between neuronal activity and local haemodynamics? *Brain.* 2004; 127:99–110. [PubMed: 14570819]
- Rother J, Knab R, Hamzei F, et al. Negative dip in BOLD fMRI is caused by blood flow–oxygen consumption uncoupling in humans. *Neuroimage.* 2002; 15:98–102. [PubMed: 11771977]
- Sakatani K, Xie Y, Lichty W, et al. Language-activated cerebral blood oxygenation and hemodynamic changes of the left prefrontal cortex in poststroke aphasic patients: a near-infrared spectroscopy study. *Stroke.* 1998; 29:1299–1304. [PubMed: 9660376]
- Sakatani K, Murata Y, Fukaya C, et al. BOLD functional MRI may overlook activation areas in the damaged brain. *Acta Neurochir Suppl.* 2003; 87:59–62. [PubMed: 14518525]
- Sandor P, Komjati K, Reivich M, et al. Major role of nitric oxide in the mediation of regional CO₂ responsiveness of the cerebral and spinal cord vessels of the cat. *J Cereb Blood Flow Metab.* 1994; 14:49–58. [PubMed: 7505282]
- Schmidt-Kastner R, Ophoff BG, Hossmann KA. Delayed recovery of CO₂ reactivity after one hour's complete ischaemia of cat brain. *J Neurol.* 1986; 233:367–369. [PubMed: 3100726]
- Schmidt-Kastner R, Hossmann KA, Ophoff BG. Pial artery pressure after one hour of global ischemia. *J Cereb Blood Flow Metab.* 1987; 7:109–117. [PubMed: 3100544]
- Schreiber WG, Guckel F, Stritzke P, et al. Cerebral blood flow and cerebrovascular reserve capacity: estimation by dynamic magnetic resonance imaging. *J Cereb Blood Flow Metab.* 1998; 18:1143–1156. [PubMed: 9778191]
- Seki H, Yoshimoto T, Ogawa A, et al. The CO₂ response in focal cerebral ischemia—sequential changes following recirculation. *Stroke.* 1984; 15:699–704. [PubMed: 6431652]
- Shen Q, Ren H, Cheng H, et al. Functional, perfusion and diffusion MRI of acute focal ischemic brain injury. *J Cereb Blood Flow Metab.* 2005; 25:1265–1279. [PubMed: 15858531]
- Shima T, Hossmann KA, Date H. Pial arterial pressure in cats following middle cerebral artery occlusion. 1. Relationship to blood flow, regulation of blood flow and electrophysiological function. *Stroke.* 1983; 14:713–719. [PubMed: 6658955]

- Shiokawa O, Sadoshima S, Kusuda K, et al. Cerebral and cerebellar blood flow autoregulation in acutely induced cerebral ischemia in spontaneously hypertensive rats—transtentorial remote effect. *Stroke*. 1986; 17:1309–1313. [PubMed: 3810734]
- Silver IA, Erecinska M. Ion homeostasis in rat brain in vivo: intra- and extracellular $[Ca^{2+}]$ and $[H^+]$ in the hippocampus during recovery from short-term, transient ischemia. *J Cereb Blood Flow Metab*. 1992; 12:759–772. [PubMed: 1324251]
- Skinhoj E, Hoedt-Rasmussen K, Paulson OB, et al. Regional cerebral blood flow and its autoregulation in patients with transient focal cerebral ischemic attacks. *Neurology*. 1970; 20:485–493. [PubMed: 5462241]
- Sokoloff L, Reivich M, Kennedy C, et al. The $[^{14}C]$ deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *J Neurochem*. 1977; 28:897–916. [PubMed: 864466]
- Soriano FG, Virag L, Szabo C. Diabetic endothelial dysfunction: role of reactive oxygen and nitrogen species production and poly(ADP-ribose) polymerase activation. *J Mol Med*. 2001; 79:437–448. [PubMed: 11511974]
- Stamler JS. Alzheimer's disease. A radical vascular connection. *Nature*. 1996; 380:108–111. [PubMed: 8600380]
- Sundt TM Jr, Waltz AG. Cerebral ischemia and reactive hyperemia. Studies of cortical blood flow and micro-circulation before, during, and after temporary occlusion of middle cerebral artery of squirrel monkeys. *Circ Res*. 1971; 28:426–433. [PubMed: 4994637]
- Symon L, Khodadad G, Montoya G. Effect of carbon dioxide inhalation on the pattern of gaseous metabolism in ischaemic zones of the primate cortex. An experimental study of the 'intracerebral steal' phenomenon in baboons. *J Neurol Neurosurg Psychiatry*. 1971; 34:481–486. [PubMed: 4999541]
- Symon L, Pasztor E, Branston NM. The distribution and density of reduced cerebral blood flow following acute middle cerebral artery occlusion: an experimental study by the technique of hydrogen clearance in baboons. *Stroke*. 1974; 5:355–364. [PubMed: 4209598]
- Symon L, Crockard HA, Dorsch NW, et al. Local cerebral blood flow and vascular reactivity in a chronic stable stroke in baboons. *Stroke*. 1975; 6:482–492. [PubMed: 1179454]
- Takano T, Nagatsuka K, Ohnishi Y, et al. Vascular response to carbon dioxide in areas with and without diaschisis in patients with small, deep hemispheric infarction. *Stroke*. 1988; 19:840–845. [PubMed: 3133840]
- Takano T, Tian GF, Peng W, et al. Astrocyte-mediated control of cerebral blood flow. *Nat Neurosci*. 2006; 9:260–267. [PubMed: 16388306]
- Tatemichi TK, Young WL, Prohovnik I, et al. Perfusion insufficiency in limb-shaking transient ischemic attacks. *Stroke*. 1990; 21:341–347. [PubMed: 2406995]
- Thompson BG, Pluta RM, Girton ME, et al. Nitric oxide mediation of chemoregulation but not autoregulation of cerebral blood flow in primates. *J Neurosurg*. 1996; 84:71–78. [PubMed: 8613839]
- Todd NV, Picozzi P, Crockard HA, et al. Reperfusion after cerebral ischemia: influence of duration of ischemia. *Stroke*. 1986; 17:460–466. [PubMed: 3715943]
- Traupe H, Kruse E, Heiss WD. Reperfusion of focal ischemia of varying duration: postischemic hyper- and hypo-perfusion. *Stroke*. 1982; 13:615–622. [PubMed: 7123593]
- Traystman RJ, Kirsch JR, Koehler RC. Oxygen radical mechanisms of brain injury following ischemia and reperfusion. *J Appl Physiol*. 1991; 71:1185–1195. [PubMed: 1757340]
- Tsubokawa T, Katayama Y, Kondo T, et al. Changes in local cerebral blood flow and neuronal activity during sensory stimulation in normal and sympathectomized cats. *Brain Res*. 1980; 190:51–64. [PubMed: 7378745]
- Ueki M, Linn F, Hossmann KA. Functional activation of cerebral blood flow and metabolism before and after global ischemia of rat brain. *J Cereb Blood Flow Metab*. 1988; 8:486–494. [PubMed: 3392113]
- Ugurbil K, Toth L, Kim DS. How accurate is magnetic resonance imaging of brain function? *Trends Neurosci*. 2003; 26:108–114. [PubMed: 12536134]

- Van den Kerckhoff W, Hossmann KA, Hossmann V. No effect of prostacyclin on blood flow, regulation of blood flow and blood coagulation following global cerebral ischemia. *Stroke*. 1983; 14:724–730. [PubMed: 6362092]
- Vignais PV. The superoxide-generating NADPH oxidase: structural aspects and activation mechanism. *Cell Mol Life Sci*. 2002; 59:1428–1459. [PubMed: 12440767]
- Von Monakow, C. Die Lokalisation im Grosshirn und der Abbau der Funktion durch kortikale Herde. JF Berg-mann; Wiesbaden: 1914.
- Vorstrup S, Brun B, Lassen NA. Evaluation of the cerebral vasodilatory capacity by the acetazolamide test before EC-IC bypass surgery in patients with occlusion of the internal carotid artery. *Stroke*. 1986; 17:1291–1298. [PubMed: 3492787]
- Walder CE, Green SP, Darbonne WC, et al. Ischemic stroke injury is reduced in mice lacking a functional NADPH oxidase. *Stroke*. 1997; 28:2252–2258. [PubMed: 9368573]
- Waltz AG. Effect of PaCO₂ on blood flow and micro-vasculature of ischemic and nonischemic cerebral cortex. *Stroke*. 1970; 1:27–37. [PubMed: 5522898]
- Wang X, Lo EH. Triggers and mediators of hemorrhagic transformation in cerebral ischemia. *Mol Neurobiol*. 2003; 28:229–244. [PubMed: 14709787]
- Wei EP, Kontos HA, Beckman JS. Mechanisms of cerebral vasodilation by superoxide, hydrogen peroxide, and peroxynitrite. *Am J Physiol*. 1996; 271:H1262–H1266. [PubMed: 8853367]
- Wellman GC, Nathan DJ, Saundry CM, et al. Ca²⁺ sparks and their function in human cerebral arteries. *Stroke*. 2002; 33:802–808. [PubMed: 11872907]
- Yamamoto M, Meyer JS, Sakai F, et al. Aging and cerebral vasodilator responses to hypercarbia: responses in normal aging and in persons with risk factors for stroke. *Arch Neurol*. 1980; 37:489–496. [PubMed: 6774700]
- Yamauchi H, Kudoh T, Sugimoto K, et al. Altered patterns of blood flow response during visual stimulation in carotid artery occlusive disease. *Neuroimage*. 2005; 25:554–560. [PubMed: 15784434]
- Zhang F, Slungaard A, Vercellotti GM, et al. Super-oxide-dependent cerebrovascular effects of homocysteine. *Am J Physiol*. 1998; 274:R1704–R1711. [PubMed: 9608025]
- Zhang J, Piantadosi CA. Prolonged production of hydroxyl radical in rat hippocampus after brain ischemiareperfusion is decreased by 21-aminosteroids. *Neurosci Lett*. 1994; 177:127–130. [PubMed: 7824163]
- Zonta M, Angulo MC, Gobbo S, et al. Neuron-toastrocyte signaling is central to the dynamic control of brain microcirculation. *Nat Neurosci*. 2003; 6:43–50. [PubMed: 12469126]
- Zou MH, Leist M, Ullrich V. Selective nitration of prostacyclin synthase and defective vasorelaxation in atherosclerotic bovine coronary arteries. *Am J Pathol*. 1999; 154:1359–1365. [PubMed: 10329589]




Fig. 14.1.

Recording of cerebral blood flow (CBF) measured by laser Doppler flowmetry in the center of the ischemic territory (ischemic core) before, during, and after occlusion of the middle cerebral artery using an intravascular filament in mice. CBF is markedly reduced during the arterial occlusion. At reperfusion, CBF first increases (post-ischemic hyperemia) and then drops below pre-ischemic levels (post-ischemic hypoperfusion).




Fig. 14.2.

Relationship between cerebral blood flow (CBF) and mean arterial pressure (MAP) in physiological conditions (gray) and following cerebral ischemia (black). In the normal state, CBF is kept constant over a relatively wide range of MAPs, a phenomenon termed cerebrovascular auto-regulation. Thus, cerebral resistance vessels dilate when MAP decreases and constrict when MAP increases and, as a result, CBF remains relatively constant. When the change in MAP exceeds this compensatory ability of cerebral vessels, CBF increases when MAP rises (upper limit) and decreases when MAP is reduced (lower limit). Following cerebral ischemia, cerebral resistance arteries are in a state of vasoparalysis and lose their ability to constrict or dilate in response to changes in MAP. Thus, autoregulation is lost and CBF follows MAP passively. Such disruption of autoregulation can amplify the damage by aggravating cerebral ischemia if MAP falls and by inducing blood-barrier damage and cerebral edema when MAP rises.

Table 14.1

Cerebral ischemia and autoregulation—selected animal studies

Ischemia model (duration)	Species	CBF measurement	Effect on autoregulation (upper–lower limit)	Reference
Global ischemia (30 and 60 min)	Cat	¹³³ Xe-injection	Preserved (LL)	Hossmann et al., 1973
3 years after pMCAO	Baboon	H ₂ clearance	Abolished in infarct, impaired in peri-infarct area (LL)	Symon et al., 1975
Global ischemia (15 min)	Dog	¹³³ Xe-injection	Partially preserved (UL, LL)	Nemoto et al., 1975
pMCAO	Cat	Heated thermocouple	Impaired (UL)	Shima et al., 1983
Global ischemia (60 min)	Rat	H ₂ clearance	Abolished (LL)	Shiokawa et al., 1986
pMCAO	Rat (SHR)	LDF	Impaired if ischemic CBF >30%, abolished if ischemic CBF <30% (UL, LL)	Dirnagl and Pulsinelli, 1990
Global ischemia (12 and 18 min)	Dog	Sagittal sinus venous outflow	Impaired (UL, LL)	Christopherson et al., 1993
pMCAO	Rat	LDF	Impaired, dependent on proximity to core (UL, LL)	Lauer et al., 1998
tMCAO (endothelin-1 administration)	Rat	IAP, SPECT	Abolished in viable tissue with normal CBF (LL)	MacGregor et al., 2000

CBF, cerebral blood flow; IAP, C¹⁴-iodoantipyrine; LDF, laser-Doppler flowmetry; LL, lower limit; pMCAO, permanent middle cerebral artery occlusion; SHR, spontaneously hypertensive rats; SPECT, single photon emission computed tomography; tMCAO, transient middle cerebral artery occlusion; UL, upper limit.

Table 14.2

Cerebral ischemia and autoregulation—selected human studies

Pathology	CBF measurement	Effect on autoregulation (upper–lower limit)	Reference
TIA (ICA territory)	¹³³ Xe-injection	Impaired/abolished in 2 of 12 patients (UL, LL)	Skinhoj et al., 1970
Stroke (ICA territory)	¹³³ Xe-injection	Abolished, restored by hypocapnia (UL)	Paulson et al., 1972
Stroke within 72 h (MCA territory)	¹³³ Xe-injection	Impaired in 3 of 16 patients (UL)	Olsen et al., 1981
Subacute stroke (MCA territory)	¹³³ Xe-injection	Impaired in collaterally perfused areas (UL)	Olsen et al., 1983
TIA (bilateral ICA stenosis/occlusion)	¹³³ Xe-inhalation, TCD	Impaired (LL)	Tatemichi et al., 1990
Stroke (ICA, MCA territory)	PET	Impaired (LL)	Ouchi et al., 2001
Stroke within 72 h (ICA, VB territories)	TCD*	Impaired (spontaneous ABP fluctuations)	Eames et al., 2002
ICA stenosis/occlusion	TCD*	Impaired (spontaneous ABP fluctuations)	Gooskens et al., 2003
ICA stenosis/occlusion	TCD*	Impaired (spontaneous ABP fluctuations)	Reinhard et al., 2003
MCA stroke, lacunar stroke (within 72 h)	TCD*	Impaired (spontaneous ABP fluctuations)	Immink et al., 2005
MCA stenosis/occlusion	TCD*	Impaired, improvement after MCA stent angioplasty (thigh cuff)	Gong et al., 2006
ICA stenosis (before/after angioplasty)	TCD*	Impaired (spontaneous ABP fluctuations), improvement after angioplasty	Haubrich et al., 2007

ABP, arterial blood pressure; CBF, cerebral blood flow; ICA, internal carotid artery; LL, lower limit; MCA, middle cerebral artery; PET, positron emission tomography; TCD, transcranial Doppler; TIA, transient ischemic attack; UL, upper limit; VB, vertebrobasilar artery.

* 'Dynamic autoregulation' was estimated based on spontaneous ABP fluctuations except for Gong et al. (2006), who used the thigh cuff technique.

Table 14.3

Cerebral ischemia and cerebral blood flow reactivity to carbon dioxide—selected animal studies

Ischemia model	Species	CBF measurement	Effect on CO ₂ reactivity	Reference
Global ischemia (30 and 60 min)	Cat	¹³³ Xe-injection	Abolished	Hossmann et al., 1973
Global ischemia (15 min)	Dog	¹³³ Xe-injection	Abolished	Nemoto et al., 1975
pMCAO	Baboon	H ₂ clearance	Impaired, steal phenomenon	Symon et al., 1974
3 years after pMCAO	Baboon	H ₂ clearance	Impaired	Symon et al., 1975
pMCAO	Baboon	¹³³ Xe-injection	Paradoxical CBF increase in ischemic area during hypocapnia	Ott et al., 1975
pMCAO	Cat	Heated thermocouple	Abolished or steal phenomenon	Shima et al., 1983
Global ischemia (60 min)	Cat	¹³³ Xe-injection	Abolished during postischemic hypoperfusion	Van den Kerckhoff et al., 1983
Global ischemia (12 min)	Dog	Microspheres	Abolished at 3 h, restored at 24 h after ischemia	Koch et al., 1984
tMCAO (30 min, 1, 2, 6 h)	Dog	H ₂ clearance	Impaired, dependent on duration and severity of stroke	Seki et al., 1984
Global ischemia (60 min)	Cat	¹³³ Xe-injection	Prolonged impairment	Schmidt-Kastner et al., 1986
Global ischemia (60 min)	Cat	¹³³ Xe-injection	Abolished during postischemic hypoperfusion	Schmidt-Kastner et al., 1987
pMCAO	Rat	IAP	Abolished in core	Jones et al., 1989
Global ischemia (20 min)	Piglet	Pial diameter	Abolished	Leffler et al., 1989
Global ischemia (12 and 18 min)	Dog	Sagittal venous outflow	Impaired	Christopherson et al., 1993
pMCAO	Baboon	Microspheres	Core: abolished or steal phenomenon Penumbra: impaired	Dettmers et al., 1993
p or tMCAO (30 and 90 min)	Rat	MRI (T2*, DWI)	Abolished, recovery of CO ₂ reactivity in viable tissue	Ono et al., 1997
tMCAO (60 min)	Rat	Perfusion MRI	Prolonged impairment	Olah et al., 2000
pMCAO	Rat	BOLD imaging	Core: abolished/reduced Borderzone: normal	Harris et al., 2001

BOLD, blood-oxygen-level-dependent; CBF, cerebral blood flow; DWI, diffusion-weighted imaging; IAP, C¹⁴-iodoantipyrine; MRI, magnetic resonance imaging; pMCAO, permanent middle cerebral artery occlusion; tMCAO, transient middle cerebral artery occlusion.

Table 14.4

Cerebral ischemia and cerebral blood flow reactivity to carbon dioxide—selected human studies

Pathology	CBF measurement	Effect on CO ₂ reactivity	Reference
Stroke (hemispheric, VB)	¹³³ Xe-inhalation	Impaired	Yamamoto et al., 1980
Stroke within 72 h (MCA territory)	¹³³ Xe-injection	Impaired	Olsen et al., 1981
TIA or stroke (MCA territory)	¹³³ Xe-inhalation	Impaired	Norrving et al., 1982
TIA, stroke in patients with ICA stenosis/occlusion	¹³³ Xe-inhalation	Impaired	Bullock et al., 1985
Uni-/bilateral ICA stenosis occlusion (TIA, stroke, no symptoms)	¹³³ Xe-injection	Impaired	Brown et al., 1986
Hemispherical stroke/TIA (ICA occlusion), follow-up after EC/IC bypass	¹³³ Xe-inhalation	Acetazolamide reactivity impaired in 9 of 18 patients, two patients with steal, improvement in two patients after EC/IC bypass	Vorstrup et al., 1986
Stroke/TIA, patients with unilateral ICA occlusion	¹³³ Xe-inhalation	Impaired	Clifton et al., 1988
ICA stenosis occlusion (TIA, stroke, no symptoms)	PET, ¹³³ Xe-injection	Impaired, correlates with CBF/CBV ratio	Herold et al., 1988
ICA/MCA stenosis occlusion, moyamoya disease	PET	Impaired, correlates with OEF	Kanno et al., 1988
TIA, stroke, ICA stenosis occlusion	¹³³ Xe-inhalation	Impaired	Keyeux et al., 1988
TIA, minor stroke (ICA territory)	PET	Impaired	Levine et al., 1988
Uni-/bilateral ICA occlusion (stroke, TIA, no symptoms)	TCD	Impaired	Ringelstein et al., 1988
Chronic capsular stroke	¹³³ Xe-injection	Vasoconstriction to hypocapnia attenuated in regions with diaschisis	Takano et al., 1988
TIA (ICA territory)	¹³³ Xe-inhalation	Acetazolamide reactivity impaired in 14 of 15 patients	Chollet et al., 1989
TIA, severe ICA disease	¹³³ Xe-inhalation, TCD	Impaired	Tatemichi et al., 1990
TIA (ICA territory), ICA stenosis occlusion	PET	Impaired	Levine et al., 1991
TIA, stroke, ICA stenosis occlusion	TCD	Impaired	Miller et al., 1992
ICA stenosis occlusion (TIA, stroke, no symptoms)	TCD	CO ₂ and acetazolamide reactivity impaired	Ringelstein et al., 1992
Stroke (ICA territory)	TCD	Impaired	Maeda et al., 1993
Uni-/bilateral ICA occlusion (stroke, TIA, no symptoms)	TCD	Impaired	Ringelstein et al., 1994
TIA, stroke (ICA/MCA territory, moyamoya disease)	Stable Xe-CT, PET	Acetazolamide reactivity impaired	Nariai et al., 1995
TIA, stroke (MCA territory)	¹³³ Xe-inhalation	Preserved reactivity; CO ₂ : 23/24pts, acetazolamide: 13/24pts	Kazumata et al., 1996
Stroke, TIA (ICA/MCA stenosis occlusion)	Dynamic MRI	Acetazolamide reactivity impaired in 5/8 patients	Schreiber et al., 1998
CADASIL	TCD	Impaired	Pfefferkorn et al., 2001
TIA (ICA territory)	TCD	Impaired	Bakker et al., 2003
Lacunar stroke (MCA territory, within 1 week)	TCD	Impaired	De Leeuw et al., 2003
ICA stenosis/occlusion	TCD	CO ₂ reactivity correlates with degree of stenosis	Gooskens et al., 2003
Minor stroke (MCA territory)	TCD	Impaired	Novak et al., 2003

Pathology	CBF measurement	Effect on CO₂ reactivity	Reference
ICA stenosis/occlusion	TCD	Impaired	Reinhard et al., 2003
ICA occlusion	PET	Correlation between reduced acetazolamide reactivity and OEF increase to acetazolamide	Nemoto et al., 2007

CADASIL, cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy; CBF, cerebral blood flow; CBV, cerebral blood volume; CT, computed tomography; EC/IC, extracranial/intracranial; ICA, internal carotid artery; MCA, middle cerebral artery; MRI, magnetic resonance imaging; OEF, oxygen extraction fraction; PET, positron emission tomography; TCD, transcranial Doppler; TIA, transient ischemic attack; Xe, Xenon; VB, vertebrobasilar.

Table 14.5

Cerebral ischemia and neurovascular coupling—selected animal studies

Ischemia model	Species	CBF measurement	Effect on CO ₂ reactivity	Reference
Global ischemia (10 min)	Rat	Mitochondrial redox state	Transient mitochondrial dysfunction	Duckrow et al., 1981
Global ischemia (30 min)	Rat	2-DG	Transient reduction of CGU	Dietrich et al., 1986
Photothrombotic frontal lesion	Rat	IAP, 2-DG	5 days post-lesion: activation of rCBF in barrel cortex suppressed	Ginsberg et al., 1989
Global ischemia (30 min)	Rat	IAP, 2-DG	Impaired neurovascular coupling	Ueki et al., 1988
Focal ischemia (30 min)	Mouse	LDF	Impaired neurovascular coupling	Kunz et al., 2007b

2-DG, 2-deoxyglucose; CGU, cerebral glucose utilization; IAP, C¹⁴-iodoantipyrine; LDF, laser-Doppler flowmetry; rCBF, regional cerebral blood flow.

Table 14.6

Cerebral ischemia and functional activation—selected human studies

Pathology	CBF measurement	Effect on functional activation	Reference
TIA, stroke (ICA territory)	PET	Motor activation: ipsilateral CBF increase Acetazolamide: no CBF increase	Inao et al., 1998
Stroke (MCA territory)	NIRS	Increased deoxyhemoglobin (five of ten patients)	Sakatani et al., 1998
ICA stenosis/occlusion	BOLD-fMRI	Reduced BOLD response in auditory cortex if ischemic borderzone lesions present	Bilecen et al., 2002
ICA/MCA stenosis/occlusion	BOLD-fMRI	Reduced BOLD response ipsilateral	Carusone et al., 2002
TIA, stroke (within 3–12 months)	BOLD-fMRI, NIRS	Reduced BOLD response ipsilaterally, increased deoxyhemoglobin	Murata et al., 2002
Lacunar stroke	BOLD-fMRI	Reduced BOLD response (ipsi-/contralateral)	Pineiro et al., 2002
TIA (ICA territory), ICA occlusion	BOLD-fMRI	Negative BOLD response in a patient with impaired CO ₂ reactivity	Rother et al., 2002
TIA (stenosis extra-/intracranial)	BOLD-fMRI	Reduced BOLD response in patients with impaired CO ₂ reactivity	Hamzei et al., 2003
Stroke, TIA (ICA territory)	BOLD-fMRI, NIRS	BOLD: limited activation, NIRS: sustained increase in deoxyhemoglobin	Sakatani et al., 2003
TIA, stroke (ICA territory)	MEG, evoked field BOLD-fMRI	VMR preserved, BOLD absent	Rossini et al., 2004
Stroke (frontal lobe)	BOLD-fMRI	Reduced BOLD response on lesion side	Krainik et al., 2005
ICA stenosis/occlusion (stroke, TIA, no symptoms)	PET	Increased CBF in primary visual cortex, decreased CBF in surrounding areas	Yamauchi et al., 2005
TIA, stroke	BOLD-fMRI, NIRS	Reduced BOLD response and increased deoxyhemoglobin after severe ischemia	Murata et al., 2006

BOLD, blood-oxygen-level-dependent; CBF, cerebral blood flow; fMRI, functional magnetic resonance imaging; ICA, internal carotid artery; MCA, middle cerebral artery; MEG, magnetoencephalography; NIRS, near-infrared spectroscopy; PET, positron emission tomography; TIA, transitory ischemic attack; VMR, vasomotor reserve.