

Increasing the Number of Adverse Drug Reactions Reporting: the Role of Clinical Pharmacy Residents

Shadi Baniasadi^a, Maryam Habibi^b, Roodabeh Haghgoo^b, Masoumeh Karimi Gamishan^c, Fatemeh Dabaghzadeh^d, Maryam Farasatinasab^e, Shadi Farsaeif, Afshin Gharekhani^g, Hamidreza Kafi^e, Iman Karimzadeh^g, Ali Kharazmkia^h, Farhad Najmeddin^g, Naemeh Nikvarz^d, Mohammad Bagher Oghazian^g, Haleh Rezaeeⁱ, Kourosh Sadeghi^g, Ali Tafazzoli^e, Nahid Shahsavari^e and Fanak Fahimi^{h,e*}

^aVirology Research Center, NRITLD, Masih Daneshvari Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^bChronic Respiratory Disease Research Center, NRITLD, Masih Daneshvari Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^cImporting and Exporting Office, Food and Drug Organization, Ministry of Health and Medical Education, Tehran, Iran. ^dFaculty of Pharmacy and Pharmaceutical Sciences, Kerman Medical University, Kerman, Iran. ^e Clinical Pharmacy Department, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran. ^fDepartment of Clinical Pharmacy and Pharmacy Practice, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, Iran. ^g Clinical Pharmacy Department, School of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran. ^hLorestan University of Medical Sciences, Khorramabad, Iran. ⁱInfectious Diseases and Tropical Medicine Research Center, Department of Clinical Pharmacy, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Abstract

Detection of adverse drug reactions (ADRs) in hospitals provides an important measure of the burden of drug related morbidity on the healthcare system. Spontaneous reporting of ADRs is scarce and several obstacles to such reporting have been identified formerly. This study aimed to determine the role of clinical pharmacy residents in ADR reporting within a hospital setting. Clinical pharmacy residents were trained to report all suspected ADRs through ADR-reporting yellow cards. The incidence, pattern, seriousness, and preventability of the reported ADRs were analyzed. During the period of 12 months, for 8559 patients, 202 ADR reports were received. The most frequently reported reactions were due to anti-infective agents (38.38%). Rifampin accounted for the highest number of the reported ADRs among anti-infective agents. The gastro-intestinal system was the most frequently affected system (21.56%) of all reactions. Fifty four of the ADRs were reported as serious reactions. Eighteen of the ADRs were classified as preventable. Clinical pharmacy residents involvement in the ADR reporting program could improve the ADR reporting system.

Keywords: Adverse drug reaction; Clinical pharmacist; Pharmacy resident; Hospital, Pharmacovigilance; Spontaneous reporting.

* Corresponding author:

E-mail: fanakfahimi@yahoo.com

Introduction

Adverse drug reactions (ADRs) are common causes of morbidity and mortality within the hospital setting. The hospital setting, with its clearly defined patient population, is an ideal setting to identify potential adverse drug reaction signals and to report them to either the drug company or the FDA (1). Reporting of ADRs is an important component of monitoring and evaluation activities performed in the hospitals. A productive hospital-based reporting program can be helpful in providing valuable information regarding potential problems of drug usage in an institution. Through these efforts, problems are identified and resolved, which results in continuous improvement in patient care (2).

Spontaneous reporting program, a common method of drug surveillance is capable of recognizing ADRs in the daily medical practice, even though underreporting and absence of information on ADRs are its disadvantages (3, 4). Although identification and documentation of ADRs are considered as all health care professionals' responsibility, pharmacists and especially clinical pharmacists are often in charge for these tasks (5).

Clinical pharmacy is relatively a new concept in Iran (6). This new breed of pharmacy practices is more patient oriented rather than drug product oriented (7). Clinical pharmacy residents are in an ideal position to increase ADR reporting rates within health systems (8). ADR review can be added to the daily responsibilities of residents participating in the hospitals and clinical rotations. In this article we tried to demonstrate the contribution of clinical pharmacy residents in the process of ADR reporting within a hospital setting and assess the role of the clinical pharmacy residents on the number and reporting rate of ADRs.

Experimental

The study was conducted in Masih Daneshvari Hospital over a one year period from March 2010 to February 2011. During the orientation session at the beginning of each new rotation, residents received a brief overview of the hospital's ADR reporting program. Residents began

prospectively and retrospectively collecting data on suspected ADRs. Prospective ADRs were identified by physicians, nurses, or residents during interdisciplinary rounds. Retrospective ADRs were identified by reviewing patient's records. The residents documented the suspected drug(s) for the event on yellow cards. Pharmaceutical care department staffs reviewed each ADR report and were responsible for the accuracy of the reported ADR.

An ADR was defined according to the World Health Organization (WHO) definition as: "a response to a drug which is noxious and unintended, and which occurs at doses normally used in man for the prophylaxis, diagnosis, or therapy of disease, or for the modification of physiological function."

Each ADR was classified according to the WHO system organ classification (9). All reported ADRs were evaluated for the causality in accordance with Naranjo's algorithm (10). The seriousness of reported adverse reactions was assessed based on the WHO definition, which included any adverse event that resulted in death, life-threatening situation, hospitalization, prolonged hospital stay, disability or birth defect. Assessment of preventability was determined using the scale developed by Schumock *et al.* (11).

All documented ADRs were electronically entered into an Excel spreadsheet for the purpose of generating annually reports. ADR forms were then sent to the national ADR monitoring centre at the Ministry of Health for further evaluation.

The number and reporting rate (number of reports/number of beds) of ADRs were compared with our previous study (12).

Results

During 12 months, a total of 8559 patients were admitted to the hospital. Sixteen clinical pharmacy residents were rotating who reported 202 ADRs from 178 patients. A total of 202 ADRs were documented in this study, compared with 112 in previous study (12).

In terms of the patient demographics in the reported ADRs, 111 were women and 91 were men.

Number of patients in pediatric (0–18 years), adult (19–60 years) and geriatric (> 60 years)

Table 1. Wards associated with the reported ADRs.

Ward	Number of ADRs	Percentage
Internal medicine	64	31.68
Cardiac Care Unit	34	16.83
Intensive Care Unit	29	14.36
Tuberculosis	28	13.86
Oncology	26	12.87
Surgery	9	4.46
Pediatric	7	3.47
Transplant	5	2.48

groups were 11, 112 and 79, respectively. ADRs were reported from 8 wards. The majority of the ADR reports were from the Department of Internal Medicine (31.68%), followed by the Coronary Care Unit (16.83%) as presented in Table 1. The most frequent reports were due to anti-infective agents (38.38%), followed by antineoplastic agents (11.45%) and central nervous system agents (10.44%). The drug class involved in the ADRs is shown in Table 2.

Rifampin was the most frequent anti-infective agent associated with the suspected ADRs such as nausea, vomiting, abdominal pain, urticaria, hepatic enzyme increased, pancytopenia, and Stevens-Johnson syndrome (SJS) (Table 3).

The gastrointestinal system was the most frequently affected system (21.56% of all reactions), followed by the skin and appendages

system (14.50%). The classification of the ADRs by system-organ class is demonstrated in Table 4. The causality assessment of ADRs revealed that 35.64% as probable and 64.36% as possible reactions. Fifty four (26.73%) of the ADRs were classified as serious according to the WHO definition and resulted in prolonged hospital stay, persistent disability or death. Table 5 shows the drugs with a serious reaction.

Eighteen (8.91%) of the ADRs were classified as preventable according to the scale developed by Schumock *et al.*(11).

Discussion

The contribution of clinical pharmacy residents in improving ADR reporting was determined by comparing current with previous ADR data. A total of 202 ADRs were documented in this study, compared with 112 in previous study (12) an increase of almost twofold. Reporting rate of ADRs in our center was increased from 40% in year 2006 to 80% in year 2010.

Sullivan *et al.* showed that pharmacy student participation in the ADR reporting significantly increases the number of ADRs documented (5).

Until recent years, colleges of pharmacy in Iran had no clinical education (13). Significant progress has been made since 1996 toward changing the emphasis of pharmaceutical

Table 2. Drug class implicated in the reported ADRs.

Drug class	Number of ADRs	Percentage
Anti-infective agents	114	38.38
Antineoplastic agents	34	11.45
Central nervous system agents	31	10.44
Cardiovascular drugs	24	8.08
Miscellaneous Therapeutic Agents	19	6.40
Blood formation and coagulation	17	5.72
Autonomic drugs	16	5.39
Hormones and synthetic substitutes	15	5.05
Electrolytic, Caloric, and Water Balance	11	3.70
Gastrointestinal drugs	9	3.03
Antitussives, Expectorants, and Mucolytic Agents	4	1.35
Antihistamine drugs	1	0.34
Vitamins	1	0.34
Eye, Ear, Nose, and Throat preparations	1	0.34

Table 3. Anti-infective agents implicated in ADRs.

Anti-infective agents	Number of ADRs	Percentage
Rifampin	23	20.35
Isoniazid	21	18.58
Pyrazinamide	18	15.93
Vancomycin	14	12.39
Clindamycin	5	4.42
Ceftriaxone	4	3.54
Co-trimoxazole	4	3.54
Clarithromycin	4	3.54
Ofloxacin	4	3.54
Ethambutol	3	2.65
Azithromycin	2	1.77
Dapsone	1	0.88
Erythromycin	1	0.88
Oseltamivir	1	0.88
Ganciclovir	1	0.88
Chloramphenicol	1	0.88
Linezolid	1	0.88
Nitrofurantoin	1	0.88
Penicillin	1	0.88
Cefalexin	1	0.88
Meropenem	1	0.88
Piperacillin	1	0.88

Table 4. Organ systems associated with the reported ADRs.

System associated with ADRs	Number of ADRs	Percentage
Gastro-intestinal system	58	21.56
Skin and appendages	39	14.50
Central & peripheral nervous system	31	11.52
Metabolic and nutritional	22	8.18
Liver and biliary system	20	7.43
Platelet, bleeding & clotting	18	6.69
Respiratory system	13	4.83
Vascular (extracardiac)	12	4.46
Body as a whole-general	10	3.72
Psychiatric	10	3.72
Vision	6	2.23
Red blood cell	5	1.86
Urinary system	4	1.49
White cell	4	1.49
Musculo- skeletal system	4	1.49
Cardiovascular	3	1.12
Resistance mechanism	3	1.12
Heart rate and rhythm	2	0.74
Application site	2	0.74
Special senses other	2	0.74
Hearing and vestibular	1	0.37

education from preparing a product-oriented pharmacist to preparing a patient-oriented individual with good basic knowledge of drugs. These pharmacists can work in close association with physicians and others on the health care teams (14). It is the pharmacist's responsibility and professional commitment to report any suspected ADRs (15).

In our study, the most frequent ADRs were related to antibiotics that are costly and highly consumed in our hospital. While in other studies, different drug classes such as antineoplastic, cardiovascular, anticoagulant, nonsteroidal anti-inflammatory, hypoglycemic, and anti-infective agents have been most frequently associated with ADRs (16-21).

Our previous study has shown ceftriaxone as the most frequent anti-infective agent associated with the suspected ADRs (12). Perhaps emphasis on the appropriate and rational use of this drug by pharmaceutical care department in the rounds,

consultations, random medication reviews, or via newsletter, (22-25), has changed drug utilization pattern and consequently ADR pattern.

In the present study rifampin has been replaced as the most frequent anti-infective agent associated with the suspected ADRs. It could be rational since our hospital is a tuberculosis referral center and therefore rifampin is highly prescribed.

Gastro-intestinal system was the most reported organ system which was affected by ADRs. Skin and appendages systems have usually been reported as the most affected organ system by ADRs (26-28). This reveals that clinical pharmacy residents have used their clinical judgments which could be superior to documenting only those visible ADRs. Pathological and laboratory data which are objective markers of ADRs should be considered by residents (12).

In this study, 8.91% of ADRs were

Table 5. Drugs reported in serious ADRs by frequency of times reported and type of reaction.

Drugs	Type of reaction	Number of reports
Rifampin	Hepatic enzyme increased	9
	Pancytopenia	1
	Stevens Johnson Syndrome	1
Isoniazid	Hepatic enzyme increased	8
Heparin	Thrombocytopenia	3
	Haematuria, Gastric ulcer hemorrhagic, Ecchymosis	1
Cyclosporine	Nephropathy toxic, Hepatic enzyme increased	2
	Vomiting, Nausea	1
Vancomycin	Renal failure	1
	Thrombocytopenia	1
Carboplatin	Pancytopenia	1
	Leucopenia, Anemia	1
Anti-thymocyte globulin	Thrombocytopenia	2
Insulin Regular	Hypoglycemia	2
Mycophenolate mofetil	Leucopenia, Neutropenia	2
Ranitidine	Thrombocytopenia	1
Allopurinol	Stevens Johnson Syndrome	1
Amiodarone	Thrombocytopenia, Anemia	1
Betaxolol	Dyspnoea, Bronchospasm	1
Carbamazepine	Dyspnoea, Dermatitis	1
Co-trimoxazole	Anemia	1
Digoxin	Thrombocytopenia	1
Cyclophosphamide	Pulmonary edema, Oedema peripheral	1
Enoxaparin	Thrombocytopenia	1
Hyoscine	Dyspnoea	1
Insulin NPH	Hypoglycemia	1
Morphine sulfate	Convulsion	1
Propranolol	Bronchospasm, Dyspnoea, Consciousness decreased	1
Pyrazinamide	BUN increased	1
Sodium valproate	Thrombocytopenia	1
Warfarin	Pulmonary haemorrhage	1
Ceftriaxone	Allergic reaction	1

preventable. Higher rate of preventable ADRs in other studies (the wide range of 30-70%) (28, 29) and 13.39% reduction in preventable ADRs in our hospital (22.30% preventable ADRs in previous study) suggest that preventive strategies have been effective in our institute.

Serious ADRs have been increased from 16.70% (in previous study) to 26.73% (in this

study). Perhaps clinical pharmacy residents are eligible to detect serious ADRs more efficiently. The assessment of whether an ADR has increased the length of stay or caused persistent disability or death can be difficult to assess because individual patient factors such as the nature and severity of the underlying disease, and social factors may also contributed to the length of stay

(30) medication errors (31) and inappropriate prescribing patterns (32).

References

- (1) Pushkin R, Frassetto L, Tsourounis C, Segal ES and Kim S. Improving the reporting of adverse drug reactions in the hospital setting. *Postgrad. Med.* (2010) 122: 154-164.
- (2) Murphy BM and Frigo LC. Development, implementation, and results of a successful multidisciplinary adverse drug reaction reporting program in a university teaching hospital. *Hosp. Pharm.* (1993) 28: 1204-1240.
- (3) Moride Y, Haramburu F, Requeyo AA and Begaud B. Underreporting of adverse drug reactions in general practice. *Br. J. Clin. Pharmacol.* (1997) 43: 177-181.
- (4) Alvarez-Requejo A, Carvajal A, Begaud B, Moride Y, and Vega T and Martin Arias LH. Under reporting of adverse drug reactions. Estimate based on a spontaneous reporting scheme and a sentinel system. *Eur. J. Clin. Pharmacol.* (1998) 54: 483-488.
- (5) Sullivan KM and Spooner LM. Adverse-drug-reaction reporting by pharmacy students in a teaching hospital. *Am. J. Health. Syst. Pharm.* (2008) 65: 1177-1179.
- (6) Salamzadeh J. Clinical pharmacy in Iran: where do we stand? *Iran. J. Pharm. Res.* (2004) 3: 1-2.
- (7) Miller RR. History of clinical pharmacy and clinical pharmacology. *J. Clin. Pharmacol.* (1981) 21: 195-197.
- (8) Bond CA and Raehl CL. Clinical pharmacy services, pharmacy staffing, and adverse drug reactions in United States hospitals. *Pharmacotherapy.* (2006) 26: 735-747.
- (9) Uppsala Monitoring Centre. Adverse Reaction Terminology. World Health Organization Collaborating Centre for International Drug Monitoring. Uppsala, Sweden (2003).
- (10) Naranjo CA, Busto U, Sellers EM, Sandor P, Ruiz I, Roberts EA and Janecek E. A method for estimating the probability of adverse drug reactions. *Clin. Pharmacol. Ther.* (1981) 30: 239-245.
- (11) Schumock GT and Thornton JP. Focusing on the preventability of adverse drug reactions. *Hosp. Pharm.* (1992) 27: 538.
- (12) Baniasadi S, Fahimi F and Shalviri G. Developing an adverse drug reaction reporting system at a teaching hospital. *Basic Clin. Pharmacol. Toxicol.* (2008) 102: 408-411.
- (13) Hayatshahi A, Javadi MR, Sarayani A and Gholami K. Clinical pharmacy residency in Iran. *ACCP International Clinical* 2013 Summer [cited 2013 Oct. 22]. Available from: URL: <http://www.accp.com/docs/international/0813.pdf>.
- (14) Hemerson P and Izakovic M. Hospitalist-pharmacist collaboration. A success story of the integration of a clinical pharmacist on the multidisciplinary hospitalist team. *Bratisl. Lek. Listy.* (2011) 112: 88-91.
- (15) Ratz Y, Shafir I, Berkovitch S, Sharristh M, Jacoby M, Kozer E, Golik A, Berkovitch M and Bar-Haim S. The importance of the pharmacist in reporting adverse drug reactions in the emergency department. *J. Clin. Pharmacol.* (2010) 50: 1217-1221.
- (16) Jose J and Rao PG. Pattern of adverse drug reactions notified by spontaneous reporting in an Indian tertiary care teaching hospital. *Pharmacol. Res.* (2006) 54: 226-233.
- (17) Runciman WB, Roughead EE, Semple SJ and Adams RJ. Adverse drug events and medication errors in Australia. *Int. J. Qual. Health Care* (2003) 15: 49-59.
- (18) Pirmohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ, Farrar K, Park BK and Breckenridge AM. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ.* (2004) 329: 15-19.
- (19) Camargo AL, Cardoso Ferreira MB and Heineck I. Adverse drug reactions: a cohort study in internal medicine units at a University hospital. *Eur. J. Clin. Pharmacol.* (2006) 62: 143-149.
- (20) Schneeweiss S, Göttinger M, Hasford J, Swoboda W, Hippus M, Hoffmann AK, Riethling AK and Krappweis J. First results from an intensified monitoring system to estimate drug-related hospital admissions. *Br. J. Clin. Pharmacol.* (2001) 52: 196-200.
- (21) Suh DC, Woodall BS, Shin SK and Hermes-De Santis ER. Clinical and economic impact of adverse drug reactions in hospitalized patients. *Ann. Pharmacother.* (2000) 34: 1373-1379.
- (22) Baniasadi S, Fahimi F and Namdar R. Development of an adverse drug reaction bulletin in a teaching hospital. *Formulary* (2009) 44: 333-335.
- (23) Fahimi F, Sahraee Z and Amini S. Evaluation of Stat Orders in a Teaching Hospital. *Clinical Drug Invest.* (2011) 31: 231-235.
- (24) Fahimi F. Implementation of a clinical pharmacy education program in a teaching hospital: Resident oriented documentation and intervention. *Iran. J. Pharm. Res.* (2010) 9: 297-302.
- (25) Allameh Z, Mehrpooya M, Baniasadi S and Fahimi F. Clinical pharmacy services in an Iranian teaching hospital: Type, severity, resolution, and accuracy. *J. Res. Pharm. Practice.* (2013) 2: 10-17.
- (26) Kanjanarat P, Winterstein AG, Johns TE, Hatton RC, Gonzalez-Rothi R and Segal R. Nature of preventable adverse drug events in hospitals: a literature review. *Am. J. Health Syst. Pharm.* (2003) 60: 1750-1759.
- (27) Baniasadi S and Fahimi F. Adverse drug reactions in a pulmonary teaching hospital: incidence, pattern, seriousness, and preventability. *Curr. Drug Saf.* (2011) 6: 230-236.
- (28) Fahimi F, Baniasadi S and Amini S. Adverse drug reactions in the post coronary care unit inpatients of a teaching hospital. *Iran. J. Pharm. Res.* (2008) 7: 223-228.
- (29) Bates DW, Cullen DJ, Laird N, Petersen LA, Small SD, Servi D, Laffel G, Schweitzer BJ, Shea BF, Hallisey R, Vander Vliet M, Nemeskal R, Leape LL, Bates D, Hojnowski-Diaz P, Petrycki S, Cotugno M, Patterson

- H, Hickey M, Kleefield S, Cooper J, Kinneally E, Demonaco HJ, Clapp MD, Gallivan T, Jeanette Ives, Porter K, Thompson BT, Hackman R and Edmondson A. ADE Prevention Group. Incidence of adverse drug events and potential adverse drug events. Implications for prevention. *JAMA*. (1995) 274: 29-34.
- (30) Ducharme MM and Boothby LA. Analysis of adverse drug reactions for preventability. *Int. J. Clin. Pract.* (2007) 61: 157-161.
- (31) Davies EC, Green CF, Taylor S, Williamson PR, Mottram DR and Pirmohamed M. Adverse drug reactions in hospital in-patients: a prospective analysis of 3695 patient-episodes. *PLoS One*. (2009) 4: 4439.
- (32) Dabaghzadeh F, Rashidian A, Torkamandi H, Alahyari S, Hanafi S, Farsaei S and Javadi MR. Medication errors in an emergency department in a large teaching hospital in Tehran. *Iran. J. Pharm. Res.* (2013) 12: 937-942.
- (33) Sadeghiana, Leila Safaeian GH, Mahdanian AR, Salami S and Kebriaee-Zadeh J. Prescribing quality in medical specialists in Isfahan, Iran. *Iran. J. Pharm. Res.* (2013) 12: 235-241.

This article is available online at <http://www.ijpr.ir>

Back issues?
Visit <http://www.ijpr.ir>
or
[http:// ijpr.sbm.ac.ir](http://ijpr.sbm.ac.ir)