

# A Link between Arabinose Utilization and Oxalotrophy in *Bradyrhizobium japonicum*

Marion Koch,<sup>a</sup> Nathanaël Delmotte,<sup>a</sup> Christian H. Ahrens,<sup>b</sup> Ulrich Omasits,<sup>b</sup> Kathrin Schneider,<sup>a</sup> Francesco Danza,<sup>a</sup> Barnali Padhi,<sup>a</sup> Valérie Murset,<sup>a</sup> Olivier Braissant,<sup>c</sup> Julia A. Vorholt,<sup>a</sup> Hauke Hennecke,<sup>a</sup> Gabriella Pessi<sup>a\*</sup>

Institute of Microbiology, ETH Zurich, Zurich, Switzerland<sup>a</sup>; Institute of Molecular Life Sciences, University of Zurich, Zurich, Switzerland<sup>b</sup>; Laboratory of Biomechanics and Biocalorimetry, University of Basel, Basel, Switzerland<sup>c</sup>

**Rhizobia have a versatile catabolism that allows them to compete successfully with other microorganisms for nutrients in the soil and in the rhizosphere of their respective host plants. In this study, *Bradyrhizobium japonicum* USDA 110 was found to be able to utilize oxalate as the sole carbon source. A proteome analysis of cells grown in minimal medium containing arabinose suggested that oxalate oxidation extends the arabinose degradation branch via glycolaldehyde. A mutant of the key pathway genes *oxc* (for oxalyl-coenzyme A decarboxylase) and *frc* (for formyl-coenzyme A transferase) was constructed and shown to be (i) impaired in growth on arabinose and (ii) unable to grow on oxalate. Oxalate was detected in roots and, at elevated levels, in root nodules of four different *B. japonicum* host plants. Mixed-inoculation experiments with wild-type and *oxc-frc* mutant cells revealed that oxalotrophy might be a beneficial trait of *B. japonicum* at some stage during legume root nodule colonization.**

Rhizobia have a broad metabolic capacity and can use a multitude of carbon and nitrogen sources which allow them to be successful and competitive in soil and in the rhizosphere of host plants (1). Within root nodules, differentiated rhizobia (bacteroids) reduce N<sub>2</sub> to ammonium, which is secreted to the plant in return for C<sub>4</sub>-dicarboxylic acids as carbon and energy sources (2). The C<sub>4</sub>-dicarboxylic acids malate, succinate, and fumarate have been shown to be the primary carbon sources of bacteroids and can actively cross the peribacteroid membrane (3–5). In fact, transport of C<sub>4</sub>-dicarboxylic acids is required for nitrogen fixation (6–8). C<sub>4</sub>-dicarboxylic acids are directly fed into the tricarboxylic acid (TCA) cycle to supply the bacteroid with enough energy to perform nitrogen fixation (3).

However, despite the fact that dicarboxylic acids were shown to be the major carbon sources for N<sub>2</sub>-fixing bacteroids, large quantities of hexose and pentose sugars are also found in nodules (9), suggesting an important role in nodule metabolism and N<sub>2</sub> fixation. Enzymes for hexose and pentose transport and metabolism have been reported to be present in bacteroids (10–12). The pentose L-arabinose is a well-known substrate of *B. japonicum* (13, 14), and this sugar is routinely used for *in vitro* growth studies. Previous work has shown that L-arabinose is degraded by a pathway that conceptually resembles the Entner-Doudoroff pathway (14). By analogy, L-arabinose is first oxidized to L-2-keto-3-deoxyarabonate (L-KDA) (15–17), which is then converted into  $\alpha$ -ketoglutarate in the case of fast-growing rhizobia (18–20) or to glycolaldehyde and pyruvate in the case of slow-growing species like *B. japonicum* (14). While pyruvate most likely is oxidized in the TCA cycle, the fate of glycolaldehyde has not been resolved yet. In this study, through proteomic analysis of *B. japonicum* cells grown either in minimal medium with L-arabinose as the sole carbon and energy source or in complete medium containing L-arabinose, we identified highly expressed products of candidate genes involved in the degradation of arabinose (17). Based on protein expression data, we suggest that glycolaldehyde is oxidized to glyoxylate and then reduced to glycerate through a glyoxylate carboligase and a tartronate semialdehyde reductase for assimilation and/or is converted to oxalate and then oxidized to formate and CO<sub>2</sub> through

the formyl-coenzyme A (CoA) transferase (Frc), oxalyl-CoA decarboxylase (Oxc), and formate dehydrogenase reactions for energy generation (Fig. 1). Results from previous transcriptome and proteome studies in *B. japonicum* nodules (12, 21) suggested that glyoxylate is preferentially degraded via oxalate oxidation during symbiosis. In fact, in contrast to the glyoxylate carboligase and tartronate semialdehyde reductase, the Frc and Oxc enzymes were detected in nodules where oxalate presence also could be measured. In agreement with these results, the  $\Delta$ *frc-oxc* mutant strain constructed in the present work is not able to grow on oxalate and is partially defective in free-living growth in minimal medium containing arabinose. We also show that the mutant has a disadvantage when competing for nodule occupancy against the wild-type strain.

## MATERIALS AND METHODS

**Bacterial strains, media, and growth conditions.** The bacterial strains and plasmids used in this work are listed in Table S1 in the supplemental material. *Escherichia coli* cells were cultivated in Luria-Bertani medium (22) at 37°C using the following concentrations of antibiotics (in  $\mu$ g/ml): ampicillin (200), kanamycin (30), chloramphenicol (20), and tetracycline (10). *Bradyrhizobium japonicum* cells were routinely cultivated at 30°C on peptone-salts-yeast extract (PSY) medium (23) supplemented with 0.1% arabinose or in defined buffered Vincent's minimal medium (BVM) (24, 25), referred to as minimal medium. Carbon sources used in defined

Received 4 October 2013 Accepted 13 January 2014

Published ahead of print 24 January 2014

Editor: G. Voordouw

Address correspondence to Gabriella Pessi, [gabriella.pessi@botinst.uzh.ch](mailto:gabriella.pessi@botinst.uzh.ch).


\* Present address: Gabriella Pessi, Institute of Plant Biology, University of Zurich, Zurich, Switzerland.

N.D. and C.H.A. contributed equally to this work.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AEM.03314-13>.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AEM.03314-13



**FIG 1** Proposed pathway for the catabolism of L-arabinose and its link to the oxalate degradation pathway (based on reference 17). Gene names and enzyme assignments were made according to amino acid sequence similarity to previously identified enzymes. The chemical structures have been created with ChemDraw.

media were filter sterilized and used at a final concentration of 20 mM arabinose or 20 mM succinate. The appropriate antibiotic concentrations (in  $\mu\text{g/ml}$ ) were added: spectinomycin (100) and kanamycin (100). Aerobic cultures for phenotypic growth analysis in PSY medium or BVM were grown in 500-ml Erlenmeyer flasks containing 25 ml medium supplemented with spectinomycin ( $100 \mu\text{g ml}^{-1}$ ) and the respective C source

on a shaker (160 rpm) at 30°C. For each strain or condition, the growth of three independent cultures was analyzed.

**Sample preparation and liquid chromatography-tandem MS (LC-MS/MS) analysis for proteomics.** Sample preparation and mass spectrometric (MS) analysis were performed as described in detail elsewhere (12). In brief, proteins extracted from three replicates of *B. japonicum* cells

grown in complex and minimal medium with L-arabinose until mid-exponential phase were separated on a Tris-HCl polyacrylamide gel. After reduction and carbamidomethylation the proteins were digested with trypsin (Promega, Madison, WI, USA), and the resulting peptides were separated by reverse-phase high-performance liquid chromatography (RP-HPLC) and analyzed by a hybrid LTQ-Orbitrap XL mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA) interfaced with a nano-electrospray source (12). Mass spectra were further processed with an in-house processing pipeline (26). In brief, fragment ion mass spectra were extracted from Thermo RAW files using msconvert (ProteoWizard, version 3.0.3831), merged in one MGF file per gel and searched against a composite *B. japonicum* USDA 110 protein database (RefSeq NC\_004463.1, 22 July 2013) containing 256 common contaminants (e.g., human keratin and trypsin). The search was performed with the powerful search engine MS-GF+ (MS-GFDB v7747) for a match to fully tryptic and semitryptic peptides with up to two missed cleavage sites and a mass tolerance of 25 ppm. Oxidation (M), deamidation (NQ), and methylation (DE) were used as variable modifications and carbamidomethylation (C) as a fixed modification. Using the decoy option of MS-GF+, we filtered the list of peptide spectrum matches (PSMs) to an estimated overall false discovery rate (FDR) of 0.2%, classified the PSMs with PeptideClassifier (27), and further considered only peptides (tryptic or semitryptic) that unambiguously imply one bacterial protein sequence. The FDR at the peptide level amounted to about 1.5%, and that at the protein level amounted to about 2.5% when requiring two peptides or three spectra for protein identification. A total of 3,390 and 3,149 proteins were identified in complex and minimal medium, respectively.


**Protein abundance calculation.** Relative protein abundance (in ppm) was estimated based on spectral counts as described by Schrimpf and colleagues (28) and extended recently (29).

**qPCR analysis.** RNA was extracted from *B. japonicum* 110*spc4* cultures grown in minimal medium, with 20 mM arabinose or 20 mM succinate as the C source, to mid-exponential phase. Cell harvest, RNA extraction, cDNA synthesis, and quantitative PCR (qPCR) analysis were done as described previously (30, 31).

**Metabolite extraction and HPLC-MS analysis.** Several intermediates of the L-arabinose degradation pathway (L-2-keto-3-deoxyarabonate, glycolaldehyde, and glyoxylate) as well as oxalate could be identified by a metabolomics approach (using nanoflow ion-pair RP-HPLC coupled with nanospray high-resolution MS with a split-free nano-LC Ultra system connected to an LTQ-Orbitrap mass spectrometer). Sampling, quenching, and central metabolite extraction were carried out as described by Schneider and colleagues (32). The data were normalized to the biomass produced in the cultures.

**Oxalotrophic growth.** The *B. japonicum* wild type and mutant strain  $\Delta$ *frc-oxc* were grown on double-layered Schlegel mineral medium plates supplemented with either Ca-oxalate or Na-oxalate (33). The first layer contains the following ingredients: Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O, 9.0 g/liter; KH<sub>2</sub>PO<sub>4</sub>, 1.5 g/liter; NH<sub>4</sub>Cl, 1.0 g/liter; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.2 g/liter; ammonium ferric citrate, 0.005 g/liter; CaCl<sub>2</sub>, 0.01 g/liter; ZnSO<sub>4</sub>·7H<sub>2</sub>O, 50 µg/liter; MnCl<sub>2</sub>·4H<sub>2</sub>O, 15 µg/liter; H<sub>3</sub>BO<sub>3</sub>, 150 µg/liter; CoCl<sub>2</sub>·6H<sub>2</sub>O, 100 µg/liter; CuCl<sub>2</sub>·2H<sub>2</sub>O, 50 µg/liter; NiCl<sub>2</sub>·6H<sub>2</sub>O, 10 µg/liter; NaMoO<sub>4</sub>·2H<sub>2</sub>O, 15 µg/liter; and agar, 15 g/liter. The second layer is composed of Schlegel mineral medium to which Ca-oxalate or Na-oxalate (7 g/liter or 2.7 g/liter; Sigma-Aldrich, Steinheim, Germany) was added (33). For this test, wild-type and mutant cultures were pregrown in minimal medium. Cells were washed and set to an optical density at 600 nm (OD<sub>600</sub>) of 4. A cotton swab was used to streak the cells on the plate. Plates were incubated at 30°C for 7 days. The assays were repeated three times.

**Isothermal calorimetry.** Isothermal calorimetry analyses were performed according to Bravo et al. (34). Briefly, solid Angle's medium (35) was supplemented with different carbon sources at a final concentration of 35 mM Ca-oxalate. The media were poured into sterile microcalorimetric ampoules to obtain slants. These slants were inoculated with *B. japonicum* wild-type and  $\Delta$ *frc-oxc* strains using a loop with an inoculum



**FIG 2** Physical map of the *B. japonicum* genomic region that harbors genes of the oxalate metabolism. Gene *blI3156* (*frc*) codes for a formyl-CoA transferase and *blI3157* (*oxc*) for an oxalyl-CoA decarboxylase. The precise structure of the *B. japonicum* deletion mutant is indicated with an arrow along with the strain number. Numbers below the vertical lines represent genome coordinates (the map is drawn to scale).

size sufficient to grow bacteria as a lawn. The ampoules were then sealed and introduced in the microcalorimeter (TAM48; Waters/TA, DE). After the thermal equilibration procedure, measurements were taken for at least 7 days. The data were recorded continuously by the microcalorimeter and resampled to obtain an effective sampling rate of 1 data point every 5 min. Samples were removed from the microcalorimeter and visually inspected in order to check that bacteria did form a lawn.

**Construction of mutant strains.** DNA was isolated from the *B. japonicum* wild-type strain 110*spc4* as previously described (36). Plasmid DNA from *E. coli* strains was obtained by using the NucleoSpin plasmid kit (Macherey-Nagel, Düren, Germany). Mutagenesis of genes was done by marker replacement. To construct a deletion mutant in the *blI3156-3157* genes, PCR fragments of the 5' and 3' flanking regions of *blI3156-3157* were amplified using the following primer pairs: *blI3156-1\_rev* (TACGGCTGGCGTCCGGCGAAC) and *blI3156-2\_for* (GTCTCTGGTCCGACCGCTTACTCGGCG) for the 3' region of *blI3156* and *blI3156-3\_for* (TGTCTCCCTGGTCTCAATACTG) and *blI3156-4\_rev* (GTCGAGCATCACCGGCTCG) for the 5' region of *blI3157*. PCR products were cloned in the pGEM-T Easy vector (Promega, Madison, WI, USA), and the correct sequence was verified by sequence analysis. Up- and downstream regions were subcloned into the pSUP202pol4 vector (37), and a kanamycin resistance cassette (*aphII*) derived from pBSL15 (38) was inserted between both regions. The resulting plasmid, pRJ6243, was used for conjugation with *B. japonicum* strain 110*spc4*. The correct genomic integration was verified by PCR. The resulting  $\Delta$ *frc-oxc* deletion mutant was named 6243 (Fig. 2).

**Plant material, inoculation, and cultivation.** Sterilization of soybean (*Glycine max* [L.] Merr. cv. Williams), mungbean (*Vigna radiata*), cowpea (*Vigna unguiculata*), and siratro (*Macroptilium atropurpureum*) seeds was done as described earlier (21). Plants were inoculated with cultures of *B. japonicum* that had been grown for 5 days in full medium and diluted to approximately 100 bacteria per plant. Nitrogenase activity was determined by using the acetylene reduction assay 21 days postinfection (dpi) for soybean, mungbean, and cowpea and 31 dpi for siratro. Bacteria were isolated from randomly selected nodules to confirm the presence of appropriate genetic markers.

**Competitiveness in symbiosis.** Soybean, mungbean, cowpea, and siratro plants were infected with a mixture of the wild type and the  $\Delta$ *frc-oxc* mutant strain containing a total of 100 bacteria per plant. Cultures of the wild type and mutant were grown and diluted to the same CFU per milliliter. A 90:10 ratio ( $\Delta$ *frc-oxc* to wild type) was chosen to assess symbiotic competition. Serial dilutions of the mixed inoculum were plated on selective agar to control the number of inoculated cells. All nodules from one plant were harvested at the peak of nitrogenase activity 21 (soybean, mungbean, and cowpea) and 31 (siratro) dpi. Nodules were surface sterilized (100% ethanol for 5 min) and rinsed in sterile distilled water. Nodules were then crushed in 1 ml PSY medium using a mortar, and this suspension was serially diluted and spotted on plates containing the appropriate selection for strain differentiation. The plates were incubated for 4 days at 30°C, and the ratio of the mutant to wild-type strain in nodule extracts was determined and compared to the initial inoculum ratio. As a

TABLE 1 Normalized protein abundance data for enzymes involved in L-arabinose degradation and oxalate catabolism in *B. japonicum*<sup>d</sup>

Enzyme <sup>c</sup>	Designation <sup>a</sup>	Normalized protein abundance <sup>b</sup> (ppm)				
		Symbiotic			Free living	
		Soybean	Cowpea	Siratiro	BVM plus arabinose	PSY plus arabinose
L-Arabinose 1-DH	Blr3205	23	38	42	522	502
L-Arabinonolactonase	Blr3207	31	55	13	555	458
Glycolate oxidase	Bll7540	114	81	91	98	68
	Bll7541	9	47	34	73	80
	Bll7543		5	11	46	45
Oxalyl-CoA decarboxylase (Oxc)	Bll3157	342	237	205	1,389	519
Formyl-CoA transferase (Frc)	Bll3156	364	394	211	800	604
Formate DH	Bll0288					
	Bll0280					
	Blr2317				63	56
	Bll3135					
	Bll3136				2	2
	Bll5475					
	Bll5476	18	66	207		
	Bll5477		19	108		2
Bll5478	27	49	145		4	
Glyoxylate carboligase	Blr3166				526	13
Tartronate semialdehyde reductase	Blr3168				774	28

<sup>a</sup> Nomenclature is according to Kaneko et al. (53).

<sup>b</sup> Calculated as described previously (28).

<sup>c</sup> DH, dehydrogenase.

<sup>d</sup> Cells were grown *in vitro* in minimal (BVM) and complex (PSY) media containing L-arabinose (the two rightmost columns) or were extracted from bacteroids of different host plants (12, 21). The proteomics data summarize expression from three independent biological replicates.

control, plants infected with either the wild type or mutant strain only were processed. At least three independent plants were processed per host. The nodule extracts were spotted in duplicates. Data were evaluated for statistical significance using Student's *t* test and SPSS 17.0 software.

**Determination of the oxalate content of roots and nodules of legumes inoculated with *B. japonicum*.** Root nodules of soybean, mungbean, cowpea, and siratro infected with *B. japonicum* as well as root-only material were collected, immediately flash-frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  until use. To assess the soluble oxalate content in nodules, 50 mg of nodule material was homogenized in 200  $\mu\text{l}$  distilled water ( $\text{dH}_2\text{O}$ ) using a TissueLyzer (Qiagen, Valencia, CA, USA) (1.3 min at 30 Hz) with a tungsten carbide bead (3 mm; Qiagen). One hundred mg of root material was ground in liquid nitrogen using a mortar and pestle. The resulting powder was collected and resuspended in 200  $\mu\text{l}$  of  $\text{dH}_2\text{O}$ . To ensure optimal destruction, the material was further homogenized in a TissueLyzer (1.3 min at 30 Hz). The supernatant of nodule and root samples was collected by a low-speed centrifugation step. The oxalate concentration in each sample was determined using a urinalysis diagnostic kit (TrinityBiotech, St. Louis, MO) according to the manufacturer's instructions. Measurements were done in triplicate on three independent plants.


**Proteomics data accession number.** Proteomics data associated with the manuscript can be downloaded from ProteomeXchange under accession number PXD000487.

## RESULTS AND DISCUSSION

**Proteome analysis of *B. japonicum* grown in minimal medium containing arabinose.** The proteome of cells grown in minimal medium containing 20 mM L-arabinose as the carbon and energy source revealed the presence of several enzymes involved in L-arabinose degradation (Fig. 1 and Table 1). Apart from detecting L-arabinose 1-dehydrogenase (Blr3205), L-arabinolactonase (Blr3207),

and glycolate oxidase (Bll7540-41, Bll7543), the enzymes glyoxylate carboligase (Blr3166) as well as the tartronate semialdehyde reductase (Blr3168) (Fig. 1) were expressed abundantly in minimal medium containing L-arabinose. Remarkably, large amounts of two enzymes involved in oxalate degradation were also detected: a formyl-CoA transferase (Frc) (bll3156) and an oxalyl-CoA decarboxylase (Oxc) (bll3157) that share 71% and 78% amino acid sequence identity with the previously studied Frc and Oxc of *Oxalobacter formigenes* (39–41). This led us to speculate that glycolaldehyde is oxidized to glyoxylate, which can be fed into at least two pathways: (i) reduction to glycerate through the activity of glyoxylate carboligase (blr3166) (42) and tartronate semialdehyde reductase (blr3168), or (ii) oxidation to oxalate followed by the stepwise, complete oxidation to formate and  $\text{CO}_2$  through the activities of Oxc and Frc (43, 44) and formate dehydrogenase. We previously showed that only the enzymes Oxc and Frc involved in the second pathway were detected in bacteroids during symbiosis with all different host plants (12, 21) (Table 1). Interestingly, using qPCR analysis, an elevated expression of bll3157 and bll3156 was not observed in *B. japonicum* cultures grown in minimal medium with succinate as the carbon source. The induction factors in arabinose-grown cells were measured as 45 for bll3157 and 116 for bll3156 compared to levels for succinate-grown cells. Thus, it appeared as if the oxalate oxidation pathway was switched on when L-arabinose was offered as the carbon source.

**Construction and growth analysis of a  $\Delta\text{frc-oxc}$  deletion mutant.** To further analyze the hypothesis that oxalate degradation is linked to arabinose utilization, a  $\Delta\text{frc-oxc}$  mutant strain was con-



**FIG 3** Growth analysis of the *B. japonicum* wild type and the  $\Delta frc-oxc$  mutant in minimal medium supplemented with two different carbon sources. (A) Twenty millimolar arabinose; (B) 20 mM succinate. Shown are mean values for all taken time points  $\pm$  standard deviations. (C) The heat production related to the metabolic activity during oxalotrophic growth of *B. japonicum* wild-type and mutant cells was monitored using isothermal microcalorimetry.

structured (strain 6243) (Fig. 2). Growth of *B. japonicum* wild-type and mutant strains was analyzed in minimal and complex medium. The minimal medium was supplemented with either 20 mM L-arabinose, 20 mM succinate, or 20 mM Na-oxalate.

First, we compared the ability of both the wild type and the  $\Delta frc-oxc$  mutant to catabolize oxalate using plate assays and microcalorimetry (see Materials and Methods). We initially aimed at monitoring oxalotrophic growth in liquid cultures using Ca- or Na-oxalate as the sole source of carbon. Even though various concentrations (5, 10, and 20 mM) of oxalate were used, these cultivation attempts failed. Therefore, we examined growth on Schlegel's minimal medium agar plates in the presence of 20 mM Na-oxalate or 35 mM Ca-oxalate as the sole carbon source (data not shown). For this test, strains were first grown in minimal medium and then streaked out on agar plates. This test revealed that the *B. japonicum* wild type was able to grow on plates supplemented with 20 mM Na-oxalate, whereas the  $\Delta frc-oxc$  mutant did not grow. Likewise, microcalorimetric analyses on Angle's medium supplemented with 35 mM Ca-oxalate revealed that the *B. japonicum* wild type was able to grow on Ca-oxalate, whereas no growth was observed for the  $\Delta frc-oxc$  mutant (Fig. 3C). The metabolic heat production resulting from the oxidation of oxalate according to the reaction  $2C_2H_2O_4 + O_2 \rightarrow 4CO_2 + 2H_2O$  with a reaction enthalpy ( $\Delta H^\circ_{rxn}$ ) of  $-499$  KJ/mol is directly proportional to the oxalate consumption rate. At this point, it must be noted that biomass production can easily be neglected (34), because biomass yield is extremely low with oxalate. Thus, each curve represents the overall metabolic activity related to the growth of *B. japonicum*. Such measurements can be compared using common metabolic assays, such as the 2,3,5-triphenyltetrazolium chloride assay, for example (45). The metabolic activity pattern for the *B. japonicum* wild-type strain was similar to what could be observed in other studies (34).

When grown on minimal medium with L-arabinose as the carbon source, the  $\Delta frc-oxc$  mutant displayed a diminished growth rate compared to that of the wild type (Fig. 3A). The mean generation time of the parental strain was 23.4 h, compared to 37.3 h for the  $\Delta frc-oxc$  mutant. Moreover, wild-type cells reached a final optical density of 5.6, while the mutant strain did not exceed an OD of 2. In contrast, the  $\Delta frc-oxc$  strain exhibited growth behavior similar to that of the wild-type strain when cultivated either in minimal medium with the dicarboxylic acid succinate (Fig. 3B) or in complex medium (data not shown). Thus, the results have shown that a deletion of two specific oxalate degradation genes caused a partial defect in growth with arabinose. This can be explained in at least two different ways: (i) in the mutant, L-arabinose may not be consumed as efficiently as in the wild type for lack of oxalate degradation and partial energy conservation through the formate dehydrogenase reaction, and (ii) the mutant may accumulate inhibitory concentrations of oxalate and its precursors. We tested the latter scenario by measuring the cellular contents of

**TABLE 2** Oxalate content of root material and root nodules of cowpea, mungbean, siratro, and soybean plants infected with the *B. japonicum* wild type

Host plant	Oxalate level ( $mg\ g^{-1}$ wet wt) in <sup>a</sup> :	
	Root	Root nodules
Cowpea	$0.10 \pm 0.01$	$0.37 \pm 0.08$
Mungbean	$0.09 \pm 0.01$	$0.31 \pm 0.05$
Siratro	$0.09 \pm 0.01$	$0.22 \pm 0.01$
Soybean	$0.03 \pm 0.01$	$0.18 \pm 0.02$

<sup>a</sup> Results are presented as means  $\pm$  standard deviations where  $n \geq 2$ .


oxalate, glyoxylate, and glycolaldehyde and found these compounds to be elevated by a factor of up to 2 in the mutant compared to the wild type (data not shown). Whether or not this slight increase is inhibitory to growth has not been further investigated. Regardless of which of the two explanations is true, the degradation of oxalate appears to be a necessary requirement for the complete utilization of arabinose for growth and biomass formation.

**Oxalate content in roots and root nodules induced by *B. japonicum*.** Since both *Frc* and *Oxc* had been detected in soybean, cowpea, and siratro bacteroids using a global proteomics approach (12, 21) (Table 1), we became interested in exploring whether oxalate is present in nodules as a potential C source for the bacteroids. Notably, it has been hypothesized previously that oxalate is a potential energy-yielding substrate sufficient to sustain nitrogen fixation in *Vicia faba* (46).

The oxalate content of roots and root nodules of four *B. japonicum* host plants (soybean, siratro, cowpea, and mungbean) was analyzed using an oxalate oxidase assay (Table 2). In general, three times higher oxalate concentrations were detected in root nodules than in the roots of uninfected plants (Table 2). This showed that oxalate is indeed present in nodules and might be available as a C source together with other compounds, such as succinate and malate.

**Plant symbiosis and competition for nodule occupancy.** The fact that *B. japonicum* host plants displayed different concentrations of soluble oxalate prompted us to investigate the role of the oxalate degradation genes in symbiosis. The symbiotic efficiency of the wild type and the  $\Delta frc-oxc$  mutant was tested on all plant hosts. Plants infected with  $\Delta frc-oxc$  grew as healthy as the wild type, displaying green leaves, normal nodule development, and wild-type-like nitrogenase activity 21 dpi for soybean, mungbean, and cowpea and 31 dpi for siratro (see Table S2 in the supplemental material). Moreover, the same number of nodules was observed in plants inoculated with the  $\Delta frc-oxc$  mutant and with the parental strain. Reisolation of bacteroids from plants used in this test revealed comparable viable cell counts for wild-type and mutant strains (data not shown).

It was previously shown that mutations in genes responsible for the catabolism of carbon sources, such as rhamnose and myo-inositol, and other used nutrients, such as mimosine, are correlated with a decreased ability to compete for nodule occupancy (47–50). Therefore, we investigated if the presence of the oxalate catabolic pathway could represent a competitive advantage for nodulation occupancy on all host plants. In order to test nodule colonization, 50 CFU/ml of *B. japonicum*  $\Delta frc-oxc$  and wild-type cells were mixed at a 90:10 ratio and used for inoculation of soybean, siratro, cowpea, and mungbean seedlings. After 21 dpi for soybean, mungbean, and cowpea and 31 dpi for siratro, all nodules from one plant were collected. In total, three independent experiments including at least three plants were performed per strain and host. To isolate bacteroids, root nodules were homogenized and the wild type-to-mutant ratio was investigated by comparing the recovered viable cell counts of  $\Delta frc-oxc$  and wild-type strains using genetic markers. In all host plants the  $\Delta frc-oxc$  mutant was affected in nodule occupancy when competing with the parental strain (Fig. 4), despite the fact that the mutant was present in 10-fold excess over the wild type in the inoculum mixture. A possible impact of the antibiotic resistance cassette (*aphII* in strain  $\Delta frc-oxc$ ) on the nodulation competition phenotype had been excluded by results presented recently (51), where an ACC deami-



**FIG 4** Symbiosis and competition for root-nodule colonization. Soybean, cowpea, siratro, and mungbean seedlings were coinoculated into 3 plants at a ratio of 10:90 *B. japonicum* wild-type (wt) to  $\Delta frc-oxc$  cells. Percentages of wild-type and mutant strains recovered from plant nodules at different inoculum ratios were compared. The tested host plants are shown along the x axis. The y axis represents mean values  $\pm$  standard deviations of percentages of bacteroids recovered from crushed nodules. Experimental data were assessed for statistical significance by means of the Student *t* test. Statistical analyses were performed using SPSS 17.0 software. Recovered percentages were significantly different between the wild type and mutant ( $P < 0.05$ ).

nase mutant containing the identical *aphII* antibiotic resistance cassette could compete as well as the wild type for nodule occupancy.

**Concluding remarks.** In this study, we demonstrated that *B. japonicum* is capable of entertaining an oxalotrophic lifestyle. Sequence analyses indicated that the *oxc* and *frc* genes involved in oxalate degradation are present and conserved in other *Bradyrhizobium* strains (52) but not in fast-growing rhizobia such as *Sinorhizobium meliloti*, *Rhizobium leguminosarum*, and *R. etli*. In this study, based on proteomics and growth analysis on different carbon sources, we showed that the degradation of L-arabinose creates intermediates that most likely are fed into the oxalate degradation pathway (Fig. 1). Disrupting the oxalate degradation branch by mutation leads to compromised growth on arabinose, because either the additive benefit from reductant formation in the formate dehydrogenase reaction is lacking or inhibitory amounts of oxalate and its precursors build up in the mutant. While the presence of functional formyl-CoA transferase (*frc*) and oxalyl-CoA decarboxylase (*oxc*) genes in *B. japonicum* is dispensable for the establishment of an effective symbiosis, these enzymes nevertheless seem to provide an advantage in the process of root-nodule colonization by *B. japonicum*. It can be speculated that at some point during rhizobial infection and nodule occupation the ability to degrade oxalate, which is present in roots and root nodules, represents a beneficial trait for *B. japonicum*.

#### ACKNOWLEDGMENTS

We thank Pilar Junier for advice on oxalotrophy analyses, Andrea Lindemann for initial work on L-arabinose degradation, Martina Lardi for help with statistical analysis, and Hans-Martin Fischer for constructive discussions.

This work was supported by ETH Zurich and by a grant from the Swiss National Foundation for Scientific Research.

We are grateful to Dulce-Nombre Rodríguez-Navarro and Francisco Temprano (Las Torres-Tomejil, Seville, Spain) for providing soybean

seeds and to William Broughton (University of Geneva, Switzerland) for cowpea and siratro seeds.

## REFERENCES

- Udvardi M, Poole PS. 2013. Transport and metabolism in legume-rhizobia symbioses. *Annu. Rev. Plant Biol.* 64:781–805. <http://dx.doi.org/10.1146/annurev-arplant-050312-120235>.
- Lodwig E, Poole P. 2003. Metabolism of *Rhizobium* bacteroids. *Crit. Rev. Plant Sci.* 22:37–78. <http://dx.doi.org/10.1080/713610850>.
- Prell J, Poole P. 2006. Metabolic changes of rhizobia in legume nodules. *Trends Microbiol.* 14:161–168. <http://dx.doi.org/10.1016/j.tim.2006.02.005>.
- Udvardi MK, Price GD, Gresshoff PM, Day DA. 1988. A dicarboxylate transporter on the peribacteroid membrane of soybean nodules. *FEBS Lett.* 231:36–40. [http://dx.doi.org/10.1016/0014-5793\(88\)80697-5](http://dx.doi.org/10.1016/0014-5793(88)80697-5).
- Udvardi MK, Day DA. 1997. Metabolite transport across symbiotic membranes of legume nodules. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 48:493–523. <http://dx.doi.org/10.1146/annurev.arplant.48.1.493>.
- Ronson CW, Lyttleton P, Robertson JG. 1981. C<sub>4</sub>-dicarboxylate transport mutants of *Rhizobium trifolii* form ineffective nodules on *Trifolium repens*. *Proc. Natl. Acad. Sci. U. S. A.* 78:4284–4288. <http://dx.doi.org/10.1073/pnas.78.7.4284>.
- Ronson CW, Astwood PM, Downie JA. 1984. Molecular cloning and genetic organization of C<sub>4</sub>-dicarboxylate transport genes from *Rhizobium leguminosarum*. *J. Bacteriol.* 160:903–909.
- Yurgel SN, Kahn ML. 2004. Dicarboxylate transport by rhizobia. *FEMS Microbiol. Rev.* 28:489–501. <http://dx.doi.org/10.1016/j.femsre.2004.04.002>.
- Streeter JG. 1980. Carbohydrates in soybean nodules. II. Distribution of compounds in seedlings during the onset of nitrogen fixation. *Plant Physiol.* 66:471–476.
- Glenn AR, McKay IA, Arwas R, Dilworth MJ. 1984. Sugar metabolism and the symbiotic properties of carbohydrate mutants of *Rhizobium leguminosarum*. *J. Gen. Microbiol.* 130:239–245.
- Salminen SO, Streeter JG. 1987. Uptake and metabolism of carbohydrates by *Bradyrhizobium japonicum* bacteroids. *Plant Physiol.* 83:535–540. <http://dx.doi.org/10.1104/pp.83.3.535>.
- Delmotte N, Ahrens CH, Knief C, Qeli E, Koch M, Fischer HM, Vorholt JA, Hennecke H, Pessi G. 2010. An integrated proteomics and transcriptomics reference data set provides new insights into the *Bradyrhizobium japonicum* bacteroid metabolism in soybean root nodules. *Proteomics* 10: 1391–1400. <http://dx.doi.org/10.1002/pmic.200900710>.
- Stowers MD. 1985. Carbon metabolism in *Rhizobium* species. *Annu. Rev. Microbiol.* 39:89–108. <http://dx.doi.org/10.1146/annurev.mi.39.100185.000513>.
- Pedrosa FO, Zancan GT. 1974. L-Arabinose metabolism in *Rhizobium japonicum*. *J. Bacteriol.* 119:336–338.
- Novick NJ, Tyler ME. 1982. L-Arabinose metabolism in *Azospirillum brasilense*. *J. Bacteriol.* 149:364–367.
- Watanabe S, Kodaki T, Makino K. 2006. Cloning, expression, and characterization of bacterial L-arabinose 1-dehydrogenase involved in an alternative pathway of L-arabinose metabolism. *J. Biol. Chem.* 281:2612–2623. <http://dx.doi.org/10.1074/jbc.M506477200>.
- Watanabe S, Shimada N, Tajima K, Kodaki T, Makino K. 2006. Identification and characterization of L-arabinose dehydratase, L-2-keto-3-deoxyarabonate dehydratase, and L-arabinolactonase involved in an alternative pathway of L-arabinose metabolism. Novel evolutionary insight into sugar metabolism. *J. Biol. Chem.* 281:33521–33536. <http://dx.doi.org/10.1074/jbc.M606727200>.
- Duncan MJ, Fraenkel DG. 1979.  $\alpha$ -Ketoglutarate dehydrogenase mutant of *Rhizobium meliloti*. *J. Bacteriol.* 137:415–419.
- Watanabe S, Kodaki T, Makino K. 2006. A novel  $\alpha$ -ketoglutaric semialdehyde dehydrogenase: evolutionary insight into an alternative pathway of bacterial L-arabinose metabolism. *J. Biol. Chem.* 281:28876–28888. <http://dx.doi.org/10.1074/jbc.M602585200>.
- Poysti NJ, Loewen ED, Wang Z, Oresnik IJ. 2007. *Sinorhizobium meliloti* pSymB carries genes necessary for arabinose transport and catabolism. *Microbiology* 153:727–736. <http://dx.doi.org/10.1099/mic.0.29148-0>.
- Koch M, Delmotte N, Rehrauer H, Vorholt JA, Pessi G, Hennecke H. 2010. Rhizobial adaptation to hosts, a new facet in the legume root-nodule symbiosis. *Mol. Plant Microbe Interact.* 23:784–790. <http://dx.doi.org/10.1094/MPMI-23-6-0784>.
- Miller J. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Regensburger B, Hennecke H. 1983. RNA polymerase from *Rhizobium japonicum*. *Arch. Microbiol.* 135:103–109. <http://dx.doi.org/10.1007/BF00408017>.
- Serventi F, Youard ZA, Murset V, Huwiler S, Bühler D, Richter M, Luchsinger R, Fischer HM, Brogioli R, Niederer M, Hennecke H. 2012. Copper starvation-inducible protein for cytochrome oxidase biogenesis in *Bradyrhizobium japonicum*. *J. Biol. Chem.* 287:38812–38823. <http://dx.doi.org/10.1074/jbc.M112.406173>.
- Vincent JM. 1970. A manual for the practical study of root nodule bacteria. Blackwell Scientific Publications, Oxford, United Kingdom.
- Omasits U, Quebatte M, Stekhoven DJ, Fortes C, Roschitzki B, Robinson MD, Dehio C, Ahrens CH. 2013. Directed shotgun proteomics guided by saturated RNA-seq identifies a complete expressed prokaryotic proteome. *Genome Res.* 23:1916–1927. <http://dx.doi.org/10.1101/gr.151035.112>.
- Qeli E, Ahrens CH. 2010. PeptideClassifier for protein inference and targeted quantitative proteomics. *Nat. Biotechnol.* 28:647–650. <http://dx.doi.org/10.1038/nbt0710-647>.
- Schrimpf SP, Weiss M, Reiter L, Ahrens CH, Jovanovic M, Malmstrom J, Brunner E, Mohanty S, Lercher MJ, Hunziker PE, Aebersold R, von Mering C, Hengartner MO. 2009. Comparative functional analysis of the *Caenorhabditis elegans* and *Drosophila melanogaster* proteomes. *PLoS Biol.* 7:e48. <http://dx.doi.org/10.1371/journal.pbio.1000048>.
- Carlier AL, Omasits U, Ahrens CH, Eberl L. 2013. Proteomics analysis of *Psychotria* leaf nodule symbiosis: improved genome annotation and metabolic predictions. *Mol. Plant Microbe Interact.* 26:1325–1333. <http://dx.doi.org/10.1094/MPMI-05-13-0152-R>.
- Pessi G, Ahrens CH, Rehrauer H, Lindemann A, Hauser F, Fischer HM, Hennecke H. 2007. Genome-wide transcript analysis of *Bradyrhizobium japonicum* bacteroids in soybean root nodules. *Mol. Plant Microbe Interact.* 20:1353–1363. <http://dx.doi.org/10.1094/MPMI-20-11-1353>.
- Lindemann A, Moser A, Pessi G, Hauser F, Friberg M, Hennecke H, Fischer HM. 2007. New target genes controlled by the *Bradyrhizobium japonicum* two-component regulatory system RegSR. *J. Bacteriol.* 189: 8928–8943. <http://dx.doi.org/10.1128/JB.01088-07>.
- Schneider K, Skovran E, Vorholt JA. 2012. Oxalyl-coenzyme A reduction to glyoxylate is the preferred route of oxalate assimilation in *Methylobacterium extorquens* AM1. *J. Bacteriol.* 194:3144–3155. <http://dx.doi.org/10.1128/JB.00288-12>.
- Aragno MS, Schlegel HG. 1991. The mesophilic hydrogen-oxidizing (Knallgas) bacteria. Springer Verlag, Berlin, Heidelberg, New York.
- Bravo D, Braissant O, Solokhina A, Clerc M, Daniels AU, Verrecchia E, Junier P. 2011. Use of an isothermal microcalorimetry assay to characterize microbial oxalotrophic activity. *FEMS Microbiol. Ecol.* 78:266–274. <http://dx.doi.org/10.1111/j.1574-6941.2011.01158.x>.
- Angle JS, McGrath SP, Chaney RL. 1991. New culture medium containing ionic concentrations of nutrients similar to concentrations found in the soil solution. *Appl. Environ. Microbiol.* 57:3674–3676.
- Hahn M, Hennecke H. 1984. Localized mutagenesis in *Rhizobium japonicum*. *Mol. Gen. Genet.* 193:46–52. <http://dx.doi.org/10.1007/BF00327412>.
- Fischer HM, Babst M, Kaspar T, Acuña T, Arigoni F, Hennecke H. 1993. One member of a *groESL*-like chaperonin multigene family in *Bradyrhizobium japonicum* is co-regulated with symbiotic nitrogen fixation genes. *EMBO J.* 12:2901–2912.
- Alexeyev MF. 1995. Three kanamycin resistance gene cassettes with different polylinkers. *Biotechniques* 18:52–56.
- Lung HY, Baetz AL, Peck AB. 1994. Molecular cloning, DNA sequence, and gene expression of the oxalyl-coenzyme A decarboxylase gene, *oxc*, from the bacterium *Oxalobacter formigenes*. *J. Bacteriol.* 176:2468–2472.
- Sidhu H, Ogden SD, Lung HY, Luttge BG, Baetz AL, Peck AB. 1997. DNA sequencing and expression of the formyl coenzyme A transferase gene, *frc*, from *Oxalobacter formigenes*. *J. Bacteriol.* 179:3378–3381.
- Sahin N. 2003. Oxalotrophic bacteria. *Res. Microbiol.* 154:399–407. [http://dx.doi.org/10.1016/S0923-2508\(03\)00112-8](http://dx.doi.org/10.1016/S0923-2508(03)00112-8).
- Chang YY, Wang AY, Cronan JE, Jr. 1993. Molecular cloning, DNA sequencing, and biochemical analyses of *Escherichia coli* glyoxylate carboxylase, an enzyme of the acetoacetyl acid synthase-pyruvate oxidase family. *J. Biol. Chem.* 268:3911–3919.
- Stewart CS, Duncan SH, Cave DR. 2004. *Oxalobacter formigenes* and its role in oxalate metabolism in the human gut. *FEMS Microbiol. Lett.* 230: 1–7. [http://dx.doi.org/10.1016/S0378-1097\(03\)00864-4](http://dx.doi.org/10.1016/S0378-1097(03)00864-4).
- Svedruzic D, Jonsson S, Toyota CG, Reinhardt LA, Ricagno S, Linqvist

- Y, Richards NGJ. 2005. The enzymes of oxalate metabolism: unexpected structures and mechanisms. *Arch. Biochem. Biophys.* 433:176–192. <http://dx.doi.org/10.1016/j.abb.2004.08.032>.
45. Braissant O, Wirz D, Gopfert B, Daniels AU. 2010. Use of isothermal microcalorimetry to monitor microbial activities. *FEMS Microbiol. Lett.* 303:1–8. <http://dx.doi.org/10.1111/j.1574-6968.2009.01819.x>.
  46. Trinchant JC, Guerin V, Rigaud J. 1994. Acetylene reduction by symbiosomes and free bacteroids from broad bean (*Vicia faba* L.) nodules. *Plant Physiol.* 105:555–561.
  47. Fry J, Wood M, Poole PS. 2001. Investigation of myo-inositol catabolism in *Rhizobium leguminosarum* bv. viciae and its effect on nodulation competitiveness. *Mol. Plant Microbe Interact.* 14:1016–1025. <http://dx.doi.org/10.1094/MPMI.2001.14.8.1016>.
  48. Jimenez-Zurdo JI, van Dillewijn P, Soto MJ, de Felipe MR, Olivares J, Toro N. 1995. Characterization of a *Rhizobium meliloti* proline dehydrogenase mutant altered in nodulation efficiency and competitiveness on alfalfa roots. *Mol. Plant Microbe Interact.* 8:492–498. <http://dx.doi.org/10.1094/MPMI-8-0492>.
  49. Oresnik IJ, Pacarynuk LA, O'Brien SAP, Yost CK, Hynes MF. 1998. Plasmid-encoded catabolic genes in *Rhizobium leguminosarum* bv. trifolii: evidence for a plant-inducible rhamnose locus involved in competition for nodulation. *Mol. Plant Microbe Interact.* 11:1175–1185. <http://dx.doi.org/10.1094/MPMI.1998.11.12.1175>.
  50. Soedarjo M, Borthakur D. 1998. Mimosine, a toxin produced by the tree-legume *Leucaena* provides a nodulation competition advantage to mimosine-degrading *Rhizobium* strains. *Soil Biol. Biochem.* 30:1605–1613. [http://dx.doi.org/10.1016/S0038-0717\(97\)00180-6](http://dx.doi.org/10.1016/S0038-0717(97)00180-6).
  51. Murset V, Hennecke H, Pessi G. 2012. Disparate role of rhizobial ACC deaminase in root-nodule symbioses. *Symbiosis* 57:43–50. <http://dx.doi.org/10.1007/s13199-012-0177-z>.
  52. Bravo D, Martin G, David MM, Cailleau G, Verrecchia E, Junier P. 2013. Identification of active oxalotrophic bacteria by bromodeoxyuridine DNA-labeling in a microcosm soil experiments. *FEMS Microbiol. Lett.* 348:103–111. <http://dx.doi.org/10.1111/1574-6968.12244>.
  53. Kaneko T, Nakamura Y, Sato S, Minamisawa K, Uchiumi T, Sasamoto S, Watanabe A, Idesawa K, Iriguchi M, Kawashima K, Kohara M, Matsumoto M, Shimpo S, Tsuruoka H, Wada T, Yamada M, Tabata S. 2002. Complete genomic sequence of nitrogen-fixing symbiotic bacterium *Bradyrhizobium japonicum* USDA110. *DNA Res.* 9:189–197. <http://dx.doi.org/10.1093/dnares/9.6.189>.