



Published in final edited form as:

*Dev Neurosci.* 2014 ; 36(1): 1–9. doi:10.1159/000356219.

## Improved Fronto-Parietal White Matter Integrity in Overweight Children is Associated with Attendance in an After-School Exercise Program

Cynthia E. Krafft<sup>a</sup>, David J. Schaeffer<sup>b</sup>, Nicolette F. Schwarz<sup>a</sup>, Lingxi Chi<sup>a</sup>, Abby L. Weinberger<sup>a</sup>, Jordan E. Pierce<sup>a</sup>, Amanda L. Rodrigue<sup>a</sup>, Jerry D. Allison<sup>c</sup>, Nathan E. Yanasak<sup>c</sup>, Tianming Liu<sup>d</sup>, Catherine L. Davis<sup>e</sup>, and Jennifer E. McDowell<sup>a</sup>

<sup>a</sup>Psychology Department, University of Georgia

<sup>b</sup>Neuroscience Department, University of Georgia

<sup>c</sup>Radiology Department, Medical College of Georgia, Georgia Regents University

<sup>d</sup>Computer Science Department, University of Georgia

<sup>e</sup>Pediatrics Department and Georgia Prevention Center, Medical College of Georgia, Georgia Regents University

### Abstract

Aerobic fitness is associated with white matter integrity (WMI) in adults as measured by diffusion tensor imaging (DTI). This study examined the effect of an 8-month exercise intervention on WMI in children. Participants were 18 sedentary, overweight (body mass index ≥85th percentile) 8- to 11-year-old children (94% Black), randomly assigned to either an aerobic exercise ( $n=10$ ) or sedentary attention control group ( $n=8$ ). Each group was offered an instructor-led after-school program every school day for approximately 8 months. Before and after the program, all subjects participated in DTI scans. Tractography was conducted to isolate the superior longitudinal fasciculus and investigate whether the exercise intervention affected WMI in this region. There was no group by time interaction for WMI in the superior longitudinal fasciculus. There was a group by time by attendance interaction, however, such that higher attendance at the exercise intervention, but not the control intervention, was associated with increased WMI. Heart rate and the total dose of exercise correlated with WMI changes in the exercise group. In the overall sample, increased WMI was associated with improved scores on a measure of attention and improved teacher ratings of executive function. This study indicates that participating in an exercise intervention improves WMI in children as compared to a sedentary after-school program.

### Keywords

aerobic exercise; children; cognition; cognitive control; diffusion tensor imaging; overweight; superior longitudinal fasciculus

Aerobic fitness is associated with cognitive performance across the lifespan. In cross-sectional studies, higher-fit individuals showed better performance than lower-fit peers on measures of cognitive control, and higher-fit children demonstrated comparatively better academic achievement and parent and teacher ratings of behavior [1; 2; 3; 4; 5]. There is evidence that aerobic exercise plays a causal role in improving cognition, as aerobic exercise training improves performance on a variety of cognitive tasks. Randomized controlled trials showed that both adults and children who were assigned to chronic exercise improved cognitive performance compared to their peers assigned to control groups [6; 7; 8] with more improvement associated with greater attendance [9]. While exercise benefits many types of cognition, there is evidence that exercise-related improvement is greater for cognitive control (higher-order processes including inhibition, working memory, and attentional control) compared to lower-level tasks [10].

Cognitive control is supported by fronto-parietal circuitry [11; 12]. Exercise interventions improved cognitive control and altered frontal and parietal activation as assessed by functional magnetic resonance imaging (fMRI). Specifically, a 6-month exercise intervention in older adults improved performance and increased prefrontal and posterior parietal activation during a flanker task in an exercise group as compared to controls [7]. Regional alterations also were found in studies of children. Our group initially found that a three-month exercise intervention in 7- to 11-year-old children improved cognitive control performance (as measured by the Planning scale of the Cognitive Assessment System) compared to controls [6]. The exercise group also increased prefrontal and decreased posterior parietal activation during an antisaccade task compared to controls. We recently extended these findings using an eight-month intervention (the current study). We found that children assigned to eight months of exercise showed altered activation during antisaccade and flanker tasks as compared to the sedentary attention control group [13].

With evidence that fronto-parietal brain activation is affected by exercise, one issue that warrants investigation is whether exercise alters fronto-parietal brain structure. Altered white matter structure may be an underlying cause of functional change, given evidence that inter-individual differences in brain activation reflect white matter integrity (WMI) [14]. WMI reflects axonal membrane structure and myelination and can be assessed by diffusion tensor imaging (DTI), which measures the anisotropy (directional dependence) of water diffusion. Fractional anisotropy (FA) is a frequent measure of interest in DTI and describes the anisotropy of water diffusion. FA values range between 0 and 1, with 1 indicating fully anisotropic diffusion. Higher values are generally interpreted as greater WMI (myelination and axonal membrane structure [15]). Another measure based on the same tensor model is radial diffusivity (RD), which measures water diffusion perpendicular to axons (with higher values indicating greater diffusion). Taken together, higher FA and lower RD values are often interpreted as primarily reflecting greater myelination [16; 17].

WMI was associated with fitness in several cross-sectional studies. Higher aerobic fitness in adults was associated with higher FA in the cingulum and corpus callosum, possibly relating to motor planning and control [18; 19; 20]. Fitness also was associated with integrity of the uncinate fasciculus, which is involved in memory [20]. In one of the first investigations to date of the impact of a randomized controlled exercise intervention on WMI, Voss et al.

randomly assigned adults to one year of either an aerobic exercise group or a flexibility and toning control group. Increased aerobic fitness was associated with increases in prefrontal, parietal, and temporal FA in the aerobic group [21]. While literature indicates that exercise affects WMI in adults, this topic has yet to be investigated in children.

Given evidence that exercise improved cognitive control and altered associated frontoparietal brain activation in prior studies, we investigated whether an exercise intervention in overweight children improves WMI in a tract that connects frontal and parietal regions: the superior longitudinal fasciculus (SLF). Only overweight participants were recruited for the current study since sedentary, obese children are likely to show the greatest benefits from exercise. Greater integrity of the SLF is related to better performance on several measures of cognitive control, such as working memory and attention [16; 22; 23]. As the SLF does not completely mature until young adulthood [24], ongoing development makes it an interesting target for investigation across the ages included in the current study (children 8 – 11 years old). The SLF may be particularly susceptible to intervention effects due to its continuing maturation and in fact was altered by another type of intervention in children. Five weeks of spelling training decreased RD in the right SLF in children 9 to 16 years of age [25]. In sum, the hypotheses were generated based on the literature indicating that exercise improves cognitive control and that fitness is associated cross-sectionally with higher WMI. Specifically, we hypothesized that a randomized controlled exercise intervention in children would increase WMI in the SLF, particularly in children with good attendance.

## Methods and Procedures

### Participants

Participants (94% Black) were a subset of children in a larger randomized trial [13], recruited from public schools around Augusta, Georgia, U.S.A. and eligible if they were 8-11 years old, overweight (BMI ≥ 85<sup>th</sup> percentile; [26]), and inactive (no regular physical activity program ≥ 1 hr/week). Exclusions included any medical condition that would limit physical activity or affect study results (including neurological or psychiatric disorders). Children and parents completed written informed assent and consent in accordance with the Human Assurance Committee of Georgia Regents University. Each child's parent or guardian reported the child's age, sex, race, and health status. Parents also reported their own educational attainment, used as an index of socioeconomic status (1 = 7 or fewer years of school; 2 = 8 – 9 years of school; 3 = 10 – 11 years of school; 4 = 12 years of school; 5 = partial college; 6 = college graduate; 7 = post-graduate). The study took place at the Georgia Prevention Center at Georgia Regents University. Magnetic resonance imaging was completed with DTI data available for 41 children at baseline and 30 at post-test. Three of the 30 children scanned at post-test were not scanned at baseline. Of the 27 children with both baseline and post-test data, 8 were excluded due to scanner artifact or excessive motion and 1 was excluded due to a neurological anomaly. Thus the present study included eighteen children (10 exercise, 8 control; see Table 1).

## Cognitive Measures

The Cognitive Assessment System (CAS), a standardized individual assessment of children's cognitive processes [27], was administered and Standard Scores analyzed for all 18 children at both baseline and post-test. The Full Scale score of the CAS takes into account the four scales of the test (Planning, Attention, Simultaneous and Successive processing), each of which include 3 subtests. For the CAS, higher scores reflect better performance. The Behavioral Rating Inventory of Executive Function (BRIEF) was completed by up to three teachers per child [28]. This questionnaire provides a standardized measure of executive function behaviors in the school environment, with higher scores reflecting worse performance. For each child, teachers completing the BRIEF were matched at baseline and post-test to maintain consistency. T-scores were obtained for 17 children at baseline (10 exercise, 7 control) and 16 at post-test (9 exercise, 7 control), with 15 children having both baseline and post-test data and thus being included in BRIEF analyses (9 exercise, 6 control). The BRIEF was completed by an average of 1.6 teachers ( $SD = 0.8$ ) per child, with no significant difference in the number of teachers between groups ( $t(13) = 1.02$ ,  $p > 0.3$ ).

## Intervention

Participants were assigned randomly to one of two conditions, both of which were offered for 40 min/day: aerobic exercise or sedentary attention control. Randomization (balanced by race, sex, and school) was performed by the study statistician and concealed until after baseline testing was completed, at which point the study coordinator informed the families. Both groups were offered an after-school program every school day for approximately 8 months (average number of days offered = 139,  $SD = 9$ ). All participants were offered daily bus transportation after school to the Georgia Prevention Center where they spent half an hour on supervised homework time and were given a snack. Lead instructors were rotated between the two groups every two weeks and assistants were rotated between the two groups every week.

The aerobic exercise group engaged in instructor-led aerobic activities (e.g., tag and jump rope). They wore heart rate monitors every day (S610i; Polar Electro, Oy, Finland) with which they could monitor their own performance and from which data were collected daily. Participants in the exercise group had an average heart rate of 161 beats per minute ( $SD = 8$ ) during the intervention. The product of each child's average heart rate with their attendance provided an index of total energy expenditure during the exercise intervention. The attention control group engaged in instructor-led sedentary activities (e.g., art and board games).

## Diffusion Tensor Imaging Procedure and Analysis

**MR image acquisition**—Images were acquired at Georgia Regents University on a 3T GE Signa Excite HDx MRI system (General Electric Medical Systems, Milwaukee, WI). During scanning, head position was stabilized with a vacuum pillow and/or foam padding. Diffusion images were acquired using an echo planar imaging (EPI) sequence (acquisition matrix =  $128 \times 128$ , 60 interleaved slices, voxel size =  $1 \times 1 \times 2.4$  mm, FOV =  $256 \times 256$  mm, TR = 15500 ms, TE = min-full, 3 B0 images, 30 diffusion weighted images,  $b = 1000$  s/mm<sup>2</sup>).

**Image analysis**—Raw diffusion images were converted from GE DICOM format to NIFTI format using the dcm2nii tool [29]. For each subject, volumes were visually inspected for motion artifacts; volumes distorted by motion were removed from the image series and b value/vector tables (volumes removed  $M = 2$ ,  $SD = 2.7$ ). To test for inhomogeneity of gradient application due to volume removal, volume gradient vectors were plotted on a sphere after motion volumes were removed. If the any surface of the sphere had a gap greater than the free surface area of 6 non-co-linear directions, the participant was excluded from analysis (two participants thus were excluded for excessive motion). Diffusion tensor image preprocessing was conducted using the FMRIB Software Library (FSL; [30]). Non-brain tissue was removed using the Brain Extraction Tool (BET; [31]). Diffusion images were corrected for eddy-current-induced distortions. Tractography was then conducted with whole brain tensors in individual space for the bilateral SLF following established anatomical markers [32] using the ExploreDTI software package [33]. Average values for each SLF (right and left) were extracted for four measures of white matter microstructure, all of which were included in analyses: fractional anisotropy (FA), radial diffusivity (RD), mean diffusivity (MD), and axial diffusivity (AD). High MD values are often explained as disorganized development, immaturity, or other structural anomalies in white matter, whereas high AD values are thought to reflect a lower number of axonal fibers or smaller axonal diameter.

Statistical analyses of white matter variables were conducted in SPSS Version 21 (IBM, Armonk, NY). Repeated measures ANOVAs were conducted for each measure of WMI to examine effects of group, attendance, measurement time, and their interactions on the outcome. Each model controlled for age and gender. Group by time interactions were the primary tests of our hypothesis. Time by group by attendance interactions tested whether attendance was associated with change in WMI differently between groups. The same analyses also were conducted for cognitive measures. Correlations between change in WMI and attendance were conducted separately for each group to examine significant group by attendance interactions. For the exercise group alone, correlations were conducted between change in WMI and average heart rate at the intervention, as well as the product of average heart rate and attendance.

An exploratory analysis was conducted to investigate whether improved white matter integrity was associated with improved cognitive performance in the overall sample. For each white matter variable that showed a significant time by group by attendance interaction, the change from baseline to post-test was calculated for each individual, controlling for age and gender. These difference scores were then entered into two separate  $k$ -means clustering analyses, one each for left and right SLF ( $k = 2$ , input: FA and RD for the hemisphere of interest; 10 iterations). T-tests evaluated differences in cognitive task performance between the resulting clusters.

## Results

### Cognitive Measures

Participants attended an average of 3.3 days per week ( $M = 65\%$  attendance,  $SD = 6\%$ ) and the groups not differ significantly in the percentage of days they attended the program out of

the number of days offered ( $t(16) = 0.95, p = .36$ ). The groups did not differ significantly at baseline on any of the characteristics listed in Table 1. There was no significant group by time interaction in any cognitive measure, indicating that the exercise intervention did not differentially affect cognition compared to the control condition. Time by group by attendance interactions were significant for BRIEF Global Executive Composite and Metacognition Index scores,  $F(1,9) = 13.17, p < .01$  and  $F(1,9) = 8.49, p = .02$ , respectively. For both interactions, attendance at the exercise intervention was associated with more improvement in scores as compared to attendance at the control condition. Partial correlations between attendance and change in BRIEF scores (controlling for age and gender) were not significant for either group alone (a trend-level correlation was found between attendance and worse Metacognition Index scores in the control group,  $r(2) = .94, p = .06$ ).

### White Matter Structure

There was no significant group by time interaction in any WMI measure, indicating that the exercise intervention did not differentially affect WMI compared to the control condition. Time by group by attendance interactions were significant in left SLF for both FA ( $F(1,12) = 7.4, p = .02$ ) and RD ( $F(1,12) = 5.3, p = .04$ ), and in right SLF for RD ( $F(1,12) = 5.4, p = .04$ ) and at the trend-level for FA ( $F(1,12) = 4.2, p = .06$ ). For all group by time by attendance interactions, the exercise group showed a significant relationship between attendance and change in WMI, with increased WMI associated with more frequent attendance ( $r(6) = .92, p < .01$  for right SLF FA and  $r(6) = -.88, p < .01$  for right SLF RD) or trend-level ( $r(6) = .64, p = .06$  for left SLF FA and  $r(6) = -.69, p = .06$  for left SLF RD). The control group did not show significant relationships between attendance and change in WMI (all  $p$  values  $> .10$ ). Group by attendance interactions were not significant for MD or AD. For further details, see Figure 1.

In addition to attendance, the intensity and dose of exercise were related to improved white matter integrity in the exercise group. Higher average heart rate in the exercise intervention was correlated with decreased MD ( $r(6) = -.71, p < .05$ ) and RD ( $r(6) = -.75, p < .05$ ) in the left SLF. A measure of the dose of exercise (the product of average heart rate and attendance) was correlated with increased FA in left and right SLF ( $r(6) = .73, p = .04$  and  $r(6) = .93, p < .01$ , respectively), decreased MD in the right SLF ( $r(6) = -.75, p = .03$ ), and decreased RD in the left and right SLF ( $r(6) = -.74, p = .04$  and  $r(6) = -.87, p < .01$ , respectively). There were no significant correlations between age and any measure of WMI at either baseline or post-test (all  $p$  values  $> .07$ ).

### Exploratory K-means Analyses

Cluster analyses were conducted to supplement the main results by illustrating the overall relationship between change in WMI and change in cognitive performance. Analyses divided all participants into two groups based on those who improved WMI (FA and RD) and those who did not for each SLF (left and right). With two groups identified for each SLF, it was then possible to compare alterations in cognitive measures (CAS and BRIEF scores) between the improved and not-improved SLF groups in order to investigate whether improved WMI was associated with improved cognition in the overall study.



**Cluster analysis 1: Left SLF**—Two clusters were identified which significantly differed in the change in WMI from baseline to post-test. Cluster membership did not significantly overlap with exercise/control group membership. The cluster that showed greater improvement in WMI also showed better cognitive outcome as indicated by decreased BRIEF scores. Specifically, Cluster 1 ( $n = 11$ ) was composed of participants with improved left SLF WMI as indicated by increased FA ( $t(16) = -6.1, p < .01$ ) and decreased RD ( $t(16) = 3.7, p < .01$ ) compared to Cluster 2 ( $n = 7$ ). Cluster 1 was associated with more cognitive improvement, significantly differing from Cluster 2 in change in BRIEF scores over time. Cluster 2 had significantly worse scores on the BRIEF as compared to Cluster 1, including the Global Executive Composite Score ( $t(13) = 3.2, p < .01$ ), as well as both the Behavioral Regulation and Metacognition Indices ( $t(13) = 2.5, p = .04$  and  $t(13) = 3.04, p = .01$ , respectively). No differences were observed between the two clusters in CAS scores. For further details, see Figure 2.

**Cluster analysis 2: Right SLF**—Two clusters were identified which significantly differed in the change in WMI from baseline to post-test. Cluster membership did not significantly overlap with exercise/control group membership. The cluster showing greater improvement in WMI also showed better cognitive outcome as indicated by increased CAS Attention Scale scores. Specifically, Cluster 1 ( $n = 11$ ) was composed of participants with improved right SLF WMI as indicated by increased FA ( $t(16) = 10, p < .01$ ) and decreased RD ( $t(16) = 2.5, p = .02$ ) compared to Cluster 2 ( $n = 7$ ). Cluster 1 was associated with more cognitive improvement, significantly differing from Cluster 2 in changes in attention scores over time. Cluster 1 significantly improved scores on the Attention Scale of the CAS compared to Cluster 2 ( $t(16) = 2.6, p = .02$ ). No differences were observed between the two clusters in BRIEF scores. While the two cluster analyses (Cluster Analysis 1 and Cluster Analysis 2) did identify similar groups of participants as indicated by a chi-square test ( $\chi^2(1, N = 18) = 5.1, p = .02$ ), they did not completely overlap with one another, with a total of four participants having different cluster membership between the two analyses. For further details, see Figure 2.

## Discussion

The current study hypothesized that SLF WMI would be improved by an exercise intervention in overweight children, particularly with greater attendance. This hypothesis was based on evidence that fronto-parietal regions and the cognitive control processes that they support are altered by exercise. While no group by time interaction for SLF was detected, a group by time by attendance interaction indicated that improved WMI in the SLF of overweight children was associated with attendance at an after-school exercise program as compared to attendance at an after-school sedentary program. Specifically, higher attendance in the exercise condition was associated with increased WMI in bilateral SLF (increased FA and decreased RD) compared to the control group, where attendance was not associated with change in WMI. A dose-response gradient was observed not only with attendance but also with aerobic intensity (heart rate) and an index of total energy expenditure (heart rate  $\times$  attendance) in the exercise group, such that better attendance, higher intensity, and greater total dose of exercise were linked with greater improvements in

SLF WMI. In addition, attendance at the exercise intervention as compared to attendance at the control group was associated with more improved teacher ratings of cognitive control as measured by the BRIEF. Finally, these results indicated that in the overall study sample, increasing WMI in the SLF benefited cognitive control. Improved WMI in the right SLF was associated with improved selective attention, while improved WMI in the left SLF was associated with better teacher ratings of classroom behavior.

Improved WMI across both groups was associated with better teacher ratings of executive function behaviors in the classroom as assessed by the BRIEF. WMI of the left SLF was associated with a global assessment of executive function and with two component scales. Behavioral Regulation includes items such as children's ability to control impulses and move freely from one situation to the next. Metacognition includes items such as children's ability to initiate activities, hold information in working memory, plan and organize future events, and monitor their own performance. The association between increased WMI and improved teacher ratings indicates that the SLF is important for cognitive control abilities in this sample, as shown in previous studies [22; 23]. It also indicates that WMI is reflected in classroom conduct and that it therefore may have implications for academic achievement and social behavior.

The other measure which was associated with WMI across both groups was the Attention Scale of the CAS, a composite measure based on three selective attention tasks: a variation of the Stroop test and two paper-and-pencil tests requiring children to find visual targets within fields of distracters. There is previous evidence to suggest that the SLF is important in attention. FA in the SLF has been associated with similar measures in children, such as the sensitivity index of a rapid visual information processing paradigm, which describes a subject's ability to detect signals independent of answering strategy or bias [34]. SLF FA also was related to visual search in adults (i.e., finding targets among distracters [16]).

Higher FA is often interpreted as reflecting coherently bundled, myelinated fibers but it is somewhat non-specific in that it can be affected by various tissue characteristics, including myelination, axon diameter, and fiber organization [35]. It is possible that decreasing RD in the current study could reflect increasing myelination more so than other alterations in axonal structure, for which AD might be a more sensitive measure [17; 36]. Therefore, the changes in WMI described in the current study may be more strongly related to myelination. This interpretation is made with caution, as the association of white matter measures with specific tissue characteristics remains a topic of debate [37]. Nevertheless, the current study indicates that FA and RD are affected by exercise rather than AD, which is supported by a previous cross-sectional study in adults showing that higher FA associated with fitness was primarily related to less RD, rather than altered AD [18]. While this study is one of the first to investigate how exercise affects white matter structure in children, it is limited by a small sample size. This limited power to detect group by time effects and may be susceptible to sample-specific results.

The current study suggests that children's attendance at an after-school exercise program, as compared to a sedentary program, improves WMI in a tract supporting cognitive control processes, as well as teacher ratings of children's cognitive control behaviors in the



classroom. This sample is unique among neuroimaging investigations of exercise in children due to the stringent nature of the control group. Because both groups attended after-school programs, several potentially beneficial effects were controlled for (e.g., social interaction, attention from adults, and supervised homework time). Additional strengths of this study include the 8-month length of the interventions and the predominantly minority sample of children. Across both groups, improved WMI was related to two different measures of cognitive control – a measure of selective attention and teacher ratings of classroom behavior. Increased integrity of the SLF with attendance at an exercise intervention may contribute to improved cognitive control, with potential benefits not only for performance on a standardized cognitive task but also for on-task behavior and interpersonal interactions in the classroom. This may be important to protect school resources for physical activity in the context of unprecedented levels of childhood obesity along with pressure on schools to improve achievement [38; 39].

## Acknowledgments


CK, DS, CD, and JM conceived and carried out experiments and analyzed data. JA, NY, and TL conceived and carried out experiments. NS, LC, AW, JP, and AR carried out experiments. All authors were involved in writing the paper and had final approval of the submitted and published versions. This research was supported by the National Institutes of Health R01 HL87923 and the National Science Foundation Graduate Research Fellowship Program.

## References

1. Davis CL, Cooper S. Fitness, fatness, cognition, behavior, and academic achievement among overweight children: Do cross-sectional associations correspond to exercise trial outcomes? *Prev Med.* 2011; 52:S65–S69. [PubMed: 21281668]
2. Voss MW, Chaddock L, Kim JS, VanPatter M, Pontifex MB, Raine LB, Cohen NJ, Hillman CH, Kramer AF. Aerobic fitness is associated with greater efficiency of the network underlying cognitive control in preadolescent children. *Neuroscience.* 2011; 199:166–176. [PubMed: 22027235]
3. Weinstein AM, Voss MW, Prakash RS, Chaddock L, Szabo A, White SM, Wojcicki TR, Mailey E, McAuley E, Kramer AF, Erickson KI. The association between aerobic fitness and executive function is mediated by prefrontal cortex volume. *Brain Behav Immun.* 2012; 26:811–819. [PubMed: 22172477]
4. Wu CT, Pontifex MB, Raine LB, Chaddock L, Voss MW, Kramer AF, Hillman CH. Aerobic fitness and response variability in preadolescent children performing a cognitive control task. *Neuropsychology.* 2011; 25:333–341. [PubMed: 21443340]
5. Chaddock L, Erickson KI, Prakash RS, VanPatter M, Voss MW, Pontifex MB, Raine LB, Hillman CH, Kramer AF. Basal ganglia volume is associated with aerobic fitness in preadolescent children. *Dev Neurosci.* 2010; 32:249–256. [PubMed: 20693803]
6. Davis CL, Tomporowski PD, McDowell JE, Austin BP, Miller PH, Yanasak NE, Allison JD, Naglieri JA. Exercise improves executive function and achievement and alters brain activation in overweight children: a randomized controlled trial. *Health Psychol.* 2011; 30:91–98. [PubMed: 21299297]
7. Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ, Webb A, Jerome GJ, Marquez DX, Elavsky S. Cardiovascular fitness, cortical plasticity, and aging. *Proc Natl Acad Sci U S A.* 2004; 101:3316–3321. [PubMed: 14978288]
8. Kamijo K, Pontifex MB, O'Leary KC, Scudder MR, Wu CT, Castelli DM, Hillman CH. The effects of an afterschool physical activity program on working memory in preadolescent children. *Dev Sci.* 2011; 14:1046–1058. [PubMed: 21884320]


9. Evers A, Klusmann V, Schwarzer R, Heuser I. Improving cognition by adherence to physical or mental exercise: A moderated mediation analysis. *Aging Ment Health*. 2011; 15:446–455. [PubMed: 21500011]
10. Colcombe SJ, Kramer AF. Fitness effects on the cognitive function of older adults: A meta-analytic study. *Psychol Sci*. 2003; 14:125–130. [PubMed: 12661673]
11. Luna B, Thulborn KR, Munoz DP, Merriam EP, Garver KE, Minshew NJ, Keshavan MS, Genovese CR, Eddy WF, Sweeney JA. Maturation of widely distributed brain function subserves cognitive development. *Neuroimage*. 2001; 13:786–793. [PubMed: 11304075]
12. Schaeffer DJ, Amlung MT, Li Q, Krafft CE, Austin BP, Dyckman KA, McDowell JE. Neural correlates of behavioral variation in healthy adults' antisaccade performance. *Psychophysiology*. 2013; 50:325–333. [PubMed: 23418930]
13. Krafft CE, Schwarz NF, Chi L, Weinberger AL, Schaeffer DJ, Pierce JE, Rodrigue AL, Yanasak NE, Miller PH, Tomporowski PD, Davis CL, McDowell JE. An eight month randomized controlled exercise trial alters brain activation during cognitive tasks in overweight children. *Obesity*. In press.
14. Gordon EM, Lee PS, Maisog JM, Foss-Feig J, Billington ME, VanMeter J, Vaidya CJ. Strength of default mode resting state connectivity relates to white matter integrity in children. *Dev Sci*. 2011; 14:738–751. [PubMed: 21676094]
15. Beaulieu C. The basis of anisotropic water diffusion in the nervous system – a technical review. *NMR in Biomedicine*. 2002; 15:435–455. [PubMed: 12489094]
16. Bennett IJ, Motes MA, Rao NK, Rypma B. White matter tract integrity predicts visual search performance in young and older adults. *Neurobiol Aging*. 2012; 33:433.e421–433.e431. [PubMed: 21402431]
17. Song SK, Sun SW, Ramsbottom MJ, Chang C, Russell J, Cross AH. Demyelination revealed through MRI as increased radial (but unchanged axial) diffusion of water. *Neuroimage*. 2002; 17:1429–1436. [PubMed: 12414282]
18. Johnson NF, Kim C, Clasey JL, Bailey A, Gold BT. Cardiorespiratory fitness is positively correlated with cerebral white matter integrity in healthy seniors. *Neuroimage*. 2012; 59:1514–1523. [PubMed: 21875674]
19. Marks B, Katz L, Styner M, Smith J. Aerobic fitness and obesity: relationship to cerebral white matter integrity in the brain of active and sedentary older adults. *Br J Sports Med*. 2011; 45:1208–1215. [PubMed: 20558529]
20. Marks B, Madden DJ, Bucur B, Provenzale JM, White LE, Cabeza R, Huettel SA. Role of aerobic fitness and aging on cerebral white matter integrity. *Ann N Y Acad Sci*. 2007; 1097:171–174. [PubMed: 17413020]
21. Voss MW, Heo S, Prakash RS, Erickson KI, Alves H, Chaddock L, Szabo AN, Mailey EL, Wójcicki TR, White SM, Gothe N, McAuley E, Sutton BP, Kramer AF. The influence of aerobic fitness on cerebral white matter integrity and cognitive function in older adults: Results of a one-year exercise intervention. *Hum Brain Mapp*. In press.
22. Østby Y, Tamnes CK, Fjell AM, Walhovd KB. Morphometry and connectivity of the fronto-parietal verbal working memory network in development. *Neuropsychologia*. 2011; 49:3854–3862. [PubMed: 22001853]
23. Vestergaard M, Madsen KS, Baaré WF, Skimminge A, Ejersbo LR, Ramsøy TZ, Gerlach C, Akeson P, Paulson OB, Jernigan TL. White matter microstructure in superior longitudinal fasciculus associated with spatial working memory performance in children. *J Cogn Neurosci*. 2010; 23:2135–2146. [PubMed: 20964591]
24. Lebel C, Walker L, Leemans A, Phillips L, Beaulieu C. Microstructural maturation of the human brain from childhood to adulthood. *Neuroimage*. 2008; 40:1044–1055. [PubMed: 18295509]
25. Gebauer D, Fink A, Filippini N, Johansen-Berg H, Reishofer G, Koschutnig K, Kargl R, Purgstaller C, Fazekas F, Enzinger C. Differences in integrity of white matter and changes with training in spelling impaired children: a diffusion tensor imaging study. *Brain Struct Funct*. 2012; 217:747–760. [PubMed: 22198594]
26. Ogden CL, Kuczmarski RJ, Flegal KM, Mei Z, Guo S, Wei R, Grummer-Strawn LM, Curtin LR, Roche AF, Johnson CL. Centers for Disease Control and Prevention 2000 growth charts for the

- United States: Improvements to the 1977 National Center for Health Statistics version. *Pediatrics*. 2002; 109:45–60. [PubMed: 11773541]
27. Naglieri, JA.; Das, JP. *Cognitive Assessment System: interpretive handbook*. Riverside Publishing; 1997.
  28. Gioia G, Isquith PK, Guy SC, Kenworthy L. Behavior rating inventory of executive function. *Child Neuropsychol*. 2000; 6:235–238. [PubMed: 11419452]
  29. Rorden C. DCM2NII (Version October 7). 2007
  30. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM. Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*. 2004; 23:S208–S219. [PubMed: 15501092]
  31. Smith SM. Fast robust automated brain extraction. *Hum Brain Mapp*. 2002; 17:143–155. [PubMed: 12391568]
  32. Wakana S, Caprihan A, Panzenboeck MM, Fallon JH, Perry M, Gollub RL, Hua K, Zhang J, Jiang H, Dubey P, Blitz A, van Zijl P, Mori S. Reproducibility of quantitative tractography methods applied to cerebral white matter. *Neuroimage*. 2007; 36:630–644.
  33. Leemans, A.; Jeurissen, B.; Sijbers, J.; Jones, DK. ExploreDTI: a graphical toolbox for processing, analyzing, and visualizing diffusion MR data; Presented at 17th Annual Meeting of the International Society for Magnetic Resonance in Medicine; Hawaii, U.S.A. 2009;
  34. Klarborg B, Madsen K, Vestergaard M, Skimminge A, Jernigan TL, Baaré WF. Sustained attention is associated with right superior longitudinal fasciculus and superior parietal white matter microstructure in children. *Hum Brain Mapp*. In press.
  35. Scholz, J.; Tomassini, V.; Johansen-Berg, H. Individual differences in white matter microstructure in the healthy brain. In: Johansen-Berg, H.; Behrens, TE., editors. *Diffusion MRI: From Quantitative Measurement to In vivo Neuroanatomy*. Academic Press; 2009. p. 105-126.
  36. Beaulieu, C. The biological basis of diffusion anisotropy. In: Johansen-Berg, H.; Behrens, TE., editors. *Diffusion MRI: From Quantitative Measurement to In vivo Neuroanatomy*. Academic Press; 2009. p. 105-126.
  37. Wheeler-Kingshott CAM, Cercignani M. About “axial” and “radial” diffusivities. *Magn Res Med*. 2009; 61:1255–1260.
  38. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999-2010. *JAMA*. 2012; 307:483–490. [PubMed: 22253364]
  39. McMurrer, J. NCLB Year 5: Instructional time in elementary schools: A closer look at changes for specific subjects. 2008. Retrieved from Center on Education Policy Website: [http://www.cep-dc.org/cfcontent\\_file.cfm?Attachment=McMurrer%5FReport%5FInstructionalTime%5F022008%2Epdf](http://www.cep-dc.org/cfcontent_file.cfm?Attachment=McMurrer%5FReport%5FInstructionalTime%5F022008%2Epdf)



**Figure 1.**

Time by group by attendance interactions. Y-axes show white matter difference scores (changes in integrity from baseline to post-test) and x-axes show attendance. Each measure shown is a residual (controlled for age and gender) which has been back-transformed to its original metric by adding its mean.



**Figure 2.**

Illustrations of *K*-means cluster analyses. The scatterplots illustrate the two clusters identified for each analysis based on the participants' alterations in white matter integrity (FA and RD) over time. These show that for both analyses, Cluster 1 had improved white matter integrity (increased FA and decreased RD) compared to Cluster 2. The bar graphs illustrate the differences in cognition between the two clusters obtained in each analysis. In Cluster Analysis 1, Cluster 2 showed worse teacher ratings of executive function as compared to Cluster 1, indicated by increased BRIEF scores. In Cluster Analysis 2, Cluster 1 showed improved attention scores as compared to Cluster 2, indicated by increased CAS Attention Scale scores. The two clusters obtained in each *k*-means analysis significantly differed in all measures shown. SLF = superior longitudinal fasciculus. FA = fractional anisotropy. RD = radial diffusivity. BRIEF = Behavioral Rating Inventory of Executive Function. CAS = Cognitive Assessment System. Measures shown are difference scores (change from baseline to post-test).

**Table 1**  
**Baseline Characteristics and Attendance of Participants Included in Analysis**

Characteristic	Exercise group	Control group	P value
<i>N</i>	10	8	
Age (years)	9.9 (0.6)	9.4 (0.8)	.17 <sup>a</sup>
Female	50%	50%	1 <sup>b</sup>
Non-white	100%	88%	.25 <sup>b</sup>
Left-handed	10%	25%	.40 <sup>b</sup>
Parental education scale	4.8 (1.0)	4.3 (1.5)	.37 <sup>a</sup>
CAS Full Scale score	94.2 (8.9)	95.6 (9.5)	.75 <sup>a</sup>
CAS Planning Scale score	88.2 (11.7)	89.5 (13.9)	.83 <sup>a</sup>
CAS Attention Scale score	98.4 (8.5)	94.8 (7.5)	.35 <sup>a</sup>
BRIEF Global Executive Composite score	56.0 (11.8)	47.9 (9.8)	.16 <sup>a</sup>
BRIEF Behavioral Regulation Index score	54.3 (11.4)	45.6 (3.7)	.07 <sup>a</sup>
BRIEF Metacognition Index score	55.9 (12.0)	49.1 (12.4)	.27 <sup>a</sup>
Attendance	60% (10%)	72% (8%)	.36 <sup>a</sup>

*Note.* Mean and SD or %. CAS = Cognitive Assessment System. BRIEF = Behavioral Rating Inventory of Executive Function.

<sup>a</sup> result of t-test

<sup>b</sup> result of  $\chi^2$  test