



CrossMark
click for updates

Research

Cite this article: Lüke L, Campbell P, Varea Sánchez M, Nachman MW, Roldan ERS. 2014 Sexual selection on protamine and transition nuclear protein expression in mouse species. *Proc. R. Soc. B* **281**: 20133359. <http://dx.doi.org/10.1098/rspb.2013.3359>

Received: 23 December 2013

Accepted: 27 February 2014

Subject Areas:

evolution

Keywords:

sexual selection, sperm competition, protamine, transition nuclear protein, sperm

Author for correspondence:

Eduardo R. S. Roldan

e-mail: roldane@mncn.csic.es

[†]Present address: Department of Zoology, Oklahoma State University, Stillwater, OK 74078, USA.

[‡]Present address: Museum of Vertebrate Zoology and Department of Integrative Biology, University of California, Berkeley, CA 94720, USA.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2013.3359> or via <http://rspb.royalsocietypublishing.org>.

Sexual selection on protamine and transition nuclear protein expression in mouse species

Lena Lüke^{1,2}, Polly Campbell^{2,†}, María Varea Sánchez¹, Michael W. Nachman^{2,‡} and Eduardo R. S. Roldan¹

¹Reproductive Ecology and Biology Group, Museo Nacional de Ciencias Naturales (CSIC), Madrid 28006, Spain

²Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721, USA

Post-copulatory sexual selection in the form of sperm competition is known to influence the evolution of male reproductive proteins in mammals. The relationship between sperm competition and regulatory evolution, however, remains to be explored. Protamines and transition nuclear proteins are involved in the condensation of sperm chromatin and are expected to affect the shape of the sperm head. A hydrodynamically efficient head allows for fast swimming velocity and, therefore, more competitive sperm. Previous comparative studies in rodents have documented a significant association between the level of sperm competition (as measured by relative testes mass) and DNA sequence evolution in both the coding and promoter sequences of protamine 2. Here, we investigate the influence of sexual selection on protamine and transition nuclear protein mRNA expression in the testes of eight mouse species that differ widely in levels of sperm competition. We also examined the relationship between relative gene expression levels and sperm head shape, assessed using geometric morphometrics. We found that species with higher levels of sperm competition express less protamine 2 in relation to protamine 1 and transition nuclear proteins. Moreover, there was a significant association between relative protamine 2 expression and sperm head shape. Reduction in the relative abundance of protamine 2 may increase the competitive ability of sperm in mice, possibly by affecting sperm head shape. Changes in gene regulatory sequences thus seem to be the basis of the evolutionary response to sexual selection in these proteins.

1. Introduction

When females mate promiscuously, the sperm of rival males compete for the fertilization of available ova [1]. Post-copulatory sexual selection mediated by sperm competition has a profound influence on male reproductive traits across a wide range of taxa (reviewed in [2–4]). In mammals, key traits affected by sperm competition include sperm quality parameters [5], processes that prepare sperm to interact with the oocyte [6], sperm design (e.g. overall size, head shape and dimensions [7–10]) and sperm swimming velocity [10–12]. Several lines of evidence suggest that sperm head shape is particularly important in competitive situations. For example, the size and curvature of the apical hook of rodent sperm heads is thought to be associated with levels of sperm competition ([9], but see [13]). Likewise, head shape may affect the hydrodynamic efficiency of spermatozoa. Head elongation, which may reduce drag, associates with faster sperm swimming velocity [10]. Faster sperm are more likely to succeed in fertilization [14].

To date, most work on the molecular evolution of male reproductive genes has focused on protein-coding regions [15,16]. A number of studies have found a positive relationship between sequence divergence of these genes and levels of sperm competition, and several such genes show evidence of positive selection in coding regions ([17–21], but see [22,23]). However, a positive correlation between sequence divergence in the promoter region of protamine 2 and

relative levels of sperm competition in house mice and their close relatives [24] suggests that regulatory changes may also contribute to species differences in sperm competitive ability. Surprisingly, despite order of magnitude differences in the absolute and relative expression levels of protamines and associated transition nuclear proteins across eutherian mammals [25], the relationship between sperm competition and gene expression remains largely unexplored.

Protamines and transition nuclear proteins are integral to chromatin remodelling and condensation during the final stages of spermatogenesis. This nuclear reshaping in postmeiotic spermatids affects the overall shape of the sperm head which, in turn, may influence hydrodynamic efficiency, resulting in an increase in sperm swimming speed and more competitive sperm. Notably, sperm from transition nuclear protein-deficient mice perform poorly in some competitive assays [26]. Whereas protamines (PRM1 and PRM2 in most eutherian mammals) bind directly to DNA in the nucleus of elongating spermatids and mature spermatozoa [27], transition nuclear proteins (TNP1 and TNP2) are involved in intermediate stages in the replacement of histones by protamines [28,29]. Protamines remain associated with sperm chromatin in the oocyte and influence the rate of nuclear decondensation, a trait associated with embryonic survival [30–33].

Protamine and transition protein mRNAs are highly co-expressed in round spermatids [34–37], and the protein products of both gene families exhibit significant overlap in elongating spermatid nuclei [28,38]. TNP1 and TNP2 seem to perform partially redundant functions: only double TNP1/TNP2 mouse knockouts are completely sterile [28]. However, deletion of either transition protein results in incomplete PRM2 processing and defective chromatin condensation [29,39]. This, together with the co-localization of mRNAs and mature proteins, strongly suggests that functional interactions between protamines and transition proteins are necessary for normal sperm development.

In mice and humans, both PRM1 and PRM2 are essential for male fertility [40]. Strikingly, although the relative abundance of PRM1 and PRM2 proteins differs widely across mammals (from 0 to 77% PRM2) [41], disruption of species-specific protamine ratios causes fertility defects comparable with gene knockouts [40,42]. In human males, for example, protamine imbalance can result in reduced sperm concentration and motility, and in abnormal head morphology, an indicator of deficits in chromatin condensation [43–45]. In particular, incomplete processing of the PRM2 precursor is associated with sperm dysfunction [45,46], and PRM2-deficient sperm are characterized by incomplete nuclear condensation and increased DNA damage [40,46,47], defects that can lead to embryonic mortality [31]. Thus, protamine ratios play a large role in sperm head morphology, a phenotype important for competitive ability both before and during fertilization. This suggests that sexual selection mediated by sperm competition should act on protamine ratios, resulting in an association between species differences in levels of sperm competition and protamine expression.

Here, we investigate the influence of sexual selection on protamine and transition nuclear protein mRNA expression in the testes of eight closely related species in the genus *Mus*. These species exhibit a wide range of relative testes mass, a robust proxy for different levels of sperm competition [2,4], and differ in sperm traits associated with competitive ability [5,7,24,48]. Moreover, evolution of the *Prm2* promoter in seven of the same species is consistent with stronger

selection in taxa with higher inferred levels of sperm competition [24]. This provides specific motivation for studying the relationship between protamine expression and sperm competition in *Mus*. Given the functional relationship between protamines and transition proteins, and the role of transition proteins in PRM2 processing, we expected that transition nuclear protein expression should covary with species differences in protamine expression. Because protamines and transition nuclear proteins are involved in the condensation of sperm chromatin and are expected to affect the shape of the sperm head, we also assessed the relationship between gene expression and sperm head shape.

2. Material and methods

(a) Species

This study included eight species in the genus *Mus*: *M. caroli*, *M. castaneus*, *M. domesticus*, *M. macedonicus*, *M. musculus*, *M. pahari*, *M. spicilegus* and *M. spretus* (four to five males per species). This group of species shows diverse levels of sperm competition, as inferred from their differences in relative testes mass (table 1). Large testes in relation to body mass (relative testes mass) is a strong predictor of high sperm competition levels in many taxa (reviewed in [2,4,50]), and relative testes mass is correlated with genetic paternity (i.e. percentages of multiple paternity) in mammals in general [51], and rodents in particular [52]. Therefore, relative testes mass is used in this study as a robust proxy for sperm competition levels.

Individuals were purchased from the Institut des Sciences de l'Evolution-Montpellier, CNRS-Universite de Montpellier II. Males were kept in our animal facilities in individual cages under standard laboratory conditions in environmentally controlled rooms (20–24°C) on a 14 L : 10 D photoperiod and were provided with food and water ad libitum. All animal handling was done following Spanish Animal Protection Regulation RD1201/2005, which conforms to European Union Regulation 2003/65.

(b) Testes collection and relative testes mass

Animals were sacrificed at an age of two to four months by cervical dislocation and were immediately weighed and dissected. Testes were removed, weighed, flash-frozen in liquid nitrogen and stored at –80°C. All dissection instruments and areas were cleaned with RNase AWAY (Molecular BioProducts, Thermo Fisher Scientific, San Diego, CA, USA) before use. Relative testes mass was calculated based on the rodent power function, following the method in Kenagy & Trombulak [49].

(c) RNA extraction and cDNA synthesis

RNA was extracted in a sterile vertical laminar flow hood using either the RNeasy Plus kit (Qiagen) or the E.Z.N.A Total RNA kit I (Omega, Madrid, Spain) following the manufacturer's recommendations. All instruments and surface areas were cleaned with RNase AWAY. RNA concentration and purity were determined using a NanoDrop 1000 spectrophotometer (Thermo Scientific, Madrid, Spain), and cDNA was synthesized the same day from 10 µg of RNA, using the Superscript III First Strand Synthesis Kit with oligo(dT) (Invitrogen, Barcelona, Spain) according to the manufacturer's recommendations. cDNA concentration and purity were determined using a NanoDrop1000 spectrophotometer, and samples were stored at –20°C.

(d) Quantitative PCR

Expression levels for *M. musculus*, *M. spretus*, *M. spicilegus* and *M. pahari* were determined at the University of Arizona in Tucson using a MyiQ2 light cycler (Bio-Rad), and expression

Table 1. Relative testes mass was calculated as described by Kenagy & Trombulak [49] followed by a calculation of the median for the species. Gene expression data are normalized, transformed median values. Species were ordered by relative testes mass (ascending).

species	relative testes mass	<i>Prm1</i> (ΔC_T)	<i>Prm2</i> (ΔC_T)	<i>Tnp1</i> (ΔC_T)	<i>Tnp2</i> (ΔC_T)	<i>Tnp1/Tnp2</i>	<i>Prm1/Prm2</i>	<i>Prm/Tnp</i>	<i>Prm2/Tnp</i>	<i>Prm2/Prm</i>	<i>Prm2/(Prm + Tnp)</i>
<i>Mus castaneus</i>	0.27	3.16	4.29	2.96	3.38	0.83	0.67	1.24	0.75	0.60	0.33
<i>Mus pahari</i>	0.27	3.80	3.69	3.01	2.26	1.13	1.01	1.38	0.68	0.50	0.29
<i>Mus domesticus</i>	0.32	2.11	3.22	1.88	2.32	0.81	0.65	1.27	0.77	0.61	0.34
<i>Mus musculus</i>	0.44	2.87	3.72	3.27	2.98	1.14	0.76	1.06	0.61	0.57	0.29
<i>Mus caroli</i>	0.46	5.83	7.28	6.71	6.18	1.07	0.78	1.00	0.56	0.56	0.28
<i>Mus spretus</i>	0.87	1.90	2.31	2.32	1.70	1.38	0.82	1.05	0.58	0.55	0.28
<i>Mus macedonicus</i>	0.95	3.54	3.49	2.48	2.14	1.05	0.99	1.49	0.74	0.50	0.30
<i>Mus spicilegus</i>	1.51	4.61	4.60	4.76	3.92	1.16	0.98	1.05	0.53	0.50	0.26
CV	0.69	0.37	0.36	0.46	0.46	0.18	0.17	0.15	0.14	0.08	0.09

levels for *M. domesticus*, *M. castaneus*, *M. macedonicus* and *M. caroli* were determined at the Museo Nacional de Ciencias Naturales in Madrid using a CFX96 Real Time System/C1000 Thermal Cycler (Bio-Rad). To check the consistency of results obtained using different cyclers, assays for the standard gene (see below) were run by the same person (L.L.) with a set of testes samples taken from the same individuals used in both Tucson and Madrid, using exactly the same protocol. Results were consistent across locations (e.g. *M. musculus* individual 1 (Tucson, right testis): average C_T (\pm s.d.) = 12.94 (0.02); *M. musculus* individual 1 (Madrid, left testis): average C_T (\pm s.d.) = 12.89 (0.07)).

Primers were designed in PRIMER3 (v. 0.4.0) to amplify a product between 70 and 150 bases across an exon–exon junction. Protamine primers were placed in sequences that are invariant across all species in this analysis. Transition protein primers were placed in sequences that are conserved between *Mus* and *Rattus*, and therefore are unlikely to vary among closely related *Mus* species. Primer sequences and amplicon sizes are provided in the electronic supplementary material, table S1. Each quantitative PCR (qPCR) run included one individual of each species with three technical replicates for the four experimental genes (*Prm1*, *Prm2*, *Tnp1* and *Tnp2*) and two technical replicates for the standard gene (*18SrRNA*). qPCR reactions were run in 96-well plates with an end volume of 16 μ l per sample containing 8 μ l SYBR green Master Mix (Invitrogen), 15 ng of each primer and 50 ng μ l⁻¹ of cDNA. The conditions of the thermocycler program consisted of an initial denaturation of 95°C for 10 min, 40 cycles of 95°C for 15 s and an annealing and elongation stage of 62°C for 1 min. Melt curve analysis was performed at the end of each run to check for multiple peaks, indicative of non-specific amplification.

(e) Analysis of expression data

Cycle threshold data (C_T) were normalized relative to *18SrRNA* for each plate (ΔC_T). To avoid statistical analysis using a dataset of mixed negative and positive values, data were transformed by adding a constant based on the lowest ΔC_T value. Expression ratios and percentages were calculated from transformed individual ΔC_T values (*M. domesticus* $n = 4$, all other species $n = 5$), and median values were obtained for each species. Because of the expectation that relative expression levels may be of greater functional significance than absolute expression levels (see above), we calculated ratios (*Prm1/Prm2*, *Tnp1/Tnp2*, *Prm/Tnp*, *Prm2/Tnp*) and proportions (*Prm2/Prm*, *Prm2/(Prm + Tnp)*, *Prm1/(Prm + Tnp)*), where *Prm* refers to the combined expression of *Prm1* and *Prm2*, and *Tnp* refers to the combined expression

of *Tnp1* and *Tnp2*. To obtain a measure of variability between individuals and species, as well as for individual genes, the coefficient of variation (CV = s.d./mean) was calculated.

(f) Phylogenetic generalized least-squares analysis

Species data may not be free of phylogenetic association because shared character values may result from common ancestry rather than independent evolution, and thus may not be truly independent. To control for this phylogenetic inertia, we used phylogenetic generalized least-squares (PGLS) analyses [53] to test for relationships between species differences in total and relative protamine and transition protein expression, and relative testes mass. PGLS analysis was implemented in COMPARE 4.6b [54], using a phylogenetic tree based on Lundrigan *et al.* [55] and Gómez Montoto *et al.* [5] (electronic supplementary material, figure S1).

(g) Geometric morphometrics analysis of sperm head shape

Geometric morphometrics methods were used to quantify head shape variation based on a set of landmarks that correspond to the spatial position of particular anatomical traits [56,57]. A total of 20 bidimensional landmark coordinates were gathered from spermatozoa of seven of the eight species used in the gene expression analysis (*M. caroli*, *M. castaneus*, *M. domesticus*, *M. macedonicus*, *M. musculus*, *M. spicilegus*, *M. spretus*; $n = 5$ males/species). Landmark data were processed as described previously [58]. All morphometric analyses were conducted with MORPHOJ [59]. An independent contrast for morphometric shape data [60] was conducted to check for phylogenetic signal in the sperm head shape dataset. This test simulates the null hypothesis of total absence of phylogenetic signal by a permutation procedure. The p -value was not significant ($p = 0.102$) for the null hypothesis of independence, which indicates a lack of phylogenetic signal and, therefore, that phylogenetic correction was not needed for this analysis.

Canonical variate analysis (CVA) [61] was used to explore the relationship between sperm head shape and relative protamine expression. Species were grouped into three categories based on well-defined differences in relative protamine expression: low, intermediate and high expression ratios (table 1; electronic supplementary material, table S2; see Results section for details). The CVA produces a set of canonical variates that are uncorrelated within and among groups and account for the maximum amount of among-group variance relative to within-group

variance. As a result of the CVA, distances in the original space are transformed to Procrustes distances. These Procrustes distances for between-category comparisons were used to test for significant differences in sperm head shapes between species with low, intermediate and high protamine expression ratios.

3. Results

(a) Expression of protamines and transition nuclear proteins

Median expression levels for each gene and species are shown in table 1. The ranges of expression medians and the CV for each gene and species are provided in the electronic supplementary material, table S2. Within species, expression levels were positively correlated in all pairwise comparisons among genes (electronic supplementary material, figure S2 and table S3), suggesting that there may be functional constraints to maintain consistent relative expression levels of these genes and/or common regulatory control. The median expression level for individual genes varied by a factor of approximately threefold among species (electronic supplementary material, table S2). *Tnp1* was expressed at a slightly higher level than *Tnp2* although both showed the same CV. Likewise, *Prm2* was expressed at a slightly higher level than *Prm1* but there was no difference in CV (table 1; electronic supplementary material, table S2).

The ratios and proportions of expression levels for different genes are shown in the electronic supplementary material, table S4. These relative levels of expression were much more constant among species (electronic supplementary material, table S4) than expression levels of individual genes (cf. electronic supplementary material, table S2). The ratio of total protamines to total transition nuclear proteins was close to one in half the species and above one in the other four species, revealing higher overall expression levels of protamines. Ratios between *Tnp1* and *Tnp2* were generally above one, in agreement with higher expression levels of *Tnp1* in comparison to *Tnp2* (see above). The reverse was true for protamines, with ratios of *Prm1/Prm2* below one (electronic supplementary material, table S4).

(b) Relationships between relative testes mass and gene expression

We tested for associations between relative testes mass and patterns of protamine and transition protein expression, both for individual genes and for ratios of expression levels among genes.

The correlation between relative testes mass and *Prm1/Prm2* or *Prm2/Prm* was not significant when all eight species were considered (figure 1a and table 2). However, we noted that *M. pahari* appears to be an outlier in this analysis. *Mus pahari* is basal to the other species included in this study and belongs to a different subgenus (*Coelomys*) [62]. When the analysis was restricted to the seven species in the subgenus *Mus*, there was a significant positive relationship between relative testes mass and *Prm1/Prm2* ($\alpha = 15.5$, CI 95% (slope) = 1.67–4.25, correlation = 0.89; figure 1a and table 2) and a significant negative relationship between relative testes mass and *Prm2/Prm* ($\alpha = 15.5$, CI 95% (slope) = -0.14 to -0.06, correlation = 0.80; table 2). By contrast, there was no




Figure 1. Relationships between relative testes mass and relative protamine 2 expression. (a) Protamine ratio (*Prm1/Prm2*): the dashed line corresponds to analyses with $N = 8$ mouse species, and the correlation is not significant. The open circle identifies *M. pahari*, a species that behaves as an outlier in these analyses. The solid line corresponds to analyses with $n = 7$ species in which *M. pahari* is not included, and this correlation is statistically significant. (b) Ratio of *Prm2* to total *Tnp* (*Prm2/Tnp*). Results of statistical analyses are given in table 2.

relationship between testes mass and transition protein ratios (data not shown).

Significant negative associations with relative testes mass were found for *Prm2/Tnp* ($\alpha = 1.56$, CI 95% (slope) = -4.64 to -0.03, correlation = -0.63; figure 1a and table 2) and *Prm2/(Prm + Tnp)* ($\alpha = 6.05$, CI 95% (slope) = -0.19 to -0.02, correlation = -0.72; table 2). By contrast, there was no association between relative testes mass and *Prm1/Tnp*, *Prm1/(Prm + Tnp)* or *Prm/Tnp* (data not shown), or between relative testes mass and any of the four genes when analysed separately (electronic supplementary material, table S5). Thus, significant relationships between testes mass and the expression of sperm condensation proteins are driven mainly by the relative expression of *Prm2*.

Together, these results indicate that species with higher inferred levels of sperm competition express proportionately less *Prm2* in relation to total transition protein and in relation to total protamine and transition protein combined. Within the subgenus *Mus*, species with higher sperm competition have a higher *Prm1/Prm2* expression ratio and therefore a lower *Prm2/Prm* proportion.

(c) Relationships between protamine expression and sperm head shape

Geometric morphometrics was employed to quantify differences in head shape between the seven species in the subgenus *Mus*.

Table 2. Relationship between relative testes mass and relative protamine or transition protein expression. Analyses were carried out with all species and excluding *Mus pahari* (see text). CI[−] and CI⁺ indicate the confidence intervals for the regression slope, lnL = log likelihood estimate of alpha, alpha = measure of evolutionary constraints acting on phenotypes, corr = the correlation value (*r*). Bold CI values indicate statistical significance.

	excluding <i>Mus pahari</i> (<i>n</i> = 7)		relationships for all species (<i>n</i> = 8)			
	<i>Prm1/Prm2</i>	<i>Prm2/Prm</i>	<i>Prm2/Tnp</i>	<i>Prm2/(Prm + Tnp)</i>	<i>Prm1/Prm2</i>	<i>Prm2/Prm</i>
CI [−]	1.67	− 0.14	− 4.64	− 0.19	−0.47	−0.12
CI ⁺	4.25	− 0.06	− 0.03	− 0.02	3.68	0.01
lnL	8.15	8.14	5.20	6.05	4.52	4.63
alpha	15.50	15.50	1.56	1.66	5.62	5.37
corr	0.89	−0.80	−0.63	−0.72	0.53	−0.54

Species were categorized as having high, intermediate or low protamine expression ratios, and Procrustes distances (*D*) calculated from CVA were used to test for between-category differences in sperm head shape.

Sperm head shapes were significantly different between species with high, intermediate and low *Prm1/Prm2* ratios (high versus intermediate: *D* = 0.08, *p* = 0.0002; high versus low: *D* = 0.1, *p* = 0.0001; intermediate versus low: *D* = 0.05, *p* = 0.05; figure 2). The same between-category differences in sperm head shape were obtained for *Prm2/Prm* ratio (high versus intermediate: *D* = 0.05, *p* = 0.0001; high versus low: *D* = 0.1, *p* = 0.0001; intermediate versus low: *D* = 0.08, *p* = 0.0001). These results support the idea that sperm head shape is influenced by relative protamine expression.

4. Discussion

Despite the long-standing debate over the relative contribution of coding versus regulatory changes to adaptive evolution [63–65], mounting empirical evidence demonstrates that regulatory evolution can play a major role in adaptive divergence, particularly between closely related lineages [65–71]. In this study, we compared protamine and transition nuclear protein mRNA expression in the testes of eight species in the genus *Mus* that share recent common ancestry but differ widely in inferred levels of sperm competition. We found that species that experience higher levels of sperm competition express less protamine 2 in relation to both transition nuclear proteins and to protamine 1. This strongly suggests that species differences in relative expression levels of these key spermiogenesis genes are influenced by variation in the strength of post-copulatory sexual selection. The fact that this pattern is driven by the relative expression of protamine 2 is consistent with evidence that the promoter region of this gene is evolving under sexual selection in *Mus* [24]. Importantly, we found that species that differ in ratios of protamine 2 expression, both in relation to protamine 1 and in relation to total protamines, also differ in sperm head shape. This suggests that regulatory changes contribute to modifications of sperm phenotype that could, ultimately, influence sperm's competitive ability. Taken together, the results of this study support the proposition that selection on regulatory regions can fine-tune adaptive phenotypes on short evolutionary timescales [72]. We discuss these results in relation to previous work on the evolution of sperm chromatin condensation genes in




Figure 2. Procrustes distances (*D*) and *p*-values for canonical variate analyses examining head shape in relation to *Prm1/Prm2* ratio. Three groups of species were defined according to their ratios of protamine expression: high (*M. macedonicus* and *M. spicilegus*), intermediate (*M. musculus*, *M. caroli* and *M. spretus*) and low (*M. castaneus* and *M. domesticus*) (see table 1). Morphometric data were taken from 35 individuals of seven species. Procrustes distances different from zero indicate shape differences between groups. Wireframe graphics show the shape associated with each group categorized according to its *Prm1/Prm2* ratio.

mammals and the genetics and functional consequences of sperm competition in rodents.

(a) Protamines and sperm competition: evolution at two levels

Sperm chromatin condensation genes, including protamines, are thought to be among the fastest evolving male reproductive proteins in eutherian mammals [73,74]. There is ample evidence from primates and rodents that selection contributes to this rapid rate of change [16,21,75,76] and sperm competition is often invoked as the driving force [15]. However, how particular substitutions might enhance sperm competitiveness remains untested, and it has been suggested that selection for protein stability is an equally parsimonious explanation for protamine-coding sequence evolution in primates [77]. Notably, in case-control studies of human males, associations between infertility and coding region SNPs in either *Prm1* or *Prm2* are rare [78–80], whereas men with imbalanced PRM1/PRM2 ratios are consistently subfertile or sterile (reviewed in [81]).

Thus, while the functional consequences of protamine-coding sequence substitutions are largely unknown, changes in protamine expression have a demonstrated impact on male fertility, and therefore might covary with the strength of post-copulatory sexual selection across species.

In the *Mus* clade comprising house mice and their close relatives, there is evidence for weak positive selection on *Prm2*-coding sequence in the three species with the highest inferred levels of sperm competition (*M. spicilegus*, *M. spretus* and *M. macedonicus*), whereas divergence in the promoter region is positively correlated with relative testes mass, and with sperm swimming speed, across the entire clade [24]. Here, using a subset of the same species, we show that the relative abundance of *Prm2* mRNA in the testes is negatively correlated with relative testes mass. These findings suggest that nucleotide substitutions in the *Prm2* promoter region influence expression and that high levels of sperm competition act to decrease the relative abundance of *Prm2* in the testes.

We emphasize, however, that our understanding of the relationship between protamine 2 regulation and sperm competition in *Mus* is far from complete. First, the functional relationship between promoter evolution and expression is not straightforward: species with higher *Prm2* promoter divergence express less *Prm2* only in relation to transition nuclear proteins and *Prm1*. Despite substantial interspecific differences in the expression levels of all four genes, there was no relationship between relative testes mass and individual gene expression. Likewise, although the *Prm1* promoter region is highly variable in *Mus*, there is no relationship between divergence and levels of sperm competition [24]. A plausible explanation for these patterns is that sexual selection for reduced PRM2 is counterbalanced by natural selection to maintain the relative proportions of protamines and transition nuclear proteins within a functional range. Potential mechanisms include compensatory evolution in the promoter regions of interacting sperm chromatin condensation proteins or a single regulatory modifier shared among genes. In mice, as in humans, *Prm1*, *Prm2* and *Tnp2* are tightly clustered in the genome. Thus, an enhancer element common to all three genes is a formal possibility. Comparative analysis of intergenic regions in the *Prm1/Prm2/Tnp2* cluster, together with the *Tnp1* and *Tnp2* promoter regions, will help to discriminate these non-mutually exclusive alternatives.

Second, the correlation between mRNA expression levels and protein abundance is often imperfect [82]. Quantification of sperm chromatin condensation proteins in mature spermatozoa will provide a direct measure of species differences in their relative abundance. Finally, evidence for selection on the *Prm2*-coding sequence in *M. spicilegus*, *M. spretus* and *M. macedonicus* is intriguing, because it suggests that high levels of sperm competition can drive coding and regulatory evolution in tandem

[24]. However, whether positively selected *Prm2* amino acid substitutions in these species affect sperm phenotypes related to competitive ability remains to be determined.

(b) The relative abundance of protamine 2: functional implications for sperm phenotypes

Why should high levels of sperm competition favour reduction in the relative abundance of PRM2? While the phenotypic effects of interspecific differences in protamine ratios are largely unstudied, there is some evidence that sperm from species that either lack PRM2, or produce very little PRM2 relative to PRM1, exhibit slower DNA decondensation in the oocyte [32,41]. Sperm with more compact heads may have higher competitive ability [10], and sperm with incomplete DNA compaction often have over-sized or less streamlined heads [34]. Thus, it is plausible that high levels of sperm competition select for higher DNA compaction, and thus proportionately less PRM2. Evaluation of this hypothesis will require comparative analyses of sperm chromatin compaction in relation to head morphology, the proportion of PRM2 and the strength of sexual selection mediated by sperm competition. Notably, the finding that relative abundance of *Prm2* is associated with differences in sperm head shape is an important first step towards revealing the functional relationship between protamine expression and sperm head morphology. Future studies will investigate the hydrodynamic consequences of these *Prm2*-associated differences in sperm head shape.

5. Conclusion

An important role of comparative studies such as this is to identify patterns that generate testable hypotheses [83]. Here, we show that species of mice with higher inferred levels of sperm competition express less protamine 2 in relation to protamine 1 and transition nuclear proteins. Based on this pattern, together with evidence for sexually selected divergence in the promoter region of protamine 2 [24], we propose that reduction in the relative abundance of protamine 2 enhances sperm competitive ability in mice by influencing sperm head shape and that regulatory evolution plays a key role in this evolutionarily rapid response to selection.

Acknowledgements. We thank François Bonhomme and Annie Orth for facilitating access to the animals used in this study.

Funding statement. L.L. holds a studentship from the Spanish Research Council (JAEpre-CSIC) co-financed by the European Social Fund (ESF). This work was supported by the Spanish Ministry of Economy and Competitiveness (grant no. CGL2011-26341) and was developed in part during a short stay at the University of Arizona with funding from CSIC ('Estancia Breve'). This work was also financially supported by NSF and NIH grants to M.W.N.

References

- Parker GA. 1970 Sperm competition and its evolutionary consequences in the insects. *Biol. Rev.* **45**, 525–567. (doi:10.1111/j.1469-185X.1970.tb01176.x)
- Birkhead TR, Møller AP. 1998 *Sperm competition and sexual selection*. London, UK: Academic Press.
- Simmons LW. 2001 *Sperm competition and its evolutionary consequences in the insects*. Princeton, NJ: Princeton University Press.
- Birkhead TR, Hosken DJ, Pitnick S. 2009 *Sperm biology: an evolutionary perspective*. Oxford, UK: Academic Press.
- Gómez Montoto L, Varea Sánchez M, Tourmente M, Martín-Coello J, Luque-Larena JJ, Gomendio M, Roldan ERS. 2011 Sperm competition differentially affects swimming velocity and size of spermatozoa from closely related muroid rodents: head first. *Reproduction* **142**, 819–830. (doi:10.1530/REP-11-0232)
- Gomendio M, Martín-Coello J, Crespo C, Magaña C, Roldan ERS. 2006 Sperm competition enhances functional capacity of mammalian spermatozoa. *Proc. Natl Acad. Sci. USA* **103**, 15 113–15 117. (doi:10.1073/pnas.0605795103)

7. Roldan ERS, Gomendio M, Vitullo AD. 1992 The evolution of eutherian spermatozoa and underlying selective forces: female selection and sperm competition. *Biol. Rev.* **67**, 551–593. (doi:10.1111/j.1469-185X.1992.tb01193.x)
8. Breed WG, Taylor J. 2000 Body mass, testes mass, and sperm size in murine rodents. *J. Mammal.* **81**, 758–768. (doi:10.1644/1545-1542(2000)081<0758:BMTMAS>2.3.CO;2)
9. Immler S, Moore HD, Breed WG, Birkhead TR. 2007 By hook or by crook? Morphometry, competition and cooperation in rodent sperm. *PLoS ONE* **2**, e170. (doi:10.1371/journal.pone.0000170)
10. Tourmente M, Gomendio M, Roldan ERS. 2011 Sperm competition and the evolution of sperm design in mammals. *BMC Evol. Biol.* **11**, 12. (doi:10.1186/1471-2148-11-12)
11. Gomendio M, Roldan ERS. 1991 Sperm competition influences sperm size in mammals. *Proc. R. Soc. Lond. B* **243**, 181–185. (doi:10.1098/rspb.1991.0029)
12. Gomendio M, Roldan ERS. 2008 Implications of diversity in sperm size and function for sperm competition and fertility. *Int. J. Dev. Biol.* **52**, 439–447. (doi:10.1387/ijdb.082595mg)
13. Firman RC, Simmons LW. 2009 Experimental evolution of sperm quality via postcopulatory sexual selection in house mice. *Evolution* **64**, 1245–1256.
14. Cummins JM, Yanagimachi R. 1982 Sperm–egg ratios and the site of the acrosome reaction during *in vivo* fertilization in the hamster. *Gamete Res.* **5**, 239–256. (doi:10.1002/mrd.1120050304)
15. Swanson WJ, Vacquier VD. 2002 The rapid evolution of reproductive proteins. *Nat. Rev. Genet.* **3**, 137–144. (doi:10.1038/nrg733)
16. Turner LM, Hoekstra HE. 2008 Causes and consequences of the evolution of reproductive proteins. *Int. J. Dev. Biol.* **52**, 769–780. (doi:10.1387/ijdb.082577lt)
17. Dorus S, Evans PD, Wyckoff GJ, Choi SS, Lahn BT. 2004 Rate of molecular evolution of the seminal protein gene SEMG2 correlates with levels of female promiscuity. *Nat. Genet.* **36**, 1326–1329. (doi:10.1038/ng1471)
18. Finn S, Civetta A. 2010 Sexual selection and the molecular evolution of ADAM proteins. *J. Mol. Evol.* **71**, 231–240. (doi:10.1007/s00239-010-9382-7)
19. Kingan SB, Tatar M, Rand DM. 2003 Reduced polymorphism in the chimpanzee semen coagulating protein, semenogelin I. *J. Mol. Evol.* **57**, 159–69. (doi:10.1007/s00239-002-2463-0)
20. Ramm SA, Oliver PL, Ponting CP, Stockley P, Emes RD. 2008 Sexual selection and the adaptive evolution of mammalian ejaculate proteins. *Mol. Biol. Evol.* **25**, 207–219. (doi:10.1093/molbev/msm242)
21. Wyckoff GJ, Wang W, Wu CI. 2000 Rapid evolution of male reproductive genes in the descent of man. *Nature* **403**, 304–309. (doi:10.1038/35002070)
22. Lüke L, Vicens A, Serra F, Luque-Larena JJ, Dopazo H, Roldan ERS, Gomendio M. 2011 Sexual selection halts the relaxation of protamine 2 among rodents. *PLoS ONE* **6**, e29247. (doi:10.1371/journal.pone.0029247)
23. Walters JR, Harrison RG. 2011 Decoupling of rapid and adaptive evolution among seminal fluid proteins in *Heliconius* butterflies with divergent mating systems. *Evolution* **65**, 2855–2871. (doi:10.1111/j.1558-5646.2011.01351.x)
24. Martin-Coello J, Dopazo H, Arbiza L, Ausió J, Roldan ERS, Gomendio M. 2009 Sexual selection drives weak positive selection in protamine genes and high promoter divergence, enhancing sperm competitiveness. *Proc. R. Soc. B* **276**, 2427–2436. (doi:10.1098/rspb.2009.0257)
25. Kleene KC, Bagarova J. 2008 Comparative genomics reveals gene-specific and shared regulatory sequences in the spermatid-expressed mammalian *Odf1*, *Prm1*, *Prm2*, *Tnp1* and *Tnp2* genes. *Genomics* **92**, 101–106. (doi:10.1016/j.ygeno.2008.05.001)
26. Nayernia K, Drabent B, Adham IM, Moschner M, Wolf S, Meinhardt A, Engel W. 2003 Mice lacking three germ cell expressed genes are fertile. *Biol. Reprod.* **69**, 1973–1978. (doi:10.1095/biolreprod.103.018564)
27. Brewer LR. 1999 Protamine-induced condensation and decondensation of the same DNA molecule. *Science* **286**, 120–123. (doi:10.1126/science.286.5437.120)
28. Meistrich ML, Mohapatra B, Shirley CR, Zhao M. 2003 Roles of transition nuclear proteins in spermiogenesis. *Chromosoma* **111**, 483–488. (doi:10.1007/s00412-002-0227-z)
29. Yu YE, Zhang Y, Unni E, Shirley CR, Deng JM, Russell LD, Weil MM, Behringer RR, Meistrich ML. 2000 Abnormal spermatogenesis and reduced fertility in transition nuclear protein 1 deficient mice. *Proc. Natl Acad. Sci. USA* **97**, 4683–4688. (doi:10.1073/pnas.97.9.4683)
30. Aoki VW, Carrell DT. 2003 Human protamines and the developing spermatid: their structure, function, expression and relationship with male infertility. *Asian J. Androl.* **5**, 315–324.
31. Cho C, Jung-Ha H, Willis WD, Goulding EH, Stein P, Xu Z, Schultz RM, Hecht NB, Eddy EM. 2003 Protamine-2 deficiency leads to sperm DNA damage and embryo death in mice. *Biol. Reprod.* **69**, 211–217. (doi:10.1095/biolreprod.102.015115)
32. Perreault SD, Barbee RR, Elstein KH, Zucker RM, Keefer CL. 1988 Interspecies differences in the stability of mammalian sperm nuclei assessed *in vivo* by sperm microinjection and *in vitro* by flow cytometry. *Biol. Reprod.* **39**, 157–167. (doi:10.1095/biolreprod39.1.157)
33. McLay DW, Clarke HJ. 2003 Remodelling the paternal chromatin at fertilization in mammals. *Reproduction* **125**, 625–633. (doi:10.1530/rep.0.1250625)
34. Balhorn R. 2007 The protamine family of sperm nuclear proteins. *Genome Biol.* **8**, 227. (doi:10.1186/gb-2007-8-9-227)
35. Hecht NB. 1993 Gene expression during male germ cell development. In *Cell and molecular biology of the testis* (eds C Desjardins, LL Ewing), pp. 400–432. New York, NY: Oxford University Press.
36. Mali P, Kaipia A, Kangasniemi M, Toppari J, Sandberg M, Hecht NB, Parvinen M. 1989 Stage-specific expression of nucleo-protein mRNAs during rat and mouse spermiogenesis. *Reprod. Fertil. Dev.* **1**, 369–382. (doi:10.1071/RD9890369)
37. Oliva R. 2006 Protamines and male infertility. *Hum. Reprod. Update* **12**, 417. (doi:10.1093/humupd/dml009)
38. Wu JY, Ribar TJ, Cummings DE, Burton KA, McKnight GS, Means AR. 2000 Spermiogenesis and exchange of basic nuclear proteins are impaired in male germ cells lacking Camk4. *Nat. Genet.* **25**, 448–452. (doi:10.1038/78153)
39. Zhao M *et al.* 2001 Targeted disruption of transition protein 2 gene subtly affects spermatogenesis in mice. *Mol. Cell Biol.* **21**, 7243–7255. (doi:10.1128/MCB.21.21.7243-7255.2001)
40. Cho C, Willis WD, Goulding EH, Haesook JH, Choi YC, Hecht NB, Eddy EM. 2001 Haploinsufficiency of protamine-1 or 2 causes infertility in mice. *Nat. Genet.* **28**, 82–86. (doi:10.1038/ng0501-82)
41. Corzett M, Mazrimas J, Balhorn R. 2002 Protamine 1, protamine 2 stoichiometry in the sperm of eutherian mammals. *Mol. Reprod. Dev.* **61**, 519–527. (doi:10.1002/mrd.10105)
42. Haueter S, Kawsumi M, Asner I, Brykczynska U, Cinelli P, Moisyadi S, Burki K, Peters AHFM, Pelczar P. 2010 Genetic vasectomy—overexpression of Prm1–EGFP fusion protein in elongating spermatids causes dominant male sterility in mice. *Genesis* **48**, 151–160.
43. Aoki VW, Liu L, Carrell DT. 2005 Identification and evaluation of a novel sperm protamine abnormality in a population of infertile males. *Hum. Reprod.* **20**, 1298–1306. (doi:10.1093/humrep/deh798)
44. Carrell DT, Liu L. 2001 Altered protamine 2 expression is uncommon in donors of known fertility, but common among men with poor fertilizing capacity, and may reflect other abnormalities of spermiogenesis. *J. Androl.* **22**, 604–610.
45. de Yebra L, Ballezá JL, Vanrell JA, Corzett M, Balhorn R, Oliva R. 1998 Detection of P2 precursors in the sperm cells of infertile patients who have reduced protamine P2 levels. *Fertil. Steril.* **69**, 755–759. (doi:10.1016/S0015-0282(98)00012-0)
46. Torregrosa N, Dominguez-Fandos D, Camejo MI, Shirley CR, Meistrich ML, Ballezá JL, Oliva R. 2006 Protamine 2 precursors, protamine 1/protamine 2 ratio, DNA integrity and other sperm parameters in infertile patients. *Hum. Reprod.* **21**, 2084–2089. (doi:10.1093/humrep/del114)
47. Carrell DT, Emery BR, Liu L. 1999 Characterization of aneuploidy rates, protamine levels, ultrastructure, and functional ability of round-headed sperm from two siblings and implications for intracytoplasmic sperm injection. *Fertil. Steril.* **71**, 511–516. (doi:10.1016/S0015-0282(98)00498-1)
48. Gómez Montoto L, Magaña C, Tourmente M, Martín-Coello J, Crespo C, Luque-Larena JJ, Gomendio M, Roldan ERS. 2011 Sperm competition, sperm numbers and sperm quality in muroid rodents. *PLoS ONE* **6**, e18173. (doi:10.1371/journal.pone.0018173)
49. Kenagy GJ, Trombulak SC. 1986 Size of mammalian testes in relation to body size. *J. Mammal.* **67**, 1–22. (doi:10.2307/1380997)

50. Gomendio M, Harcourt H, Roldan ERS. 1998 Sperm competition in mammals. In *Sperm competition and sexual selection* (eds TR Birkhead, AP Møller), pp. 667–751. London, UK: Academic Press.
51. Soulsbury CD. 2010 Genetic patterns of paternity and testes size in mammals. *PLoS ONE* **5**, e9581. (doi:10.1371/journal.pone.0009581)
52. Ramm SA, Parker GA, Stockley P. 2005 Sperm competition and the evolution of male reproductive anatomy in rodents. *Proc. R. Soc. B* **272**, 949–955. (doi:10.1098/rspb.2004.3048)
53. Felsenstein J. 1985 Phylogenies and the comparative method. *Am. Nat.* **125**, 1–15. (doi:10.1086/284325)
54. Martins EP. 2004 *COMPARE, version 46b: computer programs for the statistical analysis of comparative data*. Bloomington, IN: Department of Biology, Indiana University (<http://compare.bio.indiana.edu/>)
55. Lundrigan BL, Jansa S, Tucker PK. 2002 Phylogenetic relationships in the genus *Mus*, based on paternally, maternally, and biparentally inherited characters. *Syst. Biol.* **51**, 410–431. (doi:10.1080/10635150290069878)
56. Kendall D. 1986 The diffusion of shape. *Adv. Appl. Probab.* **9**, 428–430. (doi:10.2307/1426091)
57. Goodall C. 1991 Procrustes methods in the statistical analysis of shape. *J. R. Stat. Soc. B* **53**, 285–339.
58. Varea Sanchez M, Bastir M, Roldan ERS. 2013 Geometric morphometrics of rodent sperm head shape. *PLoS ONE* **8**, e80607. (doi:10.1371/journal.pone.0080607)
59. Klingenberg CP. 2011 MorphoJ: an integrated software package for geometric morphometrics. *Mol. Ecol. Res.* **11**, 353–357. (doi:10.1111/j.1755-0998.2010.02924.x)
60. Klingenberg CP, Gidaszewski A. 2010 Testing and quantifying phylogenetic signals and homoplasy in morphometric data. *Syst. Biol.* **59**, 245–261. (doi:10.1093/sysbio/syp106)
61. Campbell NA, Atchley WR. 1981 The geometry of canonical variate analysis. *Syst. Zool.* **30**, 268–280. (doi:10.2307/2413249)
62. Veyrunes F, Dobigny G, Yang F, O'Brien PCM, Catalan J, Robinson TJ, Britton-Davidian J. 2006 Phylogenomics of the genus *Mus* (Rodentia; Muridae): extensive genome repatterning is not restricted to the house mouse. *Proc. R. Soc. B* **273**, 2925–2934. (doi:10.1098/rspb.2006.3670)
63. Carroll SB. 2005 Evolution at two levels: on genes and form. *PLoS Biol.* **3**, 1159–1166. (doi:10.1371/journal.pbio.0030245)
64. Hoekstra HE, Coyne JA. 2007 The locus of evolution: evo devo and the genetics of adaptation. *Evolution* **61**, 995–1016. (doi:10.1111/j.1558-5646.2007.00105.x)
65. King MC, Wilson AC. 1975 Evolution at two levels in humans and chimpanzees. *Science* **188**, 107–188. (doi:10.1126/science.1090005)
66. Abzhanov A, Winston PK, Hartmann C, Grant R, Grant PR, Tabin CJ. 2006 The calmodulin pathway and evolution of elongated beak morphology in Darwin's finches. *Nature* **442**, 563–567. (doi:10.1038/nature04843)
67. Abzhanov A, Protas M, Grant R, Grant PR, Tabin CJ. 2004 *Bmp4* and morphological variation of beaks in Darwin's finches. *Science* **305**, 1462–1465. (doi:10.1126/science.1098095)
68. Carleton KL, Kocher TD. 2001 Cone opsin genes of African cichlid fishes: tuning spectral sensitivity by differential gene expression. *Mol. Biol. Evol.* **18**, 1540–1550. (doi:10.1093/oxfordjournals.molbev.a003940)
69. Jones FC *et al.* 2012 The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* **484**, 55–61. (doi:10.1038/nature10944)
70. Manceau M, Domingues VS, Mallarino R, Hoekstra HE. 2011 The developmental role of agouti in color pattern evolution. *Science* **331**, 1062–1065. (doi:10.1126/science.1200684)
71. Shapiro MD, Marks ME, Peichel CL, Blackman BK, Nereg KS, Jónsson B, Schluter D, Kingsley DM. 2004 Genetic and developmental basis of evolutionary pelvic reduction in threespine sticklebacks. *Nature* **428**, 717–723. (doi:10.1038/nature02415)
72. Wray GA. 2007 The evolutionary significance of cis-regulatory mutations. *Nat. Rev. Gen.* **8**, 206–216. (doi:10.1038/nrg2063)
73. Queralt R, Adroer R, Oliva R, Winkfein RJ, Retief JD, Dixon GH. 1995 Evolution of protamine P1 genes in mammals. *J. Mol. Evol.* **40**, 601–607. (doi:10.1007/BF00160507)
74. Su Y, Wu D, Zhou W, Irwin DM, Zhang Y. 2013 Rapid evolution of the mammalian *HILLS1* gene and the nuclear condensation process during mammalian spermiogenesis. *J. Genet. Genomics* **40**, 55–59. (doi:10.1016/j.jgg.2012.10.003)
75. Rooney AP, Zhang J. 1999 Rapid evolution of a primate sperm protein: relaxation of functional constraint or positive Darwinian selection? *Mol. Biol. Evol.* **16**, 706–710. (doi:10.1093/oxfordjournals.molbev.a026153)
76. Torgerson DG, Kulathinal RJ, Singh RS. 2002 Mammalian sperm proteins are rapidly evolving: evidence of positive selection in functionally diverse genes. *Mol. Biol. Evol.* **19**, 1973–1980. (doi:10.1093/oxfordjournals.molbev.a004021)
77. Clark AG, Civetta A. 2000 Evolutionary biology: protamine wars. *Nature* **403**, 261–263. (doi:10.1038/35002236)
78. Aoki VW, Christensen GL, Atkins JF, Carrell DT. 2006 Identification of novel polymorphisms in the nuclear protein genes and their relationship with human sperm protamine deficiency and severe male infertility. *Fertil. Steril.* **86**, 1416–1422. (doi:10.1016/j.fertnstert.2006.04.033)
79. He XJ, Ruan J, Du WD, Chen G, Zhou Y, Xu S, Zuo XB, Cao YX, Zhang XJ. 2012 Prm1 variant rs35576928. Arg > Ser is associated with defective spermatogenesis in the Chinese Han population. *Reprod. Biomed. Online* **25**, 627–634. (doi:10.1016/j.rbmo.2012.09.005)
80. Schlicker M, Schnulle V, Schnepfel L, Vorob'ev VI, Engel W. 1994 Disturbances of nuclear condensation in human spermatozoa: search for mutations in the genes for protamine 1, protamine 2 and transition nuclear protein 1. *Hum. Reprod.* **9**, 2313–2317.
81. Carrell DT, Emery BR, Hammoud S. 2007 Altered protamine expression and diminished spermatogenesis: what is the link? *Hum. Reprod. Update* **3**, 313–327. (doi:10.1093/humupd/dml057)
82. Greenbaum D, Colangelo C, Williams K, Gerstein M. 2003 Comparing protein abundance and mRNA expression levels on a genomic scale. *Genome Biol.* **4**, 117. (doi:10.1186/gb-2003-4-9-117)
83. Harvey PH, Pagel MD. 1991 *The comparative method in evolutionary biology*. New York, NY: Oxford University Press.