

Circumventing Antivector Immunity: Potential Use of Nonhuman Adenoviral Vectors

Estrella Lopez-Gordo,¹ Iva I. Podgorski,² Nicholas Downes,³ and Ramon Alemany⁴

Abstract

Adenoviruses are efficient gene delivery vectors based on their ability to transduce a wide variety of cell types and drive high-level transient transgene expression. While there have been advances in modifying human adenoviral (HAdV) vectors to increase their safety profile, there are still pitfalls that need to be further addressed. Preexisting humoral and cellular immunity against common HAdV serotypes limits the efficacy of gene transfer and duration of transgene expression. As an alternative, nonhuman AdV (NHAdV) vectors can circumvent neutralizing antibodies against HAdVs in immunized mice and monkeys and in human sera, suggesting that NHAdV vectors could circumvent preexisting humoral immunity against HAdVs in a clinical setting. Consequently, there has been an increased interest in developing NHAdV vectors for gene delivery in humans. In this review, we outline the recent advances and limitations of HAdV vectors for gene therapy and describe examples of NHAdV vectors focusing on their immunogenicity, tropism, and potential as effective gene therapy vehicles.

Introduction

ADENOVIRUSES (AdVs) ARE NONENVELOPED viruses with an icosahedral capsid containing a double-stranded linear DNA genome of 26–46 kb (Davison *et al.*, 2003). Since their discovery in 1950s in human cell cultures (Rowe *et al.*, 1953), AdVs have been identified in many species, ranging from fish to humans. Today AdVs form five accepted genera (*Mastadenovirus*, *Aviadenovirus*, *Siadenovirus*, *Atadenovirus*, and *Ichtadenovirus*) within the family *Adenoviridae*. Recently, one more genus was proposed, named *Testadenovirus* (Doszpoly *et al.*, 2013). So far 57 different AdV serotypes (Lukashev *et al.*, 2008; Walsh *et al.*, 2011) have been isolated from humans (HAdVs) and are classified within the genus *Mastadenovirus* as 7 species, HAdV A through HAdV G (Jones *et al.*, 2007). While classification was previously based on biological characteristics (erythrocyte agglutination, oncogenicity, and by specific antisera neutralization), phylogenetic relationships have been revised and clarified on the basis of genotype similarity (Bailey and Mautner, 1994; Davison *et al.*, 2003). Accordingly, individual AdV serotypes frequently share characteristics such as tropism and associated host pathologies with

other serotypes within their species. The tropism is in large part determined by the capsid proteins that interact with the host cell surface receptors to mediate cell entry. The capsid consists primarily of hexon and penton proteins that form pentameric structures at each of the 12 vertices, securing a trimeric fiber that projects outward. Other viral proteins include the cement proteins (IIIa, VI, VIII, and IX) and the core proteins (TP, V, VII, and mu) that associate with the AdV genome (Vellinga *et al.*, 2005). While the basic structure of these proteins is maintained, variations in the length and composition of the amino acid sequences of these proteins across species and genera cause them to bind to a number of receptors and host factors with variable affinity. Thus, the coxsackie and AdV receptor (CAR), CD46, desmoglein-2 (DSG2), CD80, CD86, vascular cell adhesion molecule-1 (VCAM-1), heparan sulfate proteoglycans (HSPGs), major histocompatibility complex class I- α 2 (MHC-I- α 2), sialic acid, dipalmitoylphosphatidylcholine, lactoferrin, and others can be utilized by one or more HAdV serotypes to transduce cells (Sharma *et al.*, 2009b; Arnberg, 2012).

Viral vectors remain to be the most promising gene transfer vehicles for gene therapy because of their higher

¹Institute of Cardiovascular and Medical Sciences, BHF Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow G12 8TA, United Kingdom.

²Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest 1143, Hungary.

³A.I. Virtanen Institute, Department of Biotechnology and Molecular Medicine, University of Eastern Finland, Kuopio 70211, Finland.

⁴Laboratori de Recerca Translacional, Institut Català d'Oncologia, El Instituto de Investigación Biomédica de Bellvitge, Barcelona 08907, Spain.

gene-transfer efficiencies when compared with other approaches. Currently, AdV vectors have been used in 23.5% of all clinical trials as they can efficiently transduce a wide variety of dividing and quiescent cells, they very rarely integrate into the host genome minimizing the risk of insertional mutagenesis (Stephen *et al.*, 2010), they have a large packaging capacity up to 36 kb, and they can routinely be produced at high titers. Of the 57 HAdV serotypes, HAdV-5 of species C has been the best characterized and the most widely used in clinical trials so far, mainly in the treatment of cancer but also as a vaccine (Sullivan *et al.*, 2003; Shiver and Emini, 2004).

It has, however, been estimated that over 80% of the adult population have been naturally exposed to the common HAdV serotypes (Garnett *et al.*, 2002). As a result, pre-existing humoral and cellular immunity may preclude efficient gene transfer (Kuriyama *et al.*, 1998), and local systemic immune responses against the vector and transgene product may result in a number of clinically undesirable side effects. Severe immune responses against AdV vectors can have serious consequences in humans, which have been exemplified by a fatal case of systemic inflammatory response syndrome that was directly attributed to the systemically administered AdV gene transfer vector (Raper *et al.*, 2003). Moreover, upon systemic administration of recombinant HAdV-5, transduction of immune cells in the liver and spleen leads to immune responses against the AdV vector that can limit its therapeutic efficacy. Eliminating liver tropism and the epitopes involved in viral proteins recognition by neutralizing antibodies (NAbs) has been proposed to reduce these immune responses. Taking advantage of the diversity of AdV capsid protein isoforms and their variety of ligand–receptor interactions, vectors based on low seroprevalent AdVs could potentially enable targeting to different cell types as well as overcoming pre-existing immunity. Naturally, nonhuman AdV (NHAdV) species have a low seroprevalence in all human populations and therefore represent a source of potential vectors that could fit this niche. Since the beginning of their development in 1990s (Mittal *et al.*, 1995), recombinant NHAdV vectors have been derived from many different AdV species: canine AdV-2 (CAAdV-2), bovine AdV-3 (BAAdV-3), porcine AdV-3 (PAAdV-3), ovine AdV-7 (OAdV-7), murine AdV-1 (MAAdV-1), several simian (SAdVs), and fowl (FAdVs) AdVs. Here we outline the current status of HAdV vectors, summarize the development of NHAdV vectors, and discuss the advantages and shortfalls of each.

Innate and Adaptive Immune Responses Against AdVs

Innate and adaptive immune responses play a critical role in resolving AdV infections. The presence of specific antiviral antibodies significantly increases antibody constant fraction receptor (FcR)-dependent viral internalization in macrophages, triggering an innate immune response to AdVs that leads to type I interferon (IFN) production and caspase-dependent interleukin (IL)-1 β maturation (Zaiss *et al.*, 2009). Internalized AdV DNA induces maturation of pro-IL-1 β in macrophages dependent on NALP3 and ASC, components of the innate cytosolic molecular complex called the inflammasome, but independent of Toll-like receptors (TLRs) and IFN regulatory factors (Muruve *et al.*,

2008). Later studies reported that this response requires the binding of the arginine-glycine-aspartic acid (RGD) motif in the AdV penton base with macrophage β 3 integrins (Di Paolo *et al.*, 2009). Barlan and colleagues also demonstrated that the specific AdV penetration mechanism of the endosomal membranes is required for IL-1 β release (Barlan *et al.*, 2011a,b). Macrophages and conventional dendritic cells (cDCs) can detect cytosolic AdV DNA in a TLR9-independent manner (Basner-Tschakarjan *et al.*, 2006; Zhu *et al.*, 2007), while plasmacytoid DCs (pDCs) detect non-methylated AdV CpG-containing DNA in the endosomes through TLR9. Also, internalized HAdV-5 and HAdV-6 immune complexes have been reported to induce DC maturation via TLR9 agonist motifs present in the AdV genome, whereas HAdV-35, -36, and -26 have a limited potency to induce DC maturation because of having a smaller number of motifs (Perreau *et al.*, 2012). TLR9 signaling pathway is dependent on myeloid differentiation primary response 88 (MyD88) (Yamaguchi *et al.*, 2007; Zhu *et al.*, 2007) and induces the secretion of IL-6 and IL-12 (Yamaguchi *et al.*, 2007). Heparin-sensitive receptors on DCs bind to the AdV shaft fiber region and trigger T cell immune responses (Cheng *et al.*, 2007). HAdV-5-induced natural killer (NK) cell activation relies on T cell contribution *in vitro* (Pahl *et al.*, 2012), and also appears to be related to the presence of type I IFN *in vivo* (Zhu *et al.*, 2007, 2008). In contrast to HAdV-5, in response to HAdV-35 pDCs alone are sufficient for NK cell activation in a TLR9-dependent manner, after which NK cells enhance pDC IFN- α secretion (Pahl *et al.*, 2012). This indicates that both pDCs and T cells may be necessary for NK cells' activation. Therefore, it is clear that different AdV serotypes can trigger immune responses in different ways; thus, it is essential to study immune responses in detail to ensure that efficient strategies to circumvent immune responses are developed.

The complement system also plays a critical role in AdV immune responses with effects mediated through the classical and alternative complement pathway. The complement system appears to be related to humoral immune responses to the vector capsid or the encoded transgene (Appledorn *et al.*, 2008). IgM isotype neutralization of virions activates the classical complement pathway hampering receptor–ligand interactions with the target cell (Xu *et al.*, 2013). On the other side, AdV-induced thrombocytopenia has been demonstrated to be dependent on factor B and C3 of the alternative complement pathway, and AdVs mediate induction of the nuclear factor-kappaB (NF- κ B) upon binding to C3 (Appledorn *et al.*, 2008). In addition to macrophages, DCs, and NK cells, binding of the CAR protein and α v-integrins on nonimmune cells also activates immune responses (Liu *et al.*, 2005; Schoggins *et al.*, 2005). In particular, CAR-mediated responses have been reported in epithelial cells (Tamanini *et al.*, 2006) and integrin-mediated responses in kidney epithelium-derived (REC) cells, inducing expression of CXCL10 (IP-10) through NF- κ B in an RGD motif-dependent manner (Liu *et al.*, 2005).

These anti-AdV innate immune responses are linked to adaptive immune responses through direct interactions and production of cytokines (Ginsberg, 1996; Appledorn *et al.*, 2008). When a cell is infected by AdVs, it releases type I (α and β) IFN. Plasmacytoid DCs, cDCs, and macrophages produce IFN- α upon AdV recognition. The surrounding

uninfected cells recognize IFN by IFN receptors and they induce, via the Jak-STAT signaling pathway, the expression of antiviral enzymes such as MxA, 2',5'-oligoadenylate synthase, and protein kinase R (Katze *et al.*, 2002). Also, type I IFN induces upregulation of costimulatory molecules in DCs such as CD80, CD86, and CD40, leading to DCs' maturation (Vujanovic *et al.*, 2009). Moreover, IFN takes part in B cells' activation for antibody production. Type II IFN (IFN- γ) is an immunoregulatory cytokine secreted by Th1-type T CD4+ cells, T CD8+ cells, and NK cells. IFN- γ induces the expression of major histocompatibility complex class I (MHC-I) on nearly all cells and major histocompatibility complex class II (MHC-II) on professional antigen presenting cells (APCs) promoting antigen presentation to helper T CD4+ cells, and it also activates macrophages (Gattoni *et al.*, 2006).

Dendritic cells have been shown to engulf infected cells and present antigens to naive T CD8+ cells through MHC-I (Albert *et al.*, 1998). At the same time, APCs present antigens to Th1-type T CD4+ cells through MHC-II and upon activation they produce IL-2. If the T CD8+ cell receives both antigen presence and IL-2 signal, it gets activated (Jankovic *et al.*, 2001). Once the T CD8+ cell interacts with an infected cell, mainly recognizing conserved epitopes in hexon protein (Olive *et al.*, 2002; Leen *et al.*, 2004; Tang *et al.*, 2006), it begins a cytotoxic response leading to apoptosis of the infected cell (Schumacher *et al.*, 2004). On the other hand, Th2-type T CD4+ cells induce B cell maturation and isotype switching (Jankovic *et al.*, 2001).

NAbs, mostly IgG isotype, interact with AdV capsid proteins (Gall *et al.*, 1996; Sumida *et al.*, 2005). It was reported that NAb recognize mainly the hypervariable regions (HVRs) of the hexon protein (Roberts *et al.*, 2006; Abe *et al.*, 2009; Shiratsuchi *et al.*, 2010; Bradley *et al.*, 2012b) and to a lesser extent the fiber knob domain (Myhre *et al.*, 2007; Bradley *et al.*, 2012a; Yu *et al.*, 2013) and penton base proteins (Yu *et al.*, 2013). AdV neutralization prevents cell attachment and facilitates aggregation and phagocytosis mediated by opsonization (Wohlfart, 1988). FcRs in macrophages and NK cells recognize NAb bound to AdV antigens, facilitating their phagocytic and cytolytic action, respectively (Gattoni *et al.*, 2006).

In summary, AdV gene transfer vectors are subjected to antiviral immune responses (Fig. 1) once administered to the patient. This process highly limits their efficiency for gene therapy applications. Therefore, it is critical to understand the immune response to AdVs and have strategies to circumvent antiviral immune responses. These will be discussed in the following sections, focusing on the main limiting factors: NAb against AdV proteins and CD8+ cytotoxic T lymphocytes (CTLs) against infected cells.

Implications of Antiviral Immune Responses Against HAdV Vectors and Strategies to Circumvent Them

Since the tragic death of a patient participating in a clinical gene therapy trial involving administration of a replication-incompetent HAdV-5 vector (Raper *et al.*, 2003), many efforts have been made to avoid antiviral immune responses against AdV vectors to improve their safety and efficacy. Since the AdV vector recognition by the immune system depends on the existence of antigenic epitopes

in the AdV proteins (i.e., NAb recognize AdV proteins, and CD8+ CTLs recognize infected cells), most of the strategies to circumvent the immunogenicity of HAdV vectors are based on protein modifications.

Functionally significant HAdV-5-specific NAb are directed primarily against the HAdV-5 hexon protein and to a lesser extent to fiber and penton base proteins (Gall *et al.*, 1996; Sumida *et al.*, 2005; Tian *et al.*, 2011). Despite several groups successfully generated hexon-chimeric HAdV-5 vectors that partially evaded anti-HAdV-5 immune responses (Gall *et al.*, 1998; Roy *et al.*, 1998; Ostapchuk and Hearing, 2001; Wu *et al.*, 2002; Roberts *et al.*, 2006; Bruder *et al.*, 2012), hexon pseudotyping appears to be a limited strategy as it has been reported the formation of many nonviable virions (Youil *et al.*, 2002).

The hexon protein (Rux and Burnett, 2000) contains 7 HVRs (Crawford-Miksza and Schnurr, 1996). Many groups have exchanged HAdV-5 HVRs with those from low-prevalent serotypes and demonstrated that the resultant AdV vector can evade HAdV-5 immunity (Roberts *et al.*, 2006; Abe *et al.*, 2009; Shiratsuchi *et al.*, 2010). Roberts *et al.* (2006) showed that HAdV-5 with all seven HVRs replaced with the corresponding ones from the rare serotype HAdV-48 could circumvent HAdV-5 NAb in both mice and rhesus monkeys. Further studies (Bradley *et al.*, 2012b) exchanged only some HVRs and demonstrated that HAdV-5-specific NAb target multiple HVRs, suggesting that mutation or replacement of all seven HVRs would probably be necessary to evade anti-HAdV-5 immunity. However, in 2012, Coughlan *et al.* demonstrated that, despite HAdV-5 containing all seven HVRs from HAdV-48, it displayed a decreased hepatocyte transduction and accumulation in Kupffer cells, and it triggered a robust proinflammatory response not present with the wild-type HAdV-48 (Coughlan *et al.*, 2012). In order to generate novel viable HVR-chimeric vectors able to evade immune responses, it would be necessary to use HVRs from other serotypes, and for that it is essential to determine the viability of HVR replacements in AdV vectors based on the structural and biochemical constraints of the different HVRs that limit their manipulation.

Regarding the immunogenicity of the fiber protein, little is known about the epitopes involved in fiber recognition by NAb, although it is thought that antiknob NAb represent the major population of antifiber NAb (Myhre *et al.*, 2007; Bradley *et al.*, 2012a; Yu *et al.*, 2013). It has been reported that fiber pseudotyping of AdV vectors can lead to reduced AdV-associated innate (Schoggins *et al.*, 2005) and adaptive immune responses (Parker *et al.*, 2009; Rogée *et al.*, 2010). Despite the C-terminal and shaft region heterogeneity, the N-terminal region (fiber tail) that is bound to penton base protein presents high similarity among diverse AdV species (Tarassishin *et al.*, 2000). Thus, designing chimeric AdV vectors containing all proteins from HAdV-5 but fibers from other serotypes (mainly species B and D) is a very interesting approach to retarget HAdV-5 to different tissues while circumventing antiviral preexisting immunity. Thus, several chimeric AdV vectors with tropism for different cell types such as HAdV-5/3 (Haviv *et al.*, 2002; Kanerva *et al.*, 2002; Volk *et al.*, 2003; Ulasov *et al.*, 2007), HAdV-5/11 (Havenga *et al.*, 2001; Stone *et al.*, 2005; Wang *et al.*, 2011b), HAdV-5/16 (Havenga *et al.*, 2001), and HAdV-5/35




FIG. 1. Innate and adaptive antiadenoviral immune responses. (1) Activation of macrophages and natural killer (NK) cells by recognition of opsonized adenovirus (AdV) through antibody-constant fraction receptor (FcR). (2) Interleukin (IL)-1 β maturation upon binding of arginine-glycine-aspartic acid (RGD) motif in the AdV penton base with macrophage β 3 integrins. (3) Induction of IL-6 and IL-12 secretion in plasmacytoid dendritic cell (pDC) upon AdV DNA recognition through Toll-like receptor (TLR)-9. (4) Dendritic cell (DC) maturation through internalized immune complexes (ICs). (5) Triggering of T cell immune responses upon binding of AdV fiber shaft region to heparin-sensitive receptors on DCs. (6) Interferon (IFN)- α production by activated pDCs, conventional DCs (cDCs), and macrophage and IFN- α recognition by uninfected cells leading to a cellular antiviral state. (7) Induction of CXCL10 expression by AdV penton base RGD motif binding to α_v integrins. (8) Secretion of IFN- α and β by infected cells. (9) Classical complement pathway activation by virion neutralization with IgM or recognition by IgG. (10) Thrombocytopenia related to alternative complement pathway activation on AdVs. (11) Major histocompatibility complex class II (MHC-II) induction on professional antigen presenting cells (APCs) by IFN- γ signal promoting antigen presentation to helper T CD4+ cells. (12) Induction of B cell maturation and isotype switching through AdV antigen presentation by Th2-type T CD4+ cells. (13) Activation of T CD8+ cells through AdV antigen presentation by DC via MHC-I. (14) IL-2 production by Th1-type T CD4+ cells upon AdV antigen presentation by APCs through MHC-II, which activates T CD8+ cells. BCR, B cell receptor; C3b, complement component 3b; CAR, coxsackie virus and adenovirus receptor; IFN R, interferon receptor; TCR, T cell receptor.

(Havenga *et al.*, 2001) were generated. Interestingly, HAdV-5/35 and -5/11 were assessed *in vivo* in murine and non-human primate animal models, showing reduced toxicity and limited induction of inflammatory cytokines (Ni *et al.*, 2005). However, other studies showed that HAdV-5/35 presented tropism for CD34+ human hematopoietic stem cells *in vitro* (Shayakhmetov *et al.*, 2000), monocytes, granulocytes, and blast cells of human bone marrow (Rogozhin *et al.*, 2011), and CD4+ and CD8+ T lymphocytes *in vitro* (Zhang *et al.*, 2013b). Since transduction of immune cells can lead to immune responses, extensive characterization of chimeric AdV vectors is essential.

Other retargeting strategies such as the truncation or removal of the fiber knob domain and its replacement with foreign trimerization motifs and heterologous peptides fused to the fiber shaft are also recently being assessed (fiber de-

knobbing) (Coughlan *et al.*, 2010). Despite that the removal of the fiber knob domain results in a lower number of fiber copies per virion, leading to a less efficient production of the AdV vector, it has been shown that this approach may contribute to evade NABs (Myhre *et al.*, 2007).

Regarding the adaptive immunity mediated by CD8+ CTLs, the response depends on viral antigenic epitopes loaded in the MHC of the infected cell. High-capacity, helper-dependent (HD) vectors (also called gutless vectors) bypass this problem (Kochanek *et al.*, 1996) as they are devoid of all viral genes, and therefore they are less immunogenic and allow long-term transgene expression (Maione *et al.*, 2001; Ehrhardt and Kay, 2002; Dudley *et al.*, 2004). Furthermore, as in these vectors DNA is constituted of stuffer DNA as part of the vector backbone to maintain optimum vector size, nonmethylated viral CpG-containing

DNA (recognized by DCs through TLR9) can be eliminated. However, one study showed that HD AdVs also trigger an innate immune response dependent on TLR9 (Cerullo *et al.*, 2007). Despite the advantages they offer, HD vectors are equally neutralized by preexisting antibodies against capsid proteins and also may present transgene-encoded protein immunogenicity (Tripathy *et al.*, 1996).

On the other hand, the activation of immune responses because of interactions between AdV capsid proteins and receptors in the target cell can limit AdV vector efficacy in gene therapy applications. HAdV-5 binds to the CAR protein on human erythrocytes in the bloodstream, limiting its delivery to the *in vivo* target tissue and contributing to toxicity (Seiradake *et al.*, 2009). To avoid this interaction, Nicol *et al.* (2004) described two CAR-binding mutations that abolished the previously described agglutination in human and rat erythrocytes. Moreover, HAdV-5 binding to CAR activates immune responses that can limit its potential as a gene transfer vector (Schoggins *et al.*, 2005; Tamanini *et al.*, 2006). Other interactions such as RGD with macrophage $\beta 3$ integrins (Cheng *et al.*, 2007; Di Paolo *et al.*, 2009) or the one between the shaft fiber region and the heparin-sensitive receptor from DCs (Cheng *et al.*, 2007) have also been associated with immune responses and, in the case of RGD motif, through IL-1 α production (Di Paolo *et al.*, 2009). Interestingly, the mutation of the RGD motif to arginine-glycine-glutamic acid (RGE) resulted in a significant fivefold reduction in spleen uptake, diminishing the antiviral inflammatory response after intravascular administration (Bradshaw *et al.*, 2012). These studies indicate that de-targeting from native receptors could be a useful approach to avoid immune responses against the AdV vector. Nevertheless, further research needs to be done to describe the pathways through which the AdV vectors trigger immune responses. There are more factors involved in cellular recognition that could also be implicated in promotion of immune responses. These include alternative receptors and coreceptors such as FcR (Xu *et al.*, 2008), complement-3 receptor (CR-3) (Xu *et al.*, 2008), HSPGs and low-density lipoprotein receptor-related protein (LRP) (Shayakhmetov *et al.*, 2005), complement receptor-1 (CR1) (Carlisle *et al.*, 2009), scavenging receptor-A (SR-A) (Khare *et al.*, 2012), bridging molecules such as the coagulation factor IX, X, and VII and protein C (Parker *et al.*, 2006), complement component C4-binding protein (C4BP) (Shayakhmetov *et al.*, 2005), and IgM antibodies (Xu *et al.*, 2008). Further studies will have to be done in order to describe their influence on the use of HAdV vectors for gene therapy applications.

Apart from genetic engineering of HAdV vectors, other strategies such as chemical modifications by addition of cationic polymers or lipid molecules to shield AdV vectors and prevent them from binding undesired proteins have been widely studied. Several variants of polyethylene glycol (PEG) have been used, but better shielding was found with multivalent copolymers of poly[*N*-(2-hydroxypropyl)methacrylamide] (pHPMA). Shielding of HAdV vectors with pHPMA showed an increased biological stability of the vector and a longer persistence in blood, decreased liver transduction, and therefore a lower immune response (Green *et al.*, 2004). Furthermore, coated HAdV vectors have been shown to be less vulnerable to recognition by preexisting NAbs *in vitro* (Romanczuk *et al.*, 1999; Wang *et al.*, 2011a) and by Kupffer cells

in the liver (Mok *et al.*, 2005), likely because of the hampered interaction of PEGylated AdVs and Kupffer cell SR-A (Haisma *et al.*, 2009). However, other factors could be affecting liver transduction of the coated HAdV such as the limited size of the sinusoidal endothelial cells' fenestrae (Hofherr *et al.*, 2008). The use of PEGylated HAdV vectors resulted in decreased levels of IL-6 production *in vitro* (Mok *et al.*, 2005) and of IL-6 and IL-12 *in vivo* (Croyle *et al.*, 2000). Moreover, coated HAdV vectors induce lower IL-6 levels in serum because of decreased levels of spleen AdV uptake (De Geest *et al.*, 2005). Furthermore, several groups successfully retargeted coated AdV vectors (Parker *et al.*, 2005; Stevenson *et al.*, 2007; Morrison *et al.*, 2008, 2009; Wang *et al.*, 2010). Although in some studies they presented similar levels of antitumoral efficacy (Morrison *et al.*, 2009) or transgene expression (Parker *et al.*, 2005; Stevenson *et al.*, 2007; Morrison *et al.*, 2008; Wang *et al.*, 2010) to unmodified wild-type HAdV-5, others achieved a greater therapeutic effect with negligible side effects (Eto *et al.*, 2010; Yao *et al.*, 2010). Despite the limited retargeting efficiency in comparison with unmodified HAdV-5, the use of PEGylated AdV vectors achieved lower hepatic tropism and reduction in liver and spleen toxicity *in vivo*. Moreover, evidence highlights the potential of AdV polymer coating strategies to circumvent preexisting immunity to all viral capsid proteins (Romanczuk *et al.*, 1999; Croyle *et al.*, 2000; De Geest *et al.*, 2005; Mok *et al.*, 2005; Haisma *et al.*, 2009; Zeng *et al.*, 2012). Nevertheless, the reported PEG immunogenicity could hamper PEGylated AdVs efficacy (Shimizu *et al.*, 2012).

Despite the advances made in HAdV vectors to evade anti-AdV immune responses, the current strategies are limited by the formation of nonviable virions following molecular engineering and by the limited knowledge and characterization of capsid epitopes motifs responsible for cross-reactivity. Therefore, the use of low-seroprevalent HAdV serotypes has been proposed as an alternative to hexon or fiber-chimeric AdV vectors in order to circumvent preexisting NAbs. More than 15 years ago, several studies showed that the alternate use of AdV vectors from different serotypes, even within the same AdV species, can circumvent anti-AdV humoral immunity (Kass-Eisler *et al.*, 1996; Mastrangeli *et al.*, 1996; Mack *et al.*, 1997; Parks *et al.*, 1999). A study on the presence of NAbs against different HAdV serotypes in the Belgian population reported high seroprevalence toward species A, C, and E, while low seroprevalence toward species B and D, among which, serotypes 11, 34, 35, 50 (species B), 43, and 48 (species D) showed the lowest values (Vogels *et al.*, 2003). This indicates that these serotypes could be potentially used to evade preexisting immunity. Vectors have been developed from multiple HAdV serotypes with lower seroprevalence than HAdV-5: HAdV-11 (Holterman *et al.*, 2004; Stone *et al.*, 2005; Abbink *et al.*, 2007), HAdV-35 (Gao *et al.*, 2003; Sakurai *et al.*, 2003, 2009; Seshidhar Reddy *et al.*, 2003; Vogels *et al.*, 2003; Barouch *et al.*, 2004; Wu and Tikoo, 2004; Abbink *et al.*, 2007; Brouwer *et al.*, 2007; Sakurai, 2008; McVey *et al.*, 2010; Geisbert *et al.*, 2011), HAdV-49 (Lemckert *et al.*, 2006; Abbink *et al.*, 2007), HAdV-26, -48, and -50 (Abbink *et al.*, 2007; Geisbert *et al.*, 2011), HAdV-7 (Nan *et al.*, 2003), and HAdV-6 (Capone *et al.*, 2006). However, since the presence of NAbs depends on AdV exposure, which is closely related to geographical location,

seroprevalence studies in different specific populations are very important. Moreover, better characterization of such serotypes in terms of tropism and specificity, biodistribution, and toxicity is yet required to design safe HAdV vectors.

Use of Nonhuman Adenoviral Vectors

The idea about NHAdVs as vectors appeared in 1990s, with the assumption that they would be more useful for vaccines and gene therapeutic approaches in human medicine than HAdV-based vectors. Design and characterization of NHAdV vectors was reviewed earlier (Bangari and Mittal, 2006), with a focus on vaccine vectors. In this review we focus on the progress in the development of NHAdV vectors (Table 1) for gene therapy.

Canine AdV vectors

Canine AdV (genus *Mastadenovirus*, species CAdV A)-derived vectors are currently the best described NHAdV vectors. Up to 90% of the residues involved in CAdV-2 fiber tail-penton base interactions are conserved compared with HAdV-5 (Schoehn *et al.*, 2008). CAdV-2 binds to CAR (Tomko *et al.*, 1997; Soudais *et al.*, 2000), but it does not contain an integrin-interacting RGD motif in the penton base (Chillon and Kremer, 2001) in contrast with HAdV-5 (Greber, 2002). CAdV-2 penton base appears to be ~20% shorter than that of HAdV-5 (Schoehn *et al.*, 2008). In addition, CAdV-2 has shorter protuberances on the top of the penton base, mainly where RGD motif is contained in HAdVs, and a shorter N-terminal region. The hypervariable loops in the hexon protein are shorter in CAdV-2 and the protruding structures containing the epitopes recognized by antihexon HAdV-5 antibodies are absent, while the fiber is more complex since the shaft region contains two bends (Schoehn *et al.*, 2008). Also, protein IX and IIIa are shorter in CAdV-2 (Schoehn *et al.*, 2008). Production of CAdV-2 vectors started

more than 15 years ago (Klonjowski *et al.*, 1997) in an attempt to avoid the inhibitory effect of the preexisting humoral and cellular immunity against HAdVs when used in the clinics. Later, it was shown that CAdV-2 vectors rarely cause neither the activation and proliferation of T cells nor the maturation of DCs (Perreau and Kremer, 2006). The advantage of these vectors over HAdV vectors is that CAdV-2 mainly interacts with CAR (Soudais *et al.*, 2000) and they present specific tropism for neurons in the central nervous system (Soudais *et al.*, 2001) with great capacity for axonal transport *in vivo*. Taking into account that CAdV-2 vector development was recently reviewed (Bru *et al.*, 2010), the following data will focus on the more recent trials performed with the mentioned vector. Until very recently, the production of CAdV-2 vectors was based on dog kidney (DK) cell lines (Kremer *et al.*, 2000), which is not accepted by the Food and Drug Administration (FDA) and the European Medicine Agency (EMA). To circumvent this problem, CAdV-2 E1-transcomplementing cell lines, based on Madin-Darby canine kidney (MDCK) cells accepted by the FDA and EMA for the production of vaccines, have been developed (Fernandes *et al.*, 2013a). CAdV-2 vector DNA replication in the MDCK-E1 cell line was shown to correlate with the expression levels of the transcomplementing E1A and E1B. While Cre recombinase expression in MDCK-E1-Cre cell lines (for HD CAdV-2 vectors production) can impair cell growth, controlled expression of Cre was determined not to result in negative effects on viral production (Fernandes *et al.*, 2013b). CAdV-2-based vectors are ideally suited for treating neurological disorders, but because of vector titer limitations, large volumes are necessary for applicability in the clinics. Therefore, better purification and scalable production according to good manufacturing practice (GMP) are being currently developed (Segura *et al.*, 2012; Fernandes *et al.*, 2013a), and the improvement of producer cell lines has increased vector titers (Ibanes and Kremer, 2013). In a comparative study on HD HAdV, HD CAdV-2, and a VSV-G lentiviral vector, the transduction efficiency and transgene expression in human midbrain neuroprogenitor cells was assessed. HD CAdV-2 was determined to have the most promising vector profile: equal transduction efficacy, no negative effect to neuronal development, and milder induced immune response (Piersanti *et al.*, 2013). HD CAdV-2 was also used as a vector in order to evaluate a gene therapy approach to correct the neurodegenerative lysosomal storage disorder mucopolysaccharidosis type IIIa (Lau *et al.*, 2012). It was shown to effectively transduce neurons and provide long-term transgene expression; however, its use *in vivo* was limited to discrete areas of the brain, indicating a need to find a way to increase transgene expression throughout the whole brain. Furthermore, in a clinically relevant stroke model in rats, cortical CAdV-2 vector injection transduced neurons at a greater level and covered a larger region than lentiviral vectors (Ord *et al.*, 2013), reducing infarct volume and improving neurologic recovery. For all the potential of CAdV-2 vectors, there ultimately remains the need for further understanding of the effects of CAdV-2 vectors on human cells before they are able to enter human clinical trials.

Bovine AdV vectors

Bovine AdVs (genus *Mastadenovirus*, species BAdV A, B, C; genus *Atadenovirus*, species BAdV D) were the first

TABLE 1. MOST POPULAR NONHUMAN ADENOVIRUS USED AS A VECTOR

Species	Genus	Serotype
Canine adenovirus A	<i>Mastadenovirus</i>	CAdV-2
Bovine adenovirus B	<i>Mastadenovirus</i>	BAdV-3
Porcine adenovirus A	<i>Mastadenovirus</i>	PAdV-3
Human adenovirus E	<i>Mastadenovirus</i>	ChAdV-68/Pan 9/SAdV-25 ChAdV-5/Pan 5/SAdV-22 ChAdV-6/Pan 6/SAdV-23 ChAdV-7/Pan 7/SAdV-24
Human adenovirus G	<i>Mastadenovirus</i>	SAdV-7
Murine adenovirus A	<i>Mastadenovirus</i>	MAdV-1
Fowl adenovirus A	<i>Aviadenovirus</i>	FAdV-1
Fowl adenovirus C	<i>Aviadenovirus</i>	FAdV-10
Fowl adenovirus D	<i>Aviadenovirus</i>	FAdV-9
Fowl adenovirus E	<i>Aviadenovirus</i>	FAdV-8
Ovine adenovirus D	<i>Atadenovirus</i>	OAdV-7

Different species of nonhuman adenovirus are listed, specifying the genus to which they belong and the serotypes from which the vectors have been constructed.

NHAdVs used for vectoring purposes. Mittal and colleagues substituted the nonessential E3 region of BAdV-3 (BAdV B species) with the firefly luciferase reporter gene, and luciferase expression lasted at least 6 days postinfection in human embryonic kidney (HEK)-293 cells (Mittal *et al.*, 1995). BAdV-3 might be a better option to target a specific tissue or organ, since its cell entry is independent of CAR (Bangari *et al.*, 2005b) and other primary receptors of HAdV-5 (Bangari *et al.*, 2005a). BAdV-3 uses sialic acid molecules as a primary receptor to enter into host cells (Li *et al.*, 2009) and it can efficiently transduce different cell types of various species. However, it shows limitations in transducing human cell lines (Rasmussen *et al.*, 1999; Wu and Tikoo, 2004; Bangari *et al.*, 2005b). To circumvent this problem, capsid proteins (fiber and pIX) were genetically modified to increase the ability of BAdV-3 to transduce nonbovine cells (Zakhartchouk *et al.*, 2007). Studies on BAdV-3 vectors *in vivo* biodistribution in a mouse model showed efficient transduction of the heart, kidney, and lung in addition to liver and spleen with a longer duration and higher levels of transgene expression than HAdV-5 vectors (Sharma *et al.*, 2009a). In a mouse model of breast cancer, the biodistribution of BAdV-3 vector was comparable to that of HAdV-5 vectors, the vector had the ability to efficiently transduce tumor and systemic tissues in the presence of HAdV-5 immunity, and it evaded sequestration by Kupffer cells (Tandon *et al.*, 2012). Also, it has been shown that there is minimal cross-neutralization by preexisting anti-HAdV immunity in humans (Bangari *et al.*, 2005b). These results show the suitability of BAdV-3 vectors as a delivery system for cancer gene therapy while circumventing preexisting antivector immunity. Moreover, BAdV-3 and HAdV-5 vectors have a comparable biological behavior in human, bovine, porcine, and mouse cell lines and both genomes persist at high levels in vector-permissive cell lines in the absence of immune pressure, proving BAdV-3 vectors' safety for their use as gene delivery vehicles (Sharma *et al.*, 2011).

Porcine AdV vectors

The idea of using PAdV (genus *Mastadenovirus*, species Porcine A, B, C) vectors was published over 20 years ago (Tuboly *et al.*, 1993). A few years later it was discovered that the virus can enter but is unable to replicate in dog, sheep, bovine, and human cells (Reddy *et al.*, 1999a). Soon after, PAdV-3 (PAdV A species) vectors were constructed with the long-term objective to develop replication-competent PAdV-3 as a live gene transfer vector to induce a mucosal immune response in pigs (Reddy *et al.*, 1999b). Recombinant PAdV-3 is an effective delivery system in pigs (Hammond and Johnson, 2005), and it is often studied together with BAdV-3 vectors, showing similar properties: lack of cross-neutralization with preexisting anti-HAdV immunity in humans (Bangari and Mittal, 2004; Bangari *et al.*, 2005b), CAR- and integrin-independent cell entry (Bangari and Mittal, 2005; Bangari *et al.*, 2005b), similar safety profile and biological characteristics (Sharma *et al.*, 2011), efficient transduction of several cell types of various species with PAdV-3 having an advantage in specific targeting of the breast tissue (Bangari *et al.*, 2005b), and the vector genomes remain as linear episomes (Sharma *et al.*, 2009a). In contrast with BAdV-3 vectors, the genome

levels of PAdV-3 vector were shown to be comparable or lower than those of HAdV-5 vectors *in vivo* in a mouse model, except in the case of the heart, where the levels were comparable or higher (Sharma *et al.*, 2009a). The specificity of PAdV vectors for distinct tissues is very interesting for the design of vectors for targeted gene delivery.

Simian AdV vectors

Simian AdVs are grouped into several human (certain HAdV species contain simian or ape AdVs, based on species-demarcation criteria such as phylogenetic distance and nucleotide composition) and simian AdV species within the genus *Mastadenovirus*. The first gene transfer vector derived from chimpanzee AdV-68 (ChAdV-68/Pan 9/SAHV-25; species HAdV E) was able to grow in HEK293-complementing cells without being neutralized by antibodies against HAdV serotypes (Farina *et al.*, 2001). Since neutralizing seroprevalence to ChAdV-68 in different regions of the world is significantly lower than that to HAdV-5 (Xiang *et al.*, 2006; Ersching *et al.*, 2010; Zhang *et al.*, 2013a), that makes it a good candidate for application in humans. Interestingly, a single surface loop defines a major neutralization site for the ChAdV-68 hexon (Pichla-Gollon *et al.*, 2007), which when mutated, permitted the virus to escape from neutralization by polyclonal antisera obtained from animals *in vitro*. This was actually the first neutralizing site identified for any AdV, which was later very important for the successful application of the vectors in the clinics. Gene delivery by ChAdV-68 is CAR dependent (Cohen *et al.*, 2002), and unlike the *E1* of HAdV-5, the flanking sequences of ChAdV-68 are nonhomologous with cell-derived *E1*, preventing the formation of replication-competent viruses (Xiang *et al.*, 2002). Development of vectors derived from chimpanzee AdVs Pan 5 (ChAdV-5/SAHV-22), Pan 6 (ChAdV-6/SAHV-23), and Pan 7 (ChAdV-7/SAHV-24), members of HAdV E species, started in 2004 (Roy *et al.*, 2004). The vectors transduced skeletal muscle with the same efficiency as HAdV-5 vectors, but without being neutralized by human sera, and the neutralization domains showed to be different in Pan 6 and Pan 7 both *in vitro* and *in vivo*.

One study investigated the importance of neutralizing determinants on capsid proteins by using chimeric vectors with different combinations of Pan 7 and Pan 6 penton base, and hexon and fiber proteins (Roy *et al.*, 2005), reporting that both hexon and fiber have neutralization epitopes and that *in vivo* transduction was more affected by antihexon antibodies, with no effect of antipenton base NAb. Other studies on seroprevalence rates of various ChAdVs were carried out, with Pan 6 and Pan 7 seeming promising as gene therapy vectors although depending on the region where the potential patients live (Jian *et al.*, 2013). A tropism-modified derivative of Pan 7 showed several advantages over HAdV-5 for the use in gene therapy: lower degree of inactivation, diminished liver transduction caused by poor stability of FX-Pan 7 complexes, greater gene delivery efficiency than the wild-type control Pan 7 vector *in vitro*, and higher transduction *in vivo* of the Her2 (human epidermal growth factor receptor type 2)-expressing human tumor cells injected intravenously in mice but not statistically significant compared with the control (Belousova *et al.*, 2010). The last finding indicates that there is a need to improve the vector to

achieve the desired levels of target-specific transduction. Pan 6 vector demonstrated a better capacity than HAdV-5 vector in transduction of brain tumor cells and glioma cells, indicating that it is a promising vector candidate for the brain tumor therapy (Skog *et al.*, 2007). Twenty *E1*-deleted vectors were generated from different AdVs isolated from chimpanzees, bonobos, and gorillas, of which 12 could be propagated, rescued, and expanded in HEK293 cells (Roy *et al.*, 2011b). SAdV-7 (HAdV G species) was proposed for vectoring a few years earlier (Purkayastha *et al.*, 2005b), and was subsequently constructed as an *E1*-deleted vector that could be complemented by HAdV-5 *E1* genes in HEK293 cells (Roy *et al.*, 2011a). The latter observation was surprising since SAdVs from HAdV G species are phylogenetically quite distant from HAdV C members.

Recently, it was shown that humans have low seroreactivity against simian-derived (SAdV-11, -16) or chimpanzee-derived (ChAdV-3, -63) AdV vectors compared with HAdV vectors (rHAdV-5, -28, -35) across multiple geographic regions (Quinn *et al.*, 2013). Chimpanzees and macaques elicited a systemic humoral but not systemic cellular immune response to endogenous AdVs. The structure of the *E3* locus, involved in modulating the host's response to infection, is smaller in HAdVs than in ape AdVs, which may impact the ability of the ape AdVs to evade host immune detection and elimination (Calcedo *et al.*, 2009). ChAdV-68 showed the ability to transduce immature and mature human DCs, followed by secretion of IFN- α and IL-6 but not IL-12 or tumor necrosis factor (TNF)- α , and transduced immature DCs could stimulate proliferation of autologous T lymphocytes (Varnavski *et al.*, 2003). Therefore, ChAdV-68 could be used as a vector for transduction of human DCs.

Fowl AdV vectors

Construction of FAdV (genus *Aviadenovirus*, species FAdV A to FAdV E) vectors started with the demonstration that recombinant FAdV-10 (FAdV C species) could be used to express an antigen to induce a protective immune response in pathogen-free chickens (Sheppard *et al.*, 1998). After the identification of FAdV-9 (species FAdV D) regions that are nonessential for the virus (Ojkic and Nagy, 2001, 2003; Corredor and Nagy, 2010a), FAdV-9 recombinants demonstrated wild-type growth kinetics, virus titers, cytopathic effect and plaque morphology (Corredor and Nagy, 2010a), expression in avian and mammalian cells, and lack of replication in mammalian cells, indicating their applicability as gene delivery vehicles for mammalian systems (Corredor and Nagy, 2010b). FAdV-8 (species FAdV E) vectors were also constructed and showed efficacious *in vivo* delivery of antibody fragments against pathogenic infectious bursal disease virus (Greenall *et al.*, 2010). FAdV-1 (species FAdV A) or CELO (Chicken Embryo Lethal Orphan) vectors have been widely used. The genomic regions that could be deleted were evaluated more than 10 years ago and a cosmid-based strategy was developed for generating CELO vectors (Michou *et al.*, 1999; François *et al.*, 2001). FAdV-1 has several interesting properties: it interacts with CAR and its capsid architecture allows changes in its tropism (Tan *et al.*, 2001), it transduces mammalian cells as efficiently as HAdV-5 vector, and it has a potential for mammalian gene transfer applications, es-

pecially because it has larger DNA packaging capacity and greater physical stability (Michou *et al.*, 1999). It was shown that FAdV-1 is able to deliver p53 transgene into various tissues *in vivo* restoring p53 function in human tumor cell xenografts in mice and leading therefore to the inhibition of tumor growth (Logunov *et al.*, 2004). Moreover, FAdV-1 encoding the human *IL-2* gene can successfully produce biologically active recombinant IL-2 *in vitro*, *in ovo*, and *in vivo*, and it is capable of increasing the median survival time of mice carrying melanoma tumors (Shmarov *et al.*, 2002; Cherenova *et al.*, 2004). Also, recombinant FAdV-1 encoding the HSV-1 thymidine kinase showed cytotoxicity in carcinoma cell lines and suppressed tumor growth and increase median of survival of mice with melanoma tumors (Shashkova *et al.*, 2005), demonstrating its efficacy for the gene delivery to tumor cells *in vitro* and *in vivo*. Furthermore, one study showed that FAdV-1 successfully enhanced expression levels of secreted alkaline phosphatase reporter gene in transduced mammalian cells *in vitro* and *in vivo* (Tutykhina *et al.*, 2008). Finally, PE-Glylated and retargeted with fibroblast growth factor, FAdV-1 showed increased levels of binding and internalization in a variety of human cell lines (Stevenson *et al.*, 2006), with transgene expression being greater than the unmodified version in PC-3 human prostate cells, and being fully resistant to inhibition by human serum *in vitro* (Stevenson *et al.*, 2006). All together indicates that retargeting of CELO virus to human cells via disease-specific receptors is possible, while avoiding preexisting humoral immunity.

Ovine AdV vectors

The production of OAdV (genus *Mastadenovirus*, species OAdV A, B; genus *Atadenovirus*, species OAdV D) vectors started more than 15 years ago (Xu *et al.*, 1997) after determining the nonessential regions within the genome (Vrati *et al.*, 1996). Surprisingly, much larger insertions than expected are tolerated, and portions of sequences can be deleted without affecting virus viability (Xu *et al.*, 1997). The primary receptor for OAdV-7 (OAdV D species) is not CAR, as it was shown that OAdV-7 does not compete with HAdV-5 for the entry into cells (Xu and Both, 1998; Voeks *et al.*, 2002). Recombinant OAdVs can infect a variety of nonovine cells such as rabbit, human, and murine cells but cannot replicate within them (Khatri *et al.*, 1997; Hofmann *et al.*, 1999; Löser *et al.*, 2000; Xu *et al.*, 2000; Kümin *et al.*, 2002; Lockett and Both, 2002). They are not neutralized by polyclonal serum against HAdV-5 (Xu and Both, 1998) and they can overcome preexisting humoral immunity against HAdVs *in vivo* (Hofmann *et al.*, 1999). A cosmid-based system proved useful for efficient generation of recombinant OAdVs and vectors could be rescued in an ovine fetal skin fibroblastic producer cell line (Löser *et al.*, 2003). OAdV-7 carrying purine nucleoside phosphorylase (PNP), which converts the prodrug fludarabine into its activated form 2-fluoroadenine, was effective *in vivo* against prostate cancer progression upon prodrug addition (Voeks *et al.*, 2002; Martiniello-Wilks *et al.*, 2004). Moreover, recombinant OAdV-7 expressing ovalbumin showed efficacy in inducing antitumor response in a mouse model (Tang *et al.*, 2012). OAdVs' biosafety profile and their application as gene delivery vectors were reviewed in the last decade (Both, 2004).

Murine AdV vectors

The production of MAAdV (genus *Mastadenovirus*, species Murine A, B, C) mutants started with the MAAdV-1 (MAAdV A species) E3-deleted recombinant (Beard and Spindler, 1996; Cauthen *et al.*, 1999) and E1-deleted recombinant (Ying *et al.*, 1998). MAAdV-1 presents mouse endothelial cell tropism (Charles *et al.*, 1998; Kajon *et al.*, 1998; Lenaerts *et al.*, 2005), but it can also infect human endothelial cells (Nguyen *et al.*, 1999) and it displays higher affinity for primary human smooth muscle cells than recombinant HAdV-5 (Lenaerts *et al.*, 2009). MAAdV-1 lacks the RGD motif in the penton base, but it contains it in the fiber knob domain (Raman *et al.*, 2009). The RGD motif was reported to play a role in MAAdV-1 infection through α_v integrins, which act as a receptor for the virus, and it was shown that cell surface heparan sulfate glycosaminoglycans are involved in MAAdV-1 infection (Raman *et al.*, 2009). Conversely, the primary attachment of MAAdV-1 is independent of CAR (Lenaerts *et al.*, 2006). Distribution of MAAdV-1 to the liver is markedly lower in intravenously injected immunodeficient mice than that observed with recombinant HAdV-5 (Lenaerts *et al.*, 2009). Moreover, MAAdV-1 has been used as an oncolytic vector, proving that it is a suitable murine homolog model to test the mechanism of action of murine oncolytic AdV vectors in an immunocompetent, tumor-bearing host (Robinson *et al.*, 2009). Furthermore, MAAdV-1 has preference for ovarian carcinoma cell lines, probably because of their increased expression of the enzyme involved in HSPG biosynthesis, therefore having the potential to be used in the oncolytic treatment of ovarian cancer (Lenaerts *et al.*, 2012).

Concluding Remarks and Future Perspectives

The main limitation of clinically applied HAdV-based vectors today is their failure in achieving high levels of efficient transduction in their target tissues. This is primarily because of the preexisting NABs leading to the rapid clearance of circulating vector and cellular immune responses resulting in the elimination of transduced cells, greatly limiting the expression of the transgene. These immunological responses most likely stem from the high frequency of HAdV exposure in multiple human populations across the globe, which results in immunological memory against HAdV serotypes that most of today's vectors are based on. In order to circumvent this, several strategies have been developed for HAdVs, including molecular engineering of chimeric vectors or the use of low-seroprevalent HAdV vectors.

As an alternative, NHAdV vectors have been developed and are being tested in various nonhuman and human cell lines. Most of the NHAdV vectors have two important features that make them more appropriate than HAdV vectors: there is no or very low pathogenicity for their natural host, and there is no presence of preexisting NABs, CD4+ T cells, and CD8+ CTLs against them. However, it should be noted that caution must be applied when using nonhuman primate AdVs, since their ability to infect human cells and their structural similarity could potentially result in anti-HAdV T-cell and/or NAb cross-reactivity with nonhuman primate AdV antigens. Different NHAdVs have specific tropism for different cell types, which makes them applicable in treating different diseases especially when the

targeting of specific cells is required, and therefore they could extend the range of potential target tissues. Furthermore, the inability of NHAdVs to co-replicate with wt HAdVs, as well as the fact that they are replication defective in human cells, makes their vectors much safer than HAdV vectors. Many of the NHAdVs are able to tolerate longer sequences of exogenous DNA than HAdVs. For some of the NHAdVs, there are low-cost propagation systems available, for example, chicken embryos for CELO virus (Laver *et al.*, 1971; Michou *et al.*, 1999). Despite these advantages, there are still many steps necessary before using the NHAdV vectors in the clinics. The first challenge is to find the appropriate cell lines for efficient production. There are also safety issues compared with HAdVs such as the possible occurrence of interspecies adaptation. For example, HAdV-4 is the only HAdV member of HAdV E species that is very similar to SAdVs from that group, indicating its zoonotic origin and molecular adaptation to its new host (Purkayastha *et al.*, 2005a; Dehghan *et al.*, 2013). As AdVs have co-evolved with their hosts (Benkö and Harrach, 2003; Davison *et al.*, 2003), there is a possibility that humans and monkeys could be infected with the same AdVs, a concern when considering chimpanzee or simian AdV vectors for use in humans. Nevertheless, NHAdV vectors present great potential as effective gene therapy vehicles and will hopefully be part of the successful AdV vectors used in the clinics in the coming years.

Acknowledgments

We would like to thank all members of the “ADVance” international training network (Adenoviruses as Clinical Treatments; FP7 ITN, EU grant agreement ref. 290002) and in particular Andrew Baker and Eric J. Kremer for their advice and encouragement to write this review.

Author Disclosure Statement

No competing financial interests exist.

References

- Abbinck, P., Lemckert, A.A., Ewald, B.A., *et al.* (2007). Comparative seroprevalence and immunogenicity of six rare serotype recombinant adenovirus vaccine vectors from subgroups B and D. *J. Virol.* 81, 4654–4663.
- Abe, S., Okuda, K., Ura, T., *et al.* (2009). Adenovirus type 5 with modified hexons induces robust transgene-specific immune responses in mice with pre-existing immunity against adenovirus type 5. *J. Gene Med.* 11, 570–579.
- Albert, M.L., Sauter, B., and Bhardwaj, N. (1998). Dendritic cells acquire antigen from apoptotic cells and induce class I-restricted CTLs. *Nature* 392, 86–89.
- Appledorn, D.M., McBride, A., Seregin, S., *et al.* (2008). Complex interactions with several arms of the complement system dictate innate and humoral immunity to adenoviral vectors. *Gene Ther.* 15, 1606–1617.
- Arnberg, N. (2012). Adenovirus receptors: implications for targeting of viral vectors. *Trends Pharmacol. Sci.* 33, 442–448.
- Bailey, A., and Mautner, V. (1994). Phylogenetic relationships among adenovirus serotypes. *Virology* 205, 438–452.
- Bangari, D.S., and Mittal, S.K. (2004). Porcine adenoviral vectors evade preexisting humoral immunity to adenoviruses

- and efficiently infect both human and murine cells in culture. *Virus Res.* 105, 127–136.
- Bangari, D.S., and Mittal, S.K. (2005). Porcine adenovirus serotype 3 internalization is independent of CAR and alphavbeta3 or alphavbeta5 integrin. *Virology* 332, 157–166.
- Bangari, D.S., and Mittal, S.K. (2006). Development of non-human adenoviruses as vaccine vectors. *Vaccine* 24, 849–862.
- Bangari, D.S., Sharma, A., and Mittal, S.K. (2005a). Bovine adenovirus type 3 internalization is independent of primary receptors of human adenovirus type 5 and porcine adenovirus type 3. *Biochem. Biophys. Res. Commun.* 331, 1478–1484.
- Bangari, D.S., Shukla, S., and Mittal, S.K. (2005b). Comparative transduction efficiencies of human and nonhuman adenoviral vectors in human, murine, bovine, and porcine cells in culture. *Biochem. Biophys. Res. Commun.* 327, 960–966.
- Barlan, A.U., Danthi, P., and Wiethoff, C.M. (2011a). Lysosomal localization and mechanism of membrane penetration influence nonenveloped virus activation of the NLRP3 inflammasome. *Virology* 412, 306–314.
- Barlan, A.U., Griffin, T.M., McGuire, K.A., and Wiethoff, C.M. (2011b). Adenovirus membrane penetration activates the NLRP3 inflammasome. *J. Virol.* 85, 146–155.
- Barouch, D.H., Pau, M.G., Custers, J.H., *et al.* (2004). Immunogenicity of recombinant adenovirus serotype 35 vaccine in the presence of pre-existing anti-Ad5 immunity. *J. Immunol.* 172, 6290–6297.
- Basner-Tschakarjan, E., Gaffal, E., O’Keeffe, M., *et al.* (2006). Adenovirus efficiently transduces plasmacytoid dendritic cells resulting in TLR9-dependent maturation and IFN- α production. *J. Gene Med.* 8, 1300–1306.
- Beard, C.W., and Spindler, K.R. (1996). Analysis of early region 3 mutants of mouse adenovirus type 1. *J. Virol.* 70, 5867–5874.
- Belousova, N., Mikheeva, G., Xiong, C., *et al.* (2010). Development of a targeted gene vector platform based on simian adenovirus serotype 24. *J. Virol.* 84, 10087–10101.
- Benkő, M., and Harrach, B. (2003). Molecular evolution of adenoviruses. *Curr. Top. Microbiol. Immunol.* 272, 3–35.
- Both, G.W. (2004). Ovine atadenovirus: a review of its biology, biosafety profile and application as a gene delivery vector. *Immunol. Cell Biol.* 82, 189–195.
- Bradley, R.R., Lynch, D.M., Iampietro, M.J., *et al.* (2012a). Adenovirus serotype 5 neutralizing antibodies target both hexon and fiber following vaccination and natural infection. *J. Virol.* 86, 625–629.
- Bradley, R.R., Maxfield, L.F., Lynch, D.M., *et al.* (2012b). Adenovirus serotype 5-specific neutralizing antibodies target multiple hexon hypervariable regions. *J. Virol.* 86, 1267–1272.
- Bradshaw, A.C., Coughlan, L., Miller, A.M., *et al.* (2012). Biodistribution and inflammatory profiles of novel penton and hexon double-mutant serotype 5 adenoviruses. *J. Control. Release* 164, 394–402.
- Brouwer, E., Havenga, M.J., Ophorst, O., *et al.* (2007). Human adenovirus type 35 vector for gene therapy of brain cancer: improved transduction and bypass of pre-existing anti-vector immunity in cancer patients. *Cancer Gene Ther.* 14, 211–219.
- Bru, T., Salinas, S., and Kremer, E.J. (2010). An update on canine adenovirus type 2 and its vectors. *Viruses* 2, 2134–2153.
- Bruder, J.T., Semenova, E., Chen, P., *et al.* (2012). Modification of Ad5 hexon hypervariable regions circumvents pre-existing Ad5 neutralizing antibodies and induces protective immune responses. *PLoS One* 7, e33920.
- Calcedo, R., Vandenberghe, L.H., Roy, S., *et al.* (2009). Host immune responses to chronic adenovirus infections in human and nonhuman primates. *J. Virol.* 83, 2623–2631.
- Capone, S., Meola, A., Ercole, B.B., *et al.* (2006). A novel adenovirus type 6 (Ad6)-based hepatitis C virus vector that overcomes preexisting anti-ad5 immunity and induces potent and broad cellular immune responses in rhesus macaques. *J. Virol.* 80, 1688–1699.
- Carlisle, R.C., Di, Y., Cerny, A.M., *et al.* (2009). Human erythrocytes bind and inactivate type 5 adenovirus by presenting Coxsackie virus-adenovirus receptor and complement receptor 1. *Blood* 113, 1909–1918.
- Cauthen, A.N., Brown, C.C., and Spindler, K.R. (1999). *In vitro* and *in vivo* characterization of a mouse adenovirus type 1 early region 3 null mutant. *J. Virol.* 73, 8640–8646.
- Cerullo, V., Seiler, M.P., Mane, V., *et al.* (2007). Toll-like receptor 9 triggers an innate immune response to helper-dependent adenoviral vectors. *Mol. Ther.* 15, 378–385.
- Charles, P.C., Guida, J.D., Brosnan, C.F., and Horwitz, M.S. (1998). Mouse adenovirus type-1 replication is restricted to vascular endothelium in the CNS of susceptible strains of mice. *Virology* 245, 216–228.
- Cheng, C., Gall, J.G., Kong, W.P., *et al.* (2007). Mechanism of ad5 vaccine immunity and toxicity: fiber shaft targeting of dendritic cells. *PLoS Pathog.* 3, e25.
- Cherenova, L.V., Logunov, D.Y., Shashkova, E.V., *et al.* (2004). Recombinant avian adenovirus CELO expressing the human interleukin-2: characterization *in vitro*, *in ovo* and *in vivo*. *Virus Res.* 100, 257–261.
- Chillon, M., and Kremer, E.J. (2001). Trafficking and propagation of canine adenovirus vectors lacking a known integrin-interacting motif. *Hum. Gene Ther.* 12, 1815–1823.
- Cohen, C.J., Xiang, Z.Q., Gao, G.P., *et al.* (2002). Chimpanzee adenovirus CV-68 adapted as a gene delivery vector interacts with the coxsackievirus and adenovirus receptor. *J. Gen. Virol.* 83, 151–155.
- Corredor, J.C., and Nagy, E. (2010a). A region at the left end of the fowl adenovirus 9 genome that is non-essential *in vitro* has consequences *in vivo*. *J. Gen. Virol.* 91, 51–58.
- Corredor, J.C., and Nagy, E. (2010b). The non-essential left end region of the fowl adenovirus 9 genome is suitable for foreign gene insertion/replacement. *Virus Res.* 149, 167–174.
- Coughlan, L., Alba, R., Parker, A.L., *et al.* (2010). Tropism-modification strategies for targeted gene delivery using adenoviral vectors. *Viruses* 2, 2290–2355.
- Coughlan, L., Bradshaw, A.C., Parker, A.L., *et al.* (2012). Ad5:Ad48 hexon hypervariable region substitutions lead to toxicity and increased inflammatory responses following intravenous delivery. *Mol. Ther.* 20, 2268–2281.
- Crawford-Miksza, L., and Schnurr, D.P. (1996). Analysis of 15 adenovirus hexon proteins reveals the location and structure of seven hypervariable regions containing serotype-specific residues. *J. Virol.* 70, 1836–1844.
- Croyle, M.A., Yu, Q.C., and Wilson, J.M. (2000). Development of a rapid method for the PEGylation of adenoviruses with enhanced transduction and improved stability under harsh storage conditions. *Hum. Gene Ther.* 11, 1713–1722.
- Davison, A.J., Benko, M., and Harrach, B. (2003). Genetic content and evolution of adenoviruses. *J. Gen. Virol.* 84, 2895–2908.
- De Geest, B., Snoeys, J., Van Linthout, S., *et al.* (2005). Elimination of innate immune responses and liver inflammation by

- PEGylation of adenoviral vectors and methylprednisolone. *Hum. Gene Ther.* 16, 1439–1451.
- Dehghan, S., Seto, J., Liu, E.B., *et al.* (2013). Computational analysis of four human adenovirus type 4 genomes reveals molecular evolution through two interspecies recombination events. *Virology* 443, 197–207.
- Di Paolo, N.C., Miao, E.A., Iwakura, Y., *et al.* (2009). Virus binding to a plasma membrane receptor triggers interleukin-1 alpha-mediated proinflammatory macrophage response *in vivo*. *Immunity* 31, 110–121.
- Doszpoly, A., Wellehan, J.F., Childress, A.L., *et al.* (2013). Partial characterization of a new adenovirus lineage discovered in testudinoid turtles. *Infect. Genet. Evol.* 17, 106–112.
- Dudley, R.W., Lu, Y., Gilbert, R., *et al.* (2004). Sustained improvement of muscle function one year after full-length dystrophin gene transfer into mdx mice by a gutted helper-dependent adenoviral vector. *Hum. Gene Ther.* 15, 145–156.
- Ehrhardt, A., and Kay, M.A. (2002). A new adenoviral helper-dependent vector results in long-term therapeutic levels of human coagulation factor IX at low doses *in vivo*. *Blood* 99, 3923–3930.
- Ersching, J., Hernandez, M.I., Cezarotto, F.S., *et al.* (2010). Neutralizing antibodies to human and simian adenoviruses in humans and New-World monkeys. *Virology* 407, 1–6.
- Eto, Y., Yoshioka, Y., Ishida, T., *et al.* (2010). Optimized PEGylated adenovirus vector reduces the anti-vector humoral immune response against adenovirus and induces a therapeutic effect against metastatic lung cancer. *Biol. Pharm. Bull.* 33, 1540–1544.
- Farina, S.F., Gao, G.P., Xiang, Z.Q., *et al.* (2001). Replication-defective vector based on a chimpanzee adenovirus. *J. Virol.* 75, 11603–11613.
- Fernandes, P., Peixoto, C., Santiago, V.M., *et al.* (2013a). Bioprocess development for canine adenovirus type 2 vectors. *Gene Ther.* 20, 353–360.
- Fernandes, P., Santiago, V.M., Rodrigues, A.F., *et al.* (2013b). Impact of E1 and Cre on adenovirus vector amplification: developing MDCK CAV-2-E1 and E1-Cre transcomplementing cell lines. *PLoS One* 8, e60342.
- François, A., Eterradossi, N., Delmas, B., *et al.* (2001). Construction of avian adenovirus CELO recombinants in cosmids. *J. Virol.* 75, 5288–5301.
- Gall, J., Kass-Eisler, A., Leinwand, L., and Falck-Pedersen, E. (1996). Adenovirus type 5 and 7 capsid chimera: fiber replacement alters receptor tropism without affecting primary immune neutralization epitopes. *J. Virol.* 70, 2116–2123.
- Gall, J.G., Crystal, R.G., and Falck-Pedersen, E. (1998). Construction and characterization of hexon-chimeric adenoviruses: specification of adenovirus serotype. *J. Virol.* 72, 10260–10264.
- Gao, W., Robbins, P.D., and Gambotto, A. (2003). Human adenovirus type 35: nucleotide sequence and vector development. *Gene Ther.* 10, 1941–1949.
- Garnett, C.T., Erdman, D., Xu, W., and Gooding, L.R. (2002). Prevalence and quantitation of species C adenovirus DNA in human mucosal lymphocytes. *J. Virol.* 76, 10608–10616.
- Gattoni, A., Parlato, A., Vangieri, B., *et al.* (2006). Interferon-gamma: biologic functions and HCV therapy (type I/II) (1 of 2 parts). *Clin. Ther.* 157, 377–386.
- Geisbert, T.W., Bailey, M., Hensley, L., *et al.* (2011). Recombinant adenovirus serotype 26 (Ad26) and Ad35 vaccine vectors bypass immunity to Ad5 and protect nonhuman primates against ebolavirus challenge. *J. Virol.* 85, 4222–4233.
- Ginsberg, H.S. (1996). The ups and downs of adenovirus vectors. *Bull. NY Acad. Med.* 73, 53–58.
- Greber, U.F. (2002). Signalling in viral entry. *Cell. Mol. Life Sci.* 59, 608–626.
- Green, N.K., Herbert, C.W., Hale, S.J., *et al.* (2004). Extended plasma circulation time and decreased toxicity of polymer-coated adenovirus. *Gene Ther.* 11, 1256–1263.
- Greenall, S.A., Tyack, S.G., Johnson, M.A., and Sapats, S.I. (2010). Antibody fragments, expressed by a fowl adenovirus vector, are able to neutralize infectious bursal disease virus. *Avian Pathol.* 39, 339–348.
- Haisma, H.J., Boesjes, M., Beerens, A.M., *et al.* (2009). Scavenger receptor A: a new route for adenovirus 5. *Mol. Pharm.* 6, 366–374.
- Hammond, J.M., and Johnson, M.A. (2005). Porcine adenovirus as a delivery system for swine vaccines and immunotherapeutics. *Vet. J.* 169, 17–27.
- Havenga, M.J., Lemckert, A.A., Grimbergen, J.M., *et al.* (2001). Improved adenovirus vectors for infection of cardiovascular tissues. *J. Virol.* 75, 3335–3342.
- Haviv, Y.S., Blackwell, J.L., Kanerva, A., *et al.* (2002). Adenoviral gene therapy for renal cancer requires retargeting to alternative cellular receptors. *Cancer Res.* 62, 4273–4281.
- Hofherr, S.E., Shashkova, E.V., Weaver, E.A., *et al.* (2008). Modification of adenoviral vectors with polyethylene glycol modulates *in vivo* tissue tropism and gene expression. *Mol. Ther.* 16, 1276–1282.
- Hofmann, C., Löser, P., Cichon, G., *et al.* (1999). Ovine adenovirus vectors overcome preexisting humoral immunity against human adenoviruses *in vivo*. *J. Virol.* 73, 6930–6936.
- Holterman, L., Vogels, R., van der Vlugt, R., *et al.* (2004). Novel replication-incompetent vector derived from adenovirus type 11 (Ad11) for vaccination and gene therapy: low seroprevalence and non-cross-reactivity with Ad5. *J. Virol.* 78, 13207–13215.
- Ibanes, S., and Kremer, E.J. (2013). Canine adenovirus type 2 vector generation via I-SceI-mediated intracellular genome release. *PLoS One* 8, e71032.
- Jankovic, D., Liu, Z., and Gause, W.C. (2001). Th1- and Th2-cell commitment during infectious disease: asymmetry in divergent pathways. *Trends Immunol.* 22, 450–457.
- Jian, L., Zhao, Q., Zhang, S., *et al.* (2013). The prevalence of neutralising antibodies to chimpanzee adenovirus type 6 and type 7 in healthy adult volunteers, patients with chronic hepatitis B and patients with primary hepatocellular carcinoma in China. *Arch. Virol.* DOI 10.1007/s00705-013-1828-y. [Epub ahead of print]
- Jones, M.S., Harrach, B., Ganac, R.D., *et al.* (2007). New adenovirus species found in a patient presenting with gastroenteritis. *J. Virol.* 81, 5978–5984.
- Kajon, A.E., Brown, C.C., and Spindler, K.R. (1998). Distribution of mouse adenovirus type 1 in intraperitoneally and intranasally infected adult outbred mice. *J. Virol.* 72, 1219–1223.
- Kanerva, A., Mikheeva, G.V., Krasnykh, V., *et al.* (2002). Targeting adenovirus to the serotype 3 receptor increases gene transfer efficiency to ovarian cancer cells. *Clin. Cancer Res.* 8, 275–280.
- Kass-Eisler, A., Leinwand, L., Gall, J., *et al.* (1996). Circumventing the immune response to adenovirus-mediated gene therapy. *Gene Ther.* 3, 154–162.
- Katze, M.G., He, Y., and Gale, M. (2002). Viruses and interferon: a fight for supremacy. *Nat. Rev. Immunol.* 2, 675–687.

- Khare, R., Reddy, V.S., Nemerow, G.R., and Barry, M.A. (2012). Identification of adenovirus serotype 5 hexon regions that interact with scavenger receptors. *J. Virol.* 86, 2293–2301.
- Khatri, A., Xu, Z.Z., and Both, G.W. (1997). Gene expression by atypical recombinant ovine adenovirus vectors during abortive infection of human and animal cells *in vitro*. *Virology* 239, 226–237.
- Klonjkowski, B., Gilardi-Hebenstreit, P., Hadchouel, J., *et al.* (1997). A recombinant E1-deleted canine adenoviral vector capable of transduction and expression of a transgene in human-derived cells and *in vivo*. *Hum. Gene Ther.* 8, 2103–2115.
- Kochanek, S., Clemens, P.R., Mitani, K., *et al.* (1996). A new adenoviral vector: replacement of all viral coding sequences with 28 kb of DNA independently expressing both full-length dystrophin and beta-galactosidase. *Proc. Natl. Acad. Sci. USA* 93, 5731–5736.
- Kremer, E.J., Boutin, S., Chillon, M., and Danos, O. (2000). Canine adenovirus vectors: an alternative for adenovirus-mediated gene transfer. *J. Virol.* 74, 505–512.
- Kümin, D., Hofmann, C., Rudolph, M., *et al.* (2002). Biology of ovine adenovirus infection of nonpermissive cells. *J. Virol.* 76, 10882–10893.
- Kuriyama, S., Tominaga, K., Kikukawa, M., *et al.* (1998). Inhibitory effects of human sera on adenovirus-mediated gene transfer into rat liver. *Anticancer Res.* 18, 2345–2351.
- Lau, A.A., Rozaklis, T., Ibanes, S., *et al.* (2012). Helper-dependent canine adenovirus vector-mediated transgene expression in a neurodegenerative lysosomal storage disorder. *Gene* 491, 53–57.
- Laver, W.G., Youngusband, H.B., and Wrigley, N.G. (1971). Purification and properties of chick embryo lethal orphan virus (an avian adenovirus). *Virology* 45, 598–614.
- Leen, A.M., Sili, U., Vanin, E.F., *et al.* (2004). Conserved CTL epitopes on the adenovirus hexon protein expand subgroup cross-reactive and subgroup-specific CD8+ T cells. *Blood* 104, 2432–2440.
- Lemckert, A.A., Grimbergen, J., Smits, S., *et al.* (2006). Generation of a novel replication-incompetent adenoviral vector derived from human adenovirus type 49: manufacture on PER.C6 cells, tropism and immunogenicity. *J. Gen. Virol.* 87, 2891–2899.
- Lenaerts, L., Verbeken, E., De Clercq, E., and Naesens, L. (2005). Mouse adenovirus type 1 infection in SCID mice: an experimental model for antiviral therapy of systemic adenovirus infections. *Antimicrob. Agents Chemother.* 49, 4689–4699.
- Lenaerts, L., Daelemans, D., Geukens, N., *et al.* (2006). Mouse adenovirus type 1 attachment is not mediated by the coxsackie-adenovirus receptor. *FEBS Lett.* 580, 3937–3942.
- Lenaerts, L., McVey, J.H., Baker, A.H., *et al.* (2009). Mouse adenovirus type 1 and human adenovirus type 5 differ in endothelial cell tropism and liver targeting. *J. Gene. Med.* 11, 119–127.
- Lenaerts, L., van Dam, W., Persoons, L., and Naesens, L. (2012). Interaction between mouse adenovirus type 1 and cell surface heparan sulfate proteoglycans. *PLoS One* 7, e31454.
- Li, X., Bangari, D.S., Sharma, A., and Mittal, S.K. (2009). Bovine adenovirus serotype 3 utilizes sialic acid as a cellular receptor for virus entry. *Virology* 392, 162–168.
- Liu, Q., White, L.R., Clark, S.A., *et al.* (2005). Akt/protein kinase B activation by adenovirus vectors contributes to NFkappaB-dependent CXCL10 expression. *J. Virol.* 79, 14507–14515.
- Lockett, L.J., and Both, G.W. (2002). Complementation of a defective human adenovirus by an otherwise incompatible ovine adenovirus recombinant carrying a functional E1A gene. *Virology* 294, 333–341.
- Logunov, D.Y., Ilyinskaya, G.V., Cherenova, L.V., *et al.* (2004). Restoration of p53 tumor-suppressor activity in human tumor cells *in vitro* and in their xenografts *in vivo* by recombinant avian adenovirus CELO-p53. *Gene Ther.* 11, 79–84.
- Löser, P., Hillgenberg, M., Arnold, W., *et al.* (2000). Ovine adenovirus vectors mediate efficient gene transfer to skeletal muscle. *Gene Ther.* 7, 1491–1498.
- Löser, P., Hofmann, C., Both, G.W., *et al.* (2003). Construction, rescue, and characterization of vectors derived from ovine adenovirus. *J. Virol.* 77, 11941–11951.
- Lukashev, A.N., Ivanova, O.E., Eremeeva, T.P., and Iggo, R.D. (2008). Evidence of frequent recombination among human adenoviruses. *J. Gen. Virol.* 89, 380–388.
- Mack, C.A., Song, W.R., Carpenter, H., *et al.* (1997). Circumvention of anti-adenovirus neutralizing immunity by administration of an adenoviral vector of an alternate serotype. *Hum. Gene Ther.* 8, 99–109.
- Maione, D., Della Rocca, C., Giannetti, P., *et al.* (2001). An improved helper-dependent adenoviral vector allows persistent gene expression after intramuscular delivery and overcomes preexisting immunity to adenovirus. *Proc. Natl. Acad. Sci. USA* 98, 5986–5991.
- Martiniello-Wilks, R., Dane, A., Voeks, D.J., *et al.* (2004). Gene-directed enzyme prodrug therapy for prostate cancer in a mouse model that imitates the development of human disease. *J. Gene Med.* 6, 43–54.
- Mastrangeli, A., Harvey, B.G., Yao, J., *et al.* (1996). “Sero-switch” adenovirus-mediated *in vivo* gene transfer: circumvention of anti-adenovirus humoral immune defenses against repeat adenovirus vector administration by changing the adenovirus serotype. *Hum. Gene Ther.* 7, 79–87.
- McVey, D., Zuber, M., ETTYREDDY, D., *et al.* (2010). Characterization of human adenovirus 35 and derivation of complex vectors. *Virology* 405, 267–276.
- Michou, A.I., Lehrmann, H., Saltik, M., and Cotten, M. (1999). Mutational analysis of the avian adenovirus CELO, which provides a basis for gene delivery vectors. *J. Virol.* 73, 1399–1410.
- Mittal, S.K., Prevec, L., Graham, F.L., and Babiuk, L.A. (1995). Development of a bovine adenovirus type 3-based expression vector. *J. Gen. Virol.* 76 (Pt 1), 93–102.
- Mok, H., Palmer, D.J., Ng, P., and Barry, M.A. (2005). Evaluation of polyethylene glycol modification of first-generation and helper-dependent adenoviral vectors to reduce innate immune responses. *Mol. Ther.* 11, 66–79.
- Morrison, J., Briggs, S.S., Green, N., *et al.* (2008). Virotherapy of ovarian cancer with polymer-cloaked adenovirus re-targeted to the epidermal growth factor receptor. *Mol. Ther.* 16, 244–251.
- Morrison, J., Briggs, S.S., Green, N.K., *et al.* (2009). Cetuximab retargeting of adenovirus via the epidermal growth factor receptor for treatment of intraperitoneal ovarian cancer. *Hum. Gene Ther.* 20, 239–251.
- Muruve, D.A., Pétrilli, V., Zaiss, A.K., *et al.* (2008). The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature* 452, 103–107.

- Myhre, S., Henning, P., Granio, O., *et al.* (2007). Decreased immune reactivity towards a knobless, affibody-targeted adenovirus type 5 vector. *Gene Ther.* 14, 376–381.
- Nan, X., Peng, B., Hahn, T.W., *et al.* (2003). Development of an Ad7 cosmid system and generation of an Ad7-deltaE1deltaE3HIV(MN) env/rev recombinant virus. *Gene Ther.* 10, 326–336.
- Nguyen, T., Nery, J., Joseph, S., *et al.* (1999). Mouse adenovirus (MAV-1) expression in primary human endothelial cells and generation of a full-length infectious plasmid. *Gene Ther.* 6, 1291–1297.
- Ni, S., Bernt, K., Gaggari, A., *et al.* (2005). Evaluation of bio-distribution and safety of adenovirus vectors containing group B fibers after intravenous injection into baboons. *Hum. Gene Ther.* 16, 664–677.
- Nicol, C.G., Graham, D., Miller, W.H., *et al.* (2004). Effect of adenovirus serotype 5 fiber and penton modifications on *in vivo* tropism in rats. *Mol. Ther.* 10, 344–354.
- Ojkic, D., and Nagy, E. (2001). The long repeat region is dispensable for fowl adenovirus replication *in vitro*. *Virology* 283, 197–206.
- Ojkic, D., and Nagy, E. (2003). Antibody response and virus tissue distribution in chickens inoculated with wild-type and recombinant fowl adenoviruses. *Vaccine* 22, 42–48.
- Olive, M., Eisenlohr, L., Flomenberg, N., *et al.* (2002). The adenovirus capsid protein hexon contains a highly conserved human CD4+ T-cell epitope. *Hum. Gene Ther.* 13, 1167–1178.
- Ord, E., Shirley, R., McClure, J., *et al.* (2013). Combined antiapoptotic and antioxidant approach to acute neuroprotection for stroke in hypertensive rats. *J. Cereb. Blood Flow Metab.* 33, 1215–1224.
- Ostapchuk, P., and Hearing, P. (2001). Pseudopackaging of adenovirus type 5 genomes into capsids containing the hexon proteins of adenovirus serotypes B, D, or E. *J. Virol.* 75, 45–51.
- Pahl, J.H., Verhoeven, D.H., Kwappenberg, K.M., *et al.* (2012). Adenovirus type 35, but not type 5, stimulates NK cell activation via plasmacytoid dendritic cells and TLR9 signaling. *Mol. Immunol.* 51, 91–100.
- Parker, A.L., Fisher, K.D., Oupicky, D., *et al.* (2005). Enhanced gene transfer activity of peptide-targeted gene-delivery vectors. *J. Drug Target.* 13, 39–51.
- Parker, A.L., Waddington, S.N., Nicol, C.G., *et al.* (2006). Multiple vitamin K-dependent coagulation zymogens promote adenovirus-mediated gene delivery to hepatocytes. *Blood* 108, 2554–2561.
- Parker, A.L., Waddington, S.N., Buckley, S.M., *et al.* (2009). Effect of neutralizing sera on factor x-mediated adenovirus serotype 5 gene transfer. *J. Virol.* 83, 479–483.
- Parks, R., Eveleigh, C., and Graham, F. (1999). Use of helper-dependent adenoviral vectors of alternative serotypes permits repeat vector administration. *Gene Ther.* 6, 1565–1573.
- Perreau, M., and Kremer, E.J. (2006). The conundrum between immunological memory to adenovirus and their use as vectors in clinical gene therapy. *Mol. Biotechnol.* 34, 247–256.
- Perreau, M., Welles, H.C., Pellaton, C., *et al.* (2012). The number of Toll-like receptor 9-agonist motifs in the adenovirus genome correlates with induction of dendritic cell maturation by adenovirus immune complexes. *J. Virol.* 86, 6279–6285.
- Pichla-Gollon, S.L., Drinker, M., Zhou, X., *et al.* (2007). Structure-based identification of a major neutralizing site in an adenovirus hexon. *J. Virol.* 81, 1680–1689.
- Piersanti, S., Astrologo, L., Licursi, V., *et al.* (2013). Differentiated neuroprogenitor cells incubated with human or canine adenovirus, or lentiviral vectors have distinct transcriptome profiles. *Plos One* 8, e69808.
- Purkayastha, A., Ditty, S.E., Su, J., *et al.* (2005a). Genomic and bioinformatics analysis of HAdV-4, a human adenovirus causing acute respiratory disease: implications for gene therapy and vaccine vector development. *J. Virol.* 79, 2559–2572.
- Purkayastha, A., Su, J., Carlisle, S., *et al.* (2005b). Genomic and bioinformatics analysis of HAdV-7, a human adenovirus of species B1 that causes acute respiratory disease: implications for vector development in human gene therapy. *Virology* 332, 114–129.
- Quinn, K.M., Da Costa, A., Yamamoto, A., *et al.* (2013). Comparative analysis of the magnitude, quality, phenotype, and protective capacity of simian immunodeficiency virus gag-specific CD8+ T cells following human-, simian-, and chimpanzee-derived recombinant adenoviral vector immunization. *J. Immunol.* 190, 2720–2735.
- Raman, S., Hsu, T.H., Ashley, S.L., and Spindler, K.R. (2009). Usage of integrin and heparan sulfate as receptors for mouse adenovirus type 1. *J. Virol.* 83, 2831–2838.
- Raper, S.E., Chirmule, N., Lee, F.S., *et al.* (2003). Fatal systemic inflammatory response syndrome in a ornithine transcarbamylase deficient patient following adenoviral gene transfer. *Mol. Genet. Metab.* 80, 148–158.
- Rasmussen, U.B., Benchaibi, M., Meyer, V., *et al.* (1999). Novel human gene transfer vectors: evaluation of wild-type and recombinant animal adenoviruses in human-derived cells. *Hum. Gene Ther.* 10, 2587–2599.
- Reddy, P.S., Idamakanti, N., Babiuk, L.A., *et al.* (1999a). Porcine adenovirus-3 as a helper-dependent expression vector. *J. Gen. Virol.* 80 (Pt 11), 2909–2916.
- Reddy, P.S., Idamakanti, N., Hyun, B.H., *et al.* (1999b). Development of porcine adenovirus-3 as an expression vector. *J. Gen. Virol.* 80 (Pt 3), 563–570.
- Roberts, D.M., Nanda, A., Havenga, M.J., *et al.* (2006). Hexon-chimaeric adenovirus serotype 5 vectors circumvent pre-existing anti-vector immunity. *Nature* 441, 239–243.
- Robinson, M., Li, B., Ge, Y., *et al.* (2009). Novel immunocompetent murine tumor model for evaluation of conditionally replication-competent (oncolytic) murine adenoviral vectors. *J. Virol.* 83, 3450–3462.
- Rogée, S., Grellier, E., Bernard, C., *et al.* (2010). Influence of chimeric human-bovine fibers on adenoviral uptake by liver cells and the antiviral immune response. *Gene Ther.* 17, 880–891.
- Rogozhin, V.N., Logunov, D.Y., Shchebliakov, D.V., *et al.* (2011). An efficient method for the delivery of the interleukin-2 gene to human hematopoietic cells using the fiber-modified recombinant adenovirus. *Acta Nat.* 3, 100–106.
- Romanczuk, H., Galer, C.E., Zabner, J., *et al.* (1999). Modification of an adenoviral vector with biologically selected peptides: a novel strategy for gene delivery to cells of choice. *Hum. Gene Ther.* 10, 2615–2626.
- Rowe, W.P., Huebner, R.J., Gilmore, L.K., *et al.* (1953). Isolation of a cytopathogenic agent from human adenoids undergoing spontaneous degeneration in tissue culture. *Proc. Soc. Exp. Biol. Med.* 84, 570–573.
- Roy, S., Shirley, P.S., McClelland, A., and Kaleko, M. (1998). Circumvention of immunity to the adenovirus major coat protein hexon. *J. Virol.* 72, 6875–6879.
- Roy, S., Gao, G.P., Lu, Y., *et al.* (2004). Characterization of a family of chimpanzee adenoviruses and development of

- molecular clones for gene transfer vectors. *Hum. Gene Ther.* 15, 519–530.
- Roy, S., Clawson, D.S., Calcedo, R., *et al.* (2005). Use of chimeric adenoviral vectors to assess capsid neutralization determinants. *Virology* 333, 207–214.
- Roy, S., Clawson, D.S., Adam, V.S., *et al.* (2011a). Construction of gene transfer vectors based on simian adenovirus 7. *J. Gen. Virol.* 92, 1749–1753.
- Roy, S., Medina-Jaszek, A., Wilson, M.J., *et al.* (2011b). Creation of a panel of vectors based on ape adenovirus isolates. *J. Gene Med.* 13, 17–25.
- Rux, J.J., and Burnett, R.M. (2000). Type-specific epitope locations revealed by X-ray crystallographic study of adenovirus type 5 hexon. *Mol. Ther.* 1, 18–30.
- Sakurai, F. (2008). Development of a replication-incompetent adenovirus vector derived from subgroup B adenovirus serotype 35. *Yakuga. Zasshi* 128, 1751–1761.
- Sakurai, F., Mizuguchi, H., and Hayakawa, T. (2003). Efficient gene transfer into human CD34+ cells by an adenovirus type 35 vector. *Gene Ther.* 10, 1041–1048.
- Sakurai, F., Nakamura, S.I., Akitomo, K., *et al.* (2009). Adenovirus serotype 35 vector-mediated transduction following direct administration into organs of nonhuman primates. *Gene Ther.* 16, 297–302.
- Schoehn, G., El Bakkouri, M., Fabry, C.M., *et al.* (2008). Three-dimensional structure of canine adenovirus serotype 2 capsid. *J. Virol.* 82, 3192–3203.
- Schoggins, J.W., Nociari, M., Philpott, N., and Falck-Pedersen, E. (2005). Influence of fiber detargeting on adenovirus-mediated innate and adaptive immune activation. *J. Virol.* 79, 11627–11637.
- Schumacher, L., Ribas, A., Dissette, V.B., *et al.* (2004). Human dendritic cell maturation by adenovirus transduction enhances tumor antigen-specific T-cell responses. *J. Immunother.* 27, 191–200.
- Segura, M.M., Puig, M., Monfar, M., and Chillón, M. (2012). Chromatography purification of canine adenoviral vectors. *Hum. Gene Ther. Methods* 23, 182–197.
- Seiradake, E., Henaff, D., Wodrich, H., *et al.* (2009). The cell adhesion molecule “CAR” and sialic acid on human erythrocytes influence adenovirus *in vivo* biodistribution. *PLoS Pathog.* 5, e1000277.
- Seshidhar Reddy, P., Ganesh, S., Limbach, M.P., *et al.* (2003). Development of adenovirus serotype 35 as a gene transfer vector. *Virology* 311, 384–393.
- Sharma, A., Bangari, D.S., Tandon, M., *et al.* (2009a). Comparative analysis of vector biodistribution, persistence and gene expression following intravenous delivery of bovine, porcine and human adenoviral vectors in a mouse model. *Virology* 386, 44–54.
- Sharma, A., Li, X., Bangari, D.S., and Mittal, S.K. (2009b). Adenovirus receptors and their implications in gene delivery. *Virus Res.* 143, 184–194.
- Sharma, A., Bangari, D.S., Vemula, S.V., and Mittal, S.K. (2011). Persistence and the state of bovine and porcine adenoviral vector genomes in human and nonhuman cell lines. *Virus Res.* 161, 181–187.
- Shashkova, E.V., Cherenova, L.V., Kazansky, D.B., and Doronin, K. (2005). Avian adenovirus vector CELO-TK displays anticancer activity in human cancer cells and suppresses established murine melanoma tumors. *Cancer Gene Ther.* 12, 617–626.
- Shayakhmetov, D.M., Papayannopoulou, T., Stamatoyannopoulos, G., and Lieber, A. (2000). Efficient gene transfer into human CD34(+) cells by a retargeted adenovirus vector. *J. Virol.* 74, 2567–2583.
- Shayakhmetov, D.M., Gaggari, A., Ni, S., *et al.* (2005). Adenovirus binding to blood factors results in liver cell infection and hepatotoxicity. *J. Virol.* 79, 7478–7491.
- Sheppard, M., Werner, W., Tsatas, E., *et al.* (1998). Fowl adenovirus recombinant expressing VP2 of infectious bursal disease virus induces protective immunity against bursal disease. *Arch. Virol.* 143, 915–930.
- Shimizu, T., Ichihara, M., Yoshioka, Y., *et al.* (2012). Intravenous administration of polyethylene glycol-coated (PEGylated) proteins and PEGylated adenovirus elicits an anti-PEG immunoglobulin M response. *Biol. Pharm. Bull.* 35, 1336–1342.
- Shiratsuchi, T., Rai, U., Krause, A., *et al.* (2010). Replacing adenoviral vector HVR1 with a malaria B cell epitope improves immunogenicity and circumvents preexisting immunity to adenovirus in mice. *J. Clin. Invest.* 120, 3688–3701.
- Shiver, J.W., and Emini, E.A. (2004). Recent advances in the development of HIV-1 vaccines using replication-incompetent adenovirus vectors. *Annu. Rev. Med.* 55, 355–372.
- Shmarov, M.M., Cherenova, L.V., Shashkova, E.V., *et al.* (2002). Eukaryotic vectors of Celo avian adenovirus genome, carrying GFP and human IL-2 genes. *Mol. Gen. Mikrobiol. Virusol.* 2, 30–35.
- Skog, J., Edlund, K., Bergenheim, A.T., and Wadell, G. (2007). Adenoviruses 16 and CV23 efficiently transduce human low-passage brain tumor and cancer stem cells. *Mol. Ther.* 15, 2140–2145.
- Soudais, C., Boutin, S., Hong, S.S., *et al.* (2000). Canine adenovirus type 2 attachment and internalization: coxsackievirus-adenovirus receptor, alternative receptors, and an RGD-independent pathway. *J. Virol.* 74, 10639–10649.
- Soudais, C., Laplace-Builhe, C., Kissa, K., and Kremer, E.J. (2001). Preferential transduction of neurons by canine adenovirus vectors and their efficient retrograde transport *in vivo*. *FASEB J.* 15, 2283–2285.
- Stephen, S.L., Montini, E., Sivanandam, V.G., *et al.* (2010). Chromosomal integration of adenoviral vector DNA *in vivo*. *J. Virol.* 84, 9987–9994.
- Stevenson, M., Boos, E., Herbert, C., *et al.* (2006). Chick embryo lethal orphan virus can be polymer-coated and retargeted to infect mammalian cells. *Gene Ther.* 13, 356–368.
- Stevenson, M., Hale, A.B., Hale, S.J., *et al.* (2007). Incorporation of a laminin-derived peptide (SIKVAV) on polymer-modified adenovirus permits tumor-specific targeting via alpha6-integrins. *Cancer Gene Ther.* 14, 335–345.
- Stone, D., Ni, S., Li, Z.Y., *et al.* (2005). Development and assessment of human adenovirus type 11 as a gene transfer vector. *J. Virol.* 79, 5090–5104.
- Sullivan, N.J., Geisbert, T.W., Geisbert, J.B., *et al.* (2003). Accelerated vaccination for Ebola virus haemorrhagic fever in non-human primates. *Nature* 424, 681–684.
- Sumida, S.M., Truitt, D.M., Lemckert, A.A., *et al.* (2005). Neutralizing antibodies to adenovirus serotype 5 vaccine vectors are directed primarily against the adenovirus hexon protein. *J. Immunol.* 174, 7179–7185.
- Tamanini, A., Nicolis, E., Bonizzato, A., *et al.* (2006). Interaction of adenovirus type 5 fiber with the coxsackievirus and adenovirus receptor activates inflammatory response in human respiratory cells. *J. Virol.* 80, 11241–11254.
- Tan, P.K., Michou, A.I., Bergelson, J.M., and Cotten, M. (2001). Defining CAR as a cellular receptor for the avian

- adenovirus CELO using a genetic analysis of the two viral fibre proteins. *J. Gen. Virol.* 82, 1465–1472.
- Tandon, M., Sharma, A., Vemula, S.V., *et al.* (2012). Sequential administration of bovine and human adenovirus vectors to overcome vector immunity in an immunocompetent mouse model of breast cancer. *Virus Res.* 163, 202–211.
- Tang, J., Olive, M., Pulmanusahakul, R., *et al.* (2006). Human CD8+ cytotoxic T cell responses to adenovirus capsid proteins. *Virology* 350, 312–322.
- Tang, R., Li, K., Wilson, M., *et al.* (2012). Potent antitumor immunity in mice induced by vaccination with an ovine adenovirus vector. *J. Immunother.* 35, 32–41.
- Tarassishin, L., Szawlowski, P., Kidd, A.H., and Russell, W.C. (2000). An epitope on the adenovirus fibre tail is common to all human subgroups. *Arch. Virol.* 145, 805–811.
- Tian, X., Su, X., Li, H., *et al.* (2011). Construction and characterization of human adenovirus serotype 3 packaged by serotype 7 hexon. *Virus Res.* 160, 214–220.
- Tomko, R.P., Xu, R., and Philipson, L. (1997). HCAR and MCAR: the human and mouse cellular receptors for subgroup C adenoviruses and group B coxsackieviruses. *Proc. Natl. Acad. Sci. USA* 94, 3352–3356.
- Tripathy, S.K., Black, H.B., Goldwasser, E., and Leiden, J.M. (1996). Immune responses to transgene-encoded proteins limit the stability of gene expression after injection of replication-defective adenovirus vectors. *Nat. Med.* 2, 545–550.
- Tuboly, T., Nagy, E., and Derbyshire, J.B. (1993). Potential viral vectors for the stimulation of mucosal antibody responses against enteric viral antigens in pigs. *Res. Vet. Sci.* 54, 345–350.
- Tutykhina, I.L., Shmarov, M.M., Logunov, D.I., *et al.* (2008). Construction of the vector based on the CELO avian adenovirus genome providing enhanced expression of secreted alkaline phosphatase gene in a non-permissive system *in vitro* and *in vivo*. *Mol. Gen. Mikrobiol. Virusol.* 4, 26–30.
- Ulasov, I.V., Rivera, A.A., Han, Y., *et al.* (2007). Targeting adenovirus to CD80 and CD86 receptors increases gene transfer efficiency to malignant glioma cells. *J. Neurosurg.* 107, 617–627.
- Varnavski, A.N., Schlienger, K., Bergelson, J.M., *et al.* (2003). Efficient transduction of human monocyte-derived dendritic cells by chimpanzee-derived adenoviral vector. *Hum. Gene Ther.* 14, 533–544.
- Vellinga, J., Van der Heijdt, S., and Hoeben, R.C. (2005). The adenovirus capsid: major progress in minor proteins. *J. Gen. Virol.* 86, 1581–1588.
- Voeks, D., Martiniello-Wilks, R., Madden, V., *et al.* (2002). Gene therapy for prostate cancer delivered by ovine adenovirus and mediated by purine nucleoside phosphorylase and fludarabine in mouse models. *Gene Ther.* 9, 759–768.
- Vogels, R., Zuijdgheest, D., van Rijnsoever, R., *et al.* (2003). Replication-deficient human adenovirus type 35 vectors for gene transfer and vaccination: efficient human cell infection and bypass of preexisting adenovirus immunity. *J. Virol.* 77, 8263–8271.
- Volk, A.L., Rivera, A.A., Kanerva, A., *et al.* (2003). Enhanced adenovirus infection of melanoma cells by fiber-modification: incorporation of RGD peptide or Ad5/3 chimerism. *Cancer Biol. Ther.* 2, 511–515.
- Vrati, S., Macavoy, E.S., Xu, Z.Z., *et al.* (1996). Construction and transfection of ovine adenovirus genomic clones to rescue modified viruses. *Virology* 220, 200–203.
- Vujanovic, L., Whiteside, T.L., Potter, D.M., *et al.* (2009). Regulation of antigen presentation machinery in human dendritic cells by recombinant adenovirus. *Cancer Immunol. Immunother.* 58, 121–133.
- Walsh, M.P., Seto, J., Liu, E.B., *et al.* (2011). Computational analysis of two species C human adenoviruses provides evidence of a novel virus. *J. Clin. Microbiol.* 49, 3482–3490.
- Wang, I.J., Jhuang, M.C., Chen, Y.H., *et al.* (2010). Chitosan modification of adenovirus to modify transfection efficiency in bovine corneal epithelial cells. *PLoS One* 5, e12085.
- Wang, C.H., Chan, L.W., Johnson, R.N., *et al.* (2011a). The transduction of Coxsackie and Adenovirus Receptor-negative cells and protection against neutralizing antibodies by HPMMA-co-oligolysine copolymer-coated adenovirus. *Biomaterials* 32, 9536–9545.
- Wang, D.Y., Liu, S.H., Li, X., *et al.* (2011b). Study on construction of chimeric adenovirus vector Ad5/11 carrying human eGFP and endostatin-K5 and its experimental investigation *in vitro*. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 27, 143–145.
- Wohlfart, C. (1988). Neutralization of adenoviruses: kinetics, stoichiometry, and mechanisms. *J. Virol.* 62, 2321–2328.
- Wu, Q., and Tikoo, S.K. (2004). Altered tropism of recombinant bovine adenovirus type-3 expressing chimeric fiber. *Virus Res.* 99, 9–15.
- Wu, H., Dmitriev, I., Kashentseva, E., *et al.* (2002). Construction and characterization of adenovirus serotype 5 packaged by serotype 3 hexon. *J. Virol.* 76, 12775–12782.
- Xiang, Z., Gao, G., Reyes-Sandoval, A., *et al.* (2002). Novel chimpanzee serotype 68-based adenoviral vaccine carrier for induction of antibodies to a transgene product. *J. Virol.* 76, 2667–2675.
- Xiang, Z., Li, Y., Cun, A., *et al.* (2006). Chimpanzee adenovirus antibodies in humans, sub-Saharan Africa. *Emerg. Infect. Dis.* 12, 1596–1599.
- Xu, Z.Z., and Both, G.W. (1998). Altered tropism of an ovine adenovirus carrying the fiber protein cell binding domain of human adenovirus type 5. *Virology* 248, 156–163.
- Xu, Z.Z., Hyatt, A., Boyle, D.B., and Both, G.W. (1997). Construction of ovine adenovirus recombinants by gene insertion or deletion of related terminal region sequences. *Virology* 230, 62–71.
- Xu, Z.Z., Nevels, M., MacAvoy, E.S., *et al.* (2000). An ovine adenovirus vector lacks transforming ability in cells that are transformed by AD5 E1A/B sequences. *Virology* 270, 162–172.
- Xu, Z., Tian, J., Smith, J.S., and Byrnes, A.P. (2008). Clearance of adenovirus by Kupffer cells is mediated by scavenger receptors, natural antibodies, and complement. *J. Virol.* 82, 11705–11713.
- Xu, Z., Qiu, Q., Tian, J., *et al.* (2013). Coagulation factor X shields adenovirus type 5 from attack by natural antibodies and complement. *Nat. Med.* 19, 452–457.
- Yamaguchi, T., Kawabata, K., Koizumi, N., *et al.* (2007). Role of MyD88 and TLR9 in the innate immune response elicited by serotype 5 adenoviral vectors. *Hum. Gene Ther.* 18, 753–762.
- Yao, X., Yoshioka, Y., Morishige, T., *et al.* (2010). Adenovirus vector covalently conjugated to polyethylene glycol with a cancer-specific promoter suppresses the tumor growth through systemic administration. *Biol. Pharm. Bull.* 33, 1073–1076.
- Ying, B., Smith, K., and Spindler, K.R. (1998). Mouse adenovirus type 1 early region 1A is dispensable for growth in cultured fibroblasts. *J. Virol.* 72, 6325–6331.
- Youil, R., Toner, T.J., Su, Q., *et al.* (2002). Hexon gene switch strategy for the generation of chimeric recombinant adenovirus. *Hum. Gene Ther.* 13, 311–320.

- Yu, B., Dong, J., Wang, C., *et al.* (2013). Characteristics of neutralizing antibodies to adenovirus capsid proteins in human and animal sera. *Virology* 437, 118–123.
- Zaiss, A.K., Vilaysane, A., Cotter, M.J., *et al.* (2009). Antiviral antibodies target adenovirus to phagolysosomes and amplify the innate immune response. *J. Immunol.* 182, 7058–7068.
- Zakhartchouk, A.N., Wu, Q., and Tikoo, S.K. (2007). Construction of capsid-modified recombinant bovine adenovirus type 3. *Methods Mol. Med.* 130, 91–106.
- Zeng, Q., Han, J., Zhao, D., *et al.* (2012). Protection of adenovirus from neutralizing antibody by cationic PEG derivative ionically linked to adenovirus. *Int. J. Nanomed.* 7, 985–997.
- Zhang, S., Huang, W., Zhou, X., *et al.* (2013a). Seroprevalence of neutralizing antibodies to human adenoviruses type-5 and type-26 and chimpanzee adenovirus type-68 in healthy Chinese adults. *J. Med. Virol.* 85, 1077–1084.
- Zhang, W.F., Wu, F.L., Shao, H.W., *et al.* (2013b). Chimeric adenoviral vector Ad5F35L containing the Ad5 natural long-shaft exhibits efficient gene transfer into human T lymphocytes. *J. Virol. Methods* 194, 52–59.
- Zhu, J., Huang, X., and Yang, Y. (2007). Innate immune response to adenoviral vectors is mediated by both Toll-like receptor-dependent and -independent pathways. *J. Virol.* 81, 3170–3180.
- Zhu, J., Huang, X., and Yang, Y. (2008). A critical role for type I IFN-dependent NK cell activation in innate immune elimination of adenoviral vectors *in vivo*. *Mol. Ther.* 16, 1300–1307.

Address correspondence to:

*Estrella Lopez-Gordo
Institute of Cardiovascular and Medical Sciences
BHF Glasgow Cardiovascular Research Centre
University of Glasgow
126 University Place
Glasgow G12 8TA
United Kingdom*

E-mail: l.estrella.1@research.gla.ac.uk

Received for publication December 17, 2013;
accepted after revision January 25, 2014.

Published online: February 5, 2014.