

Research article

Open Access

Xanthurenic acid translocates proapoptotic Bcl-2 family proteins into mitochondria and impairs mitochondrial function

Halina Z Malina*¹ and Otto M Hess²

Address: ¹Xanthurenic Acid Laboratory, Department Cardiology, Inselspital, CH-3010 Bern, Switzerland and ²Swiss Cardiovascular Research Center, University Hospital-Inselspital, CH-3010 Bern, Switzerland

Email: Halina Z Malina* - hmalina@hotmail.com; Otto M Hess - otto.martim.hess@insel.ch

* Corresponding author

Published: 06 April 2004

BMC Cell Biology 2004, 5:14

Received: 18 August 2003

Accepted: 06 April 2004

This article is available from: <http://www.biomedcentral.com/1471-2121/5/14>

© 2004 Malina and Hess; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Xanthurenic acid is an endogenous molecule produced by tryptophan degradation, produced in the cytoplasm and mitochondria. Its accumulation can be observed in aging-related diseases, e.g. senile cataract and infectious disease. We previously reported that xanthurenic acid provokes apoptosis, and now present a study of the response of mitochondria to xanthurenic acid.

Results: Xanthurenic acid at 10 or 20 μM in culture media of human aortic smooth muscle cells induces translocation of the proteins Bax, Bak, Bcl_{x_s}, and Bad into mitochondria. In 20 μM xanthurenic acid, Bax is also translocated to the nucleus. In isolated mitochondria xanthurenic acid leads to Bax and Bcl_{x_s} oligomerization, accumulation of Ca²⁺, and increased oxygen consumption.

Conclusion: Xanthurenic acid interacts directly with Bcl-2 family proteins, inducing mitochondrial pathways of apoptosis and impairing mitochondrial functions.

Background


Xanthurenic acid is an endogenous molecule derived from tryptophan degradation by indoleamine-2, 3-dioxygenase. Indoleamine-2, 3-dioxygenase is induced in infectious diseases by liposaccharides, interferon- α , and superoxide radicals [1]. Xanthurenic acid is enzymatically formed by transamination of 3-hydroxykynurenine by kynurenine-aminotransferase [2]. Kynurenine-aminotransferase, has been cloned from mitochondria and cytoplasm [3,4]. Xanthurenic acid accumulates with age in senile cataract [5-7]. Previously, we reported that this acid leads to apoptotic-like cell death [8-10], its BH3 domain in the cytoplasm being responsible for targeting the protein to mitochondria, resulting in cytochrome c release [11,12]. We now report that proteins with a BH3-

domain (Bax, Bak, Bcl-x_s, Bad) are transformed in the presence of xanthurenic acid to their proapoptotic-inducing state and translocated to mitochondria. *In vitro*, xanthurenic acid leads to covalent oligomerization of Bax, and Bcl-x_s. It also alters mitochondrial Ca²⁺ transport, and increases oxygen consumption.

Results

Xanthurenic acid provokes translocation of Bak, Bax, Bcl_{x_s}, and Bad to mitochondria

The fate of Bax, Bak, Bcl_{x_s}, and Bad in cultures of aortic human smooth muscle cells (HSMC) has been compared with control cells, using the antibody against Bax, which stains cells uniformly (Fig. 1A). 10 μM xanthurenic acid results in Bax being translocated to mitochondria (Fig.

**Figure 1**

Primary cell cultures human smooth muscle cells (A-C, Bax): (A) in control cells (magnification 400×), (B) in the presence of 10 μM xanthurenic acid (magnification 1260×), (C) in the presence of 20 μM xanthurenic acid (magnification 630×): mitochondria stained with MitotrackerRos (red staining), Bax (green), yellow-colocalization of Bax with mitochondria.

1B), and at 20 μM , Bax seems to be translocated from mitochondria to nuclei (Fig. 1C). However, at the latter concentration, some Bax remained detectable in mitochondria (Fig. 1C). Granular bodies strongly staining with anti-Bax antibodies (green) were detected in the nucleus (Fig. 1C). Translocation of Bax to the nucleus during apoptosis has previously been reported [13]. Proapoptotic proteins Bak, Bcl_x_s and Bad (Fig. 2A,2B,2C) at 10 and 20 μM xanthurenic acid (Fig. 2D,2E,2F) were translocated into mitochondria.

Xanthurenic acid leads to oligomerization of cytoplasmic Bcl-x_s and Bax in the rat liver mitochondrial fraction

Both Bcl-x_s and Bax were found in the 10,000 g mitochondrial fraction of rat liver. When it was incubated with xanthurenic acid at 5–10 μM (1 mg of protein/ml), cleavage of Bax with formation of p18 was observed. Both anti-Bax and anti-Bcl-x_s recognized the N-terminal part of the proteins, suggesting that the cleavage was more towards the C-terminal domain. When the concentration of xanthurenic acid was increased to 20–40 μM (1 mg protein/ml), Bax dimerized (Fig. 3A). Western blot analysis showed an increase of staining and dimerization of Bcl-x_s in the presence of xanthurenic acid (Fig. 3B). In the mitochondrial fraction, incubation of Bcl-x_s and Bax, with xanthurenic acid led to stronger attachment of the correspondent IgG antibody than in control cells, indicating some form of posttranscriptional modification. Dimers of the proapoptotic proteins in the presence of xanthurenic acid showed up on SDS-PAGE, confirming the formation of covalent crosslinking. Bax and Bcl-x_s were faintly detectable in the mitochondrial pellets. Cyto-

chrome c release was observed in the same protein extract (Fig. 3C).

Mitochondrial Ca²⁺ release

Ca²⁺ release was measured with Arsenazo III at a wavelength of 685/675 nm, using an Aminco DW-2 spectrophotometer. Xanthurenic acid at 15 μM (1 mg protein/ml) did not alter the release of Ca²⁺, but inhibition did occur at 66 μM (1 mg protein/ml). Ca²⁺ retained in mitochondria at this concentration was liberated after addition of 1 μM carbonyl cyanide-m-chlorophenylhydrazone (CCCP), which is known to provoke a loss of potential across mitochondrial membranes (Fig. 4). These results indicate that xanthurenic acid is involved in the modification of mitochondrial Ca²⁺ release, but does not lead to a loss of the mitochondrial membrane potential.

Consumption of O₂ by mitochondria in the presence of xanthurenic acid

Consumption of O₂ was measured with a Clarke-type electrode (Fig. 5). Mitochondrial respiration was supported with 2.5 mM potassium succinate prior to xanthurenic acid being added at 0.4 (Fig. 5, curve 1) or 4 μM (Fig. 5, curve 2). Xanthurenic acid stimulated oxygen consumption in a concentration-dependent manner.

Discussion

Apoptosis in the presence of xanthurenic acid is associated with cytochrome c release [8-10]. Our investigations show that xanthurenic acid induces the translocation of proapoptotic Bcl-2 proteins to mitochondria. A posttranscriptional modification of these proteins is suggested by the




Figure 2

Primary cell cultures human smooth muscle cells. (A-D, Bak): (A) Bak in control cells (magnification 800 \times), (D) Bak in presence of 20 μ M xanthurenic acid (magnification 800 \times), (B-E, Bclx₂): (B) Bclx₂ in control cells (magnification 1000 \times), (E) Bclx₂ in the presence of 10 μ M xanthurenic acid (magnification 1260 \times), (C-F, Bad): (C) Bad in control cells (magnification 800 \times), (F) Bad 20 μ M xanthurenic acid (magnification 800 \times).

direct crosslinking of Bax and Bcl-x₂ in isolated mitochondria. Previously, BID was reported to change the conformation of Bax [14]. Recently, we reported that accumulation of xanthurenic acid in cells interrupts the interaction of Bad with signalling proteins of the 14-3-3 family [10]. However, here we have shown that Bax is dimerized in isolated mitochondria, indicating that xanthurenic acid itself induces a conformational change of the proteins. Using Western blotting, Bax in mitochondria incubated with xanthurenic acid was detected more readily than in controls. This phenomenon was previously reported to indicate a change in Bax folding which results in a better exposition of the N-terminal part of Bax for the antibody after exposure to apoptotic stimuli [15]. Furthermore, xanthurenic acid appeared to modify the folding of proteins [16], which could be the reason for the increased IgG binding to proteins in the presence of xanthurenic acid.

Our results indicate that xanthurenic acid is an endogenous crosslinker of Bax and Bclx₂ proteins with a BH3 domain. Xanthurenic acid leads to translocation of BH3 proteins into mitochondria in human smooth muscle aortic cells. However, in isolated mitochondria, dimerized Bax and Bclx₂ found in the mitochondrial supernatant, are almost undetectable in the pellets, indicating that an additional cytoplasmic protein is necessary for translocation of these proteins into mitochondrial membranes. Therefore, the observed cytochrome c release in the presence of xanthurenic acid in isolated mitochondria is not mediated by Bax dimerization. At lower concentrations of xanthurenic acid, cleavage of Bax produced p18, which has been previously shown to enhance apoptosis [17]. Bax dimerization also has been shown to induce caspase-independent cell death [18]. Xanthurenic acid at 20 μ M causes Bax translocation into the nucleus, and plays a key role in apoptosis, because Bax/Bak deficient cells were




Figure 3

Western blot analysis of supernatant from rat liver mitochondria incubated for 1 h in the presence of xanthurenic acid for Bax, Bcl-x_s and cytochrome c; (A-B) Western blot analysis of Bax and Bcl-x_s; lane 1-control mitochondria; lanes 2-5: mitochondria incubated in the presence of xanthurenic acid at concentration 5; 10; 20; and 40 μM per mg of protein (A) Bax; (B) Bcl-x_s, (C) release of cytochrome c from mitochondria in the presence of xanthurenic acid at the same concentrations as in (A-B).

resistant to the apoptotic stimuli staurosporine and etoposide [19]. The fact that xanthurenic acid can directly provoke Bax and Bcl-x_s crosslinking suggests that an accumulation of xanthurenic acid in a cell is sufficient to induce the cell death. Xanthurenic acid might be oxidized *in vivo* to its semiquinone, and further to quinone-di-oxo-xanthurenic acid, as previously described [6]. The semiquinone is an oxidative stress [20], which can lead to uncoupling of mitochondria, characterized by an increase of O₂ consumption; in keeping, we noted an increased consumption of O₂ in the presence of xanthurenic acid. Previously it was reported that Bax is only toxic in the presence of active oxidative phosphorylation [21]. Our results indicate that at 20 μM xanthurenic acid, Bax is translocated into the nucleus and Ca²⁺ release from mito-

chondria is blocked. Xanthurenic acid accumulation in mitochondria does not affect mitochondrial membrane potential, because Ca²⁺ was accumulated in the mitochondria and CCCP was able to release Ca²⁺ under these conditions. Previously it was reported that Bax/Bax-induced apoptosis was associated with an uptake of mitochondrial Ca²⁺ by an unknown mechanism. An excessive Ca²⁺ load may induce apoptosis by stimulating the release of apoptosis-promoting factors from the mitochondrial membrane space to the cytoplasm, leading to impaired mitochondrial function (for review [22]).

The results presented here indicate that the accumulation of xanthurenic acid in the human aortic smooth muscle cell causes a proapoptotic conformation of the mitochondrial proteins, Bax, Bak, Bcl-x_s, and Bad. It was previously reported, that mitochondrial Bax plays an essential role in cytochrome c release, which can occur via Ca²⁺-dependent and -independent mechanisms [23]. Bax-driven apoptosis may be important for development of diabetes, atherosclerosis and neurological disorder [24-28]. We surmise that an accumulation of the endogenous aromatic amine, xanthurenic acid, may lead to organ failure.

Materials and methods

Reagents

The following polyclonal antibodies were obtained from Santa Cruz Biotechnology Inc. CA, USA: antibody against cytochrome c, against N-terminal part of Bax, Bak, Bcl-x_s, and Bad. IgG secondary antibodies-Texas Red conjugated were from Santa Cruz or secondary antibodies IgG-Alexa 488 conjugated Molecular Probes (Leiden, The Netherlands). Carbonyl cyanide-m-chlorophenylhydrazone (CCCP) and Arsenazo III were from Sigma.

Preparation of human smooth muscle cells primary culture

The normal human aortic smooth muscle cells were from PromoCell GmbH, Heidelberg, Germany.

The cells were cultivated in Smooth Muscle Cells Growth Medium 2 (SMCGM2) with an addition of Supplement Pack/Smooth Muscle Cells (PromoCell). Cells were grown under a humidified atmosphere of 5% CO₂ in air at 37°C. When confluent, they were incubated for 96 hours in the SMCGM2 or this medium supplemented with 10 and 20 μM xanthurenic acid. A 20 mM stock solution of xanthurenic acid was prepared in 0.5 M NaHCO₃, and diluted in 0.01 M PBS pH 7.4.

Western blot analysis of protein from rat liver mitochondrial fraction

Rat liver mitochondria (80-90 mg protein per ml) were isolated, as described previously [29]. The 10,000 g mitochondrial fraction of cells was used (referred herein as mitochondria). The mitochondria were in MSH buffer




Figure 4

Ca^{2+} release from mitochondria in the presence of xanthurenic acid (1) control mitochondria, (2, 3) in the presence of xanthurenic acid, and 1 mg of mitochondrial proteins: (2) 15 μM xanthurenic acid, (3) in the presence of 66 μM xanthurenic acid (4) after addition of CCCP to (3).

(210 mM mannitol, 70 mM sucrose, 5 mM Hepes, pH 7.35 (4°C, 2 M KOH). Mitochondria (~3 mg protein per 100 μl of reaction mixture) were incubated for 1 h at room temperature without xanthurenic acid or in the presence of the increasing concentrations of xanthurenic acid from 5 to 40 μM per mg protein in the presence of the following protease inhibitors: 0.1 mM phenyl-methylsulfonyl fluoride, and leupeptin, aprotinin, and pepstatin, each at 1 $\mu\text{g}/\text{ml}$. Thereafter, the sample was centrifuged for 10 min at 14,000 g, and the supernatant boiled in loading-buffer for 5 min. Proteins at 100 μg per lane were separated by SDS-PAGE containing 12.5% acrylamide. After transfer to Hybond ECL membrane (Amersham Pharmacia Biotech AB, Uppsala, Sweden), the proteins were probed with the appropriate antibodies. Chemiluminescence ECL system (Amersham Pharmacia Biotech AB, Uppsala, Sweden) was used for the detection of peroxidase-conjugated secondary antibody.

Immunofluorescence studies

Cells grown on glass coverslips were fixed for 10 min at room temperature in 4% paraformaldehyde in 0.1 M PIPES, pH 6.8, washed in PBS and permeabilized for 5 min in PIPES containing 0.05% saponin (65 μl per coverslip), washed in PBS, incubated for 10 min in cold acetone for additional fixing and permeabilisation, and again washed in PBS. Cells were incubated for 1.5 h with the first antibody diluted in PBS containing 1% bovine serum albumin, and after further washing incubated for 1.5 h with the secondary antibody. The coverslips were washed in PBS and incubated for 10 min with 65 μl of solution containing 1 μl of Hoechst 33342 dye (1 mg/ml), washed in PBS, and incubated with Antifade Kits (Molecular Probes, Leiden, The Netherlands) according to the supplier's instruction.




Figure 5
Mitochondrial oxygen consumption in presence of xanthurenic acid per 1 mg of protein in the presence of (1) 0.4 μ M xanthurenic acid, and (2) 4 μ M xanthurenic.

Measurement of calcium uptake and release and O_2 consumption in liver mitochondria in the presence of xanthurenic acid

An Aminco DW-2A spectrophotometer was used at 685–675 nm the presence of 50 μ M Arsenazo III, and mitochondrial respiration was measured using a Clarke-electrode as described previously [29]. Briefly, mitochondria (1 mg of mitochondrial protein/ml) were energized with 2.5 mM potassium succinate in the presence of rotenone. After Ca^{2+} uptake was complete, the uptake blocker, ruthenium red (RR), was added to see net release Ca^{2+} in the presence or absence of xanthurenic acid.

Competing interests

None declared.

Authors' contribution

HZM: conception, design, data collection and analysis, writing the paper.

OMH: revising the paper, research funds collection.

All authors approved the final version of the paper.

Acknowledgements

This work was supported by grant awarded to H. Z. M. by the Swiss National Science Foundation (32-59183.99), Swiss Heart Foundation, and MSE-Pharmaceutika GMBH, Bad Homburg, Germany. We thank Dr. C. Richter for proofreading the manuscript and Dr. M. Salvi for his help with Ca^{2+} release experiment.

References

- Carlin JM, Ozaki Y, Byrne GI, Brown RR, Borden EC: **Interferons and indoleamine 2,3-dioxygenase: role in antimicrobial and antitumor effects.** *Experientia* 1989, **45**:535-541.
- Tobes MC: **Kynurenine-oxoglutarate aminotransferase from rat kidney.** *Methods Enzymol* 1987, **142**:217-224.
- Mosca M, Cozzi L, Breton J, Speciale C, Okuno E, Schwarcz R, Benatti L: **Molecular cloning of rat kynurenine aminotransferase: identity with glutamine transaminase K.** *FEBS Lett* 1994, **353**:21-24.
- Malherbe P, Alberati-Giani D, Kohler C, Cesura AM: **Identification of a mitochondrial form of kynurenine aminotransferase/glutamine transaminase K from rat brain.** *FEBS Lett* 1995, **367**:141-144.
- Malina HZ, Martin XD: **Deamination of 3-hydroxykynurenine in bovine lenses: a possible mechanism of cataract formation in general.** *Graefes Arch Clin Exp Ophthalmol* 1995, **233**:38-44.
- Malina HZ, Martin XD: **Xanthurenic acid derivative formation in the lens is responsible for senile cataract in humans.** *Graefes Arch Clin Exp Ophthalmol* 1996, **234**:723-730.
- Malina HZ, Martin XD: **3-hydroxykynurenine transamination leads to the formation of the fluorescent substances in human lenses.** *Eur J Ophthalmol* 1996, **6**:250-256.
- Malina HZ, Richter C, Mehl M, Hess OM: **Pathological apoptosis by xanthurenic acid, a tryptophan metabolite: activation of cell caspases but not cytoskeleton breakdown.** *BMC Physiol* 2001, **1**:7.
- Malina H, Richter C, Frueh B, Hess OM: **Lens epithelial cell apoptosis and intracellular Ca^{2+} increase in the presence of xanthurenic acid.** *BMC Ophthalmol* 2002, **2**:1.
- Malina HZ, Frueh BE: **Abnormal signalling of 14-3-3 proteins in cells with accumulated xanthurenic acid.** *Biochem Biophys Res Commun* 2003, **310**:646-650.
- Luo X, Budihardjo I, Zou H, Slaughter C, Wang X: **Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors.** *Cell* 1998, **94**:481-490.
- Polster BM, Kinnally KW, Fiskum G: **BH3 death domain peptide induces cell type-selective mitochondrial outer membrane permeability.** *J Biol Chem* 2001, **276**:37887-37894.
- Gajkowska B, Motyl T, Olszewska-Badarczuk H, Gniadecki R, Koronkiewicz M: **Structural association of Bax with nuclear matrix and cytomatrix revealed by embedment-free immunogold electron microscopy.** *Cell Biol Int* 2000, **24**:649-656.
- Desagher S, Osen-Sand A, Nichols A, Eskes R, Montessuit S, Lauper S, Maundrell K, Antonsson B, Martinou JC: **Bid-induced conformational change of Bax is responsible for mitochondrial cytochrome c release during apoptosis.** *J Cell Biol* 1999, **144**:891-901.
- Priault M, Camougrand N, Chaudhuri B, Manon S: **Role of the C-terminal domain of Bax and Bcl-XL in their localization and function in yeast cells.** *FEBS Lett* 1999, **443**:225-228.
- Malina HZ: **Xanthurenic acid provokes formation of unfolded proteins in endoplasmic reticulum of the lens epithelial cells.** *Biochem Biophys Res Commun* 1999, **265**:600-605.
- Wood DE, Newcomb EW: **Cleavage of Bax enhances its cell death function.** *Exp Cell Res* 2000, **256**:375-382.
- Gross A, Jockel J, Wei MC, Korsmeyer SJ: **Enforced dimerization of BAX results in its translocation, mitochondrial dysfunction and apoptosis.** *Embo J* 1998, **17**:3878-3885.
- Wei MC, Zong WX, Cheng EH, Lindsten T, Panoutsakopoulou V, Ross AJ, Roth KA, MacGregor GR, Thompson CB, Korsmeyer SJ: **Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction and death.** *Science* 2001, **292**:727-730.
- Gutierrez PL: **The role of NAD(P)H oxidoreductase (DT-Diaphorase) in the bioactivation of quinone-containing antitumor agents: a review.** *Free Radic Biol Med* 2000, **29**:263-275.
- Harris MH, Vander Heiden MG, Kron SJ, Thompson CB: **Role of oxidative phosphorylation in Bax toxicity.** *Mol Cell Biol* 2000, **20**:3590-3596.
- Hajnoczky G, Davies E, Madesh M: **Calcium signaling and apoptosis.** *Biochem Biophys Res Commun* 2003, **304**:445-454.
- Gogvadze V, Robertson JD, Zhivotovsky B, Orrenius S: **Cytochrome c release occurs via Ca^{2+} -dependent and Ca^{2+} -independent mechanisms that are regulated by Bax.** *J Biol Chem* 2001, **276**:9066-9071.

24. Konstadoulakis MM, Kymionis GD, Karagiani M, Katergianakis V, Doundoulakis N, Pararas V et al.: **Evidence of apoptosis in human carotid atheroma.** *J Vasc Surg* 1998, **27**:733-739.
25. Podesta F, Romeo G, Liu WH, Krajewski S, Reed JC, Gerhardinger C, Lorenzi M: **Bax is increased in the retina of diabetic subjects and is associated with pericyte apoptosis in vivo and in vitro.** *Am J Pathol* 2000, **156**:1025-1032.
26. Snider BJ, Gottron FJ, Choi DW: **Apoptosis and necrosis in cerebrovascular disease.** *Ann N Y Acad Sci* 1999, **893**:243-253.
27. Srinivasan S, Stevens M, Wiley JW: **Diabetic peripheral neuropathy: evidence for apoptosis and associated mitochondrial dysfunction.** *Diabetes* 2000, **49**:1932-1938.
28. Hayakawa Y, Takemura G, Misao J, Kanoh M, Ohno M, Ohashi H et al.: **Apoptosis and overexpression of bax protein and bax mRNA in smooth muscle cells within intimal hyperplasia of human radial arteries: analysis with arteriovenous fistulas used for hemodialysis.** *Arterioscler Thromb Vasc Biol* 1999, **19**:2066-207.
29. Frei B, Winterhalter KH, Richter C: **Mechanism of alloxan-induced calcium release from rat liver mitochondria.** *J Biol Chem* 1985, **260**:7394-7401.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

